

Short Communication

Enhanced N-Transfer from a Soybean to Maize by Vesicular Arbuscular Mycorrhizal (VAM) Fungi¹

Received for publication June 3, 1985

CHRISTOPHER VAN KESSEL*, PAUL W. SINGLETON, AND HEINZ J. HOBEN
University of Hawaii NifTAL Project, Paia, Hawaii 96779

ABSTRACT

Using a split-root technique, roots of soybean plants were divided between two pots. In one of the two pots, two maize plants were grown and half of those pots were inoculated with the vesicular arbuscular mycorrhizal (VAM) fungus, *Glomus fasciculatus*. Fifty-two days after planting, ¹⁵N-labeled ammonium sulfate was applied to the pots which contained only soybean roots. Forty-eight hours after application, significantly higher values for atom per cent ¹⁵N excess were found in roots and leaves of VAM-infected maize plants as compared with the non-VAM-infected maize plants. Results indicated that VAM fungi did enhance N transfer from one plant to another.

ined VAM mediated transfer of N from a soybean to maize using highly labeled (¹⁵NH₄)₂SO₄ and a split-root technique.

MATERIALS AND METHODS

Surface sterilized (2 min in 3% NaOCl seeds of soybeans (*Glycine max* [L.] Merr) and maize (*Zea mays* L.) were germinated in vermiculite. The last 0.5 cm of the soybean tap roots were removed and the two seedlings were placed in two separate plastic elbows (pvc elbow, 21 mm o.d., 13 mm diameter hole) (16). At the same time, two pregerminated maize seedlings were planted in pot B (Fig 1).

Half of the B pots were inoculated with a VAM fungi, *Glomus fasciculatus*, maintained in a coarse sand pot culture with each inoculated pot receiving 30 g of inoculum which contained small

Vesicular arbuscular mycorrhizal fungi are ubiquitous and infect plant roots of most species under a wide variety of soil conditions (8). The fungi form a symbiosis with host plants in which the plant provides carbon for VAM² growth and in turn the VAM fungi provide plant nutrients, especially phosphorus, from the soil solution (11). Growth responses of host plants to infection by VAM may be dramatic in nutrient-poor environments (7). Hyphae of mycorrhizae may also spread from one infected plant and enter the roots of one or more other plants (9). It has been shown that assimilates may be transported from one plant to another through VAM hyphal connections. Transfer of ¹⁴C photosynthate from one plant to another was primarily through VAM hyphae rather than leakage from the roots of the donor plants (2, 6, 14). Similar results were obtained in a ³²P experiment where hyphal linkage between plants was the dominant factor for transferring P (3, 17).

Leguminous plants infected with both *Rhizobium* and VAM showed an increase in nodulation and N₂-fixation as compared with VAM-uninfected legumes (4, 5). The increase in total N has been explained mainly by an increase in N₂-fixation as a result of a higher P uptake through the VAM hyphae rather than increased soil N uptake (15). Although the role of VAM on N uptake and transport has been studied, the results are inconclusive. Rhodes and Gerdemann (15) stated that N translocation by VAM to the host plant would probably be of little significance, while Raven *et al.* (13) attributed a considerable role of VAM to the N nutrition of plants.

Under conditions of low N and P availability which exist in many tropical soils, the possible transfer from the host plant to another plant by VAM may well become important. We exam-




FIG. 1. A split-root system where the roots of the soybean were allowed to grow into two separated pots.

¹ Supported by Grant DAN-0613-C-00-2064-00 from the United States Agency for International Development. Conclusions of this paper do not necessarily reflect those of the granting agency.

² Abbreviation: VAM vesicular arbuscular mycorrhizal.

root fragments, hyphae, and approximately 25 spores per gram. The inoculum was spread evenly over the surface of sterile sand which filled one-half of 3-L pots. Sterile sand was added to fill the pots completely after inoculation. Nonmycorrhizal plants were treated with 50 ml suspensions of 30 g of inoculum filtered through Whatman No. 1 filter paper.

A nutrient solution containing 6 g/g of Mg as $MgSO_4 \cdot 7H_2O$, 39 g/g K as KNO_3 , 20 g/g Ca as $Ca(NO_3)_2 \cdot 4H_2O$, 1.5 g/g P as KH_2PO_4 , and 53 g/g N as KNO_3 (14 g/g), $Ca(NO_3)_2 \cdot 4H_2O$ (14 g/g), and NH_4NO_3 (25 g/g) was applied five times daily at a rate of 150 ml/plant per application. A liquid micronutrient concentrate (Montarry Chemical Co.) was added at 0.125 ml/L which provided 1.55 g/g Fe, 0.6 g/g Zn, 0.6 g/g Mn, 0.4 g/g B, 0.2 g/g Cu, 0.05 g/g Mo, and 0.04 g/g Co per L. Plants were grown in a greenhouse without additional light; treatments were replicated eight times. Fifty d after planting, pots A and B were leached with deionized H_2O followed by the addition of 0.7 mmol ^{15}N /pot as 99.99 atom % ^{15}N enriched $(NH_4)_2SO_4$ to pot A. A second application of $(^{15}NH_4)_2SO_4$ was added 24 h after the first application. Plants were harvested 48 h after the first ^{15}N application.

Soybeans and maize roots were separated as carefully as possible. Complete separation of the two root masses was not possible and therefore only roots which were still attached to the stem of the soybean or corn plants were analyzed. The rest of the root mass was discarded. Roots were thoroughly washed to remove any contaminating traces of enriched ^{15}N . All maize and soybean roots were checked for VAM infection after staining in 0.05% trypan blue in lactophenol (12). Maize leaves and roots and soybean roots were dried at 70°C until constant weight was obtained. Plants parts were ground separately in a cyclone sample mill. After grinding each sample, the mill was taken apart and thoroughly brushed and vacuum cleaned to avoid any cross-contamination between samples. Tissues were digested and analyzed for total N including NO_2^- and NO_3^- (1). Digestions were made alkaline with 13 N NaOH and steam distilled for 7 min in an all glass steam distilling apparatus. Distillates were collected in 0.02 N H_2SO_4 . To avoid cross-contamination, 20 ml of ethyl alcohol was distilled between each sample. Samples were analyzed for total N using the indophenol blue method (8). The rest of the distillate was adjusted to a pH of 4, concentrated and analyzed for ^{15}N . The analyses were carried out at the Isotope Service, Inc. in Los Alamos, NM.

The atom % ^{15}N of VAM-infected and non-VAM-infected plants in pot B, but which had not received ^{15}N -labeled $(NH_4)_2SO_4$ in pot A, were used for calculating the atom % ^{15}N excess of VAM-infected and non-VAM-infected maize and soybean plants which had received ^{15}N -labeled $(NH_4)_2SO_4$.

RESULTS AND DISCUSSION

All inoculated roots of maize and soybeans were infected with VAM fungi, whereas none of the noninoculated root systems showed any infection. Table I shows the % N of maize leaves,

Table I. Per cent N, Dry Weight and Atom per cent ^{15}N Excess of + and - VAM-infected Maize and Soybean Plants

| Plant Origin | Treatment | N | Dry Wt | ^{15}N |
|--------------------------|-----------|--------|--------|---------------|
| | | % | g | atom % excess |
| Maize leaves | -VAM | 1.62a* | 8.94a | 0.0013a |
| | +VAM | 1.40a | 11.64a | 0.0031b |
| Corn roots | -VAM | 1.39a | | 0.0032a |
| | +VAM | 1.17b | | 0.0113b |
| Soybean roots (Pot B) | -VAM | 2.17a | | 0.1966a |
| | +VAM | 2.00a | | 0.2128a |

* Means followed by the same letter are not significantly different at the P = 0.05 level by means of the *t* test.

maize roots, and soybean roots which were grown in pot B of VAM-infected and non-VAM-infected plants. VAM-infected plants did show a significantly lower % N for roots as compared with non-VAM-infected maize roots. There was a similar tendency in maize leaves. However, total N in VAM-infected and non-VAM-infected maize leaves were not different because of the higher dryweight of the former. VAM infection may have stimulated growth by increasing the availability of some other factor that limited dry matter accumulation in non-VAM plants.

Leaves and roots of VAM-infected maize plants did show an atom % ^{15}N excess significantly higher at the P = 0.05 level when compared with the leaves and roots of non-VAM-infected maize. This indicates that the VAM fungi has facilitated the transfer of labeled ^{15}N from the soybean to the maize plant. In addition to soil N-derived compounds, N_2 -derived compounds may also be transferred from legumes to nonlegumes. The mechanism how this occurred can be by direct, active transport through the mycelia from the donor plant to the receiving plant as has been shown for P (6) and for C (2) as well. However, it also may be possible that VAM-infected plants leak more N compounds out of the roots into the medium than non-VAM-infected plants. Subsequently, more ^{15}N -labeled material will enter the medium and become available for the corn plant.

We have established that a VAM mediated N-transfer from a legume to a nonlegume does occur. Additional experiments are needed before insight can be given about the nature and amount of N transferred from one plant to another through VAM fungi.

Acknowledgments—We would like to thank James Tavares and Patty Nakao for their help in the analysis of the samples. Our thanks also go to Dr. Jake Halliday for his encouragement and to Dr. Burris, University of Wisconsin, for the initial ^{15}N mass spectrometer analysis; Dr. Joann Roskoski and Dr. Ben Bohlool for their critical reading of the manuscript; Keith Avery for preparing the illustration and Mary Rohner for manuscript preparation.

LITERATURE CITED

- BREMNER JM, CS MULVANEY 1982 Nitrogen total. *Agronomy* 9: 595-624
- BROWNEE C, JA DUDDIDGE, A MALIBAN, D READ 1983 The structure and function of mycelial systems of ectomycorrhizal roots with special reference to their role in forming inter-plant connections and providing pathways for assimilation and water transport. *Plant Soil* 71: 433-443
- CHIARELLO N, JC HICKMAN, MA MOONEY 1982 Endomycorrhizal role for interspecific transfer of phosphorus in a community of annual plants. *Science* 217: 941-943
- CLUETT HC, DH BOUCHER 1984 Indirect mutualism of the legume-*Rhizobium*-mycorrhizal fungus interaction. *Oecologia* 59: 405-408
- CRUSH JR 1974 Plant growth responses to vesicular-arbuscular mycorrhiza. VII. Growth and nodulation of some herbage legumes. *New Phytol* 73: 743-749
- FRANCIS R, DJ READ 1984 Direct transfer of carbon between plants connected by vesicular-arbuscular mycorrhizal mycelium. *Nature* 307: 53-56
- GERDEMANN JW 1975 Vesicular-arbuscular mycorrhizae. In JG Torrey, DT Clarkson, eds, *The Development and Function of Roots*. Academic Press, New York, pp 545-591
- HARLEY JL, SE SMITH 1983 *Mycorrhizal Symbiosis*. Academic Press, New York
- HEAP AJ, EI NEWMAN 1980 Links between roots by hyphae of vesicular-arbuscular mycorrhizas. *New Phytol* 85: 169-171
- KEENEY DR, DW NELSON 1982 Nitrogen—inorganic forms. *Agronomy* 9: 643-698
- MOSSE B 1973 Advances in the study of vesicular arbuscular mycorrhiza. *Ann Rev Phytopathol* 11: 171-96
- PHILLIPS JM, DS HAYMAN 1970 Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans Br Mycol Soc* 55: 158-160
- RAVEN JA, SE SMITH, FA SMITH 1978 Ammonium assimilation and the role of mycorrhizas in climax communities in Scotland. *Trans Bot Soc Edinb* 43: 27-35
- REID CPP, FW WOODS 1969 Translocation of C^{14} labelled compounds in mycorrhizal and its implications in interplant nutrient cycling. *Ecology* 50: 178-187
- RHODES LH, JW GERDEMANN 1980 Nutrient translocation in vesicular-arbuscular mycorrhizae. In CB Cook, PW Pappas, ED Rudolph, eds, *Cellular Interactions in Symbiosis and Parasitism*. Ohio State University Press, Columbus, pp 173-195
- SINGLETON PW 1983 A split-root growth system for evaluating the effect of salinity on the components of the soybean *Rhizobium japonicum* symbiosis. *Crop Sci* 23: 259-262
- WHITTINGHAM J, DJ READ 1982 Vesicular arbuscular mycorrhiza in natural vegetation systems. III. Nutrient transfer between plants with mycorrhizal interconnections. *New Phytol* 90: 277-284