

<https://doi.org/10.17221/96/2023-VETMED>

An update about beneficial effects of medicinal plants in aquaculture: A review

FARANAK DADRAS*, JOSEF VELISEK, ELISKA ZUSKOVA

Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Research Institute of Fish Culture and Hydrobiology, University of South Bohemia in Ceske Budejovice, Ceske Budejovice, Czech Republic

*Corresponding author: fdadrasasiabar@frov.jcu.cz

Citation: Dadras F, Velisek J, Zuskova E (2023): An update about beneficial effects of medicinal plants in aquaculture: A review. *Vet Med-Czech* 68, 449–463.

Abstract: Aquaculture is an essential and growing component of agricultural and global ecosystems worldwide. Aquaculture provides more than 25% of the total aquatic food consumption by humans. The development of the aquaculture industry should be followed in successive industrial years, and therefore it is necessary to pay attention to the management and type of farming system that is compatible with the environment. The use of antibiotics for disease control has been criticised for their negative effects, including the emergence of antibiotic-resistant bacteria, the suppression of the immune system and the environment, and the accumulation of residue in aquatic tissues. The use of these products reduces the need for treatments, enhances the effect of vaccines, and, in turn, improves production indicators. Medicinal plants have increasingly been used in recent years as a disease control strategy in aquaculture, boosting the immune system of aquatic animals and helping to develop strong resistance to a wide range of pathogens. Therefore, this review aims to provide an overview of the recent evidence on the beneficial use of medicinal plants to promote growth and strengthen the immune system in farmed aquatic animals.

Keywords: disease; fish; growth; immunity; medicinal herb

INTRODUCTION

Aquaculture has been forecasted to increase by 62% between 2010 and 2030 to supply the growing fish and seafood demand derived from a steadily growing population, providing over two-thirds of the total fish and shellfish consumed worldwide (Bank 2013; Sofia 2018). On the other hand, in 2020, the total amount of globally farmed fish amounted

to 82.1 million tonnes, confirming aquaculture remains one of the significant contributors to animal protein sources (Stankus 2021). Despite the considerable role of aquaculture, the sector faces many challenges that prevent its expansion. The high culture density has led to environmental pollution and the emergence of diseases (Lafferty et al. 2015) due to stressed and immuno-compromised animals (Bondad-Reantaso et al. 2005; Pulkkinen

Supported by the Ministry of Agriculture of the Czech Republic, Project No. QK21010113 and by the Ministry of Education, Youth and Sports of the Czech Republic, Project Sustainable production of healthy fish in various aquaculture systems, PROFISH (CZ.02.1.01/0.0/0.0/16_019/0000869).

© The authors. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0).

<https://doi.org/10.17221/96/2023-VETMED>

et al. 2010). Other factors, such as storms, droughts, and high temperatures, adversely affect the water quality and may compromise the fish's immune system and the health of aquatic animals (Dubey et al. 2017; Abdel-Tawwab et al. 2019).

Aquatic animal diseases are among the key limiting factors for aquaculture development (Stentiford et al. 2017), which could lead to the partial or complete loss of growth and production and result in considerable economic losses, estimated at over 9.5 billion USD per year (Shinn et al. 2015; Ramesh and Souissi 2018). Therefore, antibiotics and other veterinary drugs are commonly used in aquaculture to treat disease outbreaks as well as to prevent and mitigate the economic losses resulting from sanitary shortcomings (Rico et al. 2013; Cabello et al. 2016; Miranda et al. 2018). The potential risks of using antibiotics and the associated global health threats due to the selection and emergence of antibiotic-resistant bacteria in aquaculture have been extensively studied (Cabello et al. 2016; Chen et al. 2018). Recently, in response to the risk of developing resistant genes that could potentially change the human pathogen, the use of antibiotics has been restricted in aquaculture (Binh et al. 2018; van Wyk and Prinsloo 2020). Several alternative strategies, including vaccination, probiotics, herbal plants, and immunostimulants, have been proposed to prevent disease outbreaks and limit the use of veterinary drugs in aquaculture (Liu et al. 2014; Reverter et al. 2014; Reverter et al. 2017). Vaccination is a precise technique that requires a clear disease diagnosis (Brudeseth et al. 2013) as well as the complicated development of multiple-strain vaccines (Sakai 1999; Pasnik et al. 2005). Furthermore, it is too expensive for widespread use among small-scale fish farmers (Pridgeon and Klesius 2012).

As one of the best alternatives, medicinal plants have been strongly recommended. Medicinal plants contain specific active substances modulating biological functions, such as growth promotion (Amin et al. 2019; Gharaei et al. 2020), anti-stress effects (Abd El-Gawad et al. 2020), appetite stimulation (Mo et al. 2016), immunostimulation (Rufchaei et al. 2017), disease resistance (Liu et al. 2010a; van Wyk and Prinsloo 2020), and antimicrobial activities (Ardo et al. 2008; Beltran et al. 2018). To ensure sustainable aquaculture, medicinal plants seem to be cheaper and more sustainable alternatives to chemotherapy. This supports the increasing number of publications on conceptu-

alising medicinal plant applications. The present review purposefully discussed the adoption of naturally available plants in the form of roots, leaves, flowers, and in a processed form, including active compounds, extracts, crude forms, and mixed with different promising effects in aquaculture.

Application of medicinal plants in aquaculture

Medicinal plants contain antioxidant compounds with a high affinity to trap free radicals inhibiting the normal functioning of cells (Moreno et al. 2020). Many antimicrobial substances in these plants make them an appropriate candidate to fight a wide range of pathogenic microbes (Jafarzadeh et al. 2020). Large bioactive compounds, including steroids, proteins, tannins, saponins, terpenoids, and alkaloids, are found in different plants. These compounds have shown strong resistance to a wide range of bacteria pathogens (*Salmonella typhi*, *Bordetella pertussis*, *Corynebacterium parvum*, *Klebsiella pneumoniae*, *Mycobacterium* and *Escherichia coli*), fungi (*Aspergillus flavus*, *Aspergillus fumigatus*, *Fusarium solani* and *Pseudomonas aeruginosa*) and viruses (*retrovirus*, *simian-virus*) (Ma et al. 2019).

Medicinal plants have been used in chemotherapy and as feed additives (Reverter et al. 2014; Garg et al. 2020). For a long time, they have been used by rural fish farmers (Reverter et al. 2017); in 46% of the surveyed fish farmers (Caruso et al. 2013). Medicinal plants, as sustainably used candidates, are easily accessible and readily available to be applied in intensive farming in aquaculture to ensure the improved productivity and health status of aquatic organisms (Cawthorn and Hoffman 2015). Medicinal plants can be administered in different ways, either as a whole plant or in parts (leaf, root, seed, fruit) and can either be applied fresh or as prepared herbal extracts with different solvents (water, methanol, chloroform, ethyl acetate) (Van Hai 2015). The biological activity and chemical composition of plants and extracts can vary greatly depending on the part used and type of extract, thus, knowledge of the plant's bioactive compounds is required.

For instance, some herbs, such as moringa (*Moringa oleifera*), can be used as a protein source or partial protein replacement due to its high pro-

<https://doi.org/10.17221/96/2023-VETMED>

tein content, i.e., 27.51% in the crude protein of the leaves (Oduro et al. 2008). Selection of a suitable dosage of a medicinal herb is crucial to obtain the desired effects since inappropriate doses can display toxic effects (Sambasivam et al. 2003; Kavitha et al. 2012). The high activation of the immune response without harmful and toxic effects on animals can be induced by a proper dose of the target material (Bulfinch et al. 2013). Some parts of the produced chemical compounds during regular metabolic activities of the plants, such as phytochemicals, comprise a diverse group of natural products. They may be nutritionally essential, but many of them, such as phytate, lectin, and trypsin inhibitors, have no nutritional value and, having antinutritional properties, can decrease the feed conversion efficiency and growth rate (Makkar et al. 2007; Hashemi and Davoodi 2011). A study on the toxic effect of Indian almonds (*Terminalia catappa*), leaf extracts at different concentrations (700, 800, 900, 1 000, and 1 100 mg/l) on the Nile tilapia (*Oreochromis niloticus*) juveniles under static toxicity exposure (96 h exposure) showed the 96hLC50 value was 900 mg/l (Yunus et al. 2019).

A diet supplemented with a 1% ethanol katuk extract (*Sauropus androgynus*) showed enhanced growth and improved food utilisation in the orange-spotted grouper (*Epinephelus coioides*), whereas 2.5% and 5% of the katuk extract presented lower growth levels (Santoso et al. 2013). Another critical factor in a medicinal plant application is the treatment length. Farmed barramundi (*Lates calcarifer*) fed an enriched diet of garlic (*Allium sativum*) for 30 days showed a 70% decreased *Neobenedenia* sp. infection compared to the control and the group fed garlic for 10 days (Militz et al. 2013).

Medicinal plants can be administered to fish by injection (intramuscular and intraperitoneal), oral administration, or immersion (Wu et al. 2010; Ji et al. 2012; Santoso et al. 2013). The intraperitoneal injection is considered the most rapid and efficient method of administration; however, it is expensive, laborious, and stressful for the fish (Anderson 1992; Yoshida et al. 1995); on the other hand, baths are extensively used for the treatment of ectoparasites (Whittington 2012; Forwood et al. 2013), but this method is also expensive and laborious and involves the release of exogenous molecules in the marine environment (Umeda et al. 2006). Zebrafish (*Danio rerio*) received an injection of a coumarin derivative (10 µl/fish) for 14 days

and showed improved resistance against spring viraemia of carp virus (SVCV) (Shen et al. 2018). An oral administration of berberine hydrochloride (30–50 mg/kg) in Prussian carp (*Carassius gibelio*) for 15 days showed antiviral activity against herpesvirus 2 (CyHV-2) (Su et al. 2021).

Tilapia were injected with a hot-water extract of Chinese mahogany (*Toona sinensis*) (with a 2 or 4 mg/ml solution) at doses of 4 or 8 mg/g and had significantly increased respiratory burst, phagocytic activity, and lysozyme activity towards *Aeromonas hydrophila* by 1 and 2 days post-injection (Wu et al. 2010). An improved survival rate and lysozyme activity were reported in rainbow trout (*Oncorhynchus mykiss*) orally administered with a Greek juniper extract at 8 mg/kg dose for 14 days (Bilen et al. 2021). The intraperitoneal injection, oral administration, and diffusion of a 0.1 ml extract of green chiretta (*Andrographis paniculata*) in Indian major carp (*Labeo rohita*) improved the non-specific immune system against *Aeromonas hydrophila* via enhancing the total erythrocyte count and normal haemoglobin levels (Palanikani et al. 2020).

Biological activity of medicinal plants in fish

The main target organs of fish influenced by medicinal plants are the thymus, spleen, kidney, and guts, which promote immune system development. Medicinal plants can directly improve the antibody production and the specific immune response. Many medicinal plants can promote the production of cytokines that mediate the specific/non-specific immunity, including interleukin, interferon, and the tumour necrosis factor (Tadese et al. 2021).

Medicinal plants enhance immune parameters (Dugenci et al. 2003; Yuan et al. 2007). The biological activity of medicinal plants is attributed to their secondary metabolites (SMs), such as essential oils, saponins, phenolics, tannins, alkaloids, polypeptides, and polysaccharides (Hoseinifar et al. 2020b). The SMs play a key role in stress mediation, antioxidant activity, and immunopotentiality by modulating the recognition, binding, catalytic activity, and turnover of proteins and DNA (Chong et al. 2020). Several *in vitro* and *in vivo* studies have demonstrated the beneficial effects of medicinal plants against a wide range of marine pathogens (bacteria,

<https://doi.org/10.17221/96/2023-VETMED>

viruses, fungi, and ectoparasites) (Direkbusarakom et al. 1996; Chitmanat et al. 2005; Ji et al. 2012; Su et al. 2021).

MEDICINAL PLANTS AS GROWTH PROMOTERS

As summarised in see Electronic Supplementary Material (ESM) Table S1, medicinal plants can stimulate the appetite and promote weight gain as they enhance the digestive enzyme activity (Van Hai 2015). For instance, some herbs, such as the sissoo spinach (*Alternanthera sessilis*), false daisy (*Eclipta alba*), and veldt grape (*Cissus quadrangularis*) used as appetisers, improved the activities of protease, amylase, and lipase of freshwater prawns (Radhakrishnan et al. 2014). Adding a wormwood (*Artemisia annua*) extract had a growth-promoting effect on rainbow trout (Koshinski 2018) and carp (Sarhadi et al. 2020). Similar results have been reported by He et al. (2022) in largemouth black bass (*Micropterus salmoides*).

The growth rate of freshwater prawns (*Macrobrachium rosenbergii*) has been improved using a diet supplemented with an emodin extract (Liu et al. 2010b). Greasy groupers (*Epinephelus tauvina*) fed a diet supplemented with a mixture of methanolic herb extracts of Bermuda grass (*Cynodon dactylon*), Indian long pepper (*Piper longum*), stonebreaker (*Phyllanthus niruri*), Tridax daisy (*Tridax procumbens*), and ginger (*Zingiber officinalis*) showed enhanced weight gain (41%) (Punitha et al. 2008).

The administration of numerous plant extracts to cultured fish induces appetite and increases weight gain (Raja Rajeswari et al. 2012; Hoseinifar et al. 2020c; Mohammadi et al. 2020). Several studies confirmed the promising effect of medicinal plants on the growth performance of fish species, including the administration of ferula (*Ferula asafoetida*) powder in common carp (*Cyprinus carpio*) (Safari et al. 2019); prickly chaff flower (*Achyranthes aspera*) (0.5%) in rohu (*Labeo rohita*) fry (Sharma et al. 2019); onion (*Allium cepa*) powder (10 g/kg) in white carp (*Cirrhinus mrigala*) fingerlings (Sikotariya 2019); chaff flower (*Achyranthes aspera*) (0.5%) in rohu (*Labeo rohita*) (Singh et al. 2019); dandelion (*Taraxacum officinale*) extract (0.8%) in common carp (Sirakov et al. 2019); ginger (*Zingiber officinale*) (0.8%) in rohu

(*Labeo rohita*) fingerlings (Sukumaran et al. 2016); climbing Senecio (*Senecio scandens* buch-ham) extracts (0.05–0.1%) in hybrid grouper (*Epinephelus lanceolatus* ♂ × *Epinephelus fuscoguttatus* ♀) (Sun et al. 2020); fluted pumpkin (*Telfairia occidentalis*) (1%) in African sharp-tooth catfish (*Clarias gariepinus*) (Ta et al. 2019); dandelion extract (1 g/kg) in golden pompano (*Trachinotus ovatus*) (Tan and Sun 2020); wolfberry (*Lycium barbarum*) extract (0.5–2%) in hybrid grouper (*Epinephelus lanceolatus* ♂ × *E. fuscoguttatus* ♀) (Tan et al. 2019); olive extract (1 g/kg) in common carp (Zemheri-Navruz et al. 2020); and curcumin (120 mg/kg) in common carp (Zhang et al. 2020a). In line with the reports mentioned above, Zhang et al. (2021) demonstrated an increased weight gain and specific growth rate of juvenile common carp fed 60 and 120 mg curcumin per kg for ten weeks and suggested that the observed improvement could be attributed to the enhanced immune response, which increased the growth performance.

Since regulatory functions, such as the metabolism, antioxidant, immune capacity, anti-stress, anti-virus, antibacterial, and anti-parasite activities can directly affect the growth rate, a balanced regulation among these physiological metabolisms seems to be necessary for aquatic species. Hence, these aspects of using medicinal plants in aquaculture must be taken into account.

MEDICINAL PLANTS AS ANTI-STRESSORS AND IMMUNOSTIMULANTS

In intensive aquaculture, external cues, such as poor water quality, a high environmental temperature, overcrowding, and pathogen infection, make aquatic animals vulnerable to adverse stress (Asaduzzaman et al. 2009; Chang et al. 2015). The immunomodulatory power of plants is mostly determined by the assessment of the fish immunity via classic biochemical approaches (lysozyme, phagocytic, or respiratory burst activity) or by the study of the immune gene expression (Lysine (Lys), Tumour Necrosis Factor-alpha (TNF-alpha), Interleukin-1 (IL-1), Interleukin-10 (IL-10) genes) (Harikrishnan et al. 2011; Kumar et al. 2013; Chakrabarti et al. 2014). Several studies recommend some natural compounds, including medicinal plants, to defend against stress induced by external cues (ESM Table S2).

<https://doi.org/10.17221/96/2023-VETMED>

Several herbal plants that originated from terrestrial and marine environments and are considered efficient immunostimulants against viral diseases was confirmed in a study (Chakraborty et al. 2014). In contrast to antibiotics, medical plants as natural resources have outstanding features, such as being highly effective and showing low toxicity and side effects when used correctly (Bone and Mills 2012). The antistress effects of medicinal plants have been mentioned in several studies (Ahmadniaye Motlagh et al. 2019). For instance, an anthraquinone extract from Chinese rhubarb (*Rheum officinale Bail*) improved the tolerance against hyperthermia in freshwater shrimp (*Macrobrachium nipponense*) (Song et al. 2020). Moreover, a sweet wormwood (*Artemisia annua*) leaf extract improved the antioxidant capacity of carp (Taheri Mirghaed et al. 2020) and enhanced the immunity of Nile tilapia (Soares et al. 2020). A methanolic extract of fenugreek (*Trigonella foenum-graecum*) significantly increased the immunity and antioxidative response via improvement of the superoxide dismutase, lysozyme, and phagocytic activities in Nile tilapia (Diab et al. 2023).

The anti-ammonia stress capacity of freshwater prawns was improved by the addition of a moringa (*Moringa oleifera*) leaf extract to the fish's diet (Kaleo et al. 2019). Furthermore, emodin protected Wuchang bream (*Megalobrama amblycephala*) from crowding stress (Liu et al. 2014), and bupleurum (*Radix Bupleuri*) extracts boosted tilapia against H₂O₂-induced oxidative stress (Jia et al. 2019). In common carp, the adverse effects of crowding stress were mitigated using the application of an anthraquinone extract from rhubarb (*Rheum officinale Bail*) (1–2%) (Xie et al. 2008). Also, the anti-stress effect of turmeric in common carp exposed to copper was reported by Rajabiesterabadi et al. (2020).

Several studies have focused on using plant extracts as fish immunostimulants (Prathomya et al. 2019; Rafieepour et al. 2019); according to scientific studies on fish species, the intraperitoneal injection or oral administration of plant extracts enhanced the phagocytic and lysosomal function, respiratory burst and complement activity, as well as the serum protein level (Liu et al. 2010a; Vallejos-Vidal et al. 2016; Balamurugan et al. 2018; Ramezanzadeh et al. 2019). A dietary supplemented by fenugreek (*Trigonella foenum graecum*) seeds (5%) had the highest haemolytic complement and

peroxidase activities, indicating that the metabolic and immune status of gilthead seabream (*Sparus aurata* L.) was improved (Guardiola et al. 2018). A supplement of an olive leaf extract (0.1%) in the diet of rainbow trout increased the immune-related gene expression, serum biochemistry parameters, and survival rate (Baba et al. 2018). It is supposed that herbal extracts can positively affect the innate immunity (Devi et al. 2019; de Assis and Urbinati 2020; Soares et al. 2020), adaptive immunity (Abd El-Gawad et al. 2020; Zhang et al. 2020b), mucosal immunity (Van Doan et al. 2019; Ahmadniaye Motlagh et al. 2020a; Ahmadniaye Motlagh et al. 2020b; Heydari et al. 2020; Srichaiyo et al. 2020a; Srichaiyo et al. 2020b), and their anti-stress functions have been proven in several studies (summarised in ESM Table S2).

MEDICINAL PLANTS AS AN ANTI-VIRAL AND AN ANTIBACTERIAL TREATMENT

The potential of medicinal plants against a wide range of marine pathogens (bacteria, viruses, fungus, and ectoparasites) was confirmed in *in vitro* and *in vivo* studies (Direkbusarakom et al. 1996; Chitmanat et al. 2005; Ji et al. 2012). The antiviral activity of plant species against aquatic primary pathogenic viruses was reported in different diseases, including white spot syndrome virus (WSSV), grouper iridovirus (GIV), grass carp reovirus (GCRV), spring viraemia of carp virus (SVCV) and cyprinid herpesvirus 2 or 3 (CyHV) (summarised in ESM Table S3). Giant tiger prawn (*Penaeus monodon*) treated with Bermuda grass (*Cynodon dactylon*) displayed no signs of disease and mortality when exposed to white spot syndrome virus (WSSV), while 100% mortality was observed in the control groups (Balasubramanian et al. 2008a; Balasubramanian et al. 2008b). A methanolic extract of fenugreek increased the tolerance of Nile tilapia against *A. hydrophila* (Diab et al. 2023); this could be attributed to the deactivation of the peptidoglycans of *A. hydrophila* due to the increased lysozyme activity that caused enhanced resistance (Masschalck and Michiels 2003; Brott and Clarke 2019). The extract of the plants, such as Bermuda grass (*Cynodon dactylon*), the olive (*olea europaea*), cape jasmine (*Gardenia jasminoides*), and Mexican poppy (*Argemone mexicana*) showed significant anti-virus activity against WSSV

<https://doi.org/10.17221/96/2023-VETMED>

through oral administration or injection (Ghosh et al. 2014; Palanikumar et al. 2018; Huang et al. 2019a; Huang et al. 2019b). Furthermore, a berberine hydrochloride (*Clinacanthus nutans*) extract showed a considerable preventive effect against CyHV-2 or CyHV-3 in *Gibel carp* (*Carassius auratus gibelio*) (Su et al. 2021) or koi carp (*Cyprinus carpio koi*) (Haetrakul et al. 2018). Several *in vitro* studies have reported the antibacterial activity in numerous plants against both Gram-positive and Gram-negative marine bacteria (Castro et al. 2008; Roomiani et al. 2013). As summarised in ESM Table S3, some medicinal plants presented specific antibacterial effects on pathogenic bacteria such as the Mongolian milkvetch (*Astragalus membranaceus*) (Wu 2020), fern (*Adiantum capillus-veneris*) (Hoseinifar et al. 2020a), horse mint (*Mentha longifolia*) (Heydari et al. 2020), rosemary (*Rosmarinus officinale*) (Naiel et al. 2020), and prepared foxglove root (*Radix rehmanniae preparate*) (Wu et al. 2019). A tamarind (*Tamarindus indica* L.) pulp extract (15 g/kg) promoted the growth and nutrient digestibility of Nile tilapia, and provided protection against an *A. hydrophila* infection (Adeniyi et al. 2021). The addition of a 1.5 g/kg chinaberry tree (*Melia Azedarach*) extract, a 4–6 g/kg velvet bean (*Mucuna pruriens*) extract, and a 0.5–2 g/kg jojoba extract to the diet improved the resistance against *A. hydrophila* in Catla (*Labeo catla*) (Rajeshwari et al. 2016), Mozambique tilapia (*Oreochromis mossambicus*) (Saiyad Musthafa et al. 2018), and Nile tilapia (Sarhan et al. 2019), respectively. A supplementation of 40% moringa (*Moringa oleifera*) leaf extracts to the diet of carp infected by *A. hydrophila* enhanced the growth, antioxidant and immune response of the carp (Zhang et al. 2020a). Similar promising effects of a medicinal herb supplementation against *A. hydrophila* were observed in different fish species, such as the addition of geniposide in the diet of crucian carp (He et al. 2020), a grape seed extract in the diet of common carp (Mehrinakhi et al. 2021), Mongolian wild onion (*Allium Mongolicum Regel*) (40 mg/kg) in the diet of juvenile snakehead (*Channa argus*) (Li et al. 2018), Smoketree (*Cotinus cogglyria*) in the diet of rainbow trout (Bilen and Elbeshti 2019), and marjoram (*Origanum majorana*) (Yousefi et al. 2021) and maidenhair tree (*Ginkgo biloba*) leaf (Bao et al. 2019) extracts in the diet of common carp. The antibacterial activity of several medicinal herbs against *Vibrio* spp. as an opportunistic

pathogenic bacteria has been confirmed in many fish species, such as applications of peppermint (*Mentha piperita*) extracts against a *V. harveyi* infection in Barramundi (Talpur 2014), a Chinese rhubarb (*Rheum officinale*) extract (0.1, 1.0 g/kg) against *V. parahaemolyticus* in the orange-spotted grouper (*Epinephelus coioides*) (Kuo et al. 2020), a gale of the wind (*Phyllanthus amarus*) extract (20 g/kg) against *V. alginolyticus* in pacific white shrimp (*Litopenaeus vannamei*) (Ngo et al. 2020), tears of the virgin (*Eleutherine bulbosa*) (12.5 g/kg) against *V. parahaemolyticus* in pacific white shrimp (*Litopenaeus Vannamei*) (Munaeni et al. 2020), and the Chinese herbal medicine, San-Huang-San against *V. parahaemolyticus* in pacific white shrimp (*Litopenaeus vannamei*) (Zhai and Li 2019).

Additionally, the protective roles of some medicinal herbs against *Streptococcus* spp. as an opportunistic pathogen in aquaculture systems have been mentioned in some scientific studies on Nile tilapia, such as the promising effects of a thumbai (*Leucas aspera*) extract (8 g/kg) against a *Streptococcus agalactiae* infection (Kurian et al. 2020) and an Assam tea (*Camellia sinensis*) extract against *S. agalactiae* (Van Doan et al. 2019). Likewise, the antibacterial activity of some herbs against *Yersinia ruckeri* has been confirmed by incorporating a peppermint (*Mentha Piperita*) extract (Adel et al. 2016), Greek juniper (*Juniperus excelsa*) (Bilen et al. 2021) and coriander (*Coriandrum sativum*) (Naderi Farsani et al. 2019) in the diet of rainbow trout. The functional activity of the plasma lysozyme, blood phagocytes, respiratory burst, and survival rates have improved in Nile tilapia fed a combination of honeysuckle (*Lonicera*) and milkvetch (*Astragalus*) extracts when infected with *A. hydrophila* (Ardo et al. 2008).

In addition, various active ingredients extracted from medicinal plants and essential oils were confirmed to have antibacterial activities against overwhelming bacterial species (Kacaniova et al. 2017). It was found that medicinal plants could show different levels of immune stimulation by injection, immersion, or oral administration (Awad and Awaad 2017).

MEDICINAL PLANTS AS AN ANTIPARASITIC

Medicinal plants can be considered as an effective alternative for treating ectoparasites. The

<https://doi.org/10.17221/96/2023-VETMED>

antiparasitic activities of medicinal plants, when added to water or administered orally, were confirmed by several studies (Yi et al. 2012; Huang et al. 2013; Fu et al. 2014). A significant improvement in the haematological profile in Nile tilapia challenged by flatworms (*Scutogyrus longicornis*), cichlidogyrus (*Cichlidogyrus thurstone*), cichlidogyrus (*Cichlidogyrus halli*), and monogeneans, cichlidogyrus (*Cichlidogyrus tilapia*) was observed when treated by essential oils extracted from peppermint (*Mentha piperita*) (Hashimoto et al. 2016).

Harikrishnan et al. (2012) reported decreased mortality (40%) and enhanced immunity in the olive flounder (*Paralichthys olivaceus*) infected by the protozoan ciliate *Miamiensis avidus* when a fish-fed diet was supplemented with herbaceous seepweed (*Suaeda maxima*).

Some studies on fish infected with white spot disease (*Ichthyophthirius multifiliis*) showed that ginger (*Zingiber officinale*), sweet wormwood (*Artemisia annua*), tea tree (*Melaleuca alternifolia*), English lavender (*Lavandula angustifolia*), peppermint (*Mentha piperita*), babchi (*Psoralea corylifolia*), orange climber (*Toddalia asiatica*) and Chinese gall (*Galla chinensis*) could potentially control the infection (Zhang et al. 2013; Shan et al. 2014; Song et al. 2015; Valladao et al. 2016; Wu et al. 2017; de Freitas Souza et al. 2019; Fu et al. 2019). The white spot disease (*Ichthyophthirius multifiliis*) in the sailfin molly (*Poecilia latipinna*) was also inhibited by a mother worth (*Matricaria chamomilla*) extract (Gholipourkanani et al. 2012). 100% *in vivo* efficacy against monogenean *Dactylogyrus intermedius* was reported in infected goldfish (*Carassius auratus*) treated by a methanol extract of bupleurum root (*Radix bupleuri chinensis*), aqueous and methanol extracts of cinnamon (*Cinnamomum cassia*), a methanol extract of Chinese spice bush (*Lindera aggregata*) and methanol and ethyl acetate extracts of golden larch (*Pseudolarix kaempferi*) (Wu et al. 2011; Ji et al. 2012). The oral administration of an aqueous extract of rosemary (*Rosmarinus officinalis*) could kill gill fluke (*Dactylogyrus minutus*) in common carp (Zoral et al. 2017). A bath treatment of guppies [*Poecilia reticulata* (Peters)] using an ethanolic extract of ginger (*Zingiber Officinale*) significantly decreased the salmon fluke (*Gyrodactylus turnbulli*) infection (Levy et al. 2015). More and more herbal medicines can be found to be used in parasite remedies of fish species (see *ESM Table S4*).

CONCLUSION

Medicinal plants exhibit a promising potential for the current needs of the intensive and largescale production in the aquaculture industry and are a substitute for chemotherapy in the treatment of disease outbreaks and control of farmed fish diseases. Studies have shown they are a proper candidate for disease prevention and control guidelines to develop pollution-free aquaculture and green aquatic products. Each medicinal plant is specific in its action against infectious diseases, and the dose of a specific medicinal plant is a crucial factor in the efficacy against specific infectious agents. However, the toxicological safety research of medicinal plants needs to be further studied and more research is also required to elucidate upon plant products and their modes of action and to test the plant effects on an organism's physiology in order to establish appropriate treatment strategies. In this context, researchers may benefit from the traditional knowledge of fish farmers who regularly use plants. Moreover, research is still relatively scarce on the use of some compatible herbs with a synergistic effect. Therefore, the effects of some compatible herbs on the health status of fish and shellfish might be an exciting topic in aquaculture.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Abd El-Gawad EA, El Asely AM, Soror EI, Abbass AA, Austin B. Effect of dietary *Moringa oleifera* leaf on the immune response and control of *Aeromonas hydrophila* infection in Nile tilapia (*Oreochromis niloticus*) fry. *Aquac Int.* 2020 Feb;28(1):389-402.
- Abdel-Tawwab M, Monier MN, Hoseinifar SH, Faggio C. Fish response to hypoxia stress: Growth, physiological, and immunological biomarkers. *Fish Physiol Biochem.* 2019 Jun;45(3):997-1013.
- Adel M, Pourgholam R, Zorriehzahra J, Ghiasi M. Hematological and biochemical parameters, skin antibacterial activity, and survival in rainbow trout (*Oncorhynchus mykiss*) following the diet supplemented with *Mentha piperita* against *Yersinia ruckeri*. *Fish Shellfish Immunol.* 2016 Aug;55:267-73.

<https://doi.org/10.17221/96/2023-VETMED>

- Adeniyi O, Emikpe B, Olaifa F, Ogunbanwo ST. Effects of dietary tamarind (*Tamarindus indica* L.) leaves extract on growth performance, nutrient utilization, gut physiology, and susceptibility to *Aeromonas hydrophila* infection in Nile tilapia (*Oreochromis niloticus* L.). *Int Aquat Res*. 2021 Mar;13(1):37-51.
- Ahmadniaye Motlagh H, Safari O, Paolucci M. Effect of different levels of milkweed (*Calotropis persica*) seed powder on the growth parameters, immunity and gut microbiota of *Oncorhynchus mykiss*. *Iran J Vet Sci*. 2019;11(1):43-50.
- Ahmadniaye Motlagh H, Rokhnareh Z, Safari O, Selahvarzi Y. Growth performance and intestinal microbial changes of *Carassius auratus* in response to pomegranate (*Punica granatum*) peel extract-supplemented diets. *J World Aquac Soc*. 2020a Dec;52(4):820-8.
- Ahmadniaye Motlagh H, Safari O, Selahvarzi Y, Baghalian A, Kia E. Non-specific immunity promotion in response to garlic extract supplemented diets in female Guppy (*Poecilia reticulata*). *Fish Shellfish Immunol*. 2020b Feb;97:96-9.
- Amin A, El Asely A, Abd El-Naby A, Samir F, El-Ashram A, Sudhakaran R, Dawood MAO. Growth performance, intestinal histomorphology and growth-related gene expression in response to dietary *Ziziphus mauritiana* in Nile tilapia (*Oreochromis niloticus*). *Aquaculture*. 2019 Oct;512:734301.
- Anderson DP. Immunostimulants, adjuvants, and vaccine carriers in fish: Applications to aquaculture. *Annu Rev Fish Dis*. 1992;2:281-307.
- Ardo L, Yin G, Xu P, Varadi L, Szigeti G, Jeney G. Chinese herbs (*Astragalus Membranaceus* and *Lonicera japonica*) and boron enhance the non-specific immune response of Nile tilapia (*Oreochromis niloticus*) and resistance against *Aeromonas hydrophila*. *Aquaculture*. 2008 Mar;275(1-4):26-33.
- Asaduzzaman M, Wahab MA, Verdegem MCJ, Mondal MN, Azim ME. Effects of stocking density of freshwater prawn *Macrobrachium rosenbergii* and addition of different levels of tilapia *Oreochromis niloticus* on production in C/N controlled periphyton based system. *Aquaculture*. 2009 Jan;286(1):72-9.
- Awad E, Awaad A. Role of medicinal plants on growth performance and immune status in fish. *Fish Shellfish Immunol*. 2017 Aug;67:40-54.
- Baba E, Acar U, Yilmaz S, Zemheri F, Ergun S. Dietary olive leaf (*Olea europea* L.) extract alters some immune gene expression levels and disease resistance to *Yersinia ruckeri* infection in rainbow trout *Oncorhynchus mykiss*. *Fish Shellfish Immunol*. 2018 Aug;79:28-33.
- Balamurugan S, Vijayakumar S, Prabhu S, Morvin Yabesh JE. Traditional plants used for the treatment of gynaecological disorders in Vedaranyam taluk, South India – An ethnomedicinal survey. *J Tradit Complement Med*. 2018 Jul;8(2):308-23.
- Balasubramanian G, Sarathi M, Venkatesan C, Thomas J, Hameed AS. Studies on the immunomodulatory effect of extract of *Cyanodon dactylon* in shrimp, *Penaeus monodon*, and its efficacy to protect the shrimp from white spot syndrome virus (WSSV). *Fish Shellfish Immunol*. 2008a Dec;25(6):820-8.
- Balasubramanian G, Sarathi M, Venkatesan C, Thomas J, Sahul Hameed AS. Oral administration of antiviral plant extract of *Cynodon dactylon* on a large scale production against White spot syndrome virus (WSSV) in *Penaeus monodon*. *Aquaculture*. 2008b Jul;279(1):2-5.
- Bank W. Fish to 2030 prospects for fisheries and aquaculture. Washington, DC: The World Bank Group (WBG) Report Number 83177-GLB.; 2013. 102 p.
- Bao L, Chen Y, Li H, Zhang J, Wu P, Ye K, Ai H, Chu W. Dietary *Ginkgo biloba* leaf extract alters immune-related gene expression and disease resistance to *Aeromonas hydrophila* in common carp *Cyprinus carpio*. *Fish Shellfish Immunol*. 2019 Nov;94:810-8.
- Beltran JMG, Espinosa C, Guardiola FA, Esteban M. In vitro effects of *Origanum vulgare* leaf extracts on gilthead seabream (*Sparus aurata* L.) leucocytes, cytotoxic, bactericidal and antioxidant activities. *Fish Shellfish Immunol*. 2018 Aug;79:1-10.
- Bilen S, Elbeshti H. A new potential therapeutic remedy against *Aeromonas hydrophila* infection in rainbow trout (*Oncorhynchus mykiss*) using tetra, *Cotinus coggygia*. *J Fish Dis*. 2019 Oct;42(10):1369-81.
- Bilen S, Ispir S, Kenanoglu ON, Tastan Y, Guney K, Terzi E. Effects of Greek juniper (*Juniperus excelsa*) extract on immune responses and disease resistance against *Yersinia ruckeri* in rainbow trout (*Oncorhynchus mykiss*). *J Fish Dis*. 2021 Jun;44(6):729-38.
- Binh VN, Dang N, Anh NTK, Thai PK. Antibiotics in the aquatic environment of Vietnam: Sources, concentrations, risk and control strategy. *Chemosphere*. 2018;197:438-50.
- Bondad-Reantaso MG, Subasinghe RP, Arthur JR, Ogawa K, Chinabut S, Adlard R, Tan Z, Shariff M. Disease and health management in Asian aquaculture. *Vet Parasitol*. 2005 Sep;132(3-4):249-72.
- Bone K, Mills S. Principles and practice of phytotherapy: Modern herbal medicine. Edinburgh, London, UK: Elsevier Health Sciences; 2012. 1051 p.
- Brott AS, Clarke AJ. Peptidoglycan O-Acetylation as a virulence factor: Its effect on lysozyme in the innate immune system. *Antibiotics*. 2019 Jul;8(3):94.
- Brudeseth BE, Wiulsrod R, Fredriksen BN, Lindmo K, Lokling KE, Bordevik M, Steine N, Klevan A, Gravnin-

<https://doi.org/10.17221/96/2023-VETMED>

- gen K. Status and future perspectives of vaccines for industrialised fin-fish farming. *Fish Shellfish Immunol.* 2013 Dec;35(6):1759-68.
- Bulfon C, Volpatti D, Galeotti M. Current research on the use of plant-derived products in farmed fish. *Aquac Res.* 2013 Jul;46(3):513-51.
- Cabello FC, Godfrey HP, Buschmann AH, Dolz HJ. Aquaculture as yet another environmental gateway to the development and globalisation of antimicrobial resistance. *Lancet Infect Dis.* 2016 Jul;16(7):e127-e33.
- Caruso D, Lusiastuti AM, Slembrouck J, Komarudin O, Legendre M. Traditional pharmacopeia in small scale freshwater fish farms in West Java, Indonesia: An ethnoveterinary approach. *Aquaculture.* 2013 Dec;416:334-45.
- Castro SB, Leal CA, Freire FR, Carvalho DA, Oliveira DE, Figueiredo HC. Antibacterial activity of plant extracts from Brazil against fish pathogenic bacteria. *Braz J Microbiol.* 2008 Oct;39(4):756-60.
- Cawthorn DM, Hoffman LC. The bushmeat and food security nexus: A global account of the contributions, conundrums and ethical collisions. *Food Res Int.* 2015 Oct(P1);76:906-25.
- Chakrabarti R, Srivastava PK, Verma N, Sharma J. Effect of seeds of *Achyranthes aspera* on the immune responses and expression of some immune-related genes in carp *Catla catla*. *Fish Shellfish Immunol.* 2014 Nov;41(1):64-9.
- Chakraborty S, Ghosh U, Balasubramanian T, Das P. Screening, isolation and optimization of anti-white spot syndrome virus drug derived from marine plants. *Asian Pac J Trop Biomed.* 2014 May;4(Suppl_1):S107-17.
- Chang CC, Jiang JR, Cheng W. A first insight into temperature stress-induced neuroendocrine and immunological changes in giant freshwater prawn, *Macrobrachium rosenbergii*. *Fish Shellfish Immunol.* 2015 Nov;47(1):528-34.
- Chen B, Lin L, Fang L, Yang Y, Chen E, Yuan K, Zou S, Wang X, Luan T. Complex pollution of antibiotic resistance genes due to beta-lactam and aminoglycoside use in aquaculture farming. *Water Res.* 2018 May;134:200-8.
- Chitmanat C, Tongdonmuan K, Khanom P, Pachontis P, Nunsong W. Antiparasitic, antibacterial, and antifungal activities derived from a terminalia catappa solution against some tilapia (*Oreochromis niloticus*) pathogens. *Acta Hort.* 2005;678:179-82.
- Chong CM, Murthy AVS, Choy CY, Koksong L. Phytotherapy in aquaculture: Integration of endogenous application with science. *J Environ Biol.* 2020 Sep;41:1204-14.
- de Assis RWS, Urbinati EC. Physiological activity of *Aloe vera* in pacu (*Piaractus mesopotamicus*) inoculated with *Aeromonas hydrophila*. *Fish Physiol Biochem.* 2020 Aug;46(4):1421-30.
- de Freitas Souza C, Baldissera MD, Descovi S, Zeppenfeld C, Eslava-Mocha PR, Gloria EM, Zanette RA, Baldisserotto B, da Silva AS. Melaleuca alternifolia essential oil abrogates hepatic oxidative damage in silver catfish (*Rhamdia quelen*) fed with an aflatoxin-contaminated diet. *Comp Biochem Physiol Part C.* 2019 Jul;221:10-20.
- Devi G, Harikrishnan R, Paray BA, Al-Sadoon MK, Hoseinifar SH, Balasundaram C. Effects of aloe-emodin on innate immunity, antioxidant and immune cytokines mechanisms in the head kidney leucocytes of *Labeo rohita* against *Aphanomyces invadans*. *Fish Shellfish Immunol.* 2019 Apr;87:669-78.
- Diab AM, Al-Khefa BT, Khalafallah MM, Salah AS, Farrag FA, Dawood MAO. Dietary methanolic extract of fenugreek enhanced the growth, haematobiochemical, immune responses, and resistance against *Aeromonas hydrophila* in Nile Tilapia, *Oreochromis niloticus*. *Aquac Res.* 2023 Feb;2023(8):1-13.
- Direkbusarakom S, Herunsalee A, Yoshimizu M, Ezura Y. Antiviral activity of several Thai traditional herb extracts against fish pathogenic viruses. *Fish Pathol.* 1996 Dec;31(4):209-13.
- Dubey SK, Trivedi RK, Chand BK, Mandal B, Rout SK. Farmers' perceptions of climate change, impacts on freshwater aquaculture and adaptation strategies in climatic change hotspots: A case of the Indian Sundarban delta. *Environ Dev.* 2017 Mar;21:38-51.
- Dugenci SK, Arda N, Candan A. Some medicinal plants as immunostimulant for fish. *J Ethnopharmacol.* 2003 Sep;88(1):99-106.
- Forwood JM, Harris JO, Deveney MR. Efficacy of current and alternative bath treatments for *Lepidotrema bidyana* infecting silver perch, *Bidyanus bidyanus*. *Aquaculture.* 2013 Dec;416:65-71.
- Fu Y, Zhang Q, Xu DH, Xia H, Cai X, Wang B, Liang J. Parasiticidal effects of *Morus alba* root bark extracts against *Ichthyophthirius multifiliis* infecting grass carp. *Dis Aquat Organ.* 2014 Feb;108(2):129-36.
- Fu YW, Wang B, Zhang QZ, Xu DH, Liu YM, Hou TL, Guo SQ. Efficacy and antiparasitic mechanism of 10-gingerol isolated from ginger *Zingiber officinale* against *Ichthyophthirius multifiliis* in grass carp. *Vet Parasitol.* 2019 Jan;265:74-84.
- Garg C, Sahu N, Maiti M, Nazeemashahul S, Deo A, Sardar P. Dietary *Houttuynia cordata* leaf extract and meal enhances the immunity and expression of immune genes in *Labeo rohita* (Hamilton, 1822). *Aquac Res.* 2020 Jan;52(1):381-94.
- Gharaei A, Jorjani H, Harijani JM, Miandare HK. Effects of *Tribulus terrestris* extract on masculinization, growth indices, sex determination and steroid hormones level

<https://doi.org/10.17221/96/2023-VETMED>

- in Zebra fish (*Danio rerio*). *Int Aquat Res.* 2020;12(1): 22-9.
- Gholipourkanani H, Sahandi J, Taheri A. Influence of garlic (*Allium sativum*) and mother worth (*Matricaria chamomilla*) extract on *Ichthyophthirius multifiliis* parasite treatment in sail fin molly (*Poecilia latipinna*) ornamental fish. *APCBEE Procedia.* 2012;4(3):6-11.
- Ghosh U, Chakraborty S, Balasubramanian T, Das P. Screening, isolation and optimization of anti-white spot syndrome virus drug derived from terrestrial plants. *Asian Pac J Trop Biomed.* 2014 May;4(Suppl_1):S118-28.
- Guardiola FA, Bahi A, Esteban MA. Effects of dietary administration of fenugreek seeds on metabolic parameters and immune status of gilthead seabream (*Sparus aurata* L.). *Fish Shellfish Immunol.* 2018 Mar;74:372-9.
- Haetrakul T, Dunbar SG, Chansue N. Antiviral activities of *Clinacanthus nutans* (Burm.f.) Lindau extract against Cyprinid herpesvirus 3 in koi (*Cyprinus carpio koi*). *J Fish Dis.* 2018 Apr;41(4):581-7.
- Harikrishnan R, Balasundaram C, Heo MS. Impact of plant products on innate and adaptive immune system of cultured finfish and shellfish. *Aquaculture.* 2011 Jul;317(1): 1-15.
- Harikrishnan R, Kim JS, Kim MC, Dharaneedharan S, Kim DH, Hong SH, Song CU, Balasundaram C. Effect of dietary supplementation with *Suaeda maritima* on blood physiology, innate immune response, and disease resistance in olive flounder against *Miamiensis avidus*. *Exp Parasitol.* 2012 Jun;131(2):195-203.
- Hashemi SR, Davoodi H. Herbal plants and their derivatives as growth and health promoters in animal nutrition. *Vet Res Commun.* 2011 Mar;35(3):169-80.
- Hashimoto GSO, Neto FAM, Ruiz ML, Acchile M, Chagas EC, Chaves FCM, Martins ML. Essential oils of *Lippia sidoides* and *Mentha piperita* against monogenean parasites and their influence on the hematology of Nile tilapia. *Aquaculture.* 2016 Jan;450:182-6.
- He G, Sun H, Liao R, Wei Y, Zhang T, Chen Y, Lin S. Effects of herbal extracts (*Foeniculum vulgare* and *Artemisia annua*) on growth, liver antioxidant capacity, intestinal morphology and microorganism of juvenile largemouth bass, *Micropterus salmoides*. *Aquac Rep.* 2022 Apr;23(7): 101081.
- He M, Liu G, Liu Y, Yang K, Qi X, Huang A, Liu T, Wang G, Wang E. Effects of geniposide as immunostimulant on the innate immune response and disease resistance in crucian carp. *Aquaculture.* 2020 Dec;529:735713.
- Heydari M, Firouzbaksh F, Paknejad H. Effects of *Mentha longifolia* extract on some blood and immune parameters, and disease resistance against yersiniosis in rainbow trout. *Aquaculture.* 2020 Jan2020;515:734586.
- Hoseinifar SH, Jahazi MA, Mohseni R, Raeisi M, Bayani M, Mazandarani M, Yousefi M, Van Doan H, Mozanazadeh MT. Effects of dietary fern (*Adiantum capillus-veneris*) leaves powder on serum and mucus antioxidant defence, immunological responses, antimicrobial activity and growth performance of common carp (*Cyprinus carpio*) juveniles. *Fish Shellfish Immunol.* 2020a Nov;106:959-66.
- Hoseinifar SH, Shakouri M, Doan HV, Shafiei S, Yousefi M, Raeisi M, Yousefi S, Harikrishnan R, Reverter M. Dietary supplementation of lemon verbena (*Aloysia citrodora*) improved immunity, immune-related genes expression and antioxidant enzymes in rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.* 2020b Apr;99: 379-85.
- Hoseinifar SH, Sun YZ, Zhou Z, Van Doan H, Davies SJ, Harikrishnan R. Boosting immune function and disease bio-control through environment-friendly and sustainable approaches in finfish aquaculture: Herbal therapy scenarios. *Rev Fish Sci Aquac.* 2020c;28(3):303-21.
- Huang AG, Yi YL, Ling F, Lu L, Zhang QZ, Wang GX. Screening of plant extracts for anthelmintic activity against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitol Res.* 2013 Dec;112(12): 4065-72.
- Huang AG, Tu X, Qi XZ, Ling F, Zhu B, Wang GX. *Gardenia jasminoides* Ellis inhibit white spot syndrome virus replication in red swamp crayfish *Procambarus larkia*. *Aquaculture.* 2019a Apr;504:239-47.
- Huang AG, Tan XP, Qu SY, Wang GX, Zhu B. Evaluation on the antiviral activity of genipin against white spot syndrome virus in crayfish. *Fish Shellfish Immunol.* 2019b Oct;93:380-6.
- Jafarzadeh S, Jafari SM, Salehabadi A, Nafchi AM, Uthaya Kumar US, Khalil HPSA. Biodegradable green packaging with antimicrobial functions based on the bioactive compounds from tropical plants and their by-products. *Trends Food Sci Technol.* 2020 Jun;100:262-77.
- Ji J, Lu C, Kang Y, Wang GX, Chen P. Screening of 42 medicinal plants for in vivo anthelmintic activity against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitol Res.* 2012 Jul;111(1):97-104.
- Jia R, Gu Z, He Q, Du J, Cao L, Jeney G, Xu P, Yin G. Antioxidative, anti-inflammatory and hepatoprotective effects of *Radix Bupleuri* extract against oxidative damage in tilapia (*Oreochromis niloticus*) via Nrf2 and TLRs signaling pathway. *Fish Shellfish Immunol.* 2019 Oct;93:395-405.
- Kacaniova M, Terentjeva M, Vukovic N, Puchalski C, Roychoudhury S, Kunova S, Kluga A, Tokar M, Kluz M, Ivanisova E. The antioxidant and antimicrobial activity of essential oils against *Pseudomonas* spp. isolated from fish. *Saudi Pharm J.* 2017 Dec;25(8):1108-16.

<https://doi.org/10.17221/96/2023-VETMED>

- Kaleo IV, Gao Q, Liu B, Sun C, Zhou Q, Zhang H, Shan F, Xiong Z, Bo L, Song C. Effects of *Moringa oleifera* leaf extract on growth performance, physiological and immune response, and related immune gene expression of *Macrobrachium rosenbergii* with *Vibrio anguillarum* and ammonia stress. *Fish Shellfish Immunol.* 2019 Jun; 89:603-13.
- Kavitha C, Ramesh M, Kumaran SS, Lakshmi SA. Toxicity of *Moringa oleifera* seed extract on some hematological and biochemical profiles in a freshwater fish, *Cyprinus carpio*. *Exp Toxicol Pathol.* 2012 Nov;64(7-8):681-7.
- Koshinski R. Effect of *Artemisia annua* L. extract on growth performance, biochemical blood parameters and meat quality of rainbow trout (*Oncorhynchus mykiss* W.), cultivated in recirculating system. *Agric Sci Technol.* 2018 Sep;10(3):266-70.
- Kumar S, Raman RP, Pandey PK, Mohanty S, Kumar A, Kumar K. Effect of orally administered azadirachtin on non-specific immune parameters of goldfish *Carassius auratus* (Linn. 1758) and resistance against *Aeromonas hydrophila*. *Fish Shellfish Immunol.* 2013 Feb;34(2): 564-73.
- Kuo IP, Lee PT, Nan FH. *Rheum officinale* extract promotes the innate immunity of orange-spotted grouper (*Epinephelus coioides*) and exerts strong bactericidal activity against six aquatic pathogens. *Fish Shellfish Immunol.* 2020 Jul;102:117-24.
- Kurian A, Van Doan H, Tapingkae W, Elumalai P. Modulation of mucosal parameters, innate immunity, growth and resistance against *Streptococcus agalactiae* by enrichment of Nile tilapia (*Oreochromis niloticus*) diet with *Leucas aspera*. *Fish Shellfish Immunol.* 2020 Feb;97: 165-72.
- Lafferty KD, Harvell CD, Conrad JM, Friedman CS, Kent ML, Kuris AM, Powell EN, Rondeau D, Saksida SM. Infectious diseases affect marine fisheries and aquaculture economics. *Annu Rev Mar Sci.* 2015;7:471-96.
- Levy G, Zilberg D, Paladini G, Fridman S. Efficacy of ginger-based treatments against infection with *Gyrodactylus turnbulli* in the guppy [*Poecilia reticulata* (Peters)]. *Vet Parasitol.* 2015 Apr;209(3-4):235-41.
- Li M, Zhu X, Tian J, Liu M, Wang G. Dietary flavonoids from *Allium mongolicum* Regel promotes growth, improves immune, antioxidant status, immune-related signaling molecules and disease resistance in juvenile northern snakehead fish (*Channa argus*). *Aquaculture.* 2018 Feb;501:473-81.
- Liu B, Xie J, Ge X, Xu P, Wang A, He Y, Zhou Q, Pan L, Chen R. Effects of anthraquinone extract from *Rheum officinale* Bail on the growth performance and physiological responses of *Macrobrachium rosenbergii* under high temperature stress. *Fish Shellfish Immunol.* 2010a Jul;29(1):49-57.
- Liu YT, Wang F, Wang GX, Han J, Wang Y, Wang YH. In vivo anthelmintic activity of crude extracts of *Radix angelicae pubescentis*, *Fructus bruceae*, *Caulis spatholobi*, *Semen aesculi*, and *Semen pharbitidis* against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitol Res.* 2010b Apr;106(5):1233-9.
- Liu B, Xu P, Xie J, Ge X, Xia S, Song C, Zhou Q, Miao L, Ren M, Pan L, Chen R. Effects of emodin and vitamin E on the growth and crowding stress of Wuchang bream (*Megalobrama amblycephala*). *Fish Shellfish Immunol.* 2014 Oct;40(2):595-602.
- Ma Q, Jiang JG, Yuan X, Qiu K, Zhu W. Comparative anti-tumor and anti-inflammatory effects of flavonoids, saponins, polysaccharides, essential oil, coumarin and alkaloids from *Cirsium japonicum* DC. *Food Chem Toxicol.* 2019 Mar;125:422-9.
- Makkar HP, Francis G, Becker K. Bioactivity of phytochemicals in some lesser-known plants and their effects and potential applications in livestock and aquaculture production systems. *Animal.* 2007 Oct;1(9):1371-91.
- Masschalck B, Michiels C. Antimicrobial properties of lysozyme in relation to foodborne vegetative bacteria. *Crit Rev Microbiol.* 2003;29(3):191-214.
- Mehrinakhi Z, Ahmadifar E, Sheikhzadeh N, Moghadam MS, Dawood MAO. Extract of grape seed enhances the growth performance, humoral and mucosal immunity, and resistance of common carp (*Cyprinus carpio*) against *Aeromonas hydrophila*. *Ann Anim Sci.* 2021 Jan;21(1): 217-32.
- Militz TA, Southgate PC, Carton AG, Hutson KS. Dietary supplementation of garlic (*Allium sativum*) to prevent monogenean infection in aquaculture. *Aquaculture.* 2013 Sep;408:95-9.
- Miranda CD, Godoy FA, Lee MR. Current status of the use of antibiotics and the antimicrobial resistance in the Chilean salmon farms. *Front Microbiol.* 2018 Jun;9:1284.
- Mo WY, Lun CHI, Choi WM, Man YB, Wong MH. Enhancing growth and non-specific immunity of grass carp and Nile tilapia by incorporating Chinese herbs (*Astragalus membranaceus* and *Lycium barbarum*) into food waste based pellets. *Environ Pollut.* 2016 Dec;219:475-82.
- Mohammadi G, Rashidian G, Hoseinifar SH, Naserabad SS, Doan HV. Ginger (*Zingiber officinale*) extract affects growth performance, body composition, haematology, serum and mucosal immune parameters in common carp (*Cyprinus carpio*). *Fish Shellfish Immunol.* 2020 Apr; 99:267-73.
- Moreno MA, Zampini IC, Isla MI. Antifungal, anti-inflammatory and antioxidant activity of bi-herbal mixtures

<https://doi.org/10.17221/96/2023-VETMED>

- with medicinal plants from Argentinean highlands. *J Ethnopharmacol.* 2020 May;253:112642.
- Munaeni W, Widanarni, Yuhana M, Setiawati M, Wahyudi AT. Effect in white shrimp *Litopenaeus vannamei* of *Eleutherine bulbosa* (Mill.) Urb. Powder on immune genes expression and resistance against *Vibrio parahaemolyticus* infection. *Fish Shellfish Immunol.* 2020 Jul;102:218-27.
- Naderi Farsani M, Hoseinifar SH, Rashidian G, Ghafari Farsani H, Ashouri G, Van Doan H. Dietary effects of *Coriandrum sativum* extract on growth performance, physiological and innate immune responses and resistance of rainbow trout (*Oncorhynchus mykiss*) against *Yersinia ruckeri*. *Fish Shellfish Immunol.* 2019 Aug;91:233-40.
- Naiel MAE, Ismael NEM, Negm SS, Ayyat MS, Al Sagheer AA. Rosemary leaf powder–supplemented diet enhances performance, antioxidant properties, immune status, and resistance against bacterial diseases in Nile Tilapia (*Oreochromis niloticus*). *Aquaculture.* 2020 Sep;526:735370.
- Ngo HV, Huang HT, Lee PT, Liao ZH, Chen HY, Nan FH. Effects of *Phyllanthus amarus* extract on nonspecific immune responses, growth, and resistance to *Vibrio alginolyticus* in white shrimp *Litopenaeus vannamei*. *Fish Shellfish Immunol.* 2020 Dec;107:1-8.
- Oduro I, Ellis W, Owusu D. Nutritional Potential of two leafy vegetables: *Moringa oleifera* and *Ipomoea batatas* leaves. *Sci Res Essay.* 2008;3:57-60.
- Palanikani R, Chanthini KMP, Soranam R, Thanigaivel A, Karthi S, Senthil-Nathan S, Murugesan AG. Efficacy of *Andrographis paniculata* supplements induce a non-specific immune system against the pathogenicity of *Aeromonas hydrophila* infection in Indian major carp (*Labeo rohita*). *Environ Sci Pollut Res.* 2020;27:23420-36.
- Palanikumar P, Daffni Benitta DJ, Lelin C, Thirumalaikumar E, Michaelbabu M, Citarasu T. Effect of *Argemone mexicana* active principles on inhibiting viral multiplication and stimulating immune system in Pacific white leg shrimp *Litopenaeus vannamei* against white spot syndrome virus. *Fish Shellfish Immunol.* 2018 Apr;75:243-52.
- Pasnik D, Evans J, Panangala V, Klesius P, Shelby R, Shoemaker C. Antigenicity of *Streptococcus agalactiae* extracellular products and vaccine efficacy. *J Fish Dis.* 2005 Apr;28(4):205-12.
- Prathomya P, Hussein H, Tuan T, Duan YH, Peng SH, Xu Y, Qian XQ, Wang GT. Effects of dietary supplementation of *Astragalus membranaceus*, *Codonopsis pilosula*, and *Glycyrrhiza uralensis* extract mixture on growth performance, haematological parameters and hepatopancreatic performance in juvenile Pacific white shrimp (*Litopenaeus vannamei*). *Aquac Res.* 2019;50:2707-17.
- Pridgeon JW, Klesius PH. Major bacterial diseases in aquaculture and their vaccine development. *CABI Reviews.* 2012 Oct;1-16.
- Pulkkinen K, Suomalainen LR, Read AF, Ebert D, Rintamaki P, Valtonen ET. Intensive fish farming and the evolution of pathogen virulence: The case of columnaris disease in Finland. *Proc Biol Sci.* 2010 Feb;277(1681):593-600.
- Punitha SMJ, Babu MM, Sivaram V, Shankar V, Dhas SA, Mahesh TC, GI Citarasu. Immunostimulating influence of herbal biomedicines on nonspecific immunity in Grouper *Epinephelus tauvina* juvenile against *Vibrio harveyi* infection. *Aquac Int.* 2008 Dec;16(6):511-23.
- Radhakrishnan S, Saravana Bhavan P, Seenivasan C, Shanthi R, Poongodi R. Influence of medicinal herbs (*Alternanthera sessilis*, *Eclipta alba* and *Cissus quadrangularis*) on growth and biochemical parameters of the freshwater prawn *Macrobrachium rosenbergii*. *Aquac Int.* 2014;22:551-72.
- Rafieepour A, Hajirezaee S, Rahimi R. Dietary oregano extract (*Origanum vulgare* L.) enhances the antioxidant defence in rainbow trout, *Oncorhynchus mykiss* against toxicity induced by organophosphorus pesticide, diazinon. *Tox Rev.* 2019 Jan;39(4):1-11.
- Raja Rajeswari P, Velmurugan S, Michael Babu M, Albin Dhas S, Kesavan KP, Citarasu T. A study on the influence of selected Indian herbal active principles on enhancing the immune system in *Fenneropenaeus indicus* against *Vibrio harveyi* infection. *Aquac Int.* 2012;20(5):1009-20.
- Rajabiesterabadi H, Hoseini SM, Fazelan Z, Hoseinifar SH, Doan HV. Effects of dietary turmeric administration on stress, immune, antioxidant and inflammatory responses of common carp (*Cyprinus carpio*) during copper exposure. *Aquac Nutr.* 2020;26(4):1143-53.
- Rajeshwari S, Rajan M, Pavaraj M, Sevarkodiyone S. Effect of *Melia azedarach* extract on some selected physiological parameters of (*Catla catla*). *Int J Aquac Fish Sci.* 2016 Jul;2(1):27-30.
- Ramesh D, Souissi S. Antibiotic resistance and virulence traits of bacterial pathogens from infected freshwater fish, *Labeo rohita*. *Microb Pathog.* 2018 Mar;116:113-9.
- Ramezanzadeh S, Abedian Kenari A, Esmaeili MA. Immunohematological parameters of rainbow trout (*Oncorhynchus mykiss*) fed supplemented diet with different forms of barberry root (*Berberis vulgaris*). *Comp Clin Pathol.* 2019;29:177-87.
- Reverter M, Bontemps N, Lecchini D, Banaigs B, Sasal P. Use of plant extracts in fish aquaculture as an alternative to chemotherapy: Current status and future perspectives. *Aquaculture.* 2014 Sep;433:50-61.
- Reverter M, Tapissier Bontemps N, Sasal P, Saulnier D. Use of medicinal plants in aquaculture. In: Austin B, Newaj-

<https://doi.org/10.17221/96/2023-VETMED>

- Fyzul A, editors. Diagnosis and control of diseases of fish and shellfish. Hoboken, New Jersey, USA: John Wiley & Sons Ltd.; 2017. p. 223-61.
- Rico A, Phu TM, Satapornvanit K, Min J, Shahabuddin AM, Henriksson PJ, Murray FJ, Little DC, Dalsgaard A, Van den Brink PJ. Use of veterinary medicines, feed additives and probiotics in four major internationally traded aquaculture species farmed in Asia. *Aquaculture*. 2013 Nov; 412:231-43.
- Roomiani L, Soltani M, Akhondzadeh Basti A, Mahmoodi A, Taheri A, Taher Mirghaed A, Yadollahi F. Evaluation of the chemical composition and in vitro antimicrobial activity of *Rosmarinus officinalis*, *Zataria multiflora*, *Anethum graveolens* and *Eucalyptus globulus* against *Streptococcus iniae* the cause of zoonotic disease in farmed fish. *Iran J Fish Sci*. 2013;12(3):702-16.
- Rufchaei R, Hoseinifar SH, Mirzajani A, Van Doan H. Dietary administration of *Pontogammarus maoticus* extract affects immune responses, stress resistance, feed intake and growth performance of caspian roach (*Rutilus caspicus*) fingerlings. *Fish Shellfish Immunol*. 2017 Apr; 63:196-200.
- Safari O, Sarkheil M, Paolucci M. Dietary administration of ferula (*Ferula asafoetida*) powder as a feed additive in diet of koi carp, *Cyprinus carpio* koi: Effects on hemato-immunological parameters, mucosal antibacterial activity, digestive enzymes, and growth performance. *Fish Physiol Biochem*. 2019 Aug;45(4):1277-88.
- Saiyad Musthafa M, Asgari SM, Kurian A, Elumalai P, Jawahar Ali AR, Paray BA, Al-Sadoon MK. Protective efficacy of *Mucuna pruriens* (L.) seed meal enriched diet on growth performance, innate immunity, and disease resistance in *Oreochromis mossambicus* against *Aeromonas hydrophila*. *Fish Shellfish Immunol*. 2018 Apr;75:374-80.
- Sakai M. Current research status of fish immunostimulants. *Aquaculture*. 1999 Mar;172(1-2):63-92.
- Sambasivam S, Karpagam G, Chandran R, Khan SA. Toxicity of leaf extract of yellow oleander *Thevetia nerifolia* on *Tilapia*. *J Environ Biol*. 2003 Apr;24(2):201-4.
- Santoso U, Lee MC, Nan FH. Effects of dietary katuk leaf extract on growth performance, feeding behavior and water quality of grouper *Epinephelus coioides*. *Aceh Int J Sci Tech*. 2013 Apr;2(1):17-25.
- Sarhadi I, Alizadeh E, Ahmadifar E, Adineh H, Dawood MAO. Skin mucosal, serum immunity and antioxidant capacity of common carp (*Cyprinus carpio*) fed *Artemisia* (*Artemisia annua*). *Ann Anim Sci*. 2020;20(3):1011-27.
- Sarhan IA, Abdel-Aziz SA, Said AA, Abdel-Aleim AAF, Awad SM. Effect of dietary supplementation of extracted jojoba meal on hematology, biochemical parameters and disease resistance in Nile tilapia (*Oreochromis niloticus*) infected by *Aeromonas hydrophila*. *Egypt J Aquac*. 2019 Sep;9(3):13-31.
- Shan XF, Meng QF, Kang YH, Bian Y, Gao YH, Wang WL, Ai-dong Q. Isolation of active compounds from methanol extracts of *Toddalia asiatica* against *Ichthyophthirius multifiliis* in goldfish (*Carassius auratus*). *Vet Parasitol*. 2014 Jan;199(3-4):250-4.
- Sharma J, Kumar N, Singh SP, Singh A, Harikrishna V. Evaluation of immunostimulatory properties of prickly chaff flower *Achyranthes aspera* in rohu *Labeo rohita* fry in pond conditions. *Aquaculture*. 2019 Apr;505:183-9.
- Shen YF, Liu L, Feng C-Z, Hu Y, Chen C, Wang GX, Zhu B. Synthesis and antiviral activity of a new coumarin derivative against spring viraemia of carp virus. *Fish Shellfish Immunol*. 2018 Oct;81:57-66.
- Shinn A, Pratoomyot J, Bron J, Paladini G, Brooker E, Brooker A. Economic impacts of aquatic parasites on global finfish production. *Glob Aquacult Advocate*. 2015;2015:58-61.
- Sikotariya S. Effect of *Allium cepa* (Onion) powder on the growth and survival in *Cirrhinus mrigala* fingerlings. *Int J Pure Appl Biosci*. 2019;7(5):186-96.
- Singh A, Sharma J, Paichha M, Chakrabarti R. *Achyranthes aspera* (Prickly chaff flower) leaves- and seeds-supplemented diets regulate growth, innate immunity, and oxidative stress in *Aeromonas hydrophila*-challenged *Labeo rohita*. *J Appl Aquac*. 2019;32:1-18.
- Sirakov I, Velichkova K, Stoyanova S, Zhelyazkov G, Staykov Y. The effect of diet supplemented with dandelion's (*Taraxacum Officinale*) extract on the productive and blood parameters of common carp (*Cyprinus Carpio* L.), cultivated in the recirculation system. *Maced Vet Rev*. 2019;42:131-9.
- Soares MP, Cardoso IL, Ishikawa MM, de Oliveira A, Sartoratto A, Jonsson CM, de Queiroz SCDN, Duarte MCT, Rantin FT, Sampaio FG. Effects of *Artemisia annua* alcohol extract on physiological and innate immunity of Nile tilapia (*Oreochromis niloticus*) to improve health status. *Fish Shellfish Immunol*. 2020 Oct;105:369-77.
- Sofia F. The state of world fisheries and aquaculture 2018 – Meeting the sustainable development goals. Rome: Fisheries and Aquaculture Department, Food and Agriculture Organization of the United Nations; 2018.
- Song C, Liu B, Jiang S, Xiong Y, Sun C, Zhou Q, Jiang Z, Liu B, Zhang H. Anthraquinone extract from *Rheum officinale* Bail improves growth performance and Toll-Relish signaling-regulated immunity and hyperthermia tolerance in freshwater prawn *Macrobrachium nipponense*. *3 Biotech*. 2020 Dec;10(12):526.
- Song K, Ling F, Huang A, Dong W, Liu G, Jiang C, Zhang Q, Wang G. In vitro and in vivo assessment of the effect

<https://doi.org/10.17221/96/2023-VETMED>

- of antiprotozoal compounds isolated from *Psoralea corylifolia* against *Ichthyophthirius multifiliis* in fish. *Int J Parasitol Drugs Drug Resist.* 2015 Aug;5(2):58-64.
- Srichaiyo N, Tongsiri S, Hoseinifar SH, Dawood MAO, Esteban M, Ringo E, Van Doan H. The effect of fishwort (*Houttuynia cordata*) on skin mucosal, serum immunities, and growth performance of Nile tilapia. *Fish Shellfish Immunol.* 2020a Mar;98:193-200.
- Srichaiyo N, Tongsiri S, Hoseinifar SH, Dawood MAO, Jaturasitha S, Esteban MA, Ringo E, Van Doan H. The effects of gotu kola (*Centella asiatica*) powder on growth performance, skin mucus, and serum immunity of Nile tilapia (*Oreochromis niloticus*) fingerlings. *Aquac Rep.* 2020b Mar;16:100239.
- Stankus A. State of world aquaculture 2020 and regional reviews: FAO webinar series. *FAO Aquaculture Newsletter.* 2021;63:17-8.
- Stentiford GD, Sritunyalucksana K, Flegel TW, Williams BA, Withyachumnarnkul B, Itsathitphaisarn O, Bass D. New paradigms to help solve the global aquaculture disease crisis. *PLoS Pathog.* 2017 Feb;13(2):e1006160.
- Su M, Tang R, Wang H, Lu L. Suppression effect of plant-derived berberine on cyprinid herpesvirus 2 proliferation and its pharmacokinetics in Crucian carp (*Carassius auratus gibelio*). *Antiviral Res.* 2021 Feb;186:105000.
- Sukumaran V, Park SC, Giri SS. Role of dietary ginger *Zingiber officinale* in improving growth performances and immune functions of *Labeo rohita* fingerlings. *Fish Shellfish Immunol.* 2016 Oct;57:362-70.
- Sun Z, Chen L, Liu Q, Mai K, Xu M, Zhou Y, Su N, Ye C. Effects of dietary *Senecio scandens* buch-ham extracts on growth performance, plasma biochemical, histology and the expression of immune-related genes in hybrid grouper (*Epinephelus lanceolatus* ♂ × *Epinephelus fuscoguttatus* ♀). *Fish Shellfish Immunol.* 2020 Mar;98:681-90.
- Ta I, Sa D, Ha A. Growth performance and hematological indices of *Clarias gariepinus* (Burchel, 1822) fingerlings fed varying levels of *Telfairia occidentalis* leaf meal additives. *Int J Fish Aquat Stud.* 2019;7(5):442-5.
- Tadese D, Song C, Sun CX, Bo L, Zhou Q, Xu P, Ge X, Liu M, Xu X, Tamiru M, Zhou Z, Lakew A, Kevin NT. The role of currently used medicinal plants in aquaculture and their action mechanisms: A review. *Rev Aquac.* 2021:1-32.
- Taheri Mirghaied A, Paknejad H, Mirzargar SS. Hepatoprotective effects of dietary *Artemisia* (*Artemisia annua*) leaf extract on common carp (*Cyprinus carpio*) exposed to ambient ammonia. *Aquaculture.* 2020 Oct;527:735443.
- Talpur AD. *Mentha piperita* (Peppermint) as feed additive enhanced growth performance, survival, immune response and disease resistance of Asian seabass, *Lates calcarifer* (Bloch) against *Vibrio harveyi* infection. *Aquaculture.* 2014 Jan;420-421:71-8.
- Tan X, Sun Z. Dietary dandelion extract improved growth performance, immunity, intestinal morphology and microbiota composition of golden pompano *Trachinotus ovatus*. *Aquac Rep.* 2020 Nov;18:100491.
- Tan X, Sun Z, Ye C, Lin H. The effects of dietary *Lycium barbarum* extract on growth performance, liver health and immune related genes expression in hybrid grouper (*Epinephelus lanceolatus* ♂ × *E. fuscoguttatus* ♀) fed high lipid diets. *Fish Shellfish Immunol.* 2019 Apr;87:847-52.
- Umeda N, Nibe H, Hara T, Hirazawa N. Effects of various treatments on hatching of eggs and viability of oncomiracidia of the monogenean *Pseudodactylogyrus anguillae* and *Pseudodactylogyrus bini*. *Aquaculture.* 2006 Mar;253(1-4):148-53.
- Valladao GM, Gallani SU, Ikefuti CV, da Cruz C, Levy-Pereira N, Rodrigues MV, Pilarski F. Essential oils to control ichthyophthiriasis in pacu, *Piaractus mesopotamicus* (Holmberg): Special emphasis on treatment with *Melaleuca alternifolia*. *J Fish Dis.* 2016 Oct;39(10):1143-52.
- Vallejos-Vidal E, Reyes-Lopez E, Teles M, MacKenzie S. The response of fish to immunostimulant diets. *Fish Shellfish Immunol.* 2016 Sep;56:34-69.
- Van Doan H, Hoseinifar SH, Sringarm K, Jaturasitha S, Yuangsoi B, Dawood MAO, Esteban MA, Ringo E, Faggio C. Effects of Assam tea extract on growth, skin mucus, serum immunity and disease resistance of Nile tilapia (*Oreochromis niloticus*) against *Streptococcus agalactiae*. *Fish Shellfish Immunol.* 2019 Oct;93:428-35.
- Van Hai N. The use of medicinal plants as immunostimulants in aquaculture: A review. *Aquaculture.* 2015 Sep;446:88-96.
- van Wyk AS, Prinsloo G. Health, safety and quality concerns of plant-based traditional medicines and herbal remedies. *S Afr J Bot.* 2020 Sep;133:54-62.
- Whittington ID. *Benedenia seriola* and *Neobenedenia* species. *Fish parasites: Pathobiology and protection.* UK: CABI Wallingford; 2012. p. 225-44.
- Wu CC, Liu CH, Chang YP, Hsieh SL. Effects of hot-water extract of *Toona sinensis* on immune response and resistance to *Aeromonas hydrophila* in *Oreochromis mossambicus*. *Fish Shellfish Immunol.* 2010 Aug;29(2):258-63.
- Wu ZF, Zhu B, Wang Y, Lu C, Wang GX. In vivo evaluation of anthelmintic potential of medicinal plant extracts against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitol Res.* 2011 Jun;108(6):1557-63.
- Wu Z, Ling F, Song C, Chen W, Wang G. Effects of oral administration of whole plants of *Artemisia annua* on *Ichthyophthirius multifiliis* and *Aeromonas hydrophila* after

<https://doi.org/10.17221/96/2023-VETMED>

- parasitism by *I. multifiliis*. *Parasitol Res.* 2017 Jan;116(1): 91-7.
- Wu C, Shan J, Feng J, Wang J, Qin C, Nie G, Ding C. Effects of dietary *Radix Rehmanniae Preparata* polysaccharides on the growth performance, immune response and disease resistance of *Luciobarbus capito*. *Fish Shellfish Immunol.* 2019 Jun;89:641-6.
- Wu S. Dietary *Astragalus membranaceus* polysaccharide ameliorates the growth performance and innate immunity of juvenile crucian carp (*Carassius auratus*). *Int J Biol Macromol.* 2020 Apr 15;149:877-81.
- Xie J, Bo L, Zhou Q, Su Y, He Y, Pan L, Ge X, Xu P. Effects of anthraquinone extract from rhubarb *Rheum officinale* Bail on the crowding stress response and growth of common carp *Cyprinus carpio* var. Jian. *Aquaculture.* 2008 Sep;281:5-11.
- Yi YL, Lu C, Hu XG, Ling F, Wang GX. Antiprotozoal activity of medicinal plants against *Ichthyophthirius multifiliis* in goldfish (*Carassius auratus*). *Parasitol Res.* 2012 Oct;111(4):1771-8.
- Yoshida T, Kruger R, Inglis V. Augmentation of non specific protection in African catfish, *Clarias gariepinus* (Burchell), by the long term oral administration of immunostimulants. *J Fish Dis.* 1995;18(2):195-8.
- Yousefi M, Ghafarifarsani H, Hoseinifar SH, Rashidian G, Van Doan H. Effects of dietary marjoram, *Origanum majorana* extract on growth performance, hematological, antioxidant, humoral and mucosal immune responses, and resistance of common carp, *Cyprinus carpio* against *Aeromonas hydrophila*. *Fish Shellfish Immunol.* 2021 Jan;108:127-33.
- Yuan C, Li D, Chen W, Sun F, Wu G, Gong Y, Tang J, Shen M, Han X. Administration of a herbal immunoregulation mixture enhances some immune parameters in carp (*Cyprinus carpio*). *Fish Physiol Biochem.* 2007;33:93-101.
- Yunus K, Jaafar AM, Akbar J. Acute-lethal toxicity (LC50) effect of *Terminalia Catappa* Linn. leaves extract on *Oreochromis Niloticus* (Red Nile Tilapia) juveniles under static toxicity exposure. *Orient J Chem.* 2019;35(1):270-4.
- Zemheri-Navruz F, Acar U, Yilmaz S. Dietary supplementation of olive leaf extract enhances growth performance, digestive enzyme activity and growth related genes expression in common carp *Cyprinus carpio*. *Gen Comp Endocrinol.* 2020 Sep;296:113541.
- Zhai Q, Li J. Effectiveness of traditional Chinese herbal medicine, San-Huang-San, in combination with enrofloxacin to treat AHPND-causing strain of *Vibrio parahaemolyticus* infection in *Litopenaeus vannamei*. *Fish Shellfish Immunol.* 2019 Apr;87:360-70.
- Zhang Q, Xu DH, Klesius PH. Evaluation of an antiparasitic compound extracted from *Galla chinensis* against fish parasite *Ichthyophthirius multifiliis*. *Vet Parasitol.* 2013 Nov;198(1-2):45-53.
- Zhang A, Wu Y, Tan Y, Shi J, Zhao Y, Hu Y, Yu J, Zheng W, Lai X, Zhang M, Zhu Y, Ye Y, Huang Y, Fu S, Huang H, Luo Y. Determining whether prophylactic antiviral treatment is necessary in HBsAg-negative/HBcAb-positive patients receiving allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant.* 2020a May;26(5):956-64.
- Zhang X, Sun Z, Cai J, Wang J, Wang G, Zhu Z, Cao F. Effects of dietary fish meal replacement by fermented moringa (*Moringa oleifera* Lam.) leaves on growth performance, nonspecific immunity and disease resistance against *Aeromonas hydrophila* in juvenile gibel carp (*Carassius auratus gibelio* var. CAS III). *Fish Shellfish Immunol.* 2020b Jul;102:430-9.
- Zhang Y, Song L, Guo H, Wu J, Wang X, Yao F. Effects of curcumin on growth and liver protection in common carp, *Cyprinus carpio*. *Pakistan J Zool.* 2021 Aug;53(4): 1211-20.
- Zoral MA, Futami K, Endo M, Maita M, Katagiri T. Anthelmintic activity of *Rosmarinus officinalis* against *Dactylogyrus minutus* (Monogenea) infections in *Cyprinus carpio*. *Vet Parasitol.* 2017 Nov 30;247:1-6.

Received: September 22, 2023

Accepted: November 30, 2023

Published online: December 26, 2023