

Journal section: Community and Preventive Dentistry  
 Publication Types: Research

doi:10.4317/jced.61052  
<https://doi.org/10.4317/jced.61052>

## Enamel fluorosis related with fluoride-containing water ingestion and urinary excretion in schoolchildren

Farith González-Martínez <sup>1,2,3</sup>, Ruth Gómez-Scarpetta <sup>3,4</sup>, Olga Bibiana Salcedo <sup>5</sup>, Patricia Bermúdez-Reyes <sup>6</sup>, Patricia Castro-Villamizar <sup>7</sup>, María del Pilar Cerezo <sup>8</sup>, Carmen Martínez <sup>9</sup>, Lesbia Tirado-Amador <sup>10</sup>, Andrés Salas-Zambrano <sup>11</sup>, Alexandra Saldarriaga <sup>12</sup>, Marisol Sánchez-Molina <sup>13</sup>, Luis-Armando Vila <sup>14</sup>

<sup>1</sup> DDS, MSPH, PhD. Public Health Research Group Universidad de Cartagena-Cartagena de Indias, Colombia

<sup>2</sup> Asociación Colombiana de Facultades de Odontología-Bogotá-ACFO, Colombia

<sup>3</sup> Latin American Network of Research on Fluorides and Dental Fluorosis, Cartagena 130014, Colombia

<sup>4</sup> BAM, MSc. Universidad Cooperativa de Colombia-Villavicencio, Colombia

<sup>5</sup> DDS, MSc. Universidad de Antioquia-Medellín, Colombia

<sup>6</sup> DDS, MSc. Universidad Cooperativa de Colombia-Envigado, Colombia

<sup>7</sup> DDS, MSc. Corporación Universitaria Rafael Núñez, - Cartagena, Colombia

<sup>8</sup> DDS, MSc. Universidad Autónoma de Manizales-Manizales, Colombia

<sup>9</sup> DDS, MSc. Universidad Santo Tomás-Bucaramanga, Colombia

<sup>10</sup> DDS, MSc. Universidad del Sinú-Cartagena, Colombia

<sup>11</sup> DDS, MSc. Universidad Cooperativa de Colombia-Pasto, Colombia

<sup>12</sup> DDS, MSc, PhD. Universidad CES-Medellín, Colombia

<sup>13</sup> DDS, MSc, PhD Universidad Metropolitana-Barranquilla, Colombia

<sup>14</sup> DDS, MSc, PhD. Universidad del Magdalena-Santa Marta, Colombia

### Correspondence:

University of Cartagena, Zaragocilla  
 Carrera 50 No. 24-99  
 130014 Cartagena, Colombia  
[fgonzalezml@unicartagena.edu.co](mailto:fgonzalezml@unicartagena.edu.co)

Received: 22/09/2023

Accepted: 23/10/2023

González-Martínez F, Gómez-Scarpetta R, Salcedo OB, Bermúdez-Reyes P, Castro-Villamizar P, Cerezo MP, Martínez C, Tirado-Amador L, Salas-Zambrano A, Saldarriaga A, Sánchez-Molina M, Vila LA. Enamel fluorosis related with fluoride-containing water ingestion and urinary excretion in schoolchildren. J Clin Exp Dent. 2024;16(1):e51-61.

Article Number: 61052 <http://www.medicinaoral.com/odo/indice.htm>

© Medicina Oral S. L. C.I.F. B 96689336 - eISSN: 1989-5488

eMail: [jced@jced.es](mailto:jced@jced.es)

#### Indexed in:

Pubmed  
 Pubmed Central® (PMC)  
 Scopus  
 DOI® System

### Abstract

**Background:** Natural water sources are considered as the major environmental exposure of fluoride, resulting in increased prevalence of enamel fluorosis. This type of natural exposure should be permanently monitored to avoid the interactions with other non-natural fluoride sources. We evaluated the prevalence of enamel fluorosis in Colombian schoolchildren and its relationship with fluoride-containing water ingestion exposure dose and urinary fluoride excretion.

**Material and Methods:** We included 923 schoolchildren aged 7–12 years residing in eight municipalities in Colombia. Sampling of consumption water was performed in major aquifers used for daily supply. Samples were collected in 98 polyethylene containers and refrigerated until analysis. Water and urine fluoride concentrations were measured using the fluoride selective electrode method. Enamel fluorosis was evaluated using Thylstrup and Ferjerskov Index (TFI). Demographic and anthropometric characteristics were assessed. Besides, other exposures to non-natural fluoride were also evaluated. Logistic regression was applied for multiple analyses.

Results: The median fluoride concentration in water and urine samples was 10.5 mg/L and 0.63 mg/L respectively, with the highest value found in Algarrobo-Magdalena, and the lowest value found in Manzanares-Caldas. The overall prevalence of enamel fluorosis was 86.1%, being more frequent the mild codes with TFI-1 to TFI-2. The highest prevalence was found in Margarita-Bolívar and Manzanares-Caldas, and the most severe codes (TFI-5 to TFI-9) were detected in Manzanares-Caldas. The multiple analysis revealed water ingestion exposure dose, urinary excretion, involuntary intake of toothpaste, amount of table salt consumption and sex as significant factors ( $p < 0.001$ ).

Conclusions: The fluoride ingestion exposure dose and its subsequent urinary excretion could be used as estimators of past fluoride exposure, explaining the current prevalence of enamel fluorosis in Colombian schoolchildren.

**Key words:** Fluoride, groundwater ingestion, enamel fluorosis, prevalence, severity.

## Introduction

Fluorine is a naturally occurring element with a widespread distribution, primarily in geological environments. It is also present in non-geological environments in substrates such as water, soil, air and vegetation (1). The most commonly identified compounds of environmental origin with fluorine content are fluorite (calcium fluoride), cryolite (sodium and aluminium fluoride), apatite (calcium phosphate) and phosphate rocks (2,3).

Artificial addition of fluorides to sources of intake such as water and table salt has been performed globally since 1950 as the major strategy for controlling the progression of dental caries. Although drinking water in Colombia does not contain added fluoride since 1989, when the public policy was changed by adding fluoride to table salt, the country established a control and surveillance system for the quality of water for human consumption (SIVICAP), which guarantees the monitoring of fluoride concentration levels in drinking water, with the maximum acceptable value of fluorides in water being 1 mg/L (4,5). However, this system does not have sufficient coverage to monitor all sources of water access for the Colombian population, because several communities residing in rural areas have a service coverage of 74 % (6), and this consumption is supplied by unconventional aquifers that are generally not included in the water samplings performed by territorial entities.

Since 2012, the National Institute of Health of Colombia has implemented a sentinel surveillance programme for fluoride exposure (SIVIGILA), which monitors fluoride levels in drinking water and table salt in some municipalities selected by the territorial entities. Between 2012 and 2015, the programme has identified 19 municipalities with fluoride values in water samples of  $>1$  mg/L. Similarly, during this same period, 53 municipalities were identified with fluoride values in table salt samples of  $>220$  mg F/kg/salt (7). Although the programme is still active, it has a limitation of under-records in the data due to lack of samplings that represent the water and salt consumption of all participating populations.

Excessive intake of fluoride on a regular basis during the tooth formation period causes a clinical defect in enamel development, which is characterised by greater discoloration and porosity than the normal enamel and is known as enamel fluorosis (8,9). Although the optimal levels of daily fluoride intake are inconsistent, some studies indicate that intake doses of  $>0.07$  mg F/kg w/day can cause dental fluorosis (10).

The last National Oral Health Survey performed in Colombia in 2014 (11), reported a prevalence of enamel fluorosis of 59.1%, confirming a significant increase compared with the National Oral Health Survey performed in 1998 (12), which reported a prevalence of 11.8 %. These data indicate possible failures in the implementation of public policy to control fluoride intake from different sources. It can be clearly recognised that massive access to fluoride sources has generated an effect on dental caries in the country by reducing its prevalence (11). Nevertheless, the non-beneficial effects of fluoride have increased concern at the level of government entities, to the point of being considered as a public health problem.

Chronic toxicity and the increase in the prevalence and severity of enamel fluorosis have been associated with exposure to different sources of fluoride administration, especially sources of systemic intake such as water, table salt, toothpastes, fluoride supplements and diet (13-18). Therefore, the impact on chronic toxicity of the various sources of fluorides is related to several variables, including the time of exposure, concentration and quantity and frequency of consumption, as well as the urinary fluoride excretion, which become a highly complex exposure, making it difficult to control (19-21).

The above-described background information indicates the need to explore the effects of exposure to fluorides by groundwater ingestion, including in the explanatory models all the variables that intervene in this process. Therefore, we investigated the prevalence of enamel fluorosis in Colombian schoolchildren and its relationship with fluoride-containing water ingestion exposure doses and fluoride urinary excretion.

## Material and Methods

### -Study population

This study was conducted on 923 schoolchildren aged 7–12 years residing in the following eight municipalities in Colombia with exposure to fluoride due to groundwater ingestion: Algarrobo-Magdalena, Campo de la Cruz-Atlántico, La Estrella-Antioquia, Manzanares-Caldas, Margarita-Bolívar, Nariño-Nariño, Oiba-Santander and Puerto López-Meta. Eligible participants enrolled in this study have been living in these municipalities since birth and consumed groundwater on a regular basis. Exclusion criteria were medical history of renal disease and severe malnutrition.

The sample size was determined based on a type I error of 5% and a power of 80%. Based on previous data, using the Stata® software (V12, Stata Corp. LP, College Station, TX, USA), the lowest expected frequency for the prevalence of dental fluorosis was calculated (11), resulting in a target sample size of 987 subjects. To account for possible incomplete processing of the questionnaire, the sample size was increased by 10 % (1086). Thus, a total of 1086 schoolchildren were recruited, and 163 were excluded because they did not fulfil the parameters cited earlier. The final study population consisted of 923 participants.

### -Groundwater collection

A total of 98 groundwater well sites were selected in the study municipalities (Fig. 1). Water samples were

collected in 250-mL polyethylene bottles that were previously treated with 10 % HNO<sub>3</sub> for 2 h, rinsed with ultra-pure water (18 MΩ•cm; distilled and deionised water) and dried (to avoid contamination). The containers were also rinsed three times with the same sampling water before sample collection. After sample collection, the containers were refrigerated (three Celsius degrees), transported and stored in the laboratory until analysis, (Fig. 1).

### -Urine collection

The urine samples of the participants were collected in 50 mL sterile falcon tubes, taking a standard volume of 20 mL for analysis. This sample represents a single point in time, corresponding to the first morning urine. After sample collection, the containers were refrigerated (three Celsius degrees), transported and stored in the laboratory until analysis.

### Questionnaire

A survey was performed to evaluate the demographic and anthropometric information of the study participants. Moreover, details regarding the ingestion of groundwater and other fluoride exposure sources such as consumption of packaged beverages, involuntary intake of toothpaste, amount of toothpaste used, frequency of teeth brushing and amount of table salt consumption were collected. These data were obtained through a self-administered questionnaire, with a supplementary interview. This questionnaire was evaluated before its




Fig. 1: Location map of the study municipalities (Colombia, 2018). Source: own elaboration.

application by three expert judges to confirm its appearance validity. Furthermore, a reliability analysis stability, test-retest was applied in a pilot group of volunteer parents at two time intervals to correlate the obtained two data distributions. In addition, all interviewers were trained to administer the assigned questionnaires. The questionnaires were also monitored during the data collection process through a random review to assess adherence to the protocols by the interviewers.

#### -Quantification of fluoride in groundwater and urine

Fluoride concentrations were measured using a potentiometer with fluoride ion-selective electrode (Thermo Scientific™ Orion™; Fisher cat. #13-642-285 Mfr. # 9609BNWP) (22,23). The water and urine samples were analyzed by duplicate adding TISAB II buffer with CDTA just before analysis and using standard reference material (NIST-USA, 1984: SRM 3183). The measurement procedures were performed at the Public Health Research Laboratory of the School of Dentistry, University of Cartagena, Colombia. Urinary fluoride concentrations were adjusted by dilution to eliminate variation from fluid balance. Therefore, urinary creatinine was determined by spectrophotometer Genesys 10S UV-VIS (Thermo Scientific, USA) and urinary fluoride concentrations were expressed as mg F per g creatinine.

#### -Determination of water ingestion exposure doses (IDag)

Human exposure to fluoride in groundwater was evaluated using IDag (24). This index was expressed as risk using the reference dose (<0.05 mg/kg/day) and was calculated according to the following equation:  $IDag = (C \times IR \times EF) / (BW)$ , where C is the fluoride concentration in water (mg/L), IR is the water intake rate per day (L/day), EF is the exposure frequency (days/year) (a value of 1 was used, representing daily exposure to fluoride) and BW is the body weight (kg) (the average body weight was determined as 32 kg).

#### -Clinical examination

The clinical signs of enamel fluorosis were evaluated using the TF index. These results are indicators of chronic fluoride exposure, and each indicator of clinical injury has been correlated with histological characterisation. This index classifies the severity of enamel fluorosis lesions among nine indicators as follows: (TFI-0) = absence of visible lesion and (TFI-1 to TFI-9) = presence of clinically visible lesions. Based on the severity categories, the indicators were classified as (TFI-1 to TFI-2) = mild fluorosis, (TFI-3 to TFI-4) = moderate fluorosis and (TFI-5 to TFI-9) = severe fluorosis (25). Two examiners received training for enamel fluorosis measurements using the TFI. Clinical calibration values included inter-examiner and intra-examiner comparison, using an external examiner as a reference point. The two examiners obtained agreement scores above 0.75 for the weighted kappa coefficient.

#### -Statistical analyses

Statistical analyses were performed using the R softwa-

re (26). Proportions, absolute and relative frequencies and mean and standard deviation were used for statistical description. In addition, correlation coefficients were used to evaluate the relationship among the fluoride-containing water ingestion exposure dose and quantification of fluoride in urine. The prevalence of enamel fluorosis was calculated as the proportion of individuals with a given category by dividing into the total number of participants. The study population was divided into two subgroups for the purpose of analysis, and pairwise comparisons were performed between the presence and absence of enamel fluorosis and the severity levels according to TFI using the chi-square ( $\chi^2$ ) test ( $p < 0.05$ ). A complementary analysis was performed to understand the significance of differences between the presence and absence of enamel fluorosis according to the low and high exposure dose, effect size test (27). A multivariate logistic regression analysis was performed incorporating enamel fluorosis prevalence as a dependent variable and water ingestion exposure dose, urine excretion, involuntary intake of toothpaste, amount of table salt as independent variables. Potential confounders (age, sex, body mass index (BMI) and exposures to fluoride other than groundwater ingestion) were also considered in this study. Interaction terms among the water ingestion exposure dose that provided a  $p < 0.2$  and potential confounders were evaluated.

## Results

#### -Characterisation of the study municipalities

This study was conducted in eight municipalities in Colombia, distributed by different geographical areas, especially rural areas. The latitude and longitude of the study locations are depicted in Figure 1.

#### -Characteristics of study subjects

The mean age, height, weight and BMI of the study population were  $9.5 \pm 0.04$  years,  $1.4 \pm 0.003$  m,  $31.9 \pm 0.23$  kg and  $18.7$  kg/m<sup>2</sup>, respectively. Girls comprised 50.2 % of the study population, and the most frequent (20.8%) school grade was fourth grade of elementary school. An amount of table salt consumption of  $\geq 1$  spoonful (75.4 %) and an amount of toothpaste of more than or equal to half of the toothbrush (54.8 %) showed the highest frequency among fluoride exposure factors. The other characteristics are shown in Table 1.

#### -Concentration of fluoride in drinking water

The water samples contained a fluoride concentration of 0.02 to 24.3 mg/L, with the overall mean concentration being  $4.32 \pm 8.35$  mg/L. Algarrobo-Magdalena ( $24.3 \pm 26$  mg/L) and Margarita-Bolívar ( $6.3 \pm 5.7$  mg/L) were the municipalities with the highest concentration of fluoride in drinking water (Table 2).

#### -Water ingestion exposure dose (IDag) and Fluoride concentration in urine

The IDag values ranged from 0.0008 to 0.88 mg/kg/day,

**Table 1:** Characteristics of the study population.

Characteristics	N	(%) <sup>a</sup> or mean ± SE <sup>b</sup>
Total sample	923	100
Anthropometrics		
Age (years)	923	9.5 ± 0.04
Height (cm)	923	1.4 ± 0.003
Weight (kg)	923	31.9 ± 0.23
BMI (kg/m <sup>2</sup> ) <sup>c</sup>	923	18.7 ± 1.6
Municipalities		
Algarrobo-Magdalena	132	14.3
Campo de la Cruz-Atlántico	89	9.6
La Estrella-Antioquia	93	10.1
Manzanares-Caldas	121	13.1
Margarita- Bolívar	115	12.5
Nariño-Nariño	53	5.7
Oiba-Santander	153	16.6
Puerto López-Meta	167	18.1
Age (categorical, years)		
7–9	463	50.2
10–12	460	49.8
Sex		
Male	463	49.8
Female	460	50.2
Water ingestion exposure dose		
IDag ≥0.05 mg/kg/day	246	26.7
IDag <0.05 mg/kg/day	677	73.4
Consumption of packaged beverages		
Yes	290	31.4
No	633	68.6
Involuntary intake of toothpaste		
Yes	455	49.3
No	468	50.7
Amount of toothpaste used		
≥Half of toothbrush	506	54.8
<Half of toothbrush	417	45.2
Frequency of teeth brushing		
>3 times a day	327	35.4
≤3 times a day	596	64.6
Amount of table salt consumption		
≥1 spoonful	696	75.4
<1 spoonful	227	24.6

Note. a:percentage; baverage ± standard error; cbody mass index

and the median value was 0.002 mg/kg/day (0.001–0.17 mg/kg/day). The highest IDag values were found in Algarrobo-Magdalena (0.88 mg/kg/day) and Marga-

rita-Bolívar (0.22 mg/kg/day), above reference dose (<0.05 mg/kg/day). The urine samples contained a range fluoride concentration of 0.11-2.06 mg/F/g-creatinine with the overall mean concentration being 0.63 ± 0.82 mg/F/g-creatinine. The individuals residing in the municipalities of Algarrobo-Magdalena and Margarita-Bolívar showed the highest concentrations with 2.06 and 1.86 mg/F/g-creatinine respectively. Besides, the mean fluoride-containing water ingestion exposure dose (IDag) and the mean fluoride concentration in urine showed a positive and statistically significant correlation ( $r = 0.98$ ;  $p < 0.001$ ), (Table 2).

-Prevalence and severity of enamel fluorosis in school-children

The overall prevalence of fluorosis among the 923 children was 86.1 %. The highest dental fluorosis prevalence was found in Margarita-Bolívar and Manzanares-Caldas. We also detected a significant association between enamel fluorosis prevalence and municipalities ( $p < 0.001$ ) and between enamel fluorosis prevalence and sex ( $p = 0.03$ ) (Table 3).

Regarding severity, the most frequent was mild fluorosis (42.5 %), followed by moderate fluorosis (33.6%). The most severity codes were found in Manzanares-Caldas (TFI-5 to TFI-9; 39.7%). We also found an association between TFI and municipalities ( $p < 0.001$ ), between TFI and sex ( $p = 0.03$ ) and between TFI and age ( $p = 0.009$ ) (Table 4).

-Factors associated with enamel fluorosis prevalence

The risk factors associated with enamel fluorosis were involuntary intake of toothpaste (OR = 1.96 ± 0.39;  $p < 0.001$ ) and amount of table salt consumption (OR = 3.58 ± 0.71;  $p < 0.001$ ). The other factors were not statistically significant (Table 5).

-Multiple analyses between exposure factors and enamel fluorosis prevalence

Binomial analyses were conducted to determine the best explanatory model of enamel fluorosis. The outcome showed a statistically significant likelihood ratio ( $p < 0.001$ ). The raw and adjusted models were compared. It was observed an increase in the statistical strength of the association by the effect of the interactions between the following exposure factors; water ingestion exposure dose, urinary excretion, involuntary intake of toothpaste, amount of table salt and sex, being the coefficient of determination slightly higher (R<sup>2</sup> = 30%) (Table 6).

## Discussion

In Colombia, fluoride is added to table salt as a measure of public health welfare for the prevention of tooth decay (28). However, fluoride can also occur naturally in geological environments (2,29). Therefore, basic instruments have been established for the control and surveillance of the quality of water for human consumption, because unconventional water sources used for human

**Table 2:** Fluoride water ingestion exposure dose (IDag) and fluoride concentration in water and urine by municipalities.

	All participants		Fluoride in Water	IDag	Fluoride in Urine	R <sup>2</sup>
	N	Col % <sup>a</sup>	Mean± SD (mg/L)	Mean ± SD <sup>b</sup> (mg/kg/day)	Mean ± SD (mg/F/g creat.)	
Total sample	923	100	4.32 ± 8.35	0.14 ± 0.308	0.63± 0.82	0.98 <sup>d</sup>
Municipalities						
Algarrobo-Magdalena	132	13.4	24.3 ± 26	0.88 ± 0.16	2.06±1.03	
Campo de la Cruz-Atlántico	89	9.0	0.05 ± 0.03	0.002 ± 0.0003	0.11±0.03	
La Estrella-Antioquia	93	9.4	0.03 ± 0.008	0.001 ± 0.0002	0.16±0.02	
Manzanares-Caldas	121	12.3	0.02 ± 0.02	0.0008 ± 0.0002	0.33±0.05	
Margarita-Bolívar	115	11.7	6.3 ± 5.7	0.22 ± 0.04	1.86±0.92	
Nariño-Nariño	53	11.9	0.05 ± 0.009	0.001 ± 0.0003	0.14±0.001	
Oiba-Santander	153	15.5	0.04 ± 0.06	0.002 ± 0.0003	0.18±0.003	
Puerto López-Meta	167	16.9	0.45 ± 0.15	0.014 ± 0.003	0.22±0.02	

Note. <sup>a</sup> Percentage or subjects over the total population; <sup>b</sup>SD: standard deviation; <sup>c</sup>R<sup>2</sup> : determination coefficient between IDag and fluoride concentration in urine; <sup>d</sup> p< 0.001

**Table 3:** Enamel fluorosis prevalence in schoolchildren by characteristics of the study population.

Characteristics	All participants		Enamel fluorosis				
	N	Col % <sup>a</sup>	Yes n (%)	CI <sup>b</sup> 95%	No n (%)	CI 95%	p-value
<b>Total sample</b>	923	100	795 (86.1)	83.9–88.4	128 (13.9)	11.6–16.1	
<b>Municipalities (fluoride water ± SE)<sup>c</sup></b>							
Algarrobo-Magdalena (24.3 ± 26)	132	14.3	99 (75.0)	67.6–82.4	33 (25.0)	17.6–32.4	
Campo de la Cruz-Atlántico (0.05 ± 0.03)	89	9.6	41 (46.1)	35.6–56.5	48 (53.9)	43.5–64.4	
La Estrella-Antioquia (0.03 ± 0.008)	93	10.1	91 (97.8)	94.9–1.0	2 (2.2)	-0.8–5.1	
Manzanares-Caldas (0.02 ± 0.02)	121	13.1	121 (100)	-	0 (0.0)	-	<0.001
Margarita-Bolívar (6.0 ± 5.7)	115	12.5	115 (100)	-	0 (0.0)	-	
Nariño-Nariño (0.05 ± 0.009)	53	5.7	45 (84.9)	75.2–94.6	8 (15.1)	5.4–24.8	
Oiba-Santander (0.04 ± 0.06)	153	16.6	129 (84.3)	78.5–90.1	24 (15.7)	9.9–21.4	
Puerto López-Meta (0.45 ± 0.15)	167	18.1	154 (92.2)	88.1–96.3	13 (7.8)	3.7–11.9	
<b>Age (categorical, years)</b>							
7–9	463	50.2	402 (86.8)	83.7–89.9	61 (13.2)	10.1–16.3	0.54
10–12	460	49.8	393 (85.4)	82.2–88.7	67 (14.6)	11.3–17.8	
<b>Sex</b>							
Male	463	49.8	410 (88.6)	85.6–91.5	53 (11.4)	8.5–14.4	0.03
Female	460	50.2	385 (83.7)	80.3–87.1	75 (16.3)	12.9–19.7	
<b>School grade</b>							
Second	135	14.6	115 (85.2)	79.2–91.2	20 (14.8)	8.8–20.8	0.34
Third	257	27.8	224 (87.2)	83.1–91.3	33 (12.8)	8.7–16.9	
Fourth	275	29.8	243 (88.4)	84.6–92.2	32 (11.6)	7.8–15.4	
Fifth	256	27.8	213 (83.2)	78.6–87.8	43 (16.8)	12.2–21.4	

Note. <sup>a</sup> Percentage or subjects over the total population; <sup>b</sup> CI: Confidence interval. Statistical chi-square (χ<sup>2</sup>) test (p < 0.05); <sup>c</sup> average of fluoride concentration in drinking water in the study municipalities and standard error.

**Table 4:** Enamel fluorosis severity (TFI) in schoolchildren by characteristics of the study population.

	Participants		TF Index				p-value <sup>b</sup>
			TF0	TF1-TF2	TF3-TF4	TF5-TF9	
Characteristics	N	Col % <sup>a</sup>	n (%)	n (%)	n (%)	n (%)	
<b>Total sample</b>	923	100	128 (13.9)	392 (42.5)	310 (33.6)	93 (10.1)	
<b>Municipalities (fluoride water ± SE)<sup>c</sup></b>							
Algarrobo-Magdalena (24.3 ± 26)	132	14.3	33 (25.0)	57 (43.2)	30 (22.7)	12 (9.1)	
Campo de la Cruz-Atlántico (0.05 ± 0.03)	89	9.6	47 (52.8)	38 (42.7)	3 (3.4)	1 (1.1)	
La Estrella-Antioquia (0.03 ± 0.008)	93	10.1	2 (2.2)	35 (37.6)	51 (54.8)	5 (5.4)	
Manzanares-Caldas (0.02 ± 0.02)	121	13.1	0 (0.0)	19 (15.7)	54 (44.6)	48 (39.7)	
Margarita-Bolívar (6.0 ± 5.7)	115	12.5	0 (0.0)	73 (66.5)	40 (34.8)	2 (1.7)	<0.001
Nariño-Nariño (0.05 ± 0.009)	53	5.7	9 (17.0)	22 (41.5)	19 (35.9)	3 (5.7)	
Oiba-Santander (0.04 ± 0.06)	153	16.6	24 (15.7)	48(31.4)	68 (44.4)	13 (8.5)	
Puerto López-Meta (0.45 ± 0.15)	167	18.1	13 (7.8)	100 (59.9)	45 (27.7)	9 (5.4)	
<b>Age (categorical, years)</b>							
7–9	463	50.2	61 (13.2)	204 (44.1)	166 (35.9)	32 (6.9)	0.009
10–12	460	49.8	67 (14.6)	188 (40.9)	144 (31.3)	61 (13.3)	
<b>Sex</b>							
Male	463	49.8	54 (11.7)	206 (44.5)	147 (31.8)	56 (12.1)	0.03
Female	460	50.2	75 (16.3)	307 (66.7)	67 (14.6)	11 (2.4)	
<b>School grade</b>							
Second	135	14.6	20 (14.8)	58 (43.0)	48 (35.6)	9 (6.7)	0.43
Third	257	27.8	32 (12.5)	107 (41.6)	96 (37.4)	22 (8.6)	
Fourth	275	29.8	33 (12.0)	123 (44.7)	86 (31.3)	33 (12.0)	
Fifth	256	27.8	43 (16.8)	104 (40.6)	80 (31.3)	29 (11.3)	

Note. <sup>a</sup> Percentage or subjects over the total population; <sup>b</sup> Statistical chi-square ( $\chi^2$ ) test ( $p < 0.05$ ); <sup>c</sup> average of fluoride concentration water in the study municipalities and standard error.

consumption can be considered to have high risk for the presence of enamel fluorosis. The participants of the present study consumed water from underground sources, due to which it is necessary to understand the exposure in detail.

The ingestion of fluoride within the ‘optimal range’ (0.05–0.07 mg F/kg/day) (30), provides concentrations required in the oral environment to inhibit mineral loss or promote mineralisation in the tooth enamel (31). In the present study, the results showed that the fluoride ingestion level through drinking water was below the ‘optimal range’. However, in two municipalities (Algarrobo-Magdalena and Margarita-Bolívar), the fluoride ingestion levels could be considered as a risk factor for enamel fluorosis, which are possible fluorosis-endemic areas. Whereby, the fluoride concentration by ingestion of water is an indicator reflecting ingestion of water source from environmental fluoride. As most fluoride ingestion in the municipalities of Colombia was from groundwater, it is necessary to monitor fluoride levels in drinking water in different areas and over time.

Several indicators have been reported to calculate the water ingestion exposure doses (24), although it is clear that the fluoride concentration in water has a high impact within the equation used to calculate the ingestion exposure dose to fluoride, other factors such as the frequency of water ingestion and body weight also contribute. The World Health Organisation recommended drinking water fluoride levels of 0.5–1.5 mg/L for the prevention of tooth decay (32). Furthermore, in countries as Canada and E.U., the recommended concentration is 0.7 mg/L (33,31). In our study, of the 94 selected water samples, 25.5% had fluoride concentrations greater than the recommended values for Colombia (1 mg/L) (5). Therefore, the schoolchildren with the highest exposure in terms of fluoride concentration and frequency of ingestion of water showed the highest water fluoride ingestion dose. This suggests the need to monitor environmental variables for the epidemiological surveillance of fluoride exposure.

When we examined the association between exposure to fluoride in water and the prevalence of enamel fluoro-

**Table 5:** Fluoride exposure related with enamel fluorosis prevalence in schoolchildren.

Sources of fluoride ingestion	All participants		Yes	No	OR <sup>b</sup> ± SE <sup>c</sup>	CI <sup>d</sup> (95%)	
	N	Col % <sup>a</sup>	n (%)	n (%)		Lower limit	Upper limit
<b>Total sample</b>	923	100	795 (86.1)	128 (13.9)			
<b>Water ingestion exposure dose</b>							
IDag <sup>e</sup> ≥0.05 mg/kg/day	246	26.7	214 (26.9)	32 (25.0)	1.10 ± 0.24	0.72	1.79
IDag <0.05 mg/kg/day	677	73.4	581 (73.1)	96 (75)			
<b>Fluoride urinary excretion</b>							
High	447	48.4	400 (50.3)	50 (39.1)	1.58±0.42	1.12	2.50
Low	476	51.6	395 (49.7)	78 (60.9)			
<b>Consumption of packaged beverages</b>							
Yes	290	31.4	259 (32.6)	31 (24.2)	1.51 ± 0.33	0.98	2.33
No	633	68.6	536 (67.4)	97 (75.8)			
<b>Involuntary intake of toothpaste</b>							
Yes	455	49.3	410 (51.6)	45 (35.2)	1.96 <sup>f</sup> ± 0.39	1.33	2.90
No	468	50.7	385 (48.4)	83 (64.8)			
<b>Amount of toothpaste used</b>							
≥Half of toothbrush	506	54.8	437 (54.9)	69 (53.9)	1.04 ± 0.20	0.72	1.50
<Half of toothbrush	417	45.2	358 (45.0)	59 (46.1)			
<b>Frequency of teeth brushing</b>							
>3 times a day	327	35.4	272 (34.2)	55 (42.9)	0.69 ± 0.13	0.47	1.0
≤3 times a day	596	64.6	523 (65.8)	73 (57.0)			
<b>Amount of table salt consumption</b>							
≥1 spoonful	696	75.4	630 (79.2)	66 (51.6)	3.58 <sup>g</sup> ± 0.71	2.44	5.28
<1 spoonful	227	24.6	165 (20.8)	62 (48.4)			

Note. <sup>a</sup> Percentage or subjects over the total population; <sup>b</sup> OR: Odd ratio; <sup>c</sup> SE: Standard error; <sup>d</sup> CI: Confidence interval; <sup>e</sup> IDag (<0.05 mg/kg/day): recommended total optimal daily dose for all ages (ATSDR, 2003); <sup>f</sup> p<0.001; <sup>g</sup> p<0.001. Statistical chi-square (χ<sup>2</sup>) test (p < 0.05).

**Table 6:** Binomial model between exposure to fluoride and enamel fluorosis prevalence, adjusted by covariates, in schoolchildren.

Enamel fluorosis (cat. 0 = no, 1 = yes)	OR <sup>a</sup> ± SE <sup>b</sup>	CI <sup>c</sup> 95%	p value	(R <sup>2</sup> ) <sup>d</sup>	(p>chi <sup>2</sup> ) <sup>e</sup>
<b>Variables</b>			<b>&lt;0.001</b>	<b>0.30</b>	<b>&lt;0.001</b>
Water ingestion exposure dose (cat. 0 = no risk, 1 = risk)	1.4 ± 0.33	0.86–2.2	0.03		
Fluoride urinary excretion (cat. 0 = low, 1 = high)	1.6±0.32	1.01-1.95	0.04		
Involuntary intake of toothpaste (cat. 0 = no, 1 = yes)	1.6 ± 0.33	1.1–2.4	0.02		
Amount of table salt (cat. 0 = <1 spoonful, 1 = ≥1 spoonful)	3.6 ± 0.76	2.4-5.5	<0.001		
Sex (cat. 0 = Female, 1 = Male)	1.4 ± 0.29	0.97–2.1	0.06		
Dose*Amount of table salt (interaction 1)	7.9 ± 3.7	3.2–19.7	<0.001		
Dose*Sex (interaction 2)	3.0 ± 1.4	1.2–7.5	0.01		

sis, the results were not consistent because of the higher prevalence of enamel fluorosis in the municipality Manzanaraes-Caldas, although it showed the optimal concentrations of fluoride in water, whereas in the municipality

Algarrobo-Magdalena with the highest concentrations of fluoride in water, the prevalence and severity of enamel fluorosis were lower. These findings could be explained by the environmental (temperature) and geographical



(altitude) differences between the study municipalities. According to the altitude, a greater prevalence of enamel fluorosis has been reported in populations residing in high-altitude cities (10), due to hypoxia, which decreases urinary pH and increases the retention of fluoride. Similarly, populations living in cities with lower temperatures have reduced energy demand, causing the excretion of fluoride due to an increase in the retention rate (34). This finding suggests a plausible conjecture, because Manzanera-Caldas has an average temperature of 19°C and an altitude of 1933 m above sea level, whereas Algarrobo-Magdalena has an average temperature of 27.8°C and an altitude of 24 m above sea level. However, this hypothesis must be tested in further studies.

Based on the presence of enamel fluorosis, the present study showed a higher prevalence and severity than the national data for Colombia (11). These high data may be explained by the different sources of fluoride exposure that contribute to increasing the daily ingestion dose (water and table salt, toothpastes, mouthwash and food and drinks, as well as application of dental fluoride) (35-41). Similarly, the presence of other factors can increase the individual susceptibility to the toxic effects of fluoride, such as altitude and environmental temperature, malnutrition, diet components, gastric and urinary pH, kidney failure and genetic predisposition (10,42,43,20). In the bivariate analyses, we found no association between the fluoride-containing water ingestion dose and prevalence of enamel fluorosis, but there was an association between the amount of table salt consumption and involuntary intake of toothpaste. Although a surveillance and control system was implemented in Colombia to monitor fluoride levels in table salt for compliance with optimal levels (180–220 mg/F kg/salt) (7), the issue of the amount of table salt consumed daily is evident. The global average salt consumption in the country is 10–15 g/day (44), although the recommended amount is 5 g/person/day, which is sufficient to maintain adequate concentrations of fluoride in the saliva for the prevention of tooth decay (45,46). Similarly, the National Nutritional Status Survey in Colombia reported a high amount of table salt consumption (47). The majority of participants in our study responded affirmatively to the preparation of food with the amount of table salt higher than the recommended level, which is an important risk factor when several sources of excessive fluoride intake are combined.

The involuntary intake of toothpaste during tooth brushing is a common habit in children aged <5 years in Colombia (11). In our study, a high percentage of parents did not monitor the amount of toothpaste used by children, and moreover, children aged between 2 and 4 years used adult toothpaste with fluoride concentrations above the recommended level (1450 mg/L). These data demonstrate the inappropriate use of toothpaste with high

fluoride concentrations among children. However, according to risk, some studies have reported that the use of toothpastes with high fluoride content did not increase the urinary fluoride excretion significantly (48,49). Similarly, these habits may contribute together with the fluoride content in water and table salt to increase the cumulative dose above the optimal levels (50,17). Several authors reported that the involuntary intake of toothpaste in 5-year-old children contributes 30% of the total fluoride ingestion, which is higher than the same contribution in adults (51,52). In the present study, the data for the involuntary intake of toothpaste were obtained from questionnaires that reconstructed the past exposure. It is clear that this method is not highly sensitive to evaluate this exposure; therefore, we recommend conducting longitudinal studies on urinary fluoride excretion to confirm the weight of this risk factor on the presence of enamel fluorosis.

In this regard, it is evident that the impact on chronic toxicity of the different fluoride sources, independent or in combination, is dependent on the time of exposure, concentration of sources and the amount and frequency of its consumption. Therefore, we conducted a multiple analysis to estimate the impact of the effect of the combination of various fluoride sources and demographic variables and obtained a multiple association model with statistical significance, where we tested the interactions between the dose of water intake, involuntary toothpaste intake, amount of table salt intake, fluoride urinary excretion and sex of the participants. We found that the variable sex influenced the model from the category of boy or girl, indicating that boys probably ingested greater amounts of water or table salt than girls, due to greater muscle mass that consequently increases the intake needs (53). Likewise, the strength of association of urinary fluoride excretion was increased in multiple analysis, probably due to the strong positive correlation with the water ingestion exposure dose, showing a higher urinary fluoride concentration in children with high levels of water ingestion fluoride exposure dose. These outcomes have also been reported by other studies, although with great variability (29,54-57).

Among the possible limitations inherent to the type of this study (cross-sectional design) are the evaluation of enamel fluorosis and exposure to fluoride at the same time point, reconstructing the exposure in the stage where it originated (between 1 and 4 years of age) and the possible memory bias that could incur in the parents' responses when trying to reconstruct the past exposure using a validated questionnaire. It is important to emphasise that only children born and living in the respective municipalities were included in the present study, due to which it was expected that the fluoride concentrations currently found in natural water sources would be similar to those that had been exposed in the past, consid-

ring that none of the municipalities reported substantial improvements in the water source in the past 15 years. The findings of this study provide relevant information for public policies on oral health in Colombia, focusing on the deficiencies in the current fluoride exposure control and surveillance system, which monitors and controls only two of the various sources of fluoride involved (natural water and table salt). Which shows a low efficacy of the government strategy used in Colombia to reduce the chronic toxic effects of fluoride on the dental enamel of people.

To conclude, the prevalence of enamel fluorosis in Colombian schoolchildren can be explained by some factors related to the fluoride-containing water exposure dose, other sources of exposure and some demographic factors.

## References

- Ozsvath D. Fluoride and environmental health: a review. *Rev Environ Sci Biotechnol*. 2009;8:59-79.
- Usuda K, Kono K, Dote T, Watanabe M, Shimizu H, Tanimoto Y, et al. Fluoride analysis and fluoride related health problems in clinical, experimental, occupational and environmental aspects: A narrative review. *Biomedical Research on Trace Elements*. 2009;20:274-283.
- Li P, Qian H, Wu J, Chen J, Zhang Y, Zhang H. Occurrence and hydrogeochemistry of fluoride in alluvial aquifer of Weihe River, China. *Environmental Earth Sciences*. 2014;71:3133-3145.
- U.S. Environmental Protection Agency, 2010. Fluoride: exposure and relative source contribution analysis. In: Office of Water, 820-R-10-015, pp. 210.
- Ministerio de la Protección Social y Ministerio del Medio Ambiente, Vivienda y Desarrollo Territorial, República de Colombia., 2007. Resolución 2115. Art. 6, pp 3.
- Ministerio de Vivienda, Ciudad y Territorio - Ministerio de Salud y Protección Social, República de Colombia., 2020- Informe Nacional de Calidad del Agua para Consumo Humano- INCA, Bogotá, pp. 119-120.
- Instituto Nacional de Salud, República de Colombia, 2012. Documento Técnico Sistema Nacional de Vigilancia en Salud Pública, SIVIGILA-INS, 2012-2015, Bogotá, pp 35-40.
- Fejerskov O, Larsen M, Richards A, Baelum V. Dental tissue effects of fluoride. *Advances in Dental Research*. 1994;8:15-31.
- Buzalaf MAR, Granjeiro JM, Damante CA. Fluoride content of infant formulas prepared with deionized, bottled mineral and fluoridated drinking water. *ASDC J Dent Child*. 2001;68:10-37.
- Buzalaf MAR, Massaro CS, Rodrigues MHC. Validation of fingernail fluoride concentration as a predictor of risk for dental fluorosis. *Caries Research*. 2012;46:394-400.
- Ministerio de Salud, República de Colombia., 2014. IV Estudio Nacional de Salud Bucal-ENSAB, Bogotá, pp. 65-69.
- Ministerio de Salud, República de Colombia., 1998. III Estudio Nacional de Salud Bucal-ENSAB, Bogotá, pp. 114-115.
- Norman M, Twetman S, Hultgren A, Granström E, Stecksén C. Urinary fluoride excretion in preschool children after intake of fluoridated milk and use of fluoride-containing toothpaste. *Community Dent Health*. 2017;34:27-31.
- Narsimha A, Sudarshan V. Assessment of fluoride contamination in groundwater from Basara, Adilabad District, Telangana State, India. *Appl Water Sci*. 2017;7:2717-2725.
- Bhattacharya P, Samal AC, Banerjee S, Pyne J, Santra SC. Assessment of potential health risk of fluoride consumption through rice, pulses, and vegetables in addition to consumption of fluoride contaminated drinking water of West Bengal, India. *Environ Sci Pollut Res Int*. 2017;24:0300-14.
- Sezgin BI, Onur ŞG, Menteş A, Okutan AE, Haznedaroğlu E, Vieira AR. Two-fold excess of fluoride in the drinking water has no obvious health effects other than dental fluorosis. *Journal of Trace Elements in Medicine and Biology*. 2018;50:216-222.
- Armas-Vega A, González-Martínez F, Rivera-Martínez M, Mayorga-Solórzano M, Banderas-Benítez V, Guevara-Cabrera O. Factors associated with dental fluorosis in three zones of Ecuador, a post hoc analysis. *J Clin Exp Dent*. 2019;11:e42-8.
- Till C, Green R, Flora D, Hornung R, Martínez-Mier A, Blazer M, et al. Fluoride exposure from infant formula and child IQ in a Canadian birth cohort. *Environment International*. 2020;134:105315.
- Chen S, Li B, Lin S, Huang Y, Zhao X, Zhang M, et al. Change of urinary fluoride and bone metabolism indicators in the endemic fluorosis areas of southern china after supplying low fluoride public water. *BMC Public Health*. 2013;13:156.
- Martínez EA, Spencer KL, Sanders BJ, Jones JE, Soto AE, Tomlin AM, et al. Fluoride in the diet of 2-years-old children. *Community Dent Oral Epidemiol*. 2017;45:251-7.
- Lima YBO, Gomes ARLF, Santos HKA, Arsati F, Oliveira MC, Freitas VS. Exposure to fluoride of children during the critical age for dental fluorosis in the semiarid region of Brazil. *Cien Saude Colet*. 2018;23:1045-54.
- Martínez-Mier EA, Cury JA, Heilman JR, Katz BP, Levy SM, Li Y, et al. Development of gold standard ion-selective electrode-based methods for fluoride analysis. *Caries Res*. 2011;45:3-12.
- APHA., 2012. Standard methods for the examination of water and wastewater. Washington, DC: Rice, E.W., Baird, R.B., Eaton, A.D., Clesceri, L.S. (Eds.)
- ATSDR., 2005. Public Health Assessment Guidance Manual (Update). Resource document [https://www.atsdr.cdc.gov/hac/phamanual/pdfs/phagm\\_final1-27-05.pdf](https://www.atsdr.cdc.gov/hac/phamanual/pdfs/phagm_final1-27-05.pdf). Accessed 20 October 2016.
- Thylstrup A, Fejerskov O. Clinical appearance of dental fluorosis in permanent teeth in relation to histologic changes. *Community Dentistry and Oral Epidemiology*. 1978;6:315-328.
- R Core Team. R: A Language and Environment for Statistical Computing; R Foundation for Statistical Computing: Vienna, Austria, 2020; Available online: <http://www.R-project.org/> (accessed on 2 May 2020).
- Glickman M, Rao S, Schultz M. False discovery rate control is a recommended alternative to Bonferroni-type adjustments in health studies. *J Clin Epidemiol*. 2014;67:850-7.
- Ministerio de Salud, República de Colombia., 2016. Documento técnico "Perspectiva del uso del flúor Vs Caries y Fluorosis Dental en Colombia, p.p. 20-25.
- Zohoori FV, Buzalaf MAR, Cardoso CAB, Olympio KPK, Levy FM, Grizzo LT, et al. Total fluoride intake and excretion in children up to 4 years of age living in fluoridated and non-fluoridated areas. *Eur J Oral Sci*. 2013;121:457-464.
- Mascarenhas AK. Risk factors for dental fluorosis: A review of the recent literature. *Pediatric Dentistry*. 2000;22:269-77.
- National Health and Medical Research Council (NHMRC), 2017. Water fluoridation: dental and other human health outcomes. Public Statement, pp 24-38.
- World Health Organization Expert Committee on Oral Health Status and Fluoride Use. Fluorides and oral health. WHO, 1994. Technical Report Series No. 846. World Health Organization, Geneva.
- Health Canada., 2010. Trends in Breastfeeding Practices in Canada (2001 to 2009-2010). <http://www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/prenatal/trends-tendances-eng.php#a2> (accessed February 21, 2016).
- Chankanka O, Levy SM, Warren JJ, Chalmers JM. A literature review of aesthetic perceptions of dental fluorosis and relationships with psychosocial aspects/oral health-related quality of life. *Community Dent Oral Epidemiol*. 2010;38:97-109.
- Martínez-Mier EA, Soto-Rojas AE, Ureña-Cirett JL, Stookey GK, Dunipace AJ. Fluoride intake from foods, beverages and dentifrice by children in Mexico. *Community Dentistry and Oral Epidemiology*. 2003;31: 221-230.
- Yadav AK, Kaushik CP, Haritash AK, Singh B, Raghuvanshi SP, Kansal A. Determination of exposure and probable ingestion of fluori-

- de through tea, toothpaste, tobacco and pan masala. *Journal of hazardous materials*. 2007;142:77-80.
37. Delano F, Adas S, Correia F. Urinary Fluoride Excretion in Children Exposed to Fluoride Toothpaste and to Different Water Fluoride Levels in a Tropical Area of Brazil. *Braz Dent J*. 2008;19:214-218.
38. Mehta A. Biomarkers of fluoride exposure in human body. *Indian Journal of Dentistry*. 2013;4:207-210.
39. Silva DN, Ventura PG, Amaral L, Ferreira T, Cury JA, Vieira W, et al. Sensitivity of two biomarkers for biomonitoring exposure to fluoride in children and women: A study in a volcanic area. *Chemosphere*. 2016;155:614-620.
40. Martignon S, Opazo-Gutierrez MO, Velasquez-Riaño M, Orjuela IR, Avila V, Martínez-Mier EA, et al. Geochemical characterization of fluoride in water; table salt, active sediment, rock and soil samples, and its possible relationship with the prevalence of enamel fluorosis in children in four municipalities of the department of Huila (Colombia). *Environmental and Monitoring Assessment*. 2017;189:264.
41. Saldarriaga A, Restrepo M, Rojas-Gualdrón DF, De Souza-Carvalho T, Buzalaf MA, Santos-Pinto L, et al. Dental Fluorosis according to Birth Cohort and Fluoride Markers in an Endemic Region of Colombia. *ScientificWorldJournal*. 2021;2021:6662940.
42. Jiang C, Zhang S, Liu H, Guan Z, Zeng Q, Zhang C, et al. Low glucose utilization and neurodegenerative changes caused by sodium fluoride exposure in rat's developmental brain. *Neuromolecular Med*. 2014;16:94-105.
43. Kebede A, Retta N, Abuye C, Whiting SJ, Kassaw M, Zeru T, et al. Dietary fluoride intake and associated skeletal and dental fluorosis in school age children in rural Ethiopian Rift Valley. *Int J Environ Res Public Health*. 2016;13:756.
44. World Health Organization. WHO, 2006. Reducing salt intake in populations: Report of a WHO Forum and Technical Meeting. October 2006.
45. Pollick H. Salt fluoridation: a review. *Journal of the California Dental Association*. 2013;41:395-7.
46. Wennhall I, Hajem S, Ilros S, Ridell K., Ekstrand KR, Twetman S. Fluoridated salt for caries prevention and control a 2-year field study in a disadvantaged community. *Int J Paediatr Dent*. 2014;24:161-167.
47. Ministerio de Salud, República de Colombia., 2015. Encuesta Nacional de la Situación Nutricional-ENSIN. Bogotá, pp.400-411.
48. Pessan JP, Pin MLG, Martinhon CCR, Silva SMB, Buzalaf MAR. Analysis of fingernails and urine as biomarkers of fluoride exposure from dentifrice and varnish in 4-to 7-year old children. *Caries Res*. 2005;39:363-370.
49. Oliveira MJL, Paiva SM, Martins LHPM, Ramos ML, Lima YBO, Cury JA. Fluoride intake by children at risk for the development of dental fluorosis: comparison of regular dentifrices and flavored dentifrices for children. *Caries Res*. 2007;41:460-466.
50. Wong MCM, Clarkson J, Glenn AM, Lo ECM, Marinho VCC, Tsang BWK, et al. Cochrane reviews on the benefits/risks of fluoride toothpastes. *Journal of Dental Research*. 2011;90:573-579.
51. Cury JA, Tenuta LM. Evidence-based recommendation on toothpaste use. *Braz Oral Res*. 2014;28:1-7.
52. Casaglia A, Cassini MA, Condò R, Iaculli F, Cerroni L. Dietary Fluoride Intake by Children: When to Use a Fluoride Toothpaste? *Int J Environ Res Public Health*. 2021;18:5791.
53. Fariás P, Estevez-García JA, Onofre-Pardo EN, Pérez-Humara ML, Rojas-Lima E, Álamo-Hernández U, et al. Fluoride Exposure through Different Drinking Water Sources in a Contaminated Basin in Guanajuato, Mexico: A Deterministic Human Health Risk Assessment. *Int. J. Environ. Res. Public Health*. 2021;18:11490.
54. Abuhaloob L, Maguire A, Moynihan P. Fractional Urinary Fluoride Excretion (FUFE) of 3-4 year children in the Gaza Strip. *Community Dent Health*. 2015;32:8-15.
55. Antonijevic E, Mandinic Z, Curcic M, Djukic-Cosic D, Milicevic N, Ivanovic M, et al. "Borderline" fluorotic region in Serbia: correlations among fluoride in drinking water, biomarkers of exposure and dental fluorosis in schoolchildren. *Environ Geochem Health*. 2016;38:885-96.
56. Linhares DP, Garcia PV, Amaral L, Ferreira T, Cury JA, Vieira W, et al. Sensitivity of two biomarkers for biomonitoring exposure to fluoride in children and women: A study in a volcanic area. *Chemosphere*. 2016;155:614-20.
57. Idowu OS, Azevedo LB, Valentine RA, Swan J, Vasantavada PV, Maguire A, et al. The use of urinary fluoride excretion to facilitate monitoring fluoride intake: A systematic scoping review. *PLoS One*. 2019;14:e0222260.

#### Acknowledgements

We wish to thank University of Cartagena and ACFO, for the financial Support and to S. Martignon for your role as external examiner as a reference point in the clinical calibration of TF Index.

#### Ethics

This study was approved by the Human Research Ethical Committee of the University of Cartagena, Colombia (minute # 92, 22/09/2016) and was registered on the University of Cartagena Register (#234262016, 10/10/2016).

#### Source of Funding

University of Cartagena (Project # 108-2018), and ACFO (Project # 02-2017).

#### Authors' contributions

Under a subheading of Authors' contributions.

#### Conflict of interest

The authors have declared that no conflict of interest exist.