

Philippine herpetology (Amphibia, Reptilia), 20 years on: two decades of progress towards an increasingly collaborative, equitable, and inclusive approach to the study of the archipelago's amphibians and reptiles

Camila G. Meneses^{1*}, Kier Mitchel E. Pitogo^{1*}, Christian E. Supsup^{1*}, Rafe M. Brown¹

¹ Department of Ecology and Evolutionary Biology and Biodiversity Institute, University of Kansas, Lawrence, Kansas 66045, USA
Corresponding author: Christian E. Supsup (supsupchristian@gmail.com)

Abstract

A first review of the history, status, and prospects for Philippine herpetology conducted more than two decades ago (2002) summarized the diverse topics studied and highlighted the development and achievements in research up to the year 2000. This study revisits and re-assesses what Philippine herpetology has accomplished, both as a discipline and a community, during the last two decades (2002–2022). A total of 423 herpetological publications was collated, revealing a substantial increase in annual publications, rising from approximately four per year during 2002–2008 to around 28 per year in 2009–2022. Half of the published studies focused on squamate reptiles (lizards 30.5%, snakes 21%) and 28.4% on amphibians, 5.9% on turtles, and 2.6% on crocodiles. The remaining 11.6% of studies focused simultaneously on multiple taxa (i.e., faunal inventories). Diversity and distribution (35.2%) and ecological (26.5%) studies remained popular, while studies on taxonomy (14.9%), phylogenetics and biogeography (11.8%), and conservation (11.6%) all increased. However, geographical gaps persist urging immediate surveys in many understudied regions of the country. Finally, we found a balanced representation between Filipino and foreign first authors (1.0:1.1), yet a substantial gender gap exists between male and female first authors (7.1:1.0). Nonetheless, the steep increase in publications and the diversity of people engaged in Philippine herpetology is a remarkable positive finding compared to the 20 years preceding the last review (1980–2000). Our hope is that the next decades will bring increasingly equitable, internationally collaborative, and broadly inclusive engagement in the study of amphibians and reptiles in the Philippines.

Key words: Biodiversity, conservation, distribution, Southeast Asia, systematics, taxonomy

Introduction

Sustained, regionally focused, field-based research programs focusing on the ecology and evolution of amphibians and reptiles provide insight into many conceptually intriguing, unique, and fundamental questions relating to the



Academic editor: Fedor Konstantinov
Received: 15 July 2023
Accepted: 4 January 2024
Published: 30 January 2024

ZooBank: <https://zoobank.org/3C99730E-4FF6-4189-9BCD-C435E31A36DD>

Citation: Meneses CG, Pitogo KME, Supsup CE, Brown RM (2024) Philippine herpetology (Amphibia, Reptilia), 20 years on: two decades of progress towards an increasingly collaborative, equitable, and inclusive approach to the study of the archipelago's amphibians and reptiles. ZooKeys 1190: 213–257. <https://doi.org/10.3897/zookeys.1190.109586>

Copyright: © Camila G. Meneses et al.
This is an open access article distributed under terms of the CC0 Public Domain Dedication

* These authors contributed equally to this work.

origin, accumulation, and continued survival of Earth's biodiversity (Brown and Alcala 1970a; Brown et al. 2013a; Gainsbury and Meiri 2017; Greenberg et al. 2018; Meiri et al. 2020; Sheu et al. 2020; Zimin et al. 2022; Camaiti et al. 2023). Regionally oriented research by herpetological systematists, biogeographers, and macroecologists have elevated our understanding of the global phylogenetic and evolutionary underpinnings of amphibian and reptile biodiversity by relating the distribution of this diversity to the geographical template itself (Esselstyn et al. 2010; Böhm et al. 2013; Barley et al. 2014, 2015; Roll et al. 2017; Vidan et al. 2019; Gumbs et al. 2020; Bernstein et al. 2021a). Similarly, regional studies on topics ranging from natural histories of species from particular areas to global analyses of clade-wide trait variation (Vidan et al. 2019; Camaiti et al. 2022) and organismal genomic variation (Formenti et al. 2022; Chan et al. 2022; Hutter et al. 2022) now contribute significantly to our collective understanding of conservation challenges facing highly imperiled amphibian and reptile populations (Böhm et al. 2013; González-del-Piiego et al. 2019; Tingley et al. 2019; Gumbs et al. 2020; Chapple et al. 2021; Cox et al. 2022; McDonald et al. 2022; Womack et al. 2022; Slavenko et al. 2023).

The Philippine Archipelago, which has been the focus of sustained herpetological inquiry for more than a century (Taylor 1921, 1922a, b, 1928; Leviton 1963a; Brown and Alcala 1970a, 1978, 1980; Brown et al. 2002a, 2012b, 2013a; Diesmos and Brown 2011; Diesmos et al. 2014, 2015; Leviton et al. 2014, 2018), is situated adjacent to the Southeast Asian mainland, between the Western Philippine Sea and the Western Pacific Ocean (Fig. 1). This unique archipelago is home to numerous remarkable evolutionary radiations of amphibians and reptiles (Brown and Guttman 2002; Brown et al. 2002a, 2009a, b, 2013a, 2015a; Evans et al. 2003; Brown and Diesmos 2009; Siler and Brown 2010, 2011; Siler et al. 2010a, b, c, d, 2011a, b, c, 2013, 2014a, b, c; Welton et al. 2013a; 2014a, b; Weinell and Brown 2017; Weinell et al. 2020a, b, c; Chan et al. 2021, 2022; Flury et al. 2021), many of which have only recently been characterized, and some of which remain to be studied in depth (e.g., Brown et al. 2010a, 2011a, b, 2012a; Linkem et al. 2010a, b, c; Linkem and Brown 2013; Wynn et al. 2016; Oliver et al. 2018; Davis et al. 2020; Bernstein et al. 2021b; Eliades et al. 2021). Currently, there are approximately 475 recognized species of amphibians and reptiles in the Philippines, 76.2% of which are endemic, and most of which can be characterized as (1) geographically circumscribed species, whose distributions coincide with one of the archipelago's 5–7 major faunal regions (Brown and Alcala 1970a, 1978, 1980; Brown et al. 2017); (2) range-restricted species from isolated islands (e.g., Brown and Alcala 1974; Brown et al. 1997; Ferner et al. 2001; Linkem et al. 2010a; Oliveros et al. 2011; Brown et al. 2018; Meneses et al. 2022), or (3) species limited to geologically isolated and/or upland habitats (e.g., Brown and Alcala 1961, 1970b, 1982a; Ferner et al. 1997; Brown et al. 1999a, b, 2020; Linkem et al. 2010b; Siler et al. 2010c). However, this estimate is expected to change over time with ongoing biodiversity inventories, taxonomic revisionary studies, critical reappraisals of earlier works, and the novel application of technologies (e.g., genomic data, ecological niche modeling) still emerging today (Brown et al. 2002a; Brown 2006; Brown and Siler 2013; Diesmos et al. 2015).

Just more than two decades ago, Brown et al. (2002a) conducted a comprehensive review of Philippine herpetological studies. The authors summarized the



Figure 1. Map of the Philippine archipelago, situated in Southeast Asia (inset map), showing the recognized Pleistocene Aggregate Island Complexes (PAICs) and small island groups.

history of amphibian and reptile studies and included the distinct periods which characterized the development of herpetological research in the country, highlighting the important contributions of biologists during the last decades leading up to the turn of the century. That review centered on five topical themes or areas of research focus: (1) diversity and distribution, (2) taxonomy, (3) ecology,

(4) phylogenetic systematics and biogeography, and (5) conservation. As a result of that exercise, it became abundantly clear that most of the archipelago's earlier herpetological studies (prior to the 2000s) were predominantly focused on species diversity, taxonomy, and biogeography (and, to a lesser extent, ecology, and conservation). Despite the many papers focused on single species descriptions, but also including synthetic reviews (e.g., Brown and Alcala 1978, 1980), an immense amount of taxonomic work was still needed by the early 2000s. This is because the majority of studies to that date were descriptive, based solely on traditional morphological characters (i.e., measurements, meristic data like scale counts, and comparisons of discrete character states), and only selected clades had been comprehensively reviewed with the goal of synthetic considerations of those particular faunal groups (Taylor 1921, 1922a, b; Inger 1954; Leviton and Brown 1958; Leviton 1962, 1963a, b, 1964a, b, c, d, 1965a, b, 1967, 1968, 1979, 1983; Brown and Alcala 1974, 1980, 1994). As discussed by Brown (2006, 2007) and Brown and Stuart (2012), the use of multiple data streams and integrative approaches (including phylogeny) for more robust, pluralistic, and quantitative approaches to species recognition was just in their infancy (Brown and Diesmos 2002; Brown and Guttman 2002; Diesmos et al. 2002; Brown et al. 2003).

Brown et al. (2002a) also highlighted some of the gaps in other research areas. For instance, on the subject of biogeography and conservation, they emphasized how understanding patterns of Philippine amphibian and reptile distributions would be essential to formulating effective conservation and management strategies (Brown and Alcala 1961, 1986; Alcala and Custodio 1995; Diesmos et al. 2002). However, at the time of their review, Brown et al. (2002a) emphasized that the full informative potential of a comprehensive understanding of verified species distributions had not been fully realized due to limited information from many unexplored islands groups (i.e., the Batanes, Bubuyans, Lubang, Leyte, Masbate, Mindoro, the Romblon Island Group, Siquijor, the Sulu Archipelago, and Samar), as well as numerous high-elevation mountain ranges of the archipelago's largest islands, Luzon and Mindanao. At that time (Brown et al. 2002a), documentation of patterns of species occurrences and community structure along elevational gradients was just beginning to take shape (Brown and Alcala 1961; Custodio 1986; Alcala and Custodio 1995; Alcala et al. 1995; Brown et al. 1996, 2000a; Heaney et al. 2000). Numerous other topics were highlighted, including formative areas of research that had become a focus by mid-century, but for which no follow-up investigations had been forthcoming during the last 20–40 years (Brown et al. 2002a). These included studies of reproductive biology (Alcala and Brown 1956, 1982; Alcala 1962; Brown and Alcala 1982b), physiology (Alcala and Brown 1966), development (Alcala 1962; Alcala and Brown 1982), and basic population biology and demography (Alcala 1967, 1970; Alcala and Brown 1967).

In general, Brown et al. (2002a) emphasized that these conspicuous gaps in Philippine herpetological research would most likely be addressed most effectively through collaborative efforts of teams of institutions (universities, local government units, non-governmental organizations, stakeholder communities) and the participation of diverse groups of foreign and local researchers, students, and local community representatives. Ultimately, the results accessible via open-access peer-reviewed publications—such studies reinforced by integrative analyses of multiple data types that are freely accessible through

web-based platforms (e.g., HerpNet, VertNet, etc.) and specimen-associated data provided by natural history museums—could be harnessed in such a way that a more inclusive, transparent, broadly participatory future of Philippine herpetology could be realized (e.g., Brown et al. 2012b, 2013b). With the turn of the millennium and given the inevitable shift in herpetological research that was coming, Brown et al. (2002a) reviewed and took stock of the accomplishments, discoveries, strengths, and shortcomings of what they envisioned as a global community of herpetologists interested in Philippine biodiversity. Today, we revisit that same general topic, but we do so with the sense that another, pronounced, and unique period of the historical development of herpetology in the Philippines is coming in the years ahead.

In this paper, we revisit studies conducted from 2002–2022 to re-assess the state of Philippine herpetological research, 20 years after the review of Brown et al. (2002a). We summarized the last two decades of published studies in five general areas or topical themes, as discussed in the previous review. One of our goals was to explore whether these studies addressed gaps previously identified (Brown et al. 2002a). Trends during the last two decades of taxonomic studies were examined, including attention to data types (e.g., morphology versus molecular data), modern quantitative methodology (e.g., phylogenetics, advanced statistical procedures, quantitative biogeographical inference), and how these were integrated or used in statistical species delimitation (e.g., Barley et al. 2013; Welton et al. 2013a). We also sought to obtain an explicitly geographical or spatial overview of the last twenty years of research and identify priority areas that receive little attention and now represent challenges for the future. Finally, to determine who is primarily involved in Philippine herpetological research—and to critically and transparently address the question of diversity, equity, and inclusion in our field (Have we broadened engagement? Is our community more diverse now than it was a quarter century ago?)—we classified published studies by the lead author’s nationality and gender and explored the diversity of people who study Philippine herpetology.

Methods

Literature review

We employed a systematic literature review of Philippine herpetology published from 2002–2022, following the guidelines from the updated Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (Page et al. 2021). We compiled a dataset of peer-reviewed journal articles by searching through Google Scholar and Web of Science. The following keywords were used: “Philippine amphibians,” “Philippine reptiles,” “Philippine lizards,” “Philippine snakes,” and “Philippine anurans/frogs.” The list of papers compiled was manually supplemented to include articles not captured in the preliminary search. This includes articles addressing other areas or topics, but which included substantial Philippine specimens (or data derived from Philippine studies, and now available in the public domain) to inform their results, and other articles published in locally refereed scientific journals. We made every effort to be as comprehensive as possible, but it is conceivable that a few published papers meeting our criteria for inclusion may have been missed by our search process. Nevertheless,

our objective was primarily focused on capturing and characterizing the general trends and overall patterns, which may be inferred from appropriately, ethically, and transparently communicated (i.e., subject to peer-review, properly documented, accessibly archived, and demonstrably repeatable) scientific research in topics relating to the biology of Philippine amphibians and reptiles.

For each article, we extracted the following information to determine trends and patterns in Philippine herpetological research: (1) year of publication, (2) major focal taxon (amphibians [frogs/toads or caecilians], lizards, snakes, crocodiles, turtles, and a combined category “multiple taxa” for studies involving combinations of each major group), and each paper’s (3) research theme as discussed above (see Table 1). Then, we performed a Poisson regression and local polynomial regression to test for a trend in the number of papers published during the last two decades. Because we wanted to address the technological advances of the last quarter century, we characterized the variable categories of evidence, employed by papers that included new species descriptions (see also Brown and Stuart 2012). We achieved this by classifying every taxonomic publication according to the type of data that was used (e.g., morphological, bioacoustic, molecular, and/or combinations of these data).

Table 1. The categories used to identify types of peer-reviewed papers involving Philippine herpetology published from 2002–2022 (based on the five topical themes of Brown et al. 2002a).

Category	Description
Diversity and distribution	surveys, checklists, distributions (range extensions, new island records), and measures of diversity
Taxonomy	species descriptions and taxonomic revisions
Ecology	natural history, community ecology, and population biology
Phylogenetics and biogeography	phylogenetics (excluding new species description), evolutionary biology, and biogeography
Conservation	conservation, outreach, threats, and methods

Geographical patterns

Available geographic coordinates were extracted from articles focusing on new species descriptions, species distribution records, natural history notes, and targeted herpetological inventories. To assess country-wide geographical patterns of published herpetological studies from the literature, we projected occurrence records on a map of the archipelago and georeferenced all points by referring to museum records and biodiversity information resources (below), if necessary. For articles that did not report geographic coordinates but which did include specific locality information (e.g., island, province, municipality, barangay, or other unique identifiers), we georeferenced occurrence data in Quantum GIS v. 3.22 using the Philippine gazetteer available from DIV-GIS Database (<https://www.diva-gis.org/>). We included regions, areas, or single sites, which have been recently surveyed, and for which all specimen-associated data have been properly curated (e.g., in museum databases linked to accessible voucher specimens, ensuring repeatability and transparency; and from the community of the biodiversity repository institutions which provide unrestricted access to specimen-associated data) and published in accessible, publicly available databases such as the Global Biodiversity Information Facility (GBIF; <https://www.gbif.org/>), VertNet (<http://vertnet.org/>), and iDigBio (<https://www.idigbio.org/>).

Our presentation of geographical patterns made use of the last half-century's prevailing context for biogeographical studies in the archipelago, namely the Pleistocene Aggregate Island Complexes (PAIC) model (Brown and Diesmos 2002; Brown and Guttman 2002; Brown et al. 2013a). This model attempts to capture the biogeographic terrestrial subregions of the archipelago based on earlier studies that traced underwater bathymetric contours around the archipelago's major landmasses (Inger 1954; Leviton 1963a; Brown and Alcalá 1970a), so as to reflect patterns of island amalgamations, terrestrial connections involving major landmasses and adjacent island banks, and other exposure of land caused by oscillating Pleistocene sea levels (review: Brown et al. 2013a). Together, the frequently discussed "PAIC Paradigm" illustrates the locations of simplified, reasonably accurate, inferred land connections that may have partitioned and isolated terrestrial vertebrates into the variable and distinct faunal compositions we find today, on the islands which together make up each unique faunal subregion: The Luzon PAIC, and those of Mindanao, Mindoro, West Visayan (also referred to as Negros-Panay), Palawan, and the smaller Romblon Island Group (RIG), the Sulu Island Group (SIG) (Fig. 1), and a few small islands which are associated with larger landmasses but which were never fully connected to them. This latter group includes Siquijor Island, adjacent to the West Visayan PAIC; Camiguin Sur Island, adjacent to the Mindanao PAIC; and finally, isolated island groups that form minor but unique subcenters of biodiversity, but which are not part of or strongly associated with, the major PAICs (e.g., Babuyan and Batanes Island Groups, north of Luzon).

First authorship nationality and gender gaps

Finally, as a simplified but important general first step towards exploring disparity and gender gaps in Philippine herpetology, we also determined the nationality (Filipino vs all non-Filipino nationalities/"foreign") and traditional biological gender (male vs female) of the first author of each article. Although we feel this somewhat crude, excessively binary view will not capture nuances needed to truly assess equity gaps in Philippine herpetology, we consider it a first step and a point from which we hope future discussions and steps towards broadening engagement can begin (see Ramírez-Castañeda et al. 2022). In particular, we emphasize that our use of the term "gender" and its binary assignment in this preliminary analysis is used solely for the purposes of assessing gender gaps in Philippine herpetological research and does not imply a binary nature of the term (see Rock et al. 2021). All analyses and visualizations were performed in R Studio v. 4.2.2 (R Core Team 2022). R code and documentation (R markdown HTML, Suppl. materials 1, 2) are available on GitHub (<https://github.com/csup-sup/PhilHerpsRev>).

Results

We compiled a total of 423 peer-reviewed scientific articles on Philippine herpetology, published from 2002–2022 (see Suppl. material 1). There was a significant increase in publications during the years based on our Poisson regression analysis ($X^2_{(1, n = 21)} = 131.9, p < 0.001$). Approximately four publications per year, for the majority of the first decade (2002–2008), was a general trend that

increased sharply to approximately 28 publications per year across the second decade (2009–2022; Fig. 2). The years with the highest number of publications were all during the past 11 years: 2020 ($n = 47$), 2021 ($n = 42$), 2014 ($n = 32$), 2011 ($n = 31$), and 2022 ($n = 31$). Notably, this dramatic seven-fold increase in publication rates during the last 11 years, comprises a remarkable 83.4% of all publications in Philippine herpetology since 2002.

Research themes

Of the 423 articles we reviewed, half the papers were conducted primarily on reptiles, with lizards and snakes comprising nearly equivalent proportions of the total, or 30.5% ($n = 129$), and 21% ($n = 89$), respectively (Fig. 3A). Amphibians comprised 28.4% ($n = 120$), turtles 5.9% ($n = 25$), and crocodiles 2.6% ($n = 11$). The remaining papers involved multiple taxa (11.6%, $n = 49$), with the predominant combinations of taxa most often employed by regional faunal studies: including all amphibian and reptile species recorded for a given island, region, local area, or specific site (Fig. 3A). Of these same 423 papers, 35.2% fell under the category Diversity and Distribution ($n = 149$), 26.5% were focused on Ecology ($n = 112$), 14.9% constituted Taxonomy ($n = 63$), 11.8% were classified as Phylogenetics and Biogeography ($n = 50$), and 11.6% were in the Conservation ($n = 49$) category (Fig. 3B).

A quarter of the Diversity and Distribution studies were targeted toward herpetological inventories and surveys (multiple taxa, 25.8%), while the remaining constituted brief distribution reports on amphibians (30.2%) and reptiles (45%), most of which consisted of single occurrence records, provided for a single species, not previously reported for that island, region, or province. Studies in Ecology, including short natural history notes, were almost equally represented among amphibians (31.2%), lizards (20.4%), and snakes (26.8%) but with few papers on turtles (7.1%) and crocodiles (4.5%). Notably, most publications in taxonomy during the last two decades focused on lizards (63.5%), with amphibians and snakes comprising 20.6% and 15.9%, respectively. Similarly, Phylogenetics and Biogeography publications primarily were conducted on lizards (44%), followed by amphibians (34%), and snakes (12%). All taxa were represented well in Conservation studies (10–20%).

New species descriptions

A total of 95 herpetological species (72 lizards, 14 amphibians, and 9 snakes) was reported as new to science and formally described during a prolific period of taxonomic activity spanning the last two decades (Fig. 4A). New species descriptions started to increase in 2009 with peaks in 2010 (18 new species), 2014 (10 new species), and 2020 (11 new species). Species description papers employing only morphological data were published more frequently between 2000 and 2012 but continue to decrease until the present time. Although the first use of molecular data (primarily mitochondrial DNA sequences) in systematic and biogeographical studies of Philippine herpetofauna occurred earlier, the use of molecular data as a form of evidence to justify new species recognition in taxonomic publications first began as a




Figure 2. Philippine herpetological research papers published (2002–2022) **A** with Poisson regression line and **B** local polynomial regression. The gray shading represents the 95% confidence interval of the regression models, and the light green shade indicates the trend in terms of the number of papers published per year.




Figure 3. Total of published papers compiled (2002–2022) **A** category per taxon and **B** category proportions in percentages.

practice in 2009 and steadily increased, eventually becoming nearly (but not quite) routine by 2022 (Fig. 4B). When we mapped the geographical position of origin for holotype specimens of these new species (see Suppl. material 2), we found that most of the holotypes originated on Luzon PAIC islands (43.2%) or were from the Mindanao PAIC landmasses (22.1%), whereas few came from the Palawan PAIC (9.5%), the West Visayan PAIC (8.4%), Mindoro Island/PAIC (7.4%), small islands of Romblon Province (RIG; 3.2%), the Sulu Archipelago (SIG; 2.1%), the Babuyan and/or Batanes groups of islands (2.1%) and the isolated island of Camiguin Sur (2.1%) (Fig. 5A; total number of points = 95).




Figure 4. **A** Accumulation curves depicting the relationships between the cumulative total number of species per taxon (key) and the year of description. The orange bars indicate the number of new species described annually **B** number of papers published under the Taxonomy category, grouped by data type.

Geographical patterns

Brief distribution records, focused on single occurrences for single species from areas where they had not previously been reported were published primarily in one journal (*Herpetological Review*), on the basis of records from the PAICs of Mindanao (31.5%), Luzon (27%), and the West Visayas (16.9%) (Fig. 5B; total number of points = 89; see Suppl. material 2). The majority of new distribution records from the Mindanao PAIC were from the central and southern regions of this large island, with few notes reporting occurrences of particular species from the eastern and western regions of Mindanao or from the northern Mindanao PAIC islands of Samar, Leyte, and Bohol. In contrast, single species distribution records from the Luzon PAIC were not grouped geographically and were more randomly distributed (Fig. 5B). On the West Visayan PAIC, the islands of Cebu, Negros, and Masbate were the focus of most new geographical distribution records. Records from Palawan (11.2%) primarily were from the northernmost portion of this island, as well as the Calamian group of islands, to its north. The remaining distribution records originated in the RIG (6.7%), SIG (2.2%), Mindoro Island/PAIC (2.2%), the Batanes Island Group (1.1%), and Siquijor (1.1%). Occurrence points from natural history notes (Fig. 5B; total number of points = 83) were biased towards the two largest PAICs, Luzon (45.7%) and Mindanao (37.3%).

Targeted herpetological surveys or sustained site-based studies attempting to characterize comprehensively, whole island communities, or regional faunas were concentrated during the last 20 years most heavily on the Luzon (35.6%) and Mindanao (33.1%) PAICs (Fig. 5C; total number of points = 247). Within the Luzon PAIC, targeted fieldwork was conducted mostly in several more northern regions of this large island, but a few sites were sampled well, along southern Luzon's Bicol Peninsula. Within the Mindanao PAIC, surveys were conducted primarily in the northern and northeastern regions (including several large




Figure 5. Maps showing the type localities of **A** new species described in the past two decades **B** localities of new distribution records and natural history notes, and **C** the inventory sites available from publications and curated databases. Dashed squares indicate understudied regions (**C**) where comprehensive surveys have not yet been published, but natural history notes and new geographical records have begun to fill spatial gaps.

mountains on northern Mindanao Island itself, but also on the East Visayan islands of Samar and Leyte). Only a relatively few sites have been sampled reasonably well in the southern portion of Mindanao Island during the last two decades. The West Visayan PAIC received attention in the form of a few surveys (10.1%) in recent years, most notably on Cebu, Panay, and Negros Islands. Approximately equal numbers of survey points were collated from the RIG (5.6%), Palawan (4.8%), and the island groups of Babuyan (5.3%) and Batanes (3.2%). There were also survey efforts in Camiguin Sur (1.6%) and Siquijor (0.4%).

First authorship nationality and gender gaps

We found approximately equal cases of Filipino- versus foreign-first authorships (1.0:1.1), with the number of Filipino first authors steadily increasing during the period of 2002–2022 (Fig. 6A). In the last decade (2013–2022), 63.7% of papers were first-authored by Filipinos which is a three-fold increase from the 2002–2012 estimate (21.1%). We identified a wide gender gap in Philippine herpetology, with males outnumbering females when quantified as a function of first authorship (7.1:1.0) during the past 20 years; only approximately 12.3% of all papers were first-authored by females (Fig. 6B).




Figure 6. The total number of papers grouped by the first author's **A** gender (7.13 male: 1.0 female) and **B** nationality (1.07 foreign: 1.0 Filipino).

Discussion

We are greatly encouraged by the increasingly collaborative and much more equitable and inclusive nature of Philippine herpetology. Great strides have been made during the past two decades in terms of our collective knowledge of herpetological biodiversity in the archipelago, public awareness and interest in amphibians and reptiles, and an increasingly vocal environmentalist movement—combined with the establishment of new protected areas and new laws for protecting historically underappreciated groups such as amphibians and reptiles (Diesmos and Brown 2011; Brown et al. 2012a; Diesmos et al. 2014; Gonzalez et al. 2018). The past 20 years provide us with a clearer picture of the progress made, gaps filled and unfilled, and collaborations formed, which now serve as a springboard for or guide to additional progress in the coming decade for our community.

Herpetology studies during the last two decades

Our appreciation of the diversity and distribution of amphibians and reptiles in the Philippines has increased during the last 20 years, with the continued efforts towards conducting targeted faunal inventories and resurveys, supplemented by short distribution notes reporting single species occurrences (Brown et al. 2012a, 2013a). Faunal inventories have dominated herpetological studies during the last two decades, which is beneficial for informing habitat conservation measures (Margules and Pressey 2000) and these works positively impact species conservation assessments (see Gonzalez et al. 2018; IUCN 2022). Importantly, data from these studies help address knowledge gaps in species distributions, broaden stakeholder inclusion and participation in Philippine herpetology, and improve our collective knowledge and understanding of biodiversity (Hortal et al. 2015).

Region-wide herpetological inventories in recent years covered a wide range of habitats and utilized multiple datasets like field records and museum specimens (Oliveros et al. 2011; Siler et al. 2011d, 2012b; Brown et al. 2012a, 2013b; Devan-Song and Brown 2012; Mcleod et al. 2012; Sanguila et al. 2016; Supsup et al. 2016). Complementing these regional inventories were site-based studies along different environmental gradients in inland montane habitats (Alcala et al. 2004; Plaza and Sanguila 2015; Supsup et al. 2017, 2020; Gojo-Cruz et al. 2018; Pitogo et al. 2021a; Sanguila et al. 2021) and island ecosystems (Bucol et al. 2011; Siler et al. 2012a; Venturina et al. 2020; Clores et al. 2021; Meneses et al. 2022). Species accounts resulting from these targeted surveys provide a much clearer picture of herpetological communities occurring in several critically threatened habitats, which serve as important baseline studies to track long-term changes in amphibian and reptile communities over time (e.g., Brown et al. 2000a, vs Siler et al. 2011d; Siler et al. 2012b, vs Meneses et al. 2022; Brown and Alcala 1986, vs Supsup et al. 2016; Brown 2015, vs Pitogo et al. 2021a). Continued conducting of targeted surveys and resurveys are still urgently needed to gather empirical field data and species occurrences throughout the Philippines, especially in understudied regions, small islands (see discussion on geographical patterns), and other threatened habitats (e.g., karst habitats, lowland forest fragments, and understudied protected areas). Repeated resurveys spanning time, seasonal variation, space, and disturbance (both anthropogenic [deforestation, mining, road-building] and natural [typhoons, earthquakes, tsunamis]; Peterson et al. 2017) in key areas and habitat types, whenever resources are available, are strongly recommended (Brown et al. 2012a; Sanguila et al. 2016).

Many of the ecology studies of Philippine herpetofauna are short observations on species' reproductive biology, natural history, habitat use, diet, and behavior (see Suppl. material 1); all of these biodiversity information products supplement our limited understanding of the natural history of many endemic and native species (i.e., Meneses 2020; Brown et al. 2021a, b, c; Maglangit et al. 2021; Pitogo 2021). However, more in-depth ecological and herpetological community composition studies have bolstered patterns that were apparent two decades ago (Brown et al. 2002a) and revealed new trends. Elevational studies have repeatedly characterized a positive relationship between elevation and proportional endemism (the portion of species at a given site that is endemic to the Philippines), but an inversely proportional relationship between elevation and species diversity (Brown and Alcala 1961; Brown et al. 2000a; Ferner et al. 2001; Diesmos et al. 2004a; Siler et al. 2011d; Brown et al. 2013a; Pitogo et al. 2021b; Brown et al. 2022; Meneses et al. 2022), with a potential mid-elevation peak in species diversity was detected in some areas (Gojo-Cruz et al. 2019; Supsup et al. 2022). There is evidence that elevation strongly shapes broad-scale distribution patterns of Philippine herpetological communities, but other habitat characteristics were also found to influence the fine-scale distribution of species (Siler et al. 2011c; Pitogo et al. 2021b; Seidl et al. 2019; Supsup et al. 2020, 2022). Functional traits potentially drive distribution patterns (Pitogo et al. 2021b) and may influence species' responses to environmental changes (Scheffers et al. 2013). Resource partitioning has also been recently investigated in amphibians (Shoeman-Goodier et al. 2019; Plaza et al. 2021) and reptiles (Garcia et al. 2014; see also Auffenberg and Auffenberg 1988), improving our

knowledge of trophic interactions among co-occurring species. These hypothesis-driven studies are welcomed developments in Philippine herpetology and should pave the way for a greater understanding of herpetological communities, limiting resources, and habitats critical for individual species' persistence and community resilience. Despite advances in research focusing on community ecology, there has been limited progress in studies involving reproductive biology, development (see Flores et al. 2023), physiology, and behavior (Lama and Senarillos 2023) during the past 20 years—as was also the case during the first review (Brown et al. 2002a).

Complementing field-based studies are works utilizing genetic data to understand the evolutionary history and phylogenetic relationships of species. The subdisciplines of model-based statistical phylogenetic systematics (Hillis et al. 1996) and quantitative biogeographical inference (Ronquist 1997) were just emerging at the time of the first review (Brown et al. 2002a). During the last two decades, numerous studies involving robust genetic datasets from Philippine species (and/or whole clades), from multilocus Sanger sequence datasets to genome-scale datasets, orders of magnitude more expansive, have provided inference into biogeographic patterns and a diversity of underlying processes. These studies have contributed greatly to our appreciation of true species diversity, species' distributions, routes of colonization, and the dynamic geographical template's role in contributing to evolutionary diversification within the archipelago (Brown and Diesmos 2009; Blackburn et al. 2010, 2013; Siler et al. 2012a, 2014a, b, c; Brown et al. 2013a, 2015a, 2016; Linkem et al. 2013a; Oaks et al. 2013, 2019, 2022; Brown 2016; Chan and Brown 2017). These studies also provide insights into surprising and intriguing evolutionary relationships not observed in other vertebrate groups (Weinell and Brown 2017; Meneses et al. 2020a; Weinell et al. 2020b), and patterns of cryptic speciation (Sanguila et al. 2011; Barley et al. 2013, 2021; Linkem and Brown 2013; Welton et al. 2013a, 2017; Weinell et al. 2020a, c; Abraham et al. 2021; Chan et al. 2021), many of which have informed taxonomic developments, fueling the description of nearly 100 new species during the past twenty years (see Taxonomic Progress, below). We expect an increase in this type of work in coming years, especially with the widespread availability and declining cost of genomic data (Carter et al. 2023), which increases accuracy in phylogenetic inference, while greatly clarifying previously obfuscated species boundaries by allowing nuanced insight into related phenomena, such as gene flow, admixture, hybridization, lineage sorting, and retention of ancestral polymorphism (Alexander et al. 2016; Wood et al. 2020; Chan et al. 2021).

Conservation of Philippine herpetofauna

The last two decades involved increasing numbers of conservation studies, particularly focused on large and charismatic species like crocodiles (van de Ven et al. 2009; van der Ploeg et al. 2011; Brown et al. 2021a), turtles (Abreo et al. 2016; Sy et al. 2020), monitor lizards (Welton et al. 2013b, 2020; Abaño-Sarigumba et al. 2018), and sailfin lizards (Siler et al. 2014d; Heinrich et al. 2021). These species are most often the highly traded and threatened species in the Philippines (Cruz and Lagunzad 2021). Although amphibians receive less attention in the conservation literature, several efforts were made to highlight their high

vulnerability to environmental perturbations and climatic changes (Alcala et al. 2004, 2012a, b; Brown et al. 2012a; Diesmos 2012; Diesmos et al. 2014). Recent amphibian rediscoveries of “lost” species have captured public attention (e.g., Bittel 2015; Pitogo and Saavedra 2021), which bolsters public interest and stimulates conservation research and action in overlooked taxa (e.g., Brown and Alcala 2000; Pitogo and Saavedra 2023). Overall, peer-reviewed literature on the conservation of Philippine herpetofauna has increased substantially from 20 years ago (Brown et al. 2002a). We are encouraged by these developments and hope to maintain this momentum, build further on these gains (including the development of more outreach materials in support of conservation objectives), and strive to improve the conservation attention to Philippine amphibians and reptiles (Brown et al. 2012a; Gonzalez et al. 2018).

Based on the most recent and updated IUCN assessments on Philippine herpetofauna (ca. 475 species), approximately 13.2% (63 species) and 13% (62 species) are threatened and data deficient, respectively (IUCN 2023). Upwards of 5.9% (28 species) are still unassessed. Notably, 131 of these 153 species are Philippine endemics, many of which have not been observed since their original descriptions. Many additional unrecognized and unprotected species await taxonomic descriptions and, as such, are at increased risk of extinction (McDonald et al. 2022). Continued lowland habitat loss, brought about by forest conversion into less complex habitats that are not suitable to support high levels of biodiversity is likely the most substantial threat to these taxa (Heaney and Regalado 1998; Brown et al. 2002a; Diesmos et al. 2004a; Suarez and Sajise 2010; BMB-DENR 2016; Gojo-Cruz et al. 2019; Decena et al. 2020; Supsup et al. 2020). Although there has been an increase in the establishment of protected areas to avert the threat of habitat loss, many of these protected areas (PAs; i.e., national parks, natural parks, biotic areas, protected landscapes) do not overlap with key biodiversity areas (Mallari et al. 2015; Supsup et al. 2023), have poor to fair management effectiveness (Madarang et al. 2017), have never been properly inventoried for terrestrial biodiversity despite their establishment decades ago (Brown et al. 2002b), and can only marginally reduce forest cover loss (Apan et al. 2017). Despite these challenges, we acknowledge that habitat protection through the establishment of PAs and other effective area-based conservation measures are the first steps and are still the most effective measures for long-term species conservation and survival/persistence. Additionally, public education and societal awareness concerning threats posed by emerging infectious diseases (Swei et al. 2011a, b; Diesmos et al. 2012; O’Hanlon et al. 2018; Byrne et al. 2019) and pet-trade facilitated introductions of highly competitive alien and invasive species (Diesmos et al. 2006; Pili et al. 2019, 2021) are all on the rise.

Taxonomic progress

At the time that the state of Philippine herpetology was last reviewed, Brown et al. (2002a) recognized a total of 101 species of Philippine amphibians (77% endemic) and approximately 258 species of Philippine reptiles (65% endemic). During the last 20 years, during which nearly 100 new species were added to the archipelago’s fauna, we recorded differing, taxonomically dependent proportional increases in the number of additionally recognized reptiles (258 vs 361, 28%)

versus amphibians (101 vs 114, an 11% increase) between 2002 and 2022 (Fig. 3A), indicating that the archipelago's native species diversity is far from comprehensively understood. This taxonomic shortfall warrants continued surveys and resurveys, plus targeted taxonomic revisionary attention to particular clades before we might conclude that the fauna is reasonably well characterized. As such, we can anticipate additional increases in cumulative total species diversity estimates in the coming decades, but the question remains, by how much?

Based on the numbers of suspected undescribed species known at that time, Brown et al. (2002a) suggested that herpetological species diversity might increase from 101 to possibly as many as 125–130 (~ 22%) amphibians and from 258 to approximately 275–280 (~ 8%) reptiles. Explanations for why amphibian diversity proportionally increased by less (11%) than the 22% estimated by Brown et al. (2002a) or why reptiles species diversity increased by proportionally far more (28%) than the 8% estimated by Brown et al. (2002a) appears to be a simple function of the fact that researchers (who were actively completing taxonomic studies) showed more interest in systematic and biogeographic questions for which Philippine reptiles represented preferable focal study subjects (e.g., Siler et al. 2009a, 2010a, b, c, d, 2011a, b, c, 2012a, 2014a, b, c, e; Welton et al. 2009, 2010a, b; Brown et al. 2010a, b, c; Linkem et al. 2010a, b, c, 2013; Barley et al. 2013, 2020, 2021; Weinell et al. 2020a, b, c; Eliades et al. 2021) than was the case for research topics involving amphibian study systems (Brown et al. 2009, 2015a, b; Siler et al. 2009b, c, 2010e; Fuiten et al. 2011; Brown 2015; Diesmos et al. 2020; Herr et al. 2021).

Recent efforts to conduct comprehensive herpetological surveys and resurveys have provided a near-complete estimation of the amphibian and reptile diversity and endemism of several islands (e.g., Siler et al. 2012b vs Meneses et al. 2022), mountain ranges (Brown et al. 2000a vs Siler et al. 2011d), or other conspicuous geographical subcenters of diversity in the archipelago through the years (Plaza and Sanguila 2015; Sanguila et al. 2016, 2021). The notable progress of an increased number of described species has been species descriptions (Brown et al. 2010a, c, 2011a, b; Brown 2015; Barley et al. 2021), resurrections of species (e.g., Brown et al. 2000b; Siler et al. 2020), redescrptions of poorly understood taxa (Davis et al. 2014, 2015; Wynn et al. 2016), and revisionary considerations of species boundaries within entire clades (e.g., Linkem and Brown 2013; Brown et al. 2015a, b; Barley et al. 2020). The vast majority of these studies involved the integration of traditional morphological characters (typical of the early 2000s; Brown and Stuart 2012) with molecular data, bioacoustic analyses, ecological information, or other independent data streams (Fig. 4B).

The majority of the 95 species newly described or recognized during the last twenty years are lizards (Fig. 3A); these are dominated by new species of the genera *Brachymeles* ($n = 12$) and *Pseudogekko* ($n = 5$; see Siler et al. 2020; Brown et al. 2020). The genus *Parvosincus* was expanded by 13 species (Brown et al. 2010c; Linkem and Brown 2013; Siler et al. 2014e) and the genus *Eutropis* by nine new taxa (Barley et al. 2020, 2021). Other new lizard species were described from the genera *Lepidodactylus*, *Gekko*, *Luperosaurus*, *Varanus*, *Cyrtodactylus*, and *Lygosoma*; in total, more than 72 previously unrecognized lizard species have been identified in the past two decades. In many ways, the most spectacular lizard discovery of the past two decades involved the

description of a third species of frugivorous monitor lizard, *Varanus bitatawa* from the southern Sierra Madre Mountain Range of Luzon Island (Welton et al. 2010c, 2012), which has since been confirmed from northern Luzon, including the northernmost reaches of the Cordillera Mountain Range or western Luzon (Abaño-Sarigumba et al. 2018; Meneses et al. 2020b). Five other species of Philippine-endemic monitor lizards in the genus *Varanus* were described during the last twenty years as well (Koch et al. 2010; Welton et al. 2014b).

In amphibians, the greatest taxonomic activities were associated with seven descriptions of new Ceratobatrachid frogs in the genus *Platymantis* (Siler et al. 2007, 2009a, 2010e; Brown et al. 2015a, b; Diesmos et al. 2020). Additional discoveries and descriptions of new species were assigned to the genera *Limnonectes*, *Sanguirana*, *Pulchrana*, *Leptobrachium*, and *Kaloula* (Brown et al. 2009a, 2016, 2017; Siler et al. 2009b; Fuiten et al. 2011; Abraham et al. 2021; Herr et al. 2021). These recorded a total increase of 14 amphibian species in two decades, and we suspect that many more await discovery, because many amphibian clades are in urgent need of taxonomic resolution (e.g., *Platymantis*, *Philautus*, and *Occidozyga*; Brown et al. 2015a, Chan et al. 2021, 2022; Flury et al. 2021), which is currently challenged by limited numbers of recordings (insufficient sample sizes necessary to permit quantitative analyses and statistical tests for species delimitation) or a complete lack of acoustic data for particular species or species groups (Brown and Alcalá 1994; Alcalá and Brown 1999; Hertwig et al. 2011; Herr et al. in press; Khalighifar et al. 2021).

One notable and striking recent discovery was the characterization of an ancient, archipelago-wide clade—a nearly 40 million-year-old endemic Philippine evolutionary radiation, now recognized as the archipelago's only endemic reptile family: snakes of the clade Cyclocoridae (Weinell and Brown 2017). Two surprising elements of this discovery were apparent; first, the unpredicted finding that snakes of the genera *Cyclocorus*, *Hologerrhum*, *Oxyrhabdium*, and *Myersophis* were a monophyletic group (they had previously and variably been treated as members of separate families or left incertae sedis, of unknown taxonomic affinities; McDiarmid et al. 1999). Second, it was clear from multilocus phylogenetic analyses that an unnamed, genus-level lineage had been overlooked (Weinell and Brown 2017; Weinell et al. 2020a). Description of the new miniaturized genus and species of snake of the family Cyclocoridae, *Levitonius mirus*, was based on three specimens of this secretive, fossorial snake from Samar and Leyte Islands. *Levitonius* exhibits highly distinctive morphology associated with its miniaturized body form, fossorial habitat, and unique diet, consisting solely of soil invertebrates (Weinell et al. 2020a). This discovery also used CT-scan, a novel method to characterize deep internal anatomy, together with molecular data, traditional morphological characters, diet, and ecological niche data (Weinell et al. 2020a), a novel degree of data-type integration to be associated with a taxonomic description for a Philippine species. Subsequent phylogenomic analyses (Das et al. 2023) confirmed the early-branching phylogenetic placement of Cyclocoridae, closely related to the globally distributed snake clade Elapoidea, which includes coral snakes and cobras. Other, highly unique, or unpredicted snake species discovered included a new species of blind snake phenotypically similar to *Acutotyphlops* (Wallach et al. 2007) from northern Luzon and the highly distinctive, krait-like *Calliophis salitan* (Brown et al. 2018, 2021b). The former is a genus otherwise restricted to the Solomon

Islands, which creates a conspicuously unusual and disjunct distribution (the single Philippine species has yet to be included in a phylogenetic analysis, which would be necessary to evaluate this disjunct and somewhat suspect taxonomic/geographic placement). In contrast, *Calliophis salitan* is related to the giant, long-glanded tropical coral snakes of the *C. bivirgata* group and most likely constitutes a separate, unique invasion of the archipelago, apart from other Philippine elapid snakes. In summary, nine species of snakes were described or newly recognized, in the genera *Acutotyphlops*, *Calamaria*, *Calliophis*, *Dendrelaphis*, *Hemibungarus*, *Levitonius*, *Malayotyphlops*, and *Lycodon* (Gaulke 2002, 2011; Wallach et al. 2007; Siler et al. 2013; Wynn et al. 2016; Weinell and Brown 2017; Leviton et al. 2018; Weinell et al. 2019, 2020a, b, c).

Geographical patterns of herpetological surveys

Our synthesis of available geographic data suggests that the last two decades were characterized by a significant surge of herpetological research across the archipelago. Herpetological surveys conducted on most major islands, particularly those that were not visited before (e.g., island groups of Babuyans, Batanes, and Romblon Province) have led to a stunning number of new species discoveries, elevating sharply the herpetological diversity of the country. Rediscoveries of poorly known species have also provided new insights about their population status and ignited the needed hope for conservation (e.g., Diesmos et al. 2004b; Siler et al. 2011b, c; Bittel 2015; Oliver et al. 2020; Supsup and Carestia 2020; Brown et al. 2021b; Pitogo and Saavedra 2021, 2023; Meneses et al. 2022). However, despite the highly celebrated discoveries, much work is still needed because many small islands and isolated habitats remain unexplored or have not been surveyed thoroughly (see Fig. 6). The islands of Jolo and Basilan in the Sulu Archipelago are the notably less explored areas of the country despite their zoogeographic importance (Seale 1917; Taylor 1918; Gaulke 1993, 1994, 1995). Few biologists have visited these islands during the last century due to logistical and security constraints. The only attempt at a comprehensive study of this archipelago (with specimens collected and still available for reconsideration) was the work of Taylor (1918); unfortunately, many of his specimens from the region, including holotypes of several of the Sulu Archipelago's endemic species (e.g., *Luperosaurus joloensis*, *Brachymeles vermis*, *B. suluensis*) were lost during World War II (Brown and Rabor 1967; Brown and Alcalá 1974; Gaulke 1993, 1994, 1995; Uetz et al. 2023). Because of the lack of studies on these islands, many wildlife biologists and biogeographers are still puzzled by uncertain taxonomic affinities and conservation status of the endemic biodiversity of the Sulu Archipelago (Siler et al. 2012c; Spinks et al. 2012; Brown and Siler 2013; Chan et al. 2021).

The apparent absence of peer-reviewed herpetological studies on the islands of Polillo (east of Luzon) and Siargao Island (northeast of Mindanao) is somewhat artefactual, and due to the fact that the majority of fieldwork conducted on these two small islands is only available as unpublished reports (but see Ross and Lazel 1991; Nuñez and Galorio 2015; Sanguila et al. 2016; Quibod et al. 2021); nevertheless, some collection information (specimens deposited in accessible biodiversity repositories such as Smithsonian National Museum Natural History [USNM], University of Kansas Natural History Museum

[KUNHM], Philippine National Museum of Natural History [PNMNH], and Father Saturnino Urios University [FSUU]) are readily accessible via online biodiversity repositories (e.g., GBIF, iDigBio). In addition to herpetologically unexplored regions of the country, the intact forest habitats of the central Sierra Madre Mountain Range of Luzon (in particular, higher elevations), including the relatively large but fragmented forests to the west have not been thoroughly explored (but see Brown et al. 2000a, 2007, 2010c, 2013b; Siler et al. 2011c; Gojo-Cruz et al. 2018, 2019). Similarly, isolated high-elevation forest habitats in southern Luzon along the borders of Quezon and Bicol Provinces and in Mindoro have not been explored well. Except for the relatively well-explored Caraga region in northeastern Mindanao (Sanguila et al. 2016), many of Mindanao's forests remain herpetologically underexplored, a condition which has persisted during the last two decades (and last century); this is particularly true of western, central, and southern Mindanao, from Mt. Piapayungan southward to the Mt. Latician complex (Taylor 1921, 1922a, b; Sanguila et al. 2016; Pitogo et al. 2021a; Maglangit et al. 2022). As in the Sulu Archipelago, the limited availability of published results from field studies on Mindanao is due, in part, to logistical and security challenges, as well as a lack of local regional expertise, training, and experience with field-based herpetological inventories (Sanguila et al. 2016; Pitogo et al. 2021a; Pitogo and Saavedra 2021).

Despite these gaps, we should note that during the last two decades, there have been significant field-based efforts focused on survey-resurvey studies at important, formerly incompletely understood areas. Several key studies have revisited areas that were targeted in periods before 2002, with the general goal of reassessing, completing, and/or providing a time series (before and after comparison) to enable a temporal perspective on faunal investigations conducted previously, and in light of deforestation, land use change, and global climate change: Zambales Mountains (Brown et al. 1996; Devan-Song and Brown 2012), the central Sierra Madre Mountains of eastern Luzon (Brown et al. 2000a; Siler et al. 2011d), the northern portions of the Sierra Madre Mountains (Brown et al. 2013b), the northern Cordillera Mountain Range of western Luzon (Diesmos et al. 2004a; Brown et al. 2012b), Panay Island (Ferner et al. 2001; Gaulke 2011), Cebu Island (Brown and Alcala 1986; Supsup et al. 2016), and Negros Island (Brown and Alcala 1955, 1961, 1963, 1986; Alcala 1958; Bucol et al. 2019), as well as recent resurveys focused on reassessments of the faunas of northeastern, central eastern, and southern Mindanao (Sanguila et al. 2016, 2021; Pitogo et al. 2021a; Plaza et al. 2021).

Many additional, earlier faunal studies (published before 2001; other areas that have been surveyed, but as of yet, have not been published) are now urgent priorities for survey-resurvey studies, hopefully in the near future (Fig. 5C). We recommend undertaking this work as soon as possible because such areas may contain critically important populations of endemic species, 'lost' species (i.e., species not encountered since their original descriptions and for which holotype specimens were destroyed in World War II), exceedingly rare species, and species awaiting rediscovery and discovery (currently unknown to science), all of which may already be facing cryptic extinction risk brought by habitat degradation and destruction (McDonald et al. 2022). We strongly encourage researchers to consider understudied areas as top priorities for field-based biodiversity research in the coming years (Fig. 5C). Also, for areas that

have not been explored due to logistical obstacles and/or security challenges (e.g., the Sulu Archipelago, southern and southwestern Mindanao Island, etc.), we strongly encourage capacity-building activities for local institutions, universities, and other community stakeholders (e.g., training of residents, students, and other community members) to conduct field-based biodiversity research involving amphibians and reptiles. Such an approach is most likely the safest, most cost-effective, and most feasible strategy for moving forward to address geographical hiatuses and knowledge gaps represented by unexplored, politically charged, or otherwise sensitive areas (see Ramírez-Castañeda et al. 2022). Training residents to effectively survey their local biodiversity reduces reliance on foreign institutions (or groups that are not residents of an area), builds the research capacity of stakeholders who may not have had the opportunity to engage in science, maximizes scholarly equity, and increases the potential for local governance and conservation action.

Diversity, equity, and inclusion in herpetology research

The rise of Filipino-first authorship in Philippine herpetological studies during the last two decades (Fig. 6B) is an encouraging development since the first review (Brown et al. 2002a). This substantial increase indicates an increased interest among early-career Filipino researchers in the discipline, which formerly was limited to a few Filipinos and their foreign collaborators. It is apparent that a large proportion of the last two decades of Filipino-led studies were field surveys and descriptive studies, whereas the majority of sophisticated studies utilizing genetic data relying on large molecular datasets or genomic analyses were led by non-Filipinos. This disparity reflects the relatively limited capacity for genomics in the Philippines, highlighting the importance of equitable collaborations to ensure skill and technology transfers between local and foreign researchers. A collaborative approach provides opportunities for capacity development for less experienced researchers (e.g., mentorship, writing, decision-making) and may further improve local interest in scientific research (Ramírez-Castañeda et al. 2022). We have seen many multi-national collaborations during the past two decades, which are consequential and contribute to our present understanding and appreciation of Philippine herpetofauna. However, we also acknowledge that access to these opportunities, including advancement in scientific careers, is limited by many socioeconomic factors.

Women are historically underrepresented in herpetology but there has been an increase in female authorship in research on amphibians and reptiles, potentially narrowing the gap if this positive trajectory continues (Rock et al. 2021). Nevertheless, the gender gap in Philippine herpetology (7.13 male per 1.0 female, as first authors) is far from the global average (1.95 male per 1.0 female first authors); ameliorating this disparity will require more representation from women in scientific publications. Despite the substantial gender gap, we are inspired by the continued emergence of next-generation Filipinas and greatly value their contributions to the advancement of the field during the past 20 years. We hope that many additional early-career women researchers, along with other historically underrepresented Filipino groups, will be encouraged to participate in the study of amphibians and reptiles toward a more inclusive and equitable scientific community in the Philippines. Additionally, finding ways to

communicate the results of our studies to the public while narrowing the gap between the scientific community and the lay-/citizen science community will be an important step for rendering our science more accessible to the public and policymakers, while bridging the gaps between science and policy (Young et al. 2014; Ramírez-Castañeda et al. 2022).

Future directions

Clearly at a crossroads of topical shifts in research themes, increased engagement, equity, inclusion, and representation of diversity in collaborations, Philippine herpetology has undergone a demonstrable maturation since the first review (Brown et al. 2002a). This work sets the stage for what we hope and anticipate will be an increasingly collaborative and inclusive engagement by diverse kinds of herpetologists during the coming several decades, in continuation of the rich history and development of our collective understanding of the amphibians and reptiles in the archipelago. Much progress has been made in some areas (increased general public education and lay-public interest, increased numbers of publications by a broad array of early-career herpetologists, narrowing of the first authorship gap between Filipinos versus foreigners), whereas other equity gaps (e.g., first authorship gender) still remain, requiring increased attention towards fostering diversity and encouraging the engagement of people from a broader array of backgrounds. It is our hope, in compiling this synthesis, that we can challenge the community of individuals, groups, and institutions interested in amphibians and reptiles of the Philippines to pursue some of the conspicuous gaps in research themes identified here and encourage early-career herpetologists to pursue research topics that have advanced in surrounding countries during the last quarter century, but which have not received comparable interest or attention in the Philippines (e.g., amphibian larval biology, developmental studies, etc.). Similarly, historically understudied geographical gaps identified in this review should be viewed as opportunities for increased attention and enhanced collaboration, both among Philippine institutions and between Filipino and foreign herpetologists. The challenge issued by Walter C. Brown a few years before the last review (e.g., 'The State of Philippine Herpetology,' Brown et al. 2002a) still stands: "Rather than view Philippine herpetology as something you might be tempted to divide up, why not just see how much you can accomplish, together, in collaboration?" (W. C. Brown to RMB and A. C. Alcala, personal communication 1998). In accordance with this perspective, and in light of the progress made during the last two decades, we are quite sure that the future of Philippine herpetology will profit most from increased engagement, involving a diversity of people, and embracing increasingly broad thematic research questions in collaboration, all with the common goal of understanding, appreciating, and conserving the archipelago's spectacularly unique amphibian and reptile fauna.

Acknowledgements

Dedicated to the memory of Angel C. Alcala (1929–2023) and Walter C. Brown (1913–2002) in recognition of their numerous fundamental contributions to the study of the amphibians and reptiles of the Philippines; with particular recogni-

tion for the research synergy they accomplished during the most trying of political times and with respect for the breakthrough international collaboration they forged—transcending historical, racial, societal, and political stereotypes—and all in the name of the passion they shared for the archipelago’s herpetological fauna. We thank Justin Bernstein, Marites Sanguila, an anonymous reviewer, and ZooKeys editors for constructive comments on earlier drafts of this manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

During the preparation of this manuscript, we were supported by a grant from the US National Science Foundation (DEB 1654388 + Pandemic Supplement to RMB), a KU College of Liberal Arts and Sciences Graduate Assistantship in Support of Faculty Research award, a KU Biodiversity Institute Research and Exploration Fund grant, and funding from the South, Southeast, and East Asia Fund of the KU Office of International Affairs, University of Kansas.

Author contributions

All authors participated in the design and conduct of the study. The literature search, data analysis, and preparation of tables and figures were all performed by CGM, KMEP, and CES. All authors wrote and reviewed the early drafts and approved the final version of the manuscript.

Author ORCIDs

Camila G. Meneses  <https://orcid.org/0000-0003-0594-5436>

Kier Mitchel E. Pitogo  <https://orcid.org/0000-0002-5785-8281>

Christian E. Supsup  <https://orcid.org/0000-0001-5176-8181>

Rafe M. Brown  <https://orcid.org/0000-0001-5338-0658>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Abaño-Sarigumba T, Brown RM, Welton LJ, Ibañez JC (2018) Records of the Northern Sierra Madre Forest Monitor *Varanus bitatawa* from the northern Cordillera Mountain Range of Luzon Island, Philippines. *Herpetological Review* 49: 208–209.
- Abraham RK, Herr MW, Sterkhova VV, Otterholt R, Siler CD, Sanguila MB, Brown RM (2021) Revisiting Linnaean and Wallacean shortfalls in Mindanao fanged frogs: The *Limnonectes magnus* complex consists of only two species. *Herpetological Monograph* 35(1): 112–140. <https://doi.org/10.1655/HERPMONOGRAPHS-D-20-00010>

- Abreo NAS, Macusi ED, Blatchley DD, Cuenca G (2016) Ingestion of Marine Plastic Debris by Green Turtle (*Chelonia mydas*) in Davao Gulf, Mindanao, Philippines. *Philippine Journal of Science* 145(1): 17–23.
- Alcala AC (1958) Amphibians on Negros Island, including two new records. *Silliman Journal* 5: 171–174.
- Alcala AC (1962) Breeding behavior and early development of frogs of Negros, Philippines Islands. *Copeia* 1962(4): 679–726. <https://doi.org/10.2307/1440671>
- Alcala AC (1967) Population biology of the “flying” lizards, *Draco volans*, on Negros Island, Philippines. *University of the Philippines Natural and Applied Sciences Bulletin* 20: 335–372.
- Alcala AC (1970) Notes on the population biology of the lizard *Mabuya multicarinata*. *The Philippine Biota* 3: 591–611. <https://doi.org/10.2307/1442304>
- Alcala AC, Brown WC (1956) Early life history of two Philippine frogs with notes on egg deposition. *Herpetologica* 12: 241–246. <https://www.jstor.org/stable/3889781>
- Alcala AC, Brown WC (1966) Thermal relations of two tropical island lizards on Negros Island, Philippine Islands. *Copeia* 1966(3): 593. <https://doi.org/10.2307/1441095>
- Alcala AC, Brown WC (1967) Population ecology of the tropical scincoid lizard. *Emoia atrocostata*, in the Philippines. *Copeia* 1967(3): 596–604. <https://doi.org/10.2307/1442238>
- Alcala AC, Brown WC (1982) Reproductive biology of some species of *Philautus* (Rhacophoridae) and other Philippine anurans. *Kalikasan. Philippine Journal of Biology* 11: 203–226.
- Alcala AC, Brown WC (1999) Philippine frogs of the genus *Platymantis* (Amphibia: Ranidae). *Philippine Journal of Science* 128: 281–287.
- Alcala AC, Custodio CC (1995) Status of endemic Philippine amphibian populations. *Sylvatrop: the Technical Journal of Philippine Ecosystems and Natural Resources* 5: 72–86.
- Alcala AC, Custodio CC, Diesmos AC, Gonzalez JCT (1995) List of amphibians of Mt. Makiling, Laguna, Philippines, with notes on their population status. *Sylvatrop: the Technical Journal of Philippine Ecosystems and Natural Resources* 5: 65–71.
- Alcala EL, Alcala AC, Dolino CN (2004) Amphibians and reptiles in tropical rainforest fragments on Negros Island, the Philippines. *Environmental Conservation* 31(3): 254–261. <https://doi.org/10.1017/S0376892904001407>
- Alcala AC, Bucol AA, Diesmos AC, Brown RM (2012a) Vulnerability of Philippine amphibians to climate change. *Philippine Journal of Science* 141(1): 77–87.
- Alcala AC, Bucol AA, Alcala EL, Brown RM (2012b) Decreasing population size of the Philippine Limestone Frog, *Platymantis insulatus*. *FrogLog* 104: 44–45.
- Alexander AM, Su Y-C, Oliveros CH, Olson KV, Travers SL, Brown RM (2016) Genomic data reveals potential for hybridization, introgression, and incomplete lineage sorting to confound phylogenetic relationships in an adaptive radiation of narrow-mouth frogs. *Evolution; International Journal of Organic Evolution* 71(2): 475–488. <https://doi.org/10.1111/evo.13133>
- Apan A, Suarez LA, Maraseni T, Castillo JA (2017) The rate, extent and spatial predictors of forest loss (2000–2012) in the terrestrial protected areas of the Philippines. *Applied Geography (Sevenoaks, England)* 81: 32–42. <https://doi.org/10.1016/j.apgeog.2017.02.007>
- Auffenberg W, Auffenberg T (1988) Resource partitioning in a community of Philippine skinks (Sauria: Scincidae). *Bulletin of the Florida State Museum Biological Sciences* 32: 151–219.

- Barley AJ, White J, Diesmos AC, Brown RM (2013) The challenge of species delimitation at the extremes: Diversification without morphological change in Philippine sun skinks. *Evolution; International Journal of Organic Evolution* 67(12): 3556–3572. <https://doi.org/10.1111/evo.12219>
- Barley AJ, Datta-Roy A, Karanth KP, Thompson R, Brown RM (2014) Sun skink diversification across the Indian-Southeast Asian biogeographical interface. *Journal of Biogeography* 42(2): 292–304. <https://doi.org/10.1111/jbi.12397>
- Barley AJ, Monnahan PJ, Thomson R, Grismer LL, Brown RM (2015) Sun skink landscape genomics across Southeast Asia: The evolution of genetic and morphological diversity across a heterogeneous and fragmented landscape. *Molecular Ecology* 24: 1696–1712. <https://doi.org/10.1111/mec.13151>
- Barley AJ, Diesmos AC, Siler CD, Martinez CM, Brown RM (2020) Taxonomic revision of Philippine Sun Skinks (Reptilia: Squamata: Scincidae: *Eutropis*), and descriptions of eight new species. *Herpetological Monograph* 34(1): 39–70. <https://doi.org/10.1655/HERPMONOGRAPHS-D-19-00009.1>
- Barley AJ, Sanguila MB, Brown RM (2021) A new species of Sun Skink (Reptilia: Squamata: Scincidae: *Eutropis*) from the Zamboanga Peninsula, southwestern Mindanao Island, Philippines. *Philippine Journal of Systematic Biology* 14(2): 1–18. <https://doi.org/10.26757/pjsb2020b14012>
- Bernstein JM, Murphy JC, Voris HK, Brown RM, Ruane S (2021a) Phylogenetics of Mud Snakes (Squamata: Serpentes: Homalopsidae); a paradox of both undescribed diversity and taxonomic inflation. *Molecular Phylogenetics and Evolution* 160: 107109. <https://doi.org/10.1016/j.ympev.2021.107109>
- Bernstein JM, Bautista J, Diesmos AC, Clores MA, Cuesta MA, Sanguila MB, Brown RM (2021b) Multivariate characterization of phenotypic variation from throughout the geographic ranges of Philippine false coral snakes: Two species or four? *Ichthyology & Herpetology* 109(4): 1036–1046. <https://doi.org/10.1643/h2021035>
- Bittel J (2015) [2 June] “Extinct” amphibians rediscovered after nearly half a century. *National Geographic*. <https://www.nationalgeographic.com/animals/article/150602-amphibians-extinct-species-toads-philippines-animals>
- Blackburn DC, Bickford DP, Diesmos AC, Iskandar DT, Brown RM (2010) An ancient origin for the enigmatic flat-headed frogs (Bombinatoridae: *Barbourula*) from the islands of Southeast Asia. *PLoS ONE* 5(8): e12090. <https://doi.org/10.1371/journal.pone.0012090>
- Blackburn DC, Siler CD, Diesmos AC, McGuire JA, Cannatella DC, Brown RM (2013) An adaptive radiation of frogs in a Southeast Asian island archipelago. *Evolution; International Journal of Organic Evolution* 67(9): 2631–2646. <https://doi.org/10.1111/evo.12145>
- BMB-DENR [Biodiversity Management Bureau-Department of Environment and Natural Resources] (2016) Philippine Biodiversity Strategy and Action Plan (2015–2028): Bringing Resilience to Filipino Communities. C. Cabrido (Ed.). Quezon City, Philippines: BMB-DENR, United Nations Development Programme – Global Environment Facility, Foundation for the Philippine Environment.
- Böhm M, Collen B, Baillie JEM, Chanson J, Cox N, Hammerson G, Hoffmann M, Livingstone SM, Ram M, Rhodin AGJ, Stuart SN, van Dijk PPL, Young BE, Aftuang LE, Aghasyan A, Aguayo AG, Aguilar C, Ajtic R, Akarsu F, Alencar LRV, Allison A, Ananjeva N, Anderson S, Andrés C, Ariano-Sánchez D, Arredondo JC, Auliya M, Austin CC, Avci A, Baker PJ, Barreto-Lima AF, Barrio-Amorós CL, Basu D, Bates MF, Batistella A, Bauer A, Bennett D, Böhme W, Broadley D, Brown RM, Burgess J, Captain A, Carreira S, Castaneda M, Castro F, Catenazzi A, Cedeño-Vázquez JR, Chapple D, Cheylan M, Cisneros-Heredia

- DF, Cogalniceanu D, Cogger H, Corti C, Costa GC, Couper PJ, Courtney T, Crnobrnja-Isailovic J, Crochet P-A, Crother B, Cruz F, Daltry F, Daniels RJR, Das I, de Silva A, Dirksen L, Doan T, Dodd K, Doody JS, Dorcas ME, Duarte de Barros Filho J, Egan VT, El Mouden EH, Embert D, Espinoza RE, Fallabrino A, Feng X, Feng Z-J, Fitzgerald L, Flores-Villela O, França FGR, Frost D, Gadsden H, Gamble T, Ganesh SR, Garcia MA, García-Pérez JE, Gatus J, Gaulke M, Geniez P, Georges A, Gerlach J, Goldberg S, Gonzalez JCT, Gower DJ, Grant T, Greenbaum E, Guo P, Haitao S, Hamilton AM, Hare K, Hedges B, Heideman N, Hilton-Taylor C, Hitchmough R, Hollingsworth B, Hutchinson M, Ineich I, Iverson J, Jaksic FM, Jenkins R, Joger U, Jose R, Kaska Y, Keogh JS, Köhler G, Kuchling G, Kumlutas Y, Kwet A, La Marca E, Lamar W, Lane A, Lardner B, Latta C, Latta G, Lau M, Lavin P, Lawson D, LeBreton M, Lehr E, Limpus D, Lipczynski N, Lobo AS, López-Luna MA, Luiselli L, Lukoschek V, Lundberg M, Lymberakis P, Macey R, Magnusson WE, Mahler L, Malhotra A, Mariaux J, Maritz B, Marques OAV, Márquez R, Martins M, Masterson G, Mateo JA, Mathew R, Mathews N, Mayer G, McCranie JR, Measey J, Mendoza-Quijano F, Menegon M, Métrailler S, Milton DA, Montgomery C, Morato SA, Mott T, Muñoz-Alonso A, Murphy J, Nguyen TQ, Nilson G, Nogueira C, Núñez H, Ota H, Ottenwalder J, Papenfuss T, Pasachnik S, Passos P, Pauwels OSG, Pérez Mellado V, Pérez-Buitrago N, Pianka ER, Pleguezuelos J, Pollock C, Ponce-Campos P, Powell R, Pupin F, Quintero Díaz GE, Radder R, Ramer J, Rasmussen AR, Raxworthy C, Reynolds R, Richman N, Rico EL, Riservato E, Rivas G, Rocha PLB, Rödel M-O, Rodríguez Schettino L, Roosenburg WM, Ross JP, Sadek R, Sanders K, Santos-Barrera G, Schleich HH, Schmidt B, Schmitz A, Sharifi M, Shea G, Shine R, Slimani T, Somaweera R, Spawls S, Stafford P, Stuebing R, Sweet S, Sy E, Temple H, Tognielli M, Tolley K, Tolson PJ, Tuniyev B, Tuniyev S, Üzümlü N, van Buurt G, Van Sluys M, Velasco A, Vences M, Veselý M, Vinke S, Vinke T, Vogel G, Vogrin M, Vogt RC, Wearn OR, Werner YL, Whiting MJ, Wiewandt T, Wilkinson J, Wilson B, Wren S, Zamin T, Zhou K, Zug G (2013) The conservation status of the world's reptiles. *Biological Conservation* 157: 372–385. <https://doi.org/10.1016/j.biocon.2012.07.015>
- Brown RM (2006) The use of genetic data for inferring species limits. In: Catibog-Sinha CS, Heaney LR (Eds) *Philippine Biodiversity: Principles and Practice*. Haribon Foundation for Conservation of Natural Resources, Quezon City.
- Brown RM (2007) Introduction to Robert F. Inger's *Systematics and Zoogeography of Philippine Amphibia*. Invited forward to the reprint of Inger's 1954 monograph. In: *Systematics and Zoogeography of Philippine Amphibia*. Natural History Publications, Kota Kinabalu.
- Brown RM (2015) A new species of stream frog (genus *Hylarana*) from the mountains of southern Mindanao Island, Philippines. *Herpetologica* 71(3): 223–233. <https://doi.org/10.1655/Herpetologica-D-14-00075>
- Brown RM (2016) Biogeography of Land Vertebrates In: Kliman RM (Ed.) *The Encyclopedia of Evolutionary Biology*. Oxford: Academic Press/Elsevier Inc., Vol. 1, 211–220. <https://doi.org/10.1016/B978-0-12-800049-6.00123-2>
- Brown WC, Alcala AC (1955) Observations on amphibians of Mount Halcon and Mount Canlaon areas, Philippine Islands. *Silliman Journal* 2: 93–105.
- Brown WC, Alcala AC (1961) Populations of amphibians and reptiles in submontane and montane forests of Cuernos de Negros, Philippine Islands. *Ecology* 42(4): 628–636. <https://doi.org/10.2307/1933494>
- Brown WC, Alcala AC (1963) Relationships of the herpetofauna of the non-dipterocarp communities to that of the dipterocarp forest of southern Negros Island, Philippines. *Senckenbergiana Biologica* 45: 591–611.

- Brown WC, Alcala AC (1970a) The zoogeography of the Philippine Islands, a fringing archipelago. *Proceedings of the California Academy of Sciences* 38: 105–130.
- Brown WC, Alcala AC (1970b) A new species of the genus *Platymantis* (Ranidae) with a list of the amphibians known from South Gigante Island, Philippines. *Occasional Papers of the California Academy of Science* 84: 1–7.
- Brown WC, Alcala AC (1974) A new frog of the genus *Platymantis* (Ranidae) from the Philippines. *Occasional Papers of the California Academy of Sciences* 113: 1–12. <https://doi.org/10.5962/bhl.part.2818>
- Brown WC, Alcala AC (1978) Philippine Lizards of the Family Gekkonidae. Silliman University Press, Dumaguete City, Philippines, 146 pp.
- Brown WC, Alcala AC (1980) Philippine Lizards of the Family Scincidae. Silliman University Press, Dumaguete City, Philippines, 264 pp.
- Brown WC, Alcala AC (1982a) A new cave *Platymantis* (Amphibia: Ranidae) from the Philippine Islands. *Proceedings of the Biological Society of Washington* 95: 386–391.
- Brown WC, Alcala AC (1982b) Modes of reproduction of Philippine anurans. In: Rodin AGJ, Miyata K (Eds) *Advances in Herpetology and Evolutionary Biology*. Museum of Comparative Biology, Cambridge, MA, USA, 416–428.
- Brown WC, Alcala AC (1986) Comparison of the herpetofaunal species richness on Negros and Cebu Islands, Philippines. *Silliman Journal* 33: 74–86.
- Brown WC, Alcala AC (1994) Philippine frogs of the family Rhacophoridae. *Proceedings of the California Academy of Sciences* 48: 185–220.
- Brown RM, Alcala AC (2000) Geckos, cave frogs, and small land-bridge islands in the Visayan sea. *Haring Ibon* 2: 19–22.
- Brown RM, Diesmos AC (2002) Application of lineage-based species concepts to oceanic island frog populations: The effects of differing taxonomic philosophies on the estimation of Philippine biodiversity. *Silliman Journal* 42: 133–162.
- Brown RN, Diesmos AC (2009) Philippines, Biology. In: Gillespie R, Clague D (Eds) *Encyclopedia of Islands*. University of California Press, Berkeley, 723–732. <https://doi.org/10.1525/9780520943728-170>
- Brown RM, Guttman SI (2002) Phylogenetic systematics of the *Rana signata* complex of Philippine and Bornean stream frogs: Reconsideration of Huxley's modification of Wallace's Line at the Oriental–Australian faunal zone interface. *Biological Journal of the Linnean Society*. Linnean Society of London 76(3): 393–461. <https://doi.org/10.1111/j.1095-8312.2002.tb01704.x>
- Brown WC, Rabor DS (1967) Review of the genus *Brachymeles* (Scincidae), with descriptions of new species and subspecies. *Proceedings of the California Academy of Sciences* 15: 525–548.
- Brown RM, Siler CD (2013) Spotted stream frog diversification at the Australasian faunal zone interface, mainland versus island comparisons, and a test of the Philippine 'dual-umbilicus' hypothesis. *Journal of Biogeography* 41(1): 182–195. <https://doi.org/10.1111/jbi.12192>
- Brown RM, Stuart BL (2012) Patterns of biodiversity discovery through time: an historical analysis of amphibian species discoveries in the Southeast Asian mainland and island archipelagos. In: Gower DJ, Johnson KG, Richardson JE, Rosen BR, Rüber L, Williams ST (Eds) *Biotic Evolution and Environmental Change in Southeast Asia*. Cambridge University Press, 348–389. <https://doi.org/10.1017/CBO9780511735882.016>
- Brown RM, Ferner JW, Sison RV, Gonzales PC, Kennedy RS (1996) Amphibians and reptiles of the Zambales Mountains of Luzon Island, Republic of the Philippines. *Herpetological Natural History* 4(1): 1–22.

- Brown WC, Brown RM, Alcala AC (1997) Species of the *hazela* group of *Platymantis* from the Philippines with descriptions of two new species. *Proceedings of the California Academy of Sciences* 49: 405–421.
- Brown RM, McGuire JA, Ferner JW, Diesmos AC (1999a) A new species of diminutive scincid lizard (Squamata; Lygosominae; *Sphenomorphus*) from Luzon Island, Republic of the Philippines. *Copeia* 1999(2): 362–370. <https://doi.org/10.2307/1447481>
- Brown RM, Leviton AE, Sison RV (1999b) Description of a new species of *Pseudorabdion* (Serpentes: Colubridae) from Panay Island, Philippines with a revised key to the genus. *Asiatic Herpetological Research* 8: 7–12. <https://doi.org/10.5962/bhl.part.24610>
- Brown RM, McGuire JA, Ferner JW, Icarangal Jr N, Kennedy RS (2000a) Amphibians and reptiles of Luzon Island, II: Preliminary report on the herpetofauna of Aurora Memorial National Park, Philippines. *Hamadryad* 25: 175–195.
- Brown RM, McGuire JA, Diesmos AC (2000b) Status of some Philippine frogs related to *Rana everetti* (Anura: Ranidae), description of a new species, and resurrection of *Rana igorota* Taylor 1922. *Herpetologica* 56: 81–104. <https://www.jstor.org/stable/3893130>
- Brown RM, Diesmos AC, Alcala AC (2002a) The state of Philippine herpetology and the challenges for the next decade. *Silliman Journal* 42(1): 18–87.
- Brown RM, Fernandez R, Rivero C, Buenviaje R, Diesmos AC (2002b) Mt. Isarog's herpetological wonders. *Haring Ibon* 3: 12–16.
- Brown RM, Dolino CN, Alcala E, Diesmos AC, Alcala AC (2003) The advertisement calls of two endangered species of endemic Philippine frogs: *Platymantis spelaeus* and *P. insulatus* (Anura; Ranidae). *Silliman Journal* 43: 91–109.
- Brown RM, Diesmos AC, Duya MV (2007) A new species of *Luperosaurus* (Squamata: Gekkonidae) from the Sierra Madre mountain range of northern Luzon Island, Philippines. *The Raffles Bulletin of Zoology* 55: 153–160.
- Brown RM, Siler CD, Diesmos AC, Alcala AC (2009a) The Philippine frogs of the genus *Leptobrachium* (Anura; Megophryidae): Phylogeny-based species delimitation, taxonomic revision, and descriptions of three new species. *Herpetological Monograph* 23(1): 1–44. <https://doi.org/10.1655/09-037.1>
- Brown RM, Oliveros C, Siler CD, Diesmos AC (2009b) Phylogeny of *Gekko* from the northern Philippines, and description of a new species from Calayan Island. *Journal of Herpetology* 43(4): 620–635. <https://doi.org/10.1670/08-207.1>
- Brown RM, Diesmos AC, Duya MV, Garcia HJD, Rico EL (2010a) A new forest gecko (Squamata; Gekkonidae; Genus *Luperosaurus*) from Mt. Mantalingajan, southern Palawan Island, Philippines. *Journal of Herpetology* 44(1): 37–48. <https://doi.org/10.1670/08-316.1>
- Brown RM, Linkem CW, Siler CD, Sukumaran J, Esselstyn JA, Diesmos AC, Iskandar DT, Bickford D, Evans BJ, McGuire JA, Grismer L, Supriatna J, Andayani N (2010b) Phylogeography and historical demography of *Polypedates leucomystax* in the islands of Indonesia and the Philippines: Evidence for recent human-mediated range expansion? *Molecular Phylogenetics and Evolution* 57(2): 598–619. <https://doi.org/10.1016/j.ympev.2010.06.015>
- Brown RM, Linkem CW, Balete D, Duya MV, Diesmos AC, Ferner JW (2010c) Species boundaries in Philippine montane forest skinks (Genus *Sphenomorphus*): Three new species from the mountains of Luzon and clarification of the status of the poorly known *S. beyeri*, *S. knollmanae*, and *S. laterimaculatus*. *Scientific Papers of the Natural History Museum of the University of Kansas* 42: 1–27.

- Brown RM, Diesmos AC, Oliveros C (2011a) A new flap-legged forest gecko (Genus *Luperosaurus*) from the northern Philippines. *Journal of Herpetology* 45(2): 202–210. <https://doi.org/10.1670/10-123.1>
- Brown RM, Siler CD, Oliveros CH, Diesmos AC, Alcalá AC (2011b) A new *Gekko* from Sibuyan Island, central Philippines. *Herpetologica* 67(4): 460–476. <https://doi.org/10.1655/HERPETOLOGICA-D-11-00025.1>
- Brown RM, Diesmos AC, Sanguila MB, Siler CD, Diesmos MLL, Alcalá AC (2012a) Amphibian Conservation in the Philippines. *FrogLog* 20(5): 40–43.
- Brown RM, Oliveros CH, Siler CD, Fernandez JB, Welton LJ, Buenavente PC, Diesmos MLD, Diesmos AC (2012b) Amphibians and Reptiles of Luzon Island (Philippines), VII: Herpetofauna of Ilocos Norte Province, Northern Cordillera Mountain Range. *Check List* 8(3): 469–490. <https://doi.org/10.15560/8.3.469>
- Brown RM, Siler CD, Oliveros CH, Esselstyn JA, Diesmos AC, Hosner PA, Linkem CW, Barley AJ, Oaks JR, Sanguila MB, Welton LJ, Blackburn DC, Moyle RG, Peterson AT, Alcalá AC (2013a) Evolutionary processes of diversification in a model island archipelago. *Annual Review of Ecology, Evolution, and Systematics* 44(1): 411–435. <https://doi.org/10.1146/annurev-ecolsys-110411-160323>
- Brown RM, Siler CD, Oliveros CH, Welton LJ, Rock A, Swab J, van Weerd M, Rodriguez JE, Diesmos AC (2013b) The amphibians and reptiles of Luzon Island, Philippines, VIII: The herpetofauna of Cagayan and Isabela Provinces, northern Sierra Madre Mountain Range. *ZooKeys* 266: 1–120. <https://doi.org/10.3897/zookeys.266.3982>
- Brown RM, Siler CD, Richards S, Diesmos AC, Cannatella DC (2015a) Multilocus phylogeny and a new classification for Southeast Asian and Melanesian forest frogs (family Ceratobatrachidae). *Zoological Journal of the Linnean Society* 174(1): 130–168. <https://doi.org/10.1111/zoj.12232>
- Brown RM, de Layola AL, Lorenzo All, Diesmos MLL, Diesmos AC (2015b) A new species of limestone karst inhabiting forest frog, genus *Platymantis* (Amphibia: Anura: Ceratobatrachidae: subgenus *Lupacolus*) from southern Luzon Island, Philippines. *Zootaxa* 4048(2): 191–210. <https://doi.org/10.11646/zootaxa.4048.2.3>
- Brown RM, Su Y-C, Barger B, Siler CD, Sanguila MB, Diesmos AC, Blackburn DC (2016) Phylogeny of the island archipelago frog genus *Sanguirana*: Another endemic Philippine radiation that diversified ‘Out-of-Palawan’. *Molecular Phylogenetics and Evolution* 94: 531–536. <https://doi.org/10.1016/j.ympev.2015.10.010>
- Brown RM, Prue A, Chan KO, Gaulke M, Sanguila MB, Siler CD (2017) Taxonomic reappraisal of the northeast Mindanao Stream Frog, *Sanguirana albotuberculata* (Inger 1954), validation of *Rana mearnsi* Stejneger 1905, and description of a new species from the central Philippines. *Herpetological Monograph* 31(1): 183–203. <https://doi.org/10.1655/HERPMONOGRAPHS-D-16-00009.1>
- Brown RM, Smart U, Leviton AE, Smith EN (2018) A new species of long-glanded coral-snake of the genus *Calliophis* (Squamata: Elapidae) from Dinagat Island, with notes on the biogeography and species diversity Philippine *Calliophis* and *Hemibungarus*. *Herpetologica* 74(1): 89–104. <https://doi.org/10.1655/Herpetologica-D-17-00008>
- Brown RM, Meneses CG, Wood Jr PS, Fernandez JB, Cuesta MA, Clores MA, Tracy C, Buehler M, Siler CD (2020) Unexpected discovery of another new species of Philippine False Gecko (Gekkonidae; *Pseudogekko*) from the Bicol Peninsula of Luzon Island. *Herpetologica* 76(3): 315–329. <https://doi.org/10.1655/Herpetologica-D-19-00029.1>
- Brown JC, Shirley MH, Yog-yog A, van Weerd M, Balbas MG, Tarun BA, Siler CD (2021a) Use of diet and body condition assessments as intermediate indicators of translocation

- success in the Critically Endangered Philippine crocodile (*Crocodylus mindorensis*). Aquatic Conservation 31(10): 2817–2829. <https://doi.org/10.1002/aqc.3700>
- Brown RM, Calvo-Revuelta M, Goyes Vallejos J, Sanchez-Vialas A, De La Riva I (2021b) On the second (or, rather, the first) specimen of the recently described *Calliophis salitan* (Squamata; Elapidae), with the first report of the species from Mindanao Island, southern Philippines. Herpetology Notes 14: 1027–1035.
- Brown RM, Bautista JB, Clores MA, Cuesta MA (2021c) *Dendrelaphis marenae* (Gaulke's Bronze-back Tree Snake): Diet. Herpetological Review 52: 153–154.
- Brown RM, Bernstein J, Fernandez JB, Sanguila MB, Meneses CG, Siler CD, Chin K, Welton LJ, Gaulke M (2022) Juvenile colouration, ontogenetic colour shifts, defensive behaviour, and frontal “hooding” threat display in Philippine false coral snakes (Squamata: Elapidae: Hemibungarus). Herpetology Notes 15: 183–192.
- Bucol AA, Alcala AC, Averia LT, Alcala EL, Alcala MLR (2011) Checklist of the herpetofauna of Siquijor Island, Philippines. Philippine Scientist 48: 100–122.
- Bucol LA, Bucol AA, Arche NC (2019) Abundance and microhabitat use by the frog *Platymantis hazelae* (Anura: Ceratobatrachidae). Revista de Biología Tropical 67: 518–533.
- Byrne AQ, Vredenburg VT, Martel A, Pasmans F, Bell RC, Blackburn DC, Bletz MC, Bosch J, Briggs CJ, Brown RM, Figueroa-Valenzuela R, Ghose SL, Jani AJ, Jaeger JR, Jirku M, Knapp RA, Portik DM, Richards-Zawacki CL, Rockney H, Stark T, Sulaeman H, Thien Thao N, Voyles J, Waddle AW, Yuan Z-Y, Rosenblum EB (2019) Cryptic diversity of a widespread global pathogen reveals new threats for amphibian conservation. Proceedings of the National Academy of Sciences of the United States of America 116(41): 20382–20387. <https://doi.org/10.1073/pnas.1908289116>
- Camaiti M, Evans AR, Hipsley CA, Meiri S, Hutchinson MN, Anderson RO, Slavenko A, Chapple DG (2022) A database of the morphology, ecology and literature of the world's limb-reduced skinks. Journal of Biogeography 49(7): 1397–1406. <https://doi.org/10.1111/jbi.14392>
- Camaiti M, Evans AR, Hipsley CA, Hutchinson MN, Meiri S, Anderson RO, Slavenko A, Chapple DG (2023) Macroecological and biogeographical patterns of limb reduction in the world's skinks. Journal of Biogeography 50(2): 428–440. <https://doi.org/10.1111/jbi.14547>
- Carter JK, Kimball RT, Funk ER, Kane NC, Schield DR, Spellman GM, Safran RJ (2023) Estimating phylogenies from genomes: A beginner's review of commonly used genomic data in vertebrate phylogenomics. The Journal of Heredity 114(1): 1–13. <https://doi.org/10.1093/jhered/esac061>
- Chan KO, Brown RM (2017) Did True Frogs ‘dispersify’? Biology Letters 13(20179299): 1–6. <https://doi.org/10.1098/rsbl.2017.0299>
- Chan KO, Schoppe S, Rico ELB, Brown RM (2021) Molecular systematic investigation of Philippine puddle frogs (Anura: Dicroglossidae: *Occidozyga* Kuhl and Van Hasselt) reveals new candidate species and a novel pattern of species dyads. Philippine Journal of Systematic Biology 14(2): 1–14. <https://doi.org/10.26757/pjsb2020b14007>
- Chan KO, Hutter CR, Wood Jr PL, Su Y-C, Brown RM (2022) Gene flow increases phylogenetic structure and inflates cryptic species estimations: A case study on widespread Philippine puddle frogs (*Occidozyga laevis*). Systematic Biology 71(1): 40–57. <https://doi.org/10.1093/sysbio/syab034>
- Chapple DG, Roll U, Böhm M, Aguilar R, Amey AP, Austin CC, Baling M, Barley AJ, Bates MF, Bauer AM, Blackburn DG, Bowles P, Brown RM, Chandramouli SR, Chirio L, Cogger H, Colli GR, Conradie W, Coupe PJ, Cowan MA, Craig MD, Das I, Datta-Roy A, Dickman CR, Ellis RJ, Fenner AL, Ford S, Ganesh SR, Gardner MG, Geissler P, Gillespie GR,

- Glaw F, Greenlees MJ, Griffith OW, Grismer LL, Haines ML, Harris DJ, Hedges SB, Hitchmough RA, Hoskin CJ, Hutchinson MN, Ineich I, Janssen J, Johnston GR, Karin BR, Keogh JS, Kraus F, LeBreton M, Lymberakis P, Masroor R, McDonald PJ, Mecke S, Melville J, Melzer S, Michael DR, Miralles A, Mitchell NJ, Nelson NJ, Nguyen TQ, de Campos Nogueira C, Ota H, Pafilis P, Pauwels OSG, Perera A, Pincheira-Donoso D, Reed RN, Ribeiro-Júnior MA, Riley JL, Rocha S, Rutherford PL, Sadlier RA, Shacham B, Shea GM, Shine R, Slavenko A, Stow A, Sumner J, Tallwin OJS, Teale R, Torres-Carvajal O, Francois Trape J, Uetz P, Ukuwela KDB, Valentine L, Van Dyke JU, van Winkel D, Vasconcelos R, Vences M, Wagner P, Wapstra E, While GM, Whiting MJ, Whittington CM, Wilson S, Ziegler T, Tingley R, Meiri S (2021) Conservation status of the world's skinks (Scincidae): Taxonomic and geographic patterns in extinction risk. *Biological Conservation* 257: 1–12. <https://doi.org/10.1016/j.biocon.2021.109101>
- Clores MA, Bautista JB, Fernandez JB, Cuesta MA, Brown RM (2021) Diversity and distribution of amphibians and reptiles in the Caramoan Island Group, Maqueda Channel, Southern Luzon, Philippines. *Journal of Asia-Pacific Biodiversity* 14(1): 1–14. <https://doi.org/10.1016/j.japb.2020.11.005>
- Cox N, Young BE, Bowles P, Fernandez M, Marin J, Rapacciuolo G, Böhm M, Brooks TM, Hedges SB, Hilton-Taylor C, Hoffmann M, Jenkins RKB, Tognelli MF, Alexander GJ, Allison A, Ananjeva NB, Auliya M, Avila LJ, Chapple DG, Cisneros-Heredia DF, Cogger HG, Colli GR, de Silva A, Eisemberg CC, Els J, Fong AG, Grant TD, Hitchmough RA, Iskandar DT, Kidera N, Martins M, Meiri S, Mitchell NJ, Molur S, Nogueira CC, Ortiz JC, Penner J, Rhodin AGJ, Rivas G, Rödel M-O, Roll U, Sanders KL, Santos-Barrera G, Shea GM, Spawls S, Stuart BL, Tolley KA, Trape J-F, Vidal MA, Wagner P, Wallace BP, Xie Y (2022) Global reptile assessment shows commonality of tetrapod conservation needs. *Nature* 605: 285–290. <https://doi.org/10.1038/s41586-022-04664-7>
- Cruz RAL, Lagunzad CGB (2021) The big picture: Consolidating national government and CITES records of animal trade in the Philippines from 1975 to 2019. *Philippine Science Letters* 14(1): 79–100.
- Custodio CC (1986) Altitudinal distribution of lizards of the Scincidae in Mt. Makiling, Laguna. *Sylvatrop: the Technical Journal of Philippine Ecosystems and Natural Resources* 1: 181–202.
- Das S, Greenbaum E, Meiri S, Bauer AM, Burbrink FT, Raxworthy CJ, Weinell JL, Brown RM, Brecko J, Olivier PM, Pauwels SG, Merilä J (2023) Ultraconserved elements-based phylogenomic systematics of snake superfamily Elapoidea, with a description of a new Afro-Asian family. *Molecular Phylogenetics and Evolution* 180: 107700. <https://doi.org/10.1016/j.ympev.2022.107700>
- Davis DR, Feller KD, Brown RM, Siler CD (2014) Evaluating the diversity of Philippine Slender Skinks of the *Brachymeles bonitae* Complex (Reptilia: Squamata: Scincidae): Redescription of *B. tridactylus* and descriptions of two new species. *Journal of Herpetology* 48(4): 480–494. <https://doi.org/10.1670/13-173>
- Davis DR, Watters JL, Köhler G, Whitsett C, Huron NA, Brown RM, Diesmos AC, Siler CD (2015) Redescription of the rare Philippine false gecko *Pseudogekko brevipes* (Reptilia: Squamata: Gekkonidae) and description of a new species. *Zootaxa* 4020(2): 357–374. <https://doi.org/10.11646/zootaxa.4020.2.7>
- Davis DR, Chan KO, Das I, Brennan IG, Karin BR, Jackman TR, Brown RM, Iskandar DT, Nashriq I, Grismer LL, Bauer AM (2020) Multi-locus phylogeny of Bornean Bent-Toed Geckos (Gekkonidae: *Cyrtodactylus*) reveals hidden diversity, taxonomic disarray, and novel biogeographic patterns. *Molecular Phylogenetics and Evolution* 147: 106785. <https://doi.org/10.1016/j.ympev.2020.106785>

- Decena SCP, Avorque CA, Decena ICP, Asis PD, Pacle B (2020) Impact of habitat alteration on amphibian diversity and species composition in a lowland tropical rainforest in Northeastern Leyte, Philippines. *Scientific Reports* 10(1): 1–15. <https://doi.org/10.1038/s41598-020-67512-6>
- Devan-Song A, Brown RM (2012) Amphibians and Reptiles of Luzon Island, Philippines, VI: The Herpetofauna of the Subic Bay Area. *Asian Herpetological Research* 3(1): 1–20. <https://doi.org/10.3724/SP.J.1245.2012.00001>
- Diesmos AC (2012) The Status of Philippine Caecilians (Amphibia: Ichthyophiidae). *FrogLog*, 46 pp.
- Diesmos AC, Watters JL, Huron NA, Davis DR, Alcala AC, Crombie RI, Afuang LE, Gee-Das G, Sison RV, Sanguila MB, Penrod ML, Labonte MJ, Davey CS, Leone EA, Diesmos ML, Sy EY, Welton LJ, Brown RM, Siler CD (2015) Amphibians of the Philippines, Part I: Checklist of the Species. *Proceedings of the California Academy of Sciences* 62: 451–531.
- Diesmos AC, Brown RM (2011) Diversity, Biogeography, and Conservation of Philippine Amphibians. In: *Biology and Conservation of Tropical Asian Amphibians. Proceedings of the Conference “Biology of the Amphibians in the Sunda Region, South-east Asia.”* In: Das I, Haas A, Tuen AA (Eds) Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak, Malaysia, 26–49.
- Diesmos AC, Brown RM, Alcala AC (2002) A new species of narrow-mouthed frog (Amphibia; Anura; Microhylidae: Genus *Kaloula*) from the mountains of southern Luzon and Polillo islands, Philippines. *Copeia* 2002(4): 1037–1051. [https://doi.org/10.1643/0045-8511\(2002\)002\[1037:NSONMF\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2002)002[1037:NSONMF]2.0.CO;2)
- Diesmos AC, Brown RM, Gee GVA (2004a) Preliminary report on the amphibians and reptiles of Balbalasang-Balbalan National Park, Luzon Island, Philippines. *Sylvatrop. Technical Journal of Philippine Ecosystems and Natural Resources* 13: 63–80.
- Diesmos AC, Gee GVA, Brown RM, Widmann PJ, Dimalibot JC (2004b) Rediscovery of the Philippine forest turtle, *Heosemys leytensis* (Chelonia; Bataguridae), from Palawan Island, Philippines. *Asiatic Herpetological Research* 10: 22–27.
- Diesmos AC, Diesmos ML, Brown RM (2006) Status and distribution of alien invasive frogs in the Philippines. *Journal of Environmental Science and Management, Philippines* 9: 41–53.
- Diesmos MLD, Diesmos AC, Siler CD, Vredenburg VT, Brown RM (2012) Detecting the distribution of the chytrid fungus in the Philippines. *FrogLog* 104: 48–49.
- Diesmos AC, Alcala AC, Siler CD, Brown RM (2014) Status and conservation of Philippine amphibians. In: Heatwole H, Das I (Eds) *Conservation Biology of Amphibians of Asia. Status and Decline of Amphibians: Eastern Hemisphere*. Natural History Publications, Malaysia, 310–336.
- Diesmos AC, Watters JL, Huron NA, Davis DR, Alcala AC, Crombie RI, Afuang LE, Gee-Das G, Sison RV, Sanguila MB, Labonte MJ, Davey CS, Penrod ML, Leone EA, Diesmos ML, Sy EY, Welton LJ, Brown RM, Siler CD (2015) Amphibians of the Philippines, Part I: Checklist of the Species. *Proceedings of the California Academy of Sciences* 62(20): 457–539.
- Diesmos AC, Scheffers BR, Mallari NAD, Siler CD, Brown RM (2020) A new forest frog of the genus *Platymantis* (Amphibia: Anura: Ceratobatrachidae: subgenus *Tirahanulap*) from Leyte and Samar islands, eastern Philippines. *Zootaxa* 4830(3): 573–591. <https://doi.org/10.11646/zootaxa.4830.3.6>
- Eliades S, Brown RM, Huang W-S, Siler CD (2021) Taxonomic revision of scaly-toed geckos (Reptilia: Gekkonidae: *Lepidodactylus*) in the northern Philippines, with descriptions of four new species. *Herpetological Monograph* 35(1): 90–111. <https://doi.org/10.1655/HERPMONOGRAPHS-D-19-00014>

- Esselstyn JA, Oliveros CH, Moyle RG, Peterson AT, McGuire JC, Brown RM (2010) Integrating phylogenetic and taxonomic evidence illuminates complex biogeographic patterns along Huxley's modification of Wallace's Line. *Journal of Biogeography* 37(11): 2054–2066. <https://doi.org/10.1111/j.1365-2699.2010.02378.x>
- Evans BJ, Brown RM, McGuire JA, Supriatna J, Andayani N, Diesmos AC, Iskandar D, Melnick DJ, Cannatella DC (2003) Phylogenetics of fanged frogs: Testing biogeographical hypotheses at the interface of the Asian and Australian faunal zones. *Systematic Biology* 52(6): 794–819. <https://doi.org/10.1080/10635150390251063>
- Ferner JW, Brown RM, Greer AE (1997) A new genus and species of closed canopy forest skinks from the Philippine islands. *Journal of Herpetology* 31(2): 187–192. <https://doi.org/10.2307/1565386>
- Ferner JW, Brown RM, Sison RV, Kennedy RS (2001) The amphibians and reptiles of Panay Island, Philippines. *Asiatic Herpetological Research* 9: 34–70. <https://doi.org/10.5962/bhl.part.15561>
- Flores ABA, Diesmos AC, Nuñez OM (2023) Morphological description and life history of the Philippine endemic *Limnonectes leytensis* Leyte swamp frog in Rogongon, Iligan City, Philippines. *Biodiversitas (Surakarta)* 24(1). <https://doi.org/10.13057/biodiv/d240117>
- Flury J, Haas A, Brown RM, Das I, Min PY, Boon-Hee K, Scheidt U, Iskandar DT, Jankowski A, Hertwig ST (2021) Unexpectedly high levels of lineage diversity in Sundaland Puddle Frogs (Dicroglossidae: *Occidozyga* Kuhl and Van Hasselt, 1822). *Molecular Phylogenetics and Evolution* 163: 107210. <https://doi.org/10.1016/j.ympev.2021.107210>
- Formenti G, Theissinger K, Fernandes C, Bista I, Bombarely A, Bleidorn C, Ciofi C, Crottini A, Godoy JA, Höglund J, Malukiewicz J, Mouton A, Oomen RA, Paez S, Palsbøll PJ, Pampoulie C, Ruiz-Lopez MJ, Svardal H, Theofanopoulou C, de Vries J, Waldvogel A-M, Zhang G, Mazzoni CJ, Jarvis ED, Bálint M, Formenti G, Theissinger K, Fernandes C, Bista I, Bombarely A, Bleidorn C, Čiampor F, Ciofi C, Crottini A, Godoy JA, Höglund J, Malukiewicz J, Mouton A, Oomen RA, Paez S, Palsbøll P, Pampoulie C, Ruiz-López MJ, Svardal H, Theofanopoulou C, de Vries J, Waldvogel A-M, Zhang G, Mazzoni CJ, Jarvis E, Bálint M, Aghayan SA, Alioto TS, Almudi I, Alvarez N, Alves PC, Amorim IR, Antunes A, Arribas P, Baldrian P, Berg PR, Bertorelle G, Böhne A, Bonisoli-Alquati A, Boštjančić LL, Boussau B, Breton CM, Buzan E, Campos PF, Carreras C, Castro LFI, Chueca LJ, Conti E, Cook-Deegan R, Croll D, Cunha MV, Delsuc F, Dennis AB, Dimitrov D, Faria R, Favre A, Fedrigo OD, Fernández R, Ficetola GF, Flot J-F, Gabaldón T, Galea Agius DR, Gallo GR, Giani AM, Gilbert MTP, Grebenc T, Guschanski K, Guyot R, Hausdorf B, Hawlitschek O, Heintzman PD, Heinze B, Hiller M, Husemann M, Iannucci A, Irisarri I, Jakobsen KS, Jentoft S, Klinga P, Kloch A, Kratochwil CF, Kusche H, Layton KKS, Leonard JA, Lerat E, Liti G, Manousaki T, Marques-Bonet T, Matos-Maraví P, Matschiner M, Maumus F, McCartney AM, Meiri S, Melo-Ferreira J, Mengual X, Monaghan MT, Montagna M, Mysłajek RW, Neiber MT, Nicolas V, Novo M, Ozretić P, Palero F, Pârvulescu L, Pascual M, Paulo OS, Pavlek M, Pegueroles C, Pellissier L, Pesole G, Primmer CR, Riesgo A, Rüber L, Rubolini D, Salvi D, Seehausen O, Seidel M, Secomandi S, Studer B, Theodoridis S, Thines M, Urban L, Vasemägi A, Vella A, Vella N, Vernes SC, Vernesi C, Vieites DR, Waterhouse RM, Wheat CW, Wörheide G, Wurm Y, Zammit G, European Reference Genome Atlas (ERGA) Consortium (2022) The era of reference genomes in conservation genomics. *Trends in Ecology & Evolution* 2940(3): 197–202. <https://doi.org/10.1016/j.tree.2021.11.008>
- Fuiten A, Diesmos AC, Welton LJ, Barley A, Oberheide B, Rico ELB, Brown RM (2011) New species of stream frog from the mountains of Luzon Island, Philippines. *Herpetologica* 67(1): 89–103. <https://doi.org/10.1655/HERPETOLOGICA-D-10-00042.1>

- Gainsbury A, Meiri S (2017) Latitudinal diversity gradient and interspecific competition: No global relationship between niche breadth and species richness. *Global Ecology and Biogeography* 26(5): 563–572. <https://doi.org/10.1111/geb.12560>
- Garcia VOS, Papa RDS, Briones JCA, Mendoza N, Okuda N, Diesmos AC (2014) Food habits and distribution of the Lake Taal sea snake (*Hydrophis semperi* Garman 1881) and the sympatric Little File Snake (*Acrochordus granulatus* Schneider 1799) in Lake Taal, Philippines. *Asian Herpetological Research* 5(4): 255–262. <https://doi.org/10.3724/SP.J.1245.2014.00255>
- Gaulke M (1993) Beobachtungen an Flugdrachen auf dem Sulu-Archipel. *Salamaandra* 28: 251–257.
- Gaulke M (1994) Contribution to the snake fauna of the Sulu archipelago with the description of a new subspecies of *Dendrelaphis caudolineatus* (Gray, 1834). *The Herpetological Journal* 4: 136–144.
- Gaulke M (1995) Der Sulu-Archipel–Besiedlungsgeschichte, geologie und herpetofauna. *Natur und Museum* 125: 217–226.
- Gaulke M (2002) A new species of *Lycodon* from Panay Island, Philippines (Reptilia, Serpentes, Colubridae). *Spixiana* 25(1): 85–92.
- Gaulke M (2011) The herpetofauna of Panay Island, Philippines: an illustrated field guide. Ed. Chimaira.
- Gojo-Cruz PH, Afuang LE, Gonzalez JCT, Gruezo WS (2018) Amphibians and reptiles of Luzon Island, Philippines: The herpetofauna of Pantabangan-Carranglan watershed, Nueva Ecija Province, Caraballo Mountain range. *Asian Herpetological Research* 9(4): 201–223. <https://doi.org/10.16373/j.cnki.ahr.180050>
- Gojo-Cruz PH, Afuang LE, Gonzalez JCT, Gruezo WS (2019) Distribution and diversity patterns of herpetofauna in the Pantabangan-Carranglan Watershed, Nueva Ecija, Caraballo Mountain Range, Philippines. *Biodiversity Data Journal* 7:e31638. <https://doi.org/10.3897/BDJ.7.E31638>
- Gonzalez JCT, Layusa CA, Afuang LE, Duya MM, Heaney LR, Balete DS, Tabaranza GE, Española CP, Van de Ven WC, Diesmos AC, Causaren RM, Diesmos ML, Lagat RT, Realubit ND, Sy ES, Lit IL, Naredo JC, Lastica-Ternura EA, Pasicolan SA, Tagtag AM, De Leon JL, Lim TM, Ong PS (2018) Review and update of the 2004 National List of Threatened Terrestrial Fauna of the Philippines. *Sylvatrop. The Technical Journal of Philippine Ecosystems and Natural Resources* 28: 73–144.
- González-del-Pliego P, Freckleton RP, Edwards DP, Koo MS, Scheffers BR, Pyron RA, Jetz W (2019) Phylogenetic and trait-based prediction of extinction risk for data-deficient amphibians. *Current Biology* 29(9): 1557–1563. <https://doi.org/10.1016/j.cub.2019.04.005>
- Greenberg DA, Palen WJ, Chan KC, Jetz W, Mooers AO (2018) Evolutionary distinct amphibians are disproportionately lost from human-modified ecosystems. *Ecology Letters* 21(10): 1530–1540. <https://doi.org/10.1111/ele.13133>
- Gumbs R, Gray CL, Böhm MMH, Grenyer R, Jetz W, Meiri S, Roll U, Owen NR, Rosindell J (2020) Global priorities for conservation of reptilian phylogenetic diversity in the face of human impacts. *Nature Communications* 11(1): 2616. <https://doi.org/10.1038/s41467-020-16410-6>
- Heaney LR, Regalado JC (1998) Vanishing treasures of the Philippine rain forest. The Field Museum, Chicago. IL, USA, viii + 88 pp.
- Heaney LR, Diesmos A, Tabaranza B, Mallari A, Brown R, Gee G (2000) Beacon of hope; a first report from Kalinga Province in the northern Central Cordillera. *Haring Ibon* 2: 14–18.

- Heinrich S, Toomes A, Janssen J (2021) Legal or unenforceable? Violations of trade regulations and the case of the Philippine Sailfin Lizard *Hydrosaurus pustulatus* (Reptilia: Squamata: Agamidae). *Journal of Threatened Taxa* 13(6): 18532–18543. <https://doi.org/10.11609/jott.7269.13.6.18532-18543>
- Herr MW, Goyes Vallejos J, Meneses CG, Abraham RK, Otterholt R, Siler CD, Rico EL, Brown RM (2021) A new, morphologically cryptic species of fanged frog, genus *Limnonectes* (Amphibia: Anura: Dicroglossidae), from Mindoro Island, central Philippines. *Ichthyology & Herpetology* 109(1): 188–210. <https://doi.org/10.1643/h2020095>
- Herr MW, Som HE, Brown RM (in press) A long overlooked new species of Fanged Frog, genus *Limnonectes* (Amphibia: Anura: Dicroglossidae), from Luzon Island northern Philippines. *Ichthyology & Herpetology*.
- Hertwig ST, Das I, Schweizer M, Brown RM, Haas A (2011) Phylogenetic relationships of the *Rhacophorus everetti*-group and implications for the evolution of reproductive modes in *Philautus* (Amphibia: Anura: Rhacophoridae). *Zoologica Scripta* 41(1): 29–46. <https://doi.org/10.1111/j.1463-6409.2011.00499.x>
- Hillis DM, Moritz C, Mable BK (1996) *Molecular Systematics*. 2nd edn. Sinauer, Sunderland, Massachusetts 1996(4): 1058–1059. <https://doi.org/10.2307/1447682>
- Hortal J, de Bello F, Diniz-Filho JAF, Lewinsohn TM, Lobo JM, Ladle RJ (2015) Seven shortfalls that beset large-scale knowledge of biodiversity. *Annual Review of Ecology, Evolution, and Systematics* 46(1): 523–549. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>
- Hutter CR, Cobb KA, Portik DM, Travers SL, Wood Jr PL, Brown RM (2022) FrogCap: A modular sequence capture probe-set for phylogenomics and population genetics for all frogs, assessed across multiple phylogenetic scales. *Molecular Ecology Resources* 2021(3): 1110–1119. <https://doi.org/10.1111/1755-0998.13517>
- Inger RF (1954) Systematics and zoogeography of Philippine Amphibia. *Fieldiana* 33: 181–531. <https://doi.org/10.5962/bhl.title.5571>
- IUCN [International Union for Conservation of Nature] (2022) IUCN Red List Categories and Criteria: Version 3.1. 2nd edn. IUCN, Gland, Switzerland and Cambridge, United Kingdom, 32 pp.
- IUCN [International Union for Conservation of Nature] (2023) The IUCN Red List of Threatened Species. Version 2023-1. <https://www.iucnredlist.org> [Accessed on 30 December 2023]
- Khalighifar A, Brown RM, Goyes Vallejos J, Peterson AT (2021) Deep learning improves acoustic biodiversity monitoring and new candidate species identification (genus *Platymantis*) in the Philippines. *Biodiversity and Conservation* 30(3): 643–657. <https://doi.org/10.1007/s10531-020-02107-1>
- Koch A, Gaulke M, Böhme W (2010) Unravelling the underestimated diversity of Philippine water monitor lizards (Squamata: *Varanus salvator* complex), with the description of two new species and a new subspecies. *Zootaxa* 2446(1): 1–54. <https://doi.org/10.11646/zootaxa.2446.1.1>
- Lama JM, Senarillos TLP (2023) Tadpole transport behavior of the Philippine Small-disked Frog, *Limnonectes parvus* (Taylor 1920), from Western Mindanao, Philippines. *Reptiles & Amphibians : Conservation and Natural History* 30(1): e20354–e20354. <https://doi.org/10.17161/randa.v30i1.20354>
- Leviton AE, Brown WC (1958) A review of the snakes of the genus *Pseudorabdion* with remarks on the status of the genera *Agrophis* and *Typhlogeophis* (Serpentes: Colubridae). *Proceedings of the California Academy of Sciences* 14: 475–508.
- Leviton AE (1962) Contributions to a review of Philippine snakes, I. The snakes of the genus *Oligodon*. *Philippine Journal of Science* 91: 365–381.

- Leviton AE (1963a) Remarks on the zoogeography of Philippine terrestrial snakes. *Proceedings of the California Academy of Sciences* 42: 112–145.
- Leviton AE (1963b) Contributions to a review of Philippine Snakes, III. The genera *Maticora* and *Calliophis*. *Philippine Journal of Science* 92: 523–550.
- Leviton AE (1964a) Contributions to a review of Philippine snakes, IV. The genera *Chrysopelea* and *Dryophiops*. *Philippine Journal of Science* 93: 131–145.
- Leviton AE (1964b) Contributions to a review of Philippine snakes, V. The snakes of the genus *Trimeresurus*. *Philippine Journal of Science* 93: 250–2760.
- Leviton AE (1964c) Contributions to a review of Philippine snakes, VI. The snakes of the genus *Oxyrhabdion*. *Philippine Journal of Science* 93: 407–422.
- Leviton AE (1964d) Contributions to a review of Philippine snakes, VII. The snakes of the genera *Naja* and *Ophiophagus*. *Philippine Journal of Science* 93: 530–550.
- Leviton AE (1965a) Contributions to a review of Philippine Snakes, VIII. The snakes of the genus *Lycodon* H. Boie. *Philippine Journal of Science* 94: 117–140.
- Leviton AE (1965b) Contributions to a review of Philippine Snakes, IX. The snakes of the genus *Cyclocorus*. *Philippine Journal of Science* 94: 519–533.
- Leviton AE (1967) Contributions to a review of Philippine Snakes, X. The snakes of the genus *Ahaetulla*. *Philippine Journal of Science* 96: 73–90.
- Leviton AE (1968) Contributions to a review of Philippine snakes, VII. The Philippine snakes of the genus *Dendrelaphis* (Serpentes: Colubridae). *Philippine Journal of Science* 97: 371–396.
- Leviton AE (1979) Contributions to a review of Philippine snakes, XIII. The snakes of the Genus *Elaphe*. *Philippine Journal of Science* 106: 99–128.
- Leviton AE (1983) Contributions to a review of Philippine snakes, XIV. The Snakes of the Genus *Xenopeltis*, *Zaocys*, *Psammodynastes*, and *Myersophis*. *Philippine Journal of Science* 112: 195–223.
- Leviton AE, Brown RM, Siler CD (2014) The dangerously venomous snakes of the Philippine archipelago, with identification keys and species accounts. In: Williams GC, Gosliner GC (Eds) *The Coral Triangle: the 2011 Hearts Philippine Biodiversity Expedition*. California Academy of Sciences, San Francisco, CA, 473–530 pp.
- Leviton AE, Siler CD, Weinell JL, Brown RM (2018) A synopsis of the snakes of the Philippines: A synthesis of data from biodiversity repositories, field studies, and the literature. *Proceedings of the California Academy of Sciences* 64: 399–568.
- Linkem CW, Brown RM (2013) Systematic revision of the *Parvosцинcus decipiens* (Boulenger 1894) complex of Philippine forest skinks (Squamata: Scincidae: Lygosominae) with descriptions of seven new species. *Zootaxa* 3700(4): 501–533. <https://doi.org/10.11646/zootaxa.3700.4.1>
- Linkem CA, Hesed K, Diesmos AC, Brown RM (2010a) Species boundaries and cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata; Scincidae: Lygosominae). *Molecular Phylogenetics and Evolution* 56(2): 572–585. <https://doi.org/10.1016/j.ympev.2010.03.043>
- Linkem CA, Diesmos AC, Brown RM (2010b) A new scincid lizard (Genus *Sphenomorphus*) from Palawan Island, Philippines. *Herpetologica* 66(1): 67–79. <https://doi.org/10.1655/08-074.1>
- Linkem CW, Siler DC, Diesmos AC, Brown RM (2010c) A new species of Gekko (Squamata: Gekkonidae) from central Luzon Island, Philippines. *Zootaxa* 2396(1): 37–94. <https://doi.org/10.11646/zootaxa.2396.1.3>
- Linkem CW, Diesmos AC, Brown RM (2011) Molecular systematics of the Philippine forest skinks (Reptilia: Scincidae: *Sphenomorphus*): testing morphological and

- biogeographic hypotheses of interspecific relationships. *Zoological Journal of the Linnean Society* 163: 1217–1243. <https://doi.org/10.1111/j.1096-3642.2011.00747.x>
- Linkem CW, Brown RM, Siler CD, Evans BJ, Austin CC, Iskandar DT, Diesmos AC, Supriatna J, Andayani N, McGuire JA (2013) Stochastic faunal exchanges drive diversification in widespread Wallacean and Pacific Island lizards (Squamata: Scincidae: *Lamprolepis smaragdina*). *Journal of Biogeography* 40(3): 507–520. <https://doi.org/10.1111/jbi.12022>
- Madarang RC, Uychiaoco AJ, Schirm B, Mallari NAD, Quitariano E (2017) Report on the Management Effectiveness and Capacity of Protected Areas in the Philippines. Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ) GmbH. Philippines: Manila, 78 pp.
- Maglangit EPT, Paraguya JJJB, Maglangit RMT, Nuñez OM, Diesmos MLL, Diesmos AC (2021) Novel cave habitat used by the cryptic lizard *Pinoyscincus abdictus abdictus* (Squamata: Scincidae) on Dinagat Islands, Philippines. *Phyllomedusa* 20(1): 99–104. <https://doi.org/10.11606/issn.2316-9079.v20i1p99-104>
- Maglangit EPT, Nuñez OM, Coritico FP, Medecilo-Guiang MMP, Mohagan AB, Patano Jr RR, Amoroso VB (2022) Richness and distribution of reptiles and amphibians in the tropical lowland habitats of Mt. Agad-Agad, Iligan City, Southern Philippines. *Reptiles & Amphibians : Conservation and Natural History* 29(1): 413–425. <https://doi.org/10.17161/randa.v29i1.18130>
- Mallari NAD, Collar NJ, McGowan PJ, Marsden SJ (2015) Philippine protected areas are not meeting the biodiversity coverage and management effectiveness requirements of Aichi Target 11. *Ambio* 45(3): 313–322. <https://doi.org/10.1007/s13280-015-0740-y>
- Margules CR, Pressey RL (2000) Systematic conservation planning. *Nature* 405(6783): 243–253. <https://doi.org/10.1038/35012251>
- McDiarmid RW, Campbell JA, Touré TA (1999) Snake Species of the World: A Taxonomic and Geographic Reference. The Herpetologists' League, Washington DC, USA, 511 pp.
- McDonald PJ, Brown RM, Kraus F, Bowles P, Arifin U, Eliades SJ, Fisher RN, Gaulke M, Grismer LL, Ineich I, Karin BR, Meneses CG, Richards SR, Sanguila MB, Siler CD, Oliver PM (2022) Cryptic extinction risk in a western Pacific lizard radiation. *Biodiversity and Conservation* 31(8–9): 2045–2062. <https://doi.org/10.1007/s10531-022-02412-x>
- McLeod DS, Siler CD, Diesmos AC, Diesmos MLL, Garcia VS, Arkonco AS, Balaquit KL, Uy CC, Vilaseran MM, Yarra EC, Brown RM (2012) Amphibians and Reptiles of Luzon Island, V: The Herpetofauna of Angat Dam Watershed, Bulacan Province, Luzon Island, Philippines. *Asian Herpetological Research* 2(4): 177–198. <https://doi.org/10.3724/SP.J.1245.2011.00177>
- Meiri S, Avila L, Bauer AM, Chapple DG, Das I, Doan TM, Doughty P, Ellis R, Grismer LL, Kraus F, Morando M, Oliver PM, Pincheira-Donoso D, Ribeiro-Junior MA, Shea G, Torres-Carvajal O, Slavenko A, Roll U (2020) The global diversity and distribution of lizard clutch sizes. *Global Ecology and Biogeography* 29(9): 1515–1530. <https://doi.org/10.1111/geb.13124>
- Meneses CG (2020) Ophiophagy in the Philippine endemic Luzon False Coral Snake, *Hemibungarus calligaster* (Weigmann 1834) on Gervais' Reed Snake *Calamaria gervaisii* Dumeril, Bibron, and Dumeril 1854 and the importance of the Rabor collection in Filipino natural history. *Laksambuhay* 7: 18–21.
- Meneses CG, Siler CD, Gonzalez JCT, Wood Jr PL, Brown RM (2020a) Molecular phylogenetic estimates of evolutionary affinities and the first reports of phenotypic variation in two secretive, endemic reptiles from the Romblon Island Group, central Philippines) from Luzon Island, the Philippines. *Philippine Journal of Systematic Biology* 14(2): 1–20. <https://doi.org/10.26757/pjsb2020b14002>

- Meneses CG, Alviola P, Parcon JA, Cosico EA, Lumibao JD, Gonzalez JCT, Brown RM (2020b) *Varanus bitatawa* (Northern Luzon Golden-spotted Forest Monitor): Geographical Distribution. *Herpetological Review* 51: 779.
- Meneses CG, Siler CD, Alviola PA, Gonzalez JCT, Balatibat JB, Natividad CA, Brown RM (2022) Amphibian and reptile diversity along a ridge-to-reef elevational gradient on a small isolated oceanic island of the central Philippines. *Check List* 18(5): 941–948. <https://doi.org/10.15560/18.5.941>
- Nuñez OM, Galorio AHN (2015) Cave Herpetofauna of Siargao Island Protected Landscape and Seascape, Philippines. *World Journal of Environmental Biosciences* 4: 26–35.
- O’Hanlon SJ, Rieux A, Farrer RA, Rosa GM, Waldman B, Bataille A, Kosch TA, Murray KA, Brankovics B, Fumagalli M, Martin MD, Wales N, Alvarado-Rybak M, Bates KA, Berger L, Böll S, Brookes L, Clare F, Courtois EA, Cunningham AA, Doherty-Bone TM, Ghosh P, Gower DJ, Hintz WE, Höglund J, Jenkinson TS, Lin CF, Laurila A, Loyau A, Martel A, Meurling S, Miaud C, Minting P, Pasmans F, Schmeller DS, Schmidt BR, Shelton JMG, Skerratt LF, Smith F, Soto-Azat C, Spagnoletti M, Tessa G, Toledo LF, Valenzuela-Sánchez A, Verster R, Vörös J, Webb RJ, Wierzbicki C, Wombwell E, Zamudio KR, Aanensen DM, James TY, Gilbert MTP, Weldon C, Bosch J, Balloux F, Garner TWJ, Fisher MC (2018) Recent Asian origin of chytrid fungi causing global amphibian declines. *Science* 360(6389): 621–627. <https://doi.org/10.1126/science.aar1965>
- Oaks J, Sukumaran JR, Esselstyn JA, Linkem CW, Siler CD, Holder MT, Brown RM (2013) Evidence for Pleistocene-driven diversification? A caution for interpreting ABC inferences of simultaneous historical events. *Evolution; International Journal of Organic Evolution* 67(4): 991–1010. <https://doi.org/10.1111/j.1558-5646.2012.01840.x>
- Oaks JR, Siler CD, Brown RM (2019) The comparative biogeography of Philippine geckos challenges predictions from a paradigm of climate-driven vicariant diversification across an island archipelago. *Evolution; International Journal of Organic Evolution* 73–6(6): 1151–1167. <https://doi.org/10.1111/evo.13754>
- Oaks JR, Wood Jr PL, Siler CD, Brown RM (2022) Generalizing Bayesian phylogenetics to infer shared evolutionary events. *Proceedings of the National Academy of Sciences of the United States of America* 119(29): 1–11. <https://doi.org/10.1073/pnas.2121036119>
- Oliver PM, Brown RM, Kraus F, Rittmeyer E, Travers SL, Siler CD (2018) Lizards of the lost arcs: Mid-Cenozoic diversification, persistence and ecological marginalization in the west pacific. *Proceedings of the Royal Society B: Biological Sciences* 285(1871) pii: 20171760. <https://doi.org/10.1098/rspb.2017.1760>
- Oliver PM, Brown RM, Karin B, Grismer LL (2020) Mountain endemism in Malaysian geckos: Are biotic interactions driving lizards up hills? *Israel Journal of Ecology & Evolution* 66(3–4): 190–20. <https://doi.org/10.1163/22244662-20191089>
- Oliveros CH, Ota H, Crombie RI, Brown RM (2011) The herpetofauna of the Babuyan Islands, northern Philippines. *Scientific Papers: Natural History Museum of the University of Kansas* 43: 1–20.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Gromshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA (2021) The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *International Journal of Surgery* 88: 105906. <https://doi.org/10.1016/j.ijsu.2021.105906>

- Peterson AT, Campbell PL, Brown RM (2017) Typhoon frequency and intensity across the western North Pacific Ocean, 1951–2014. *China Scientific Data* 2: 1–11.
- Pili AN, Sy EY, Diesmos MLL, Diesmos AC (2019) Island hopping in a biodiversity hotspot archipelago: Reconstructed invasion history and updated status and distribution of alien frogs in the Philippines. *Pacific Science* 73(3): 321–343. <https://doi.org/10.2984/73.3.2>
- Pili AN, Tingley R, Sy EY, Diesmos ML, Diesmos AC (2021) Niche shifts and environmental non-equilibrium undermine the usefulness of ecological niche models for invasion risk assessment. *Scientific Reports* 10(1): 7972. <https://doi.org/10.1038/s41598-020-64568-2>
- Pitogo KME (2021) Natural history notes on the elusive Taylor's Burrowing Snake, *Pseudorabdion taylori* Leviton and Brown 1959, from southern Mindanao, Philippines. *Herpetology Notes* 14: 759–763.
- Pitogo KME, Saavedra AJL (2021) Rediscovery of Guttman's Stream Frog, *Pulchrana guttmani* (Brown, 2015) from the mountains of southern Mindanao, Philippines. *Herpetology Notes* 14: 163–167.
- Pitogo KME, Saavedra AJL (2023) Conservation assessment of the recently rediscovered Guttman's Stream Frog, *Pulchrana guttmani* (Brown, 2015), reveals the need for its uplisting into a higher threat category. *Herpetology Notes* 16: 717–721.
- Pitogo KME, Saavedra AJL, Afuang LE (2021a) Amphibians and Reptiles of Mount Busa, Sarangani Province: A Glimpse of the Herpetological Community of Southern Mindanao, Philippines. *Philippine Journal of Science* 150(5): 1279–1306. <https://doi.org/10.56899/150.05.37>
- Pitogo KME, Saavedra AJL, Aurellado MEB, de Guia APO, Afuang LE (2021b) Functional traits and environment drive montane amphibian distribution in the southern Philippines. *Biodiversity and Conservation* 30(14): 4177–4197. <https://doi.org/10.1007/s10531-021-02299-0>
- Plaza JL, Sanguila MB (2015) Preliminary Report on the Anurans of Mount Hilong-hilong, Agusan Del Norte, Eastern Mindanao, Philippines. *Asian Herpetological Research* 6(1): 18–33. <https://doi.org/10.16373/j.cnki.ahr.140037>
- Plaza JL, Metillo EB, Sanguila MB (2021) Trophic ecology of syntopic anurans of tropical stream communities. *Journal of Tropical Ecology* 37(3): 109–117. <https://doi.org/10.1017/S0266467421000158>
- Quibod MNRM, Alcantara KNL, Bechayda NA, Estropia CJC, Guntinas JB, Obin MAHA, Raymundo RM, Soniega EP (2021) Terrestrial vertebrates in modified landscapes in northeastern Mindanao, Philippines. *Journal of Animal Diversity* 3(3): 72–85. <https://doi.org/10.52547/JAD.2021.3.3.6>
- R Core Team (2022) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Ramírez-Castañeda V, Westeen EP, Frederick J, Amini S, Wait DR, Achmadi AS, Andayani N, Arida E, Arifin U, Bernal MA, Bonaccorso E, Sanguila MB, Brown RM, Che J, Condori FP, Hartiningtias D, Hiller AE, Iskandar DT, Jimenez RA, Khelifa R, Marquez R, Martinez-Fonseca JG, Parra JL, Peñalba JV, Pinto-Garcia L, Razafindratsima OH, Ron SR, Souza S, Supriatna J, Bowje RCK, Cicero C, McGuire JA, Tarvin RD (2022) A set of principles and practical suggestions for equitable fieldwork in biology. *Proceedings of the National Academy of Sciences of the United States of America* 119(34): e2122667119. <https://doi.org/10.1073/pnas.2122667119>

- Rock KN, Barnes IN, Deyski MS, Glynn KA, Milstead BN, Rottenborn ME, Andre NS, Dekhtyar A, Dekhtyar O, Taylor EN (2021) Quantifying the gender gap in authorship in herpetology. *Herpetologica* 77(1): 1–13. <https://doi.org/10.1655/0018-0831-77.1.1>
- Roll U, Feldman A, Novosolov M, Allison A, Bauer A, Bernard R, Bohm M, Chirio L, Collen B, Colli GR, Dabul L, Das I, Doan T, Grismer L, Herrera FC, Hoogmoed M, Itescu Y, Kraus F, LeBreton M, Lewin A, Martins M, Maza E, Meirte D, Nagy Z, Nogueira CC, Pauwels OSG, Pincheira-Donoso D, Powney G, Sindaco R, Tallowin O, Torres-Carvajal O, Trape JF, Uetz P, Vidan E, Wagner P, Wang YZ, Orme D, Grenyer R, Meiri S (2017) The global distribution of tetrapods reveals a need for targeted reptile conservation. *Nature Ecology & Evolution* 1(11): 1677–1682. <https://doi.org/10.1038/s41559-017-0332-2>
- Ronquist F (1997) Dispersal-Vicariance Analysis: A new approach to the quantification of historical biogeography. *Systematic Biology* 46(1): 195–203. <https://doi.org/10.1093/sysbio/46.1.195>
- Ross CA, Lazell J (1991) Amphibians and reptiles of Dinagat and Siargao islands, Philippines. *Philippine Journal of Science* 119: 257–286.
- Sanguila MB, Siler CD, Diesmos AC, Nuñez O, Brown RM (2011) Phylogeography, geographic structure, genetic variation, and potential species boundaries in Philippine slender toads. *Molecular Phylogenetics and Evolution* 61(2): 333–350. <https://doi.org/10.1016/j.ympev.2011.06.019>
- Sanguila MB, Cobb KA, Siler CD, Diesmos AC, Alcala AC, Brown RM (2016) The amphibians and reptiles of Mindanao Island, southern Philippines, II: The herpetofauna of northeast Mindanao and adjacent islands. *ZooKeys* 624: 1–132. <https://doi.org/10.3897/zookeys.624.9814>
- Sanguila MB, Plaza JL, Mahinay MY, Edma Jr RC, Brown RM (2021) Herpetological assemblages in tropical forests of the Taguibo Watershed, Butuan City, eastern Mindanao, Philippines. *Philippine Journal of Science* 150(S1): 415–431. <https://doi.org/10.56899/150.S1.31>
- Scheffers BR, Brunner RM, Ramirez SD, Shoo LP, Diesmos AC, Williams SE (2013) Thermal Buffering of Microhabitats is a Critical Factor Mediating Warming Vulnerability of Frogs in the Philippine Biodiversity Hotspot. *Biotropica* 45(5): 628–635. <https://doi.org/10.1111/btp.12042>
- Seale A (1917) Sea products of Mindanao and Sulu III: Sponges, tortoiseshell, corals, and trepang. *Philippine Journal of Science* 12: 191–213.
- Seidl CM, Basham EW, Andriamahohatra LR, Scheffers BR (2019) Bird's nest fern epiphytes facilitate herpetofaunal arboreality and climate refuge in two paleotropical canopies. *Oecologia* 192(2): 297–309. <https://doi.org/10.1007/s00442-019-04570-2>
- Setiadi MI, McGuire JA, Brown RM, Zubairi M, Iskandar DT, Andayani N, Supriatna J, Evans BJ (2011) Adaptive radiation and ecological opportunity in Sulawesi and Philippine fanged frog (*Limnonectes*) communities. *American Naturalist* 178(2): 221–240. <https://doi.org/10.1086/660830>
- Sheu Y, Zurano JP, Ribeiro-Junior MA, Avila-Pires TC, Rodrigues MT, Colli GR, Werneck FP (2020) The combined role of dispersal and niche evolution in the diversification of Neotropical lizards. *Ecology and Evolution* 10(5): 2608–2625. <https://doi.org/10.1002/ece3.6091>
- Shooman-Goodier ME, Diaz MI, Almazan ML, Singleton GR, Hadi BAR, Propper CR (2019) Ecosystem hero and villain: Native frog consumes rice pests, while the invasive cane toad feasts on beneficial arthropods. *Agriculture, Ecosystems & Environment* 279: 100–108. <https://doi.org/10.1016/j.agee.2019.04.008>

- Siler CD, Brown RM (2010) Phylogeny-based species delimitation in Philippine slender skinks (Reptilia: Squamata: Scincidae: *Brachymeles*): taxonomic revision of the pentadactyl species groups and description of three new species. *Herpetological Monograph* 24(1): 1–54. <https://doi.org/10.1655/HERPMONOGRAPHS-D-10-00003.1>
- Siler CD, Brown RM (2011) Evidence for repeated acquisition and loss of complex body form characters in an insular clade of Southeast Asian semi-fossorial skinks. *Evolution; International Journal of Organic Evolution* 65(9): 2641–2663. <https://doi.org/10.1111/j.1558-5646.2011.01315.x>
- Siler CD, Linkem CW, Diesmos AC, Alcalá AC (2007) A new species of *Platymantis* (Amphibia: Anura: Ranidae) from Panay Island, Philippines. *Herpetologica* 63(3): 351–364. [https://doi.org/10.1655/0018-0831\(2007\)63\[351:ANSOPA\]2.0.CO;2](https://doi.org/10.1655/0018-0831(2007)63[351:ANSOPA]2.0.CO;2)
- Siler CD, Rico EL, Duya MR, Brown RM (2009a) A new limb-reduced skink, genus *Brachymeles* (Reptilia: Squamata: Scincidae), from central Luzon Island, Philippines. *Herpetologica* 65(4): 449–459. <https://doi.org/10.1655/08-076.1>
- Siler CD, McVay J, Diesmos AC, Brown RM (2009b) A new species of fanged frog (Dicroglossidae; genus *Limnonectes*) from southern Mindanao Island, Philippines. *Herpetologica* 65(1): 105–114. <https://doi.org/10.1655/08-041R1.1>
- Siler CD, Diesmos AC, Alcalá AC, Brown RM (2009c) A new species of limestone forest frogs, genus *Platymantis* (Amphibia; Anura; Ceratobatrachidae) from Samar Island, Philippines. *Herpetologica* 65(1): 92–104. <https://doi.org/10.1655/08-040R.1>
- Siler CD, Oaks JR, Esselstyn JA, Diesmos AC, Brown RM (2010a) Phylogeny and biogeography of Philippine bent-toed geckos (Gekkonidae: *Cyrtodactylus*) contradict a prevailing model of Pleistocene diversification. *Molecular Phylogenetics and Evolution* 55(2): 699–710. <https://doi.org/10.1016/j.ympev.2010.01.027>
- Siler CD, Balete DS, Diesmos AC, Brown RM (2010b) A new legless loam-swimming lizard (Reptilia: Squamata: Scincidae: Genus *Brachymeles*) from the Bicol Peninsula, Luzon Island, Philippines. *Copeia* 2010(1): 114–122. <https://doi.org/10.1643/CH-08-231>
- Siler CD, Diesmos AC, Brown R (2010c) A new loam-swimming skink, genus *Brachymeles* (Reptilia: Squamata: Scincidae) from the Bicol faunal region, Luzon and Catanduanes islands, Philippines. *Journal of Herpetology* 44(1): 49–60. <https://doi.org/10.1670/08-318.1>
- Siler CD, Diesmos AC, Linkem CW, Diesmos ML, Brown RM (2010d) A new species of limestone-forest frog, genus *Platymantis* (Amphibia: Anura: Ceratobatrachidae) from central Luzon Island, Philippines. *Zootaxa* 2482(1): 49–63. <https://doi.org/10.11646/zootaxa.2482.1.3>
- Siler CD, Fuiten AM, Jones RM, Alcalá AC, Brown RM (2011a) Phylogeny-based species delimitation in Philippine slender skinks (Reptilia: Squamata: Scincidae) II: taxonomic revision of *Brachymeles samarensis* and description of five new species. *Herpetological Monograph* 25(1): 76–112. <https://doi.org/10.1655/HERPMONOGRAPHS-D-10-00013.1>
- Siler CD, Crombie RI, Diesmos AC, Brown RM (2011b) Redescription of two poorly known loam-swimming skinks, *Brachymeles bicolor* and *Brachymeles pathfinderi* (Reptilia: Squamata: Scincidae) from the Philippines. *Journal of Herpetology* 45(3): 355–369. <https://doi.org/10.1670/10-103.1>
- Siler CD, Jones RM, Welton LJ, Brown RM (2011c) Redescription of tetradactyl, Philippine slender skinks (genus *Brachymeles*). *Herpetologica* 67(3): 300–317. <https://doi.org/10.1655/HERPETOLOGICA-D-10-00071.1>
- Siler CD, Welton LJ, Siler JM, Brown J, Bucol A, Diesmos AC, Brown RM (2011d) Amphibians and Reptiles, Luzon Island, Aurora Province and Aurora Memorial National Park, Northern Philippines: New island distribution records. *Check List* 7(2): 182–195. <https://doi.org/10.15560/7.2.182>

- Siler CD, Oaks JR, Welton LJ, Linkem CW, Swab J, Diesmos AC, Brown RM (2012a) Did geckos ride the Palawan raft to the Philippines? *Journal of Biogeography* 39(7): 1217–1234. <https://doi.org/10.1111/j.1365-2699.2011.02680.x>
- Siler CD, Swab JC, Oliveros CH, Diesmos AC, Averia L, Alcalá AC, Brown RM (2012b) Amphibians and reptiles, Romblon Island group, central Philippines: Comprehensive herpetofaunal inventory. *Check List* 8(3): 443–462. <https://doi.org/10.15560/8.3.443>
- Siler CD, Jones RM, Diesmos AC, Diesmos ML, Brown RM (2012c) Phylogeny-based species delimitation in Philippine slender skinks (Reptilia: Squamata: Scincidae) III: taxonomic revision of the *Brachymeles gracilis* complex, with description of three new species. *Herpetological Monograph* 26(1): 135–172. <https://doi.org/10.1655/HERPMONOGRAPHS-D-11-00006.1>
- Siler CS, Oliveros CH, Santannen A, Brown RM (2013) Multilocus phylogeny reveals unexpected patterns of regional and morphological diversification in Asian Wolf Snakes (genus *Lycodon*). *Zoologica Scripta* 42: 263–277. <https://doi.org/10.1111/zsc.12007>
- Siler CD, Dececchi AA, Merkord CL, Davis DR, Christiani TJ, Brown RM (2014a) Cryptic diversity and population genetic structure in the rare, endemic, forest-obligate, geckos of the Philippines. *Molecular Phylogenetics and Evolution* 70: 204–209. <https://doi.org/10.1016/j.ympev.2013.09.014>
- Siler CD, Oaks JR, Cobb K, Ota H, Brown RM (2014b) Critically endangered island endemic or peripheral population of a widespread species? Conservation genetics of Kikuchi's gecko and the global challenge of protecting peripheral oceanic island endemic vertebrates. *Diversity & Distributions* 20(7): 756–772. <https://doi.org/10.1111/ddi.12169>
- Siler CD, Welton LJ, Davis DR, Watters JL, Davey CS, Diesmos AC, Diesmos ML, Brown RM (2014c) Taxonomic revision of the *Pseudogekko compresicorpus* Complex (Reptilia: Squamata: Gekkonidae), with descriptions of three new species. *Herpetological Monograph* 28(1): 110–139. <https://doi.org/10.1655/HERPMONOGRAPHS-D-14-00005>
- Siler CD, Lira-Noriega A, Brown RM (2014d) Conservation genetics of Australasian sailfin lizards: Flagship species threatened by coastal development and insufficient protected area coverage. *Biological Conservation* 169: 100–108. <https://doi.org/10.1016/j.biocon.2013.10.014>
- Siler CD, Linkem CW, Cobb K, Watters JL, Cumings S, Diesmos AC, Brown RM (2014e) Taxonomic revision of the semi-aquatic skink *Parvosцинus leucospilos* (Reptilia: Squamata: Scincidae), with description of three new species. *Zootaxa* 3847(3): 388–412. <https://doi.org/10.11646/zootaxa.3847.3.4>
- Siler CD, Freitas ES, Sheridan J, Maguire SN, Davis DR, Watters JL, Wang K, Diesmos AC, Brown RM (2020) Additions to the Philippine slender skinks of the *Brachymeles bonitae* complex (Reptilia: Squamata: Scincidae) IV: Resurrection and redescription of *Brachymeles burksi*. *Philippine Journal of Systematic Biology* 14(2): 1–21. <https://doi.org/10.26757/pjsb2020b14005>
- Slavenko A, Allison A, Austin CC, Bauer AM, Brown RM, Fisher RN, Ineich I, Iova B, Karin BR, Kraus F, O'Shea M, Mecke S, Meiri S, Morrison C, Oliver PM, Richmond JQ, Shea GM, Tallowin OJS, Chapple DG (2023) Skinks of Oceania, New Guinea, and Eastern Wallacea: An underexplored biodiversity hotspot. *Pacific Conservation Biology* 2023(6): 1–18. <https://doi.org/10.1071/PC22034>
- Spinks PQ, Thomson RC, Hughes B, Moxley B, Brown RM, Diesmos AC, Shaffer HB (2012) Cryptic variation and the tragedy of unrecognized taxa: the case of international trade in the spiny turtle *Heosemys spinosa* (Testudines: Geoemydidae). *Zoological Journal of the Linnean Society* 164(4): 811–824. <https://doi.org/10.1111/j.1096-3642.2011.00788.x>

- Suarez RK, Sajise PE (2010) Deforestation, swidden agriculture and Philippine biodiversity. *Philippine Science Letters* 3(1): 91–99. <https://doi.org/10.54645/YZAZ96884>
- Supsup CE, Carestia Jr UV (2020) Rediscovery, new island record, and clarification of the geographic distribution of *Oligodon perkinsi* (Taylor, 1925) (Squamata, Colubridae), a poorly known endemic snake from the Palawan faunal region of western Philippines. *Check List* 16(4): 877–881. <https://doi.org/10.15560/16.4.877>
- Supsup CE, Puna NM, Asis AA, Redoblado BR, Panaguinit MFG, Guinto FM, Rico EB, Diesmos AC, Brown RM, Mallari NAD (2016) Amphibians and reptiles of Cebu, Philippines: The poorly understood herpetofauna of an island with very little remaining natural habitat. *Asian Herpetological Research* 7(3): 151–179. <https://doi.org/10.16373/j.cnki.ahr.150049>
- Supsup CE, Guinto FM, Redoblado BR, Gomez RS (2017) Amphibians and reptiles from the Mt. Hamiguitan Range of eastern Mindanao Island, Philippines: New distribution records. *Check List* 13(3): 2121. <https://doi.org/10.15560/13.3.2121>
- Supsup CE, Asis AA, Carestia Jr UV, Diesmos AC, Mallari NAD, Brown RM (2020) Variation in species richness, composition and herpetological community structure across a tropical habitat gradient of Palawan Island, Philippines. *Herpetozoa* (Wien) 33: 95–111. <https://doi.org/10.3897/herpetozoa.33.e47293>
- Supsup CE, Asis AA, Edaña JWB, Mallari NAD (2022) Anuran assemblages in western Philippines: Unraveling the effects of habitat types, water availability, and elevation. *Acta Oecologica* 117: 103869. <https://doi.org/10.1016/j.actao.2022.103869>
- Supsup CE, Asis AA, Eslava MRR, Domingo JPS, Amarga AKS, Carestia Jr UV, Cantil JA, delos Angeles MD, Acosta-Lagrada LS (2023) Revisiting environmental management zones toward conserving globally important species in western Philippines. *Journal for Nature Conservation* 73: 126415. <https://doi.org/10.1016/j.jnc.2023.126415>
- Swei A, Rowley JJJ, Rödder D, Diesmos MLL, Diesmos AC, Briggs CJ, Brown RM, Cao TT, Cheng TL, Chong RA, Han B, Hero JM, Hoang HD, Kusri MD, Le DTT, McGuire JA, Meegaskumbura M, Min MS, Mulcahy DG, Neang T, Phimmachak S, Rao DQ, Reeder NM, Schoville SD, Sivongxay N, Srei N, Stöck M, Stuart BL, Torres LS, Tran DTA, Tunstall TS, Vietes D, Vredenburg VT (2011a) Prevalence and distribution of chytridiomycosis throughout Asia. *FrogLog* 98: 33–34.
- Swei A, Rowley JJJ, Rödder D, Diesmos MLL, Diesmos AC, Briggs CJ, Brown RM, Cao TT, Cheng TL, Chong RA, Han B, Hero JM, Hoang HD, Kusri MD, Le DTT, McGuire JA, Meegaskumbura M, Min MS, Mulcahy DG, Neang T, Phimmachak S, Rao DQ, Reeder NM, Schoville SD, Sivongxay N, Srei N, Stöck M, Stuart BL, Torres LS, Tran DTA, Tunstall TS, Vietes D, Vredenburg VT (2011b) Is chytridiomycosis an emerging disease in Asia? *PLOS ONE* 6(8): e23179. <https://doi.org/10.1371/journal.pone.0023179>
- Sy EY, Schoppe S, Diesmos MLL, Lim TMS, Diesmos AC (2020) Endangered by trade: Seizure analysis of the critically endangered Philippine Forest Turtle *Siebenrockiella leytensis* from 2004–2018. *Philippine Journal of Systematic Biology* 14(2): 1–8. <https://doi.org/10.26757/pjsb2020b14003>
- Taylor EH (1918) Reptiles of the Sulu archipelago. *Philippine Journal of Science* 13: 233–267.
- Taylor EH (1921) Amphibians and Turtles of the Philippine Islands. Department of Agriculture and Natural Resources, Philippine Bureau of Science, Monograph, Manila, Philippines, 1–193.
- Taylor EH (1922a) The Lizards of the Philippine Islands. Department of Agriculture and Natural Resources, Philippine Bureau of Science, Monograph, Manila, Philippines, 17 pp.
- Taylor EH (1922b) The snakes of the Philippine Islands. Department of Agriculture and Natural Resources, Bureau of Science, Manila, Philippines, 312.

- Taylor EH (1928) Amphibians, lizards, and snakes of the Philippines. In: Dickerson R, (Ed.) Distribution of Life in the Philippines. Philippine Bureau of Science, Monograph 21, Manila, Philippines, 214–241.
- Tingley R, Macdonald SL, Mitchell NJ, Woinarski JCZ, Meiri S, Bowles P, Cox NA, Shea GM, Bohm M, Chanson J, Tognelli MF, Harris J, Walke C, Harrison N, Victor S, Woods C, Amey AP, Bamford M, Catt G, Clemann N, Couper PJ, Cogger H, Cowan M, Craig MD, Dickman CR, Doughty P, Ellis R, Fenner A, Ford S, Gaikhorst G, Gillespie GR, Greenlees MJ, Hobson R, Hoskin CJ, How R, Hutchinson MN, Lloyd R, McDonald P, Melville J, Michael DR, Moritz C, Oliver PM, Peterson G, Robertson P, Sanderson C, Somaweera R, Teale R, Valentine L, Vanderduys E, Venz M, Wapstra E, Wilson S, Chapple DG (2019) Geographic and taxonomic patterns of extinction risk in Australian squamates. *Biological Conservation* 238: 108–203. <https://doi.org/10.1016/j.biocon.2019.108203>
- Uetz P, Freed P, Aguilar R, Reyes F, Hošek J [Eds] (2023) The Reptile Database. <http://reptile-database.reptarium.cz/>
- van de Ven WA, Guerrero JP, Rodriguez DG, Telan SP, Balbas MG, Tarun BA, Van Weerd M, van der Ploeg J, Wijtten Z, Lindeyer FE (2009) Effectiveness of head-starting to bolster Philippine crocodile *Crocodylus mindorensis* populations in San Mariano Municipality, Luzon, Philippines. *Conservation Evidence* 6: 111–116.
- van der Ploeg J, Arano RR, van Weerd M (2011) What Local People Think About Crocodiles: Challenging Environmental Policy Narratives in the Philippines. *Journal of Environment & Development* 20(3): 303–328. <https://doi.org/10.1177/1070496511416743>
- Venturina REL, Del Prado YLC, Kamir RAC, Balmores MN, Diesmos AC (2020) A revised checklist of amphibians and reptiles in Camiguin Sur, Misamis Oriental, Mindanao, Philippines. *Asian Herpetological Research* 11(1): 28–43. <https://doi.org/10.16373/j.cnki.ahr.190036>
- Vidan E, Bauer AM, Herrera FC, Chirio L, Nogueira CC, Doan TM, Lewin A, Meirte D, Nagy ZT, Novosolov M, Pincheira-Donoso D, Tallowin OJ, Torres-Carvajal O, Uetz P, Wagner P, Wang Y, Belmaker J, Meiri S (2019) The global biogeography of lizard functional groups. *Journal of Biogeography* 46(10): 2147–2158. <https://doi.org/10.1111/jbi.13667>
- Wallach V, Brown RM, Diesmos AC, Gee GVA (2007) An enigmatic new species of blind snake from Luzon Island, with a Synopsis of the Genus *Acutotyphlops* northern Philippines (Serpentes: Typhlopidae). *Journal of Herpetology* 41: 690–702. <https://doi.org/10.1670/206-5.1>
- Weinell JL, Brown RM (2017) Discovery of an old, archipelago-wide, endemic radiation of Philippine snakes. *Molecular Phylogenetics and Evolution* 119: 144–150. <https://doi.org/10.1016/j.ympev.2017.11.004>
- Weinell JL, Hooper E, Leviton AE, Brown RM (2019) Illustrated key to the snakes of the Philippines. *Proceedings of the California Academy of Sciences* 66(1): 1–49.
- Weinell JL, Paluh DJ, Siler CD, Brown RM (2020a) A new, miniaturized genus and species of snake (Cyclocoridae) from the Philippines. *Copeia* 108(4): 907–923. <https://doi.org/10.1643/CH2020110>
- Weinell J, Barley AJ, Siler CD, Orlov NL, Ananjeva NB, Oaks JR, Burbrink F, Brown RM (2020b) Phylogenetic relationships and biogeographic range evolution in Cat-eyed Snakes *Boiga* (Serpentes: Colubridae). *Zoological Journal of the Linnean Society* 2020: 1–16. <https://doi.org/10.1093/zoolinnean/zlaa090>
- Weinell JL, Leviton AE, Brown RM (2020c) A new species of Reed Snake, genus *Calamaria* (Colubridae: Calamariinae), from Mindoro Island. *Philippine Journal of Systematic Biology* 14(2): 1–14. <https://doi.org/10.26757/pjsb2020b14006>

- Welton LJ, Siler CD, Diesmos AC, Brown RM (2009) A new bent-toed gecko (Genus *Cyrtodactylus*) from southern Palawan Island, Philippines, and clarification of the taxonomic status of *C. annulatus*. *Herpetologica* 65(3): 323–343. <https://doi.org/10.1655/08-057R1.1>
- Welton LJ, Siler CD, Diesmos AC, Brown RM (2010a) Phylogeny-based species delimitation of southern Philippine bent-toed geckos and a new species of *Cyrtodactylus* (Squamata: Gekkonidae) from western Mindanao and the Sulu Archipelago. *Zootaxa* 2390(1): 49–68. <https://doi.org/10.11646/zootaxa.2390.1.3>
- Welton LJ, Siler CD, Diesmos AC, Brown RM (2010b) Philippine bent-toed geckos of the *Cyrtodactylus agusanensis* complex: Multilocus phylogeny, morphological diversity, and description of three new species. *Herpetological Monograph* 24(1): 55–85. <https://doi.org/10.1655/HERPMONOGRAPHS-D-10-00005.1>
- Welton LJ, Siler CD, Bennet D, Diesmos AC, Duya MR, Dugay R, Rico EL, van Weerd M, Brown RM (2010c) A spectacular new Philippine monitor lizard reveals a hidden biogeographic boundary and a novel flagship species for conservation. *Biology Letters* 6(5): 654–658. <https://doi.org/10.1098/rsbl.2010.0119>
- Welton LJ, Siler CD, Diesmos AC, Diesmos MLD, Lagat R, Causaren R, Brown RM (2012) Genetic identity, geographic range, and major distribution records for frugivorous monitor lizards of Luzon Island, Philippines. *Herpetological Review* 43: 226–230.
- Welton LJ, Siler CD, Oaks JR, Diesmos AC, Brown RM (2013a) Multilocus phylogeny and Bayesian estimates of species boundaries reveal hidden evolutionary relationships and cryptic diversity in Southeast Asian monitor lizards. *Molecular Ecology* 22(13): 3495–3510. <https://doi.org/10.1111/mec.12324>
- Welton LJ, Siler CD, Linkem CW, Diesmos AC, Diesmos ML, Sy E, Brown RM (2013b) Dragons in our midst: Phyloforensics of illegally traded Southeast Asian monitor lizards. *Biological Conservation* 159: 7–15. <https://doi.org/10.1016/j.biocon.2012.10.013>
- Welton LJ, Wood Jr PL, Oaks JR, Siler CD, Brown RM (2014a) Fossil-calibrated phylogeny and historical biogeography of Southeast Asian water monitors (*Varanus salvator* Complex). *Molecular Phylogenetics and Evolution* 74: 29–37. <https://doi.org/10.1016/j.ympev.2014.01.016>
- Welton LJ, Siler CD, Travers SN, Brown RM (2014b) Integrative taxonomy and phylogeny-based species delimitation of Philippine water monitor lizards (*Varanus salvator* Complex) with descriptions of two new cryptic species. *Zootaxa* 3881(3): 201–227. <https://doi.org/10.11646/zootaxa.3881.3.1>
- Welton LJ, Siler CD, Grismer LL, Diesmos AC, Sites JW, Brown RM (2017) Archipelago-wide survey of Philippine forest dragons (Agamidae: *Gonocephalus*): multilocus phylogeny uncovers unprecedented levels of genetic diversity in a biodiversity hotspot. *Biological Journal of the Linnean Society*. *Linnean Society of London* 120(2): 410–426. <https://doi.org/10.1111/bj.12878>
- Welton LJ, Rechhio I, Wood Jr PL, Brown RM (2020) Phyloforensics in Action: Genetic Identity and Island Provenance of an Illegally Trafficked Philippine Monitor Lizard. *Herpetological Review* 51: 215–220.
- Womack MC, Steigerwald E, Blackburn DC, Cannatella DC, Catenazzi A, Che J, Koo MS, McGuire JA, Ron SR, Spencer CL, Vredenburg VT, Tarvin RD (2022) State of the Amphibia 2020: A Review of Five Years of Amphibian Research and Existing Resources. *Ichthyology & Herpetology* 110(4): 638–661. <https://doi.org/10.1643/h2022005>
- Wood Jr PL, Guo X, Travers SL, Su YC, Olson KV, Bauer AM, Grismer LL, Siler CD, Moyle RG, Andersen MJ, Brown RM (2020) Parachute geckos free fall into synonymy: *Gekko* phylogeny, and a new subgeneric classification, inferred from thousands

- of ultraconserved elements. *Molecular Phylogenetics and Evolution* 146: 106731. <https://doi.org/10.1016/j.ympev.2020.106731>
- Wynn A, Diesmos AC, Brown RM (2016) Two new species of *Malayotyphlops* from the northern Philippines, with redescription of *M. luzonensis* (Taylor) and *M. ruber* (Boettger). *Journal of Herpetology* 50: 157–168. <https://doi.org/10.1670/14-104>
- Young JC, Waylen KA, Sarkki S, Albon S, Bainbridge I, Balian E, Davidson J, Edwards D, Fairley R, Margerison C, McCracken D, Owen R, Quine CP, Stewart-Roper C, Thompson D, Tinch R, Van den Hove S, Watt A (2014) Improving the science-policy dialogue to meet the challenges of biodiversity conservation: Having conversations rather than talking at one-another. *Biodiversity and Conservation* 23(2): 387–404. <https://doi.org/10.1007/s10531-013-0607-0>
- Zimin A, Zimin SV, Shine R, Bauer A, Böhm M, Brown RM, de Oliveira Caetano GH, Herrera FC, Chapple DG, Chirio L, Colli GR, Doan TM, Glaw F, Grismer LL, Itescu Y, Kraus F, LeBreton M, Martins M, Morando M, Murali G, Nagy ZT, Novosolov M, Oliver P, Passos P, Pauwels OSG, Pincheira-Donoso D, Junior MAR, Shea G, Tingley R, Torres-Carvajal O, Trape JF, Uetz P, Wagner P, Roll U, Meiri S (2022) A global analysis of viviparity in squamates highlights its prevalence in cold climates. *Global Ecology and Biogeography* 2022(31): 2437–2452. <https://doi.org/10.1111/geb.13598>

Supplementary material 1

Scientific articles on Philippine herpetology, published from 2002–2022

Authors: Camila G. Meneses, Kier Mitchel E. Pitogo, Christian E. Supsup, Rafe M. Brown
Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1190.109586.suppl1>

Supplementary material 2

Brief distribution records

Authors: Camila G. Meneses, Kier Mitchel E. Pitogo, Christian E. Supsup, Rafe M. Brown
Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1190.109586.suppl2>