

MINIREVIEW

Evolution of Host Adaptation in *Salmonella enterica*

ANDREAS J. BÄUMLER,^{1*} RENÉE M. TSOLIS,² THOMAS A. FICHT,² AND L. GARRY ADAMS²

Department of Medical Microbiology and Immunology, College of Medicine,¹ and Department of Veterinary Pathobiology, College of Veterinary Medicine,² Texas A&M University, College Station, Texas 77843-4467

INTRODUCTION

The question of how bacteria are able to overcome species barriers and adapt to new hosts is central to the understanding of both the origin of infectious diseases and the emergence of new pathogens. The analysis of virulence factors used by different *Salmonella* serotypes can serve as a powerful model for studying mechanisms of host adaptation because these pathogens are physiologically well characterized and lend themselves to genetic analysis. However, they differ greatly with regard to host range and their degree of host adaptation. *Salmonella* serotypes are closely related as shown by analysis of orthologous genes. Divergence in the nucleotide sequence of orthologous genes ranges between 3.8 and 4.6% and differences in their deduced amino acid sequences range between 0.7 and 1.3% (108). This close DNA relatedness among *Salmonella* serotypes is evidence for their clonal origin, and based on the degree of sequence divergence, it can be estimated that a common ancestor of the genus existed about 25 to 40 million years ago. Which factors contributed to the clonal divergence of the genus *Salmonella* from its common ancestor to give rise to serotypes that differ with regard to their host range?

It has been postulated recently that in the genus *Salmonella* virulence evolved in three phases (11) (Fig. 1). The first phase involved acquisition of *Salmonella* pathogenicity island 1 (SPI 1) by plasmid- or phage-mediated horizontal gene transfer. SPI 1 was likely obtained by a lineage ancestral to all *Salmonella* serotypes, since it is present in all phylogenetic lineages of the genus *Salmonella* but absent from *Escherichia coli* and other related organisms (70, 79, 85). SPI 1 encodes virulence factors that mediate mechanisms used by *Salmonella* serotypes during the intestinal phase of infection, including invasion of intestinal epithelial cells (48, 62, 120), induction of neutrophil recruitment (49, 77), and secretion of intestinal fluid (49). Thus, it is likely that these or related virulence mechanisms may have existed early in the evolution of *Salmonella* serotypes.

Multilocus enzyme electrophoresis and comparative sequence analysis of housekeeping and rRNA genes revealed that the genus *Salmonella* contains two lineages that have diverged considerably from each other during evolution (24, 31, 97). By using genetic distance determined by multilocus enzyme electrophoresis and results of DNA-DNA hybridization studies as criteria, it has been proposed that these lineages represent two distinct species, designated *Salmonella enterica* and *Salmonella bongori* (Fig. 1) (69, 97). The formation of these two species could be considered a second phase in the

evolution of virulence in the genus *Salmonella*, since it involved not only divergence of their lineages by point mutation but also acquisition of new virulence determinants by horizontal gene transfer. Serotypes belonging to *S. enterica* possess a second pathogenicity island, designated SPI 2, that is not present in *S. bongori* serotypes (55, 84). The virulence genes present on SPI 2 have an average G+C content of 41%, much lower than the overall G+C content of the *S. enterica* genome, which averages 52% (55). The limited phylogenetic distribution and the atypical G+C content of SPI 2 imply that this pathogenicity island was acquired horizontally after *S. enterica* branched from the *S. bongori* lineage (55, 84). A possible mechanism for acquisition of SPI 2 by horizontal transfer is suggested by its insertion into the *S. enterica* gene encoding tRNA^{Val}, a DNA region which may facilitate integration of newly acquired genetic material because tRNA genes serve as an attachment site for bacteriophage (55, 59). Experiments performed with mice revealed that mutations in SPI 1 attenuate *S. enterica* serotype Typhimurium between 15- and 50-fold after oral injection but have no attenuating phenotype when the intestinal phase of infection is bypassed by intraperitoneal injection (17, 48). In contrast, mutations in SPI 2 attenuate *S. enterica* serotype Typhimurium more than 10,000-fold even after intraperitoneal inoculation (86, 109). Although the mechanism by which SPI 2 contributes to pathogenicity has not yet been elucidated, these data imply that virulence genes located on SPI 1 and SPI 2 are required at different stages, namely the intestinal and the systemic phases of infection, respectively.

Finally, the lineage of *S. enterica* is postulated to have branched into several distinct phylogenetic groups, which by current nomenclature are considered subspecies. The formation of one of these groups, *S. enterica* subspecies I, involved a dramatic expansion in host range: while *S. bongori* and *S. enterica* subspecies II, IIIa, IIIb, IV, VI, and VII are mainly associated with cold-blooded vertebrates, members of *S. enterica* subspecies I are most frequently isolated from avian and mammalian hosts (2, 93). The host adaptation of *S. enterica* subspecies I to warm-blooded vertebrates characterized a third phase in the evolution of virulence in the genus *Salmonella* and is the focus of this review.

LYMPH NODES, A NEW BARRIER ENCOUNTERED IN MAMMALS

Which new barriers did *S. enterica* subspecies I serotypes encounter in birds and mammals? One obstacle may have been the immune system of higher vertebrates, which is more highly developed and shows a higher level of organization than that of cold-blooded vertebrates. For instance, lymphoid aggregates located in the gut lamina propria of cold-blooded vertebrates appear only as solitary units (82) whereas gut-associated lymphoid nodules of mammals and birds are organized in complex

* Corresponding author. Mailing address: Department of Medical Microbiology and Immunology, College of Medicine, Texas A&M University Health Science Center, 407 Reynolds Medical Building, College Station, TX 77843-1114. Phone: (409) 862-7756. Fax: (409) 845-3479. E-mail: abaumler@tamu.edu.




FIG. 1. Model for the evolution of virulence in the genus *Salmonella*. The three phases in which virulence evolved in the genus *Salmonella* since its divergence from the *E. coli* lineage have been proposed previously (11). The phylogenetic tree is not drawn to scale. For explanation see the text.

organs, such as Peyer's patches, tonsils, appendix, or the avian bursa of Fabricius (51, 76). In addition, in birds and mammals B-lymphocyte variants which arise after somatic hypermutation are selected in the germinal centers of lymphoid organs for improving the affinity to their antigen. In contrast, because the lymphoid organs of cold-blooded vertebrates lack germinal centers selection of B cells is impeded, and consequently, antibody affinity does not increase during the immune response (40, 123). Furthermore, the antibody repertoire of lower vertebrates, such as fish, amphibians, and reptiles, is much less diverse than that of mammals (39). B cells selected in germinal centers of higher vertebrates show isotopic switching and are accumulated as memory cells. In cold-blooded vertebrates, on the other hand, immunological memory is poorly developed (39) and repeated immunization with *S. enterica* induces only the production of immunoglobulin M antibodies in lizards (82), suggesting that isotype switching does not occur during infection with this pathogen.

Finally, the presence of more highly developed peripheral lymphoid filter organs constitutes yet another challenge during infection of warm-blooded hosts. Pathogens that are able to penetrate the intestinal mucosa in fish can spread unchecked to central parts of the body until they are filtered from the blood by phagocytes located in sinusoids of the spleen. During the adaptation to life on land vertebrates developed peripheral lymphoid organs which function as local filtering systems (65). Although less-developed lymph node-like structures have been reported in some reptiles (82), true lymph nodes are present only in mammals and some bird species (41, 42, 51, 123). In mammals, a large number of lymph nodes located at peripheral positions form an highly effective lymph filter system (41, 42). Thus, upon penetration of the intestinal mucosae of mammals, *Salmonella* serotypes are confronted by an effective barrier to further spread, namely macrophages that line the lymphatic

sinuses of regional lymph nodes. In mammals this host defense mechanism can successfully limit bacterial expansion to the intestine, the gut-associated lymphoid tissue, and the mesenteric lymph node. For instance, humans infected with nontyphoidal *Salmonella* serotypes usually develop an acute gastroenteritis, but in only 1 to 7% of clinical cases do bacteria manage to pass through the mesenteric lymph node and cause bacteremia (19). Therefore, in order to produce systemic disease in warm-blooded hosts, microbial intruders must be able to breach the local defense formed by macrophages of regional lymph nodes.

The capability of *S. bongori* or *S. enterica* subspecies II, IIIa, IIIb, IV, VI, and VII serotypes to survive in macrophages of poikilothermic animals has not been studied. Experimental oral exposure of snakes, turtles, or lizards to isolates of *S. enterica* subspecies I, II, or III resulted in all cases in no overt signs of disease and no colonization of organs other than the intestinal tract (35–37, 111). It has therefore been speculated that *Salmonella* serotypes evolved in the alimentary tract of reptiles, where they developed from pathogens into commensal organisms (87, 102). In contrast, *S. enterica* subspecies I serotypes frequently colonize internal organs of warm-blooded animals and the ability to survive and multiply in cells of the reticuloendothelial system correlates with their capability to cause systemic disease in these hosts (8, 45). Since macrophages from distinct homeothermic animal species differ greatly in their abilities to neutralize a particular *S. enterica* serotype, adaptation to a new host may require adaptation to its mononuclear phagocytes. For instance, the human-adapted *S. enterica* serotype Typhi is able to survive in vitro in human macrophages but not in murine macrophages, whereas *S. enterica* serotype Typhimurium, which causes a systemic disease in mice, survives well in vitro in murine macrophages but not in human macrophages (117). Furthermore, these two *S. enterica*

TABLE 1. Diseases caused by *Salmonella* subspecies I serotypes in humans and higher vertebrates

Host species	Disease	<i>S. enterica</i> subspecies I serotype(s) most frequently encountered	Most susceptible age groups	Typical symptoms or sign(s) of disease	Reference
Humans	Salmonella enteritis	Typhimurium, Enteritidis	Children (<4 yr)	Diarrhea, dysentery, fever	78
	Typhoid fever	Typhi ^c	Children and adults	Septicemia, fever ^a	78
	Paratyphoid fever	Sendai; Paratyphi A, B, and C ^c	Children and adults	Septicemia, fever ^a	78
Cattle	Salmonellosis	Typhimurium Dublin	Calves (<8 wk)	Diarrhea, dysentery, septicemia, fever	104
			Calves and adult cattle	Diarrhea, dysentery, septicemia, abortion, fever	99, 104
Poultry	Pullorum disease	Pullorum ^{c,d}	Newly hatched birds	Diarrhea, septicemia	27
	Fowl typhoid	Gallinarum ^{c,d}	Growing stock and adults	Diarrhea, comb discoloration, septicemia	27
	Avian paratyphoid	Enteritidis, Typhimurium	Newly hatched birds	Diarrhea, septicemia	27
Sheep	Salmonellosis	Abortusovis ^c	Adult sheep	Septicemia, abortion, vaginal discharge	90
			Lambs	Diarrhea, dysentery, septicemia	90
		Typhimurium	Lambs	Diarrhea, dysentery, septicemia	90
Pigs	Pig paratyphoid	Choleraesuis ^c	Weaned and adult pigs	Skin discoloration, septicemia, fever ^b	106
	Salmonellosis	Typhimurium	Weaned pigs (<4 mo)	Diarrhea	106
	Chronic paratyphoid	Typhisuis		Intermittent diarrhea	7
Horses	Salmonellosis	Abortusequi ^c	Adult horses	Septicemia, abortion	110
			Foals	Diarrhea, septicemia	110
		Typhimurium	Foals	Diarrhea, septicemia	110
Wild rodents	Murine typhoid	Typhimurium, Enteritidis		Septicemia, fever	28

^a Diarrhea develops only in about one third of typhoid fever patients and usually several days after the onset of fever.

^b Diarrhea is not a typical sign of pig paratyphoid but may develop by the third or fourth day of disease.

^c These serotypes have been most frequently associated with illness in the preantibiotic era but are now rare or have been eradicated in most developed countries.

^d Gallinarum and Pullorum are considered biotypes that belong to the same serotype.

serotypes differ in their mechanism of bacterial internalization and their intracellular trafficking in human and mouse mononuclear phagocytes (3, 60, 61, 91). These data imply that the ability of *S. enterica* serotypes to cause systemic disease is directly related to the capability to withstand an assault by the macrophages of a given host. Thus it appears that mononuclear phagocytes are an important barrier that restricts the host range of *Salmonella* serotypes.

COEVOLUTIONARY COURSE TOWARD INCREASED VIRULENCE

Although *S. enterica* subspecies I contains 1,367 different serotypes (93), only one or a few are associated with the majority of cases of illness in a particular avian or mammalian species (Table 1). These serotypes show different degrees of host adaptation. Pathogens that lack host specificity, such as *S. enterica* serotype Typhimurium and *S. enterica* serotype Enteritidis, tend to be more frequently associated with disease in young animals than in adults, suggesting that they are not optimally adapted to cope with a fully mature immune system. Serotypes that are host specific, on the other hand, have acquired the ability to breach defense mechanisms in full-grown animals, as shown by their association, at similar rates, with illness in all age groups (Table 1). Furthermore, host-specific serotypes tend to be more virulent, as illustrated by the fact that they cause higher mortality rates.

It has been proposed that vertebrate pathogens which cause high mortality rates are new arrivals that are not well adapted to their hosts, because the death of an animal would destroy their habitat, thereby reducing both their transmissibility and

fitness. According to this theory, these pathogens will evolve over time into less-virulent forms, which will reflect better adaptation to their hosts (26). However, in *S. enterica* subspecies I evolution has apparently driven host-pathogen interactions in the opposite direction, since strongly host-adapted serotypes tend to cause higher mortality rates than those with a broad host range. For instance, *S. enterica* serotype Typhi exhibits host specificity for humans in whom it causes typhoid fever, a disease which results in 12 to 32% mortality (78). In contrast, death occurs in less than 0.5% of cases of human illness caused by zoonotic *Salmonella* serotypes, such as *S. enterica* serotype Typhimurium or *S. enterica* serotype Enteritidis (78). What selective forces were responsible for the increase in virulence that accompanied the adaptation of *Salmonella* serotypes to humans?

On the basis of a mathematical model, Anderson and May (5) predicted that if transmissibility in a particular host-pathogen association is linked to a high level of virulence, then the coevolutionary course will be toward high virulence. During coevolution of *Salmonella* serotypes and their human host such a link between transmissibility and high virulence may have been provided by the ability to develop chronic carriage. A chronic carrier has the potential to infect a large number of individuals, thereby increasing transmissibility. Chronic carriage develops more frequently following systemic infection, such as that caused by the host-adapted *S. enterica* serotype Typhi. For instance, 1 to 4% of patients that recover from typhoid fever but only 0.2 to 0.6% of patients with nontyphoidal salmonellosis become chronic carriers (56, 81). Thus, the link between systemic infection and increased transmissibility




FIG. 2. Phylogenetic distribution of virulence genes and virulence of different *S. enterica* subspecies I serotypes in mice and humans. The bacterial isolates shown in this figure have been described previously (23). The left side shows a dendrogram reflecting the phylogenetic relatedness of these strains as reported by Selander and coworkers (23). ET, enzyme type determined by multilocus enzyme electrophoresis (23). The distribution of the *spv* (*Salmonella* plasmid virulence), *lpf* (long polar fimbriae), *sef* (*S. enterica* serotype Enteritidis fimbriae), *agf* (thin aggregative fimbriae), *fim* (type 1 fimbriae), and *pef* (plasmid-encoded fimbriae) operons among these isolates has been determined previously (12, 17). The presence in these strains of sequences related to *viaB* (Vi capsular antigen) is described elsewhere (107). Virulence of these isolates in humans or mice has been reported recently (17, 107). a, although missing in this strain, virulence plasmids are present in most isolates of this serotype.

resulting from chronic carriage may have exerted evolutionary pressure on the development of a high degree of virulence during the adaptation of typhoidal *Salmonella* serotypes (causing typhoid or paratyphoid fever in humans) to the human host.

THE VIRULENCE PLASMID IS IMPORTANT FOR SYSTEMIC DISEASE CAUSED BY NONTYPHOIDAL *S. ENTERICA* SEROTYPES

Although nontyphoidal *S. enterica* serotypes do not cause enteric fever in humans, some do cause similar systemic infections in warm-blooded animals. In *S. enterica* subspecies I, the *spv* (*Salmonella* plasmid virulence [54]) operon is found in only a few serotypes, including *S. enterica* serotypes Typhimurium, Enteritidis, Choleraesuis, Gallinarum/Pullorum, Abortusovis, Paratyphi C, and Dublin (17, 75, 94, 95, 103, 122) (Fig. 2). Interestingly, these serotypes are also among those most frequently associated with disease in homeothermic vertebrates (Table 1), suggesting a role for *spv* during infection of these hosts. The *spv* operon is required for the systemic phase of disease caused by *S. enterica* serotype Choleraesuis in pigs (34), *S. enterica* serotype Gallinarum/Pullorum in chickens (9, 10), *S. enterica* serotype Dublin in cattle (72, 118), and *S. enterica* serotypes Typhimurium and Enteritidis in mice (53, 63, 83). Epidemiological evidence provides support for the idea that the *spv* operon is also important for the pathogenesis of extra intestinal infections associated with nontyphoidal *Salmonella* serotypes in humans (46). Therefore, in a first approximation it appears that the *spv* operon is required for systemic infections caused by nontyphoidal serotypes in warm-blooded animals, including humans (9, 10, 34, 53, 63, 72, 83, 103, 118). Mechanisms for systemic infection that are *spv*-dependent may, how-

ever, not be restricted to nontyphoidal serotypes since the *spv* operon is present in *S. enterica* serotype Paratyphi C (17, 94). Typhoidal serotypes which lack the *spv* genes, such as *S. enterica* serotypes Typhi, Paratyphi A, Paratyphi B, and Sendai, produce enteric fever by an *spv*-independent mechanism. The virulence determinants responsible have not yet been identified (94, 100, 122).

Although all members of *S. enterica* subspecies I that are able to produce lethal infection in mice possess the *spv* operon, its introduction into *S. enterica* serotype Typhi does not confer mouse virulence to this host-restricted pathogen (103). Furthermore, the *spv* operon is present in most isolates of *S. enterica* serotype Gallinarum/Pullorum, which does not cause disease in mice. These data suggest that host-restricted *S. enterica* serotypes lack additional virulence factors that are encoded on the chromosome of serotypes with a broad host range. Furthermore, in addition to its presence in several serotypes of *S. enterica* subspecies I, the *spv* operon has also been detected in some isolates of *S. bongori* and *S. enterica* subspecies IIIa and IV (98), and thus the expansion in host range observed for subspecies I cannot be explained merely by the presence of these virulence genes. Therefore, the ability to cause systemic disease in a warm-blooded host is apparently a complex phenotype that cannot be attributed to acquisition of a single virulence determinant. It has been shown recently by subtractive hybridization analysis that 20% of the genome of *S. enterica* serotype Typhimurium is not present in *S. enterica* serotype Typhi and vice versa (67). By comparison with the similarly sized *E. coli* genome, the genome of *S. enterica* serotype Typhimurium can be estimated to contain approximately 4,400 genes. Serotype-specific DNA may thus encode as many as 880 genes, some of which may contribute to determining the




FIG. 3. Linear genetic maps of virulence plasmids of *S. enterica* subspecies I serotypes. Plasmids were linearized at their conserved *par* (partitioning) region (29, 113, 114). Regions of homology shared among plasmids are shown as closed bars, which are positioned according to their locations on the *S. enterica* serotype Typhimurium virulence plasmid. The distribution among virulence plasmids of *spv* (*Salmonella* plasmid virulence) (6, 66, 122), *repC* (replicon C) (113), *pef* (plasmid encoded fimbriae) (12, 47), *repB* (replicon B) (113), and the *tra* (conjugative plasmid transfer) region (66, 101) have been reported recently. Areas of the plasmid of *S. enterica* serotype Dublin that are not present in *S. enterica* serotype Typhimurium have been described previously and are shown as open bars (68, 80). The presence on the *S. enterica* serotype Dublin plasmid of sequences with homology to *feaH* and *feaI*, two genes involved in biosynthesis of K88 fimbriae in *E. coli*, have been reported by Barrow and coworkers (105). The location and size of deletions that may account for the smaller size of plasmids from *S. enterica* serotypes Choleraesuis and Enteritidis relative to the *S. enterica* serotype Typhimurium plasmid are reported elsewhere (25, 80, 101). Dashed lines indicate the position of two DNA regions, namely the *pef* and *spv* operons, which are shown in more detail above the *S. enterica* serotype Typhimurium plasmid (1, 47, 52). The positions of genes are indicated by arrows.

host range. A combination of some of these genes, which are likely to be distributed throughout the chromosome, might determine the ability to infect a particular host. This idea may explain why attempts to extend the host range of host-restricted *S. enterica* serotypes by transferring small segments of the genome from a broad-host-range serotype have not succeeded.

Although the distribution of the *spv* operon has not been determined for all phylogenetic lineages of the genus *Salmonella*, its localization on large serotype-specific plasmids in *S. enterica* subspecies I suggests that this operon was obtained by horizontal gene transfer. Additional regions of virulence plasmids, such as the origin of transfer and the *tra* genes, hint at conjugation as a possible mechanism by which the *spv* operon was horizontally transmitted (25, 89, 101). Evidence for the presence of the *spv* operon on a predecessor plasmid that gave rise to the virulence plasmids found among extant *S. enterica* serotypes comes from analysis of their patterns of homology. For instance, the virulence plasmids of *S. enterica* serotypes Enteritidis and Choleraesuis could be seen as variants of the virulence plasmid of *S. enterica* serotype Typhimurium that were generated by deletion events which occurred during their divergence from a common predecessor (Fig. 3) (25, 80, 101, 120). Similarly, a 25-kb DNA region containing the *spv* operon is conserved between virulence plasmids of *S. enterica* serotypes Dublin and Typhimurium, suggesting a common ancestry (18, 68). But where did the virulence plasmids of *Salmonella* serotypes originate?

The genes *finO*, *traY*, and *repA* are present on the *E. coli* F plasmid and the virulence plasmids of *S. enterica* serotypes Enteritidis and Typhimurium (101). Comparison of the nucleotide sequences determined for the *finO*, *traY*, and *repA* genes of several natural isolates of *E. coli* and *S. enterica* revealed similar, average, pairwise differences within and between species (21). These data are indicative of one or more recent horizontal transfer events between *E. coli* and *S. enterica* and suggest that during formation of *S. enterica* subspecies I the *spv*

operon was obtained from a F-like plasmid pool (21) shared with ancestral *E. coli* isolates and possibly other organisms. As proposed for ancestral *E. coli* strains (87), some of the organisms that contributed to this F-like plasmid pool may have been associated with higher vertebrates long before host adaptations to mammals or birds developed in *S. enterica* subspecies I. For example, the presence on *S. enterica* virulence plasmids of *tlpA*, a gene encoding a regulatory protein that is sensitive to changes from 28°C to physiological temperatures of 37 to 42°C, implies that the donor was already able to sense temperature differences encountered upon entry into warm-blooded vertebrates (58). The identity of this ancestral virulence plasmid donor organism, if extant, remains to be discovered.

ADHESION TO THE MUCOSAL SURFACE AS A MECHANISM FOR HOST ADAPTATION

Salmonella serotypes initiate infection by attaching to the intestinal mucosa of the host. Recent evidence suggests that the recognition of intestinal surfaces by adhesins may contribute to the host adaptation of *Salmonella* serotypes. A nonfimbrial adhesin which may play a role in host adaptation is encoded by *invH*, a gene which is present within SPI 1 in *S. enterica* (22). The idea that *invH* is involved in adherence rests on the finding that mutational inactivation reduces both attachment to and invasion of cultured epithelial cells by *S. enterica* serotypes (4). In contrast, mutations in *invA* and *invE*, two genes that are part of the type III secretory apparatus encoded on SPI 1, render *S. enterica* serotypes noninvasive but have no effect on adhesion to cultured epithelial cells (48, 50). The observation that SPI 1-mediated adhesion and invasion are independent events which can be genetically separated suggests that entry of *S. enterica* into epithelial cells is a two-step process and that InvH may function as an adhesin which mediates an initial attachment step required for invasion (4, 20, 44).

A comparison of the effect on virulence of a mutation in *invH* with those in genes encoding the invasion-associated type III export apparatus illustrates the importance of adherence during interaction with different hosts. *S. enterica* serotype Dublin secretes SopB, a virulence factor which is required for eliciting inflammation and fluid secretion in calves, and its translocation into host cells can be abolished by a polar mutation in *sipB*, a gene located on SPI 1 (49, 121). Similarly, mutations in *invA*, *invB*, *invC*, and *sipC* render *S. enterica* serotypes Enteritidis and Typhimurium avirulent in newly hatched chicks (96, 115). Thus, the type III exporter encoded by SPI 1 is apparently required for *S. enterica* virulence in both chickens and cattle. However, a mutation in the *S. enterica* serotype Typhimurium *invH* gene has no effect on oral virulence or colonization of chicks but significantly reduces the severity of enteritis in calves, suggesting a role of the encoded adhesin in adaptation to bovine but not avian hosts (96, 119, 120).

In chicks, the role of InvH in mediating an initial attachment step required for invasion is likely fulfilled by an alternate adhesin. The various fimbrial operons present in *S. enterica* are possible candidates for encoding such alternate attachment factors involved in host recognition (15, 32). Like binding mediated by InvH, attachment through fimbrial adhesins has been shown to affect entry of *S. enterica* serotype Typhimurium into some epithelial cell lines (4, 14, 43, 57, 64, 112). In a murine intestinal organ culture model, adhesins encoded by the *pef* (plasmid-encoded fimbriae) and *lpf* (long polar fimbriae) operons mediate tissue tropism of *S. enterica* serotype Typhimurium to villous small intestine and Peyer's patches, respectively (13, 16). These data suggest that adhesins help to select which epithelial surfaces are colonized by *S. enterica* in a given animal, a mechanism which is likely to be of importance for determining host range. Analysis of bacterial binding to the mucosa may, in some cases, be complicated by the large number of apparently redundant adhesins. For instance, in the mouse the finding that multiple fimbrial operons are required for full virulence of *S. enterica* serotype Typhimurium suggests that alternative attachment factors can compensate for the loss of a single adhesin (116).

The repertoire of adhesins expressed by a pathogen is thought to determine which structures are recognized and bound on the surface of intestinal epithelial cells. *S. enterica* serotype Typhi utilizes the cystic fibrosis transmembrane conductance regulator as a receptor for internalization by intestinal epithelial cells. In contrast, *S. enterica* serotype Typhimurium appears to use a different epithelial cell receptor for mucosal translocation, because mutations in the cystic fibrosis transmembrane conductance regulator do not reduce the invasiveness of this pathogen for epithelial cell lines or the intestinal wall of transgenic mice (92). This use of different receptors during infection implies the involvement of distinct adhesins which may contribute to differences in host range observed for *S. enterica* serotypes Typhi and Typhimurium. New combinations of adhesion determinants, generated through horizontal gene transfer and deletion events, may thus have contributed to shifts in host range during evolution of *S. enterica* serotypes (12).

THE HOST SPECIFICITY GAMBIT

The process of adapting to a host may involve not only acquisition of virulence determinants, such as adhesins or the virulence plasmid, but also loss of gene function. The *S. enterica* biotypes Gallinarum and Pullorum are both members of the same *S. enterica* subspecies I serotype (antigen formula

1,9,12:-:-) and show host specificity for poultry and aquatic birds. Multilocus enzyme electrophoresis and comparative sequence analysis revealed that *S. enterica* biotypes Gallinarum and Pullorum are closely related (71). It has been speculated that their lineage evolved from an *S. enterica* serotype Enteritidis-like ancestor that had a broad range of hosts, including birds. Since divergence from this ancestor, the ability both to mediate mannose-sensitive hemagglutination (MSHA) and to express flagella (and hence motility) was lost in the *S. enterica* serotype Gallinarum/Pullorum lineage as a result of point mutation (33, 71, 73, 88). Interestingly, strains of *S. enterica* serotype Enteritidis and *S. enterica* serotype Typhimurium that are isolated from cases of avian disease are also frequently non-motile and lack MSHA (30, 38), suggesting that the niche occupied in birds may select against type 1 fimbriation and flagellation. Although it is arguably an advantage during infection of avian hosts, the simultaneous loss of motility and MSHA results in 100-fold-reduced virulence of *S. enterica* serotype Typhimurium in mice (74). It is tempting to speculate that the selection for point mutations in type 1 fimbrial and flagellar biosynthesis genes that occurred during the adaptation to avian hosts may account in part for the loss of virulence in mice in the *S. enterica* serotype Gallinarum/Pullorum lineage (Fig. 2). Thus, in *S. enterica* subspecies I more-complete adaptation to a particular vertebrate host appears to have evolved, in some cases, at the expense of virulence traits that were important for infection of a wider spectrum of animals, resulting in the loss of virulence for some species. The result of this adaptation was a narrowing of the host range and the development of host specificity.

However, the development of host specificity in *S. enterica* biotype Gallinarum cannot be explained by these point mutations alone. Other important changes which may have reduced virulence in mice have also been found. For instance, *S. enterica* serotype Gallinarum/Pullorum appears to be incapable of entering the follicle-associated epithelium of murine Peyer's patches, is internalized by murine macrophages by a mechanism different from that of *S. enterica* serotype Typhimurium, and is unable to survive and multiply in cells of the mouse reticuloendothelial system in vivo and in vitro (3, 8, 91). Thus, loss of virulence is a complex phenotype which involved multiple changes that accumulated in the lineage of *S. enterica* serotype Gallinarum/Pullorum during its evolution toward host specificity.

PERSPECTIVE

Despite the rapid progress in identifying some of the key elements of *S. enterica* virulence by studying the pathogenesis of murine typhoid, our knowledge about genes and mechanisms involved in the expression of host specificity is still very limited. Studies on the distribution of pathogenicity islands, fimbrial operons, and capsular biosynthesis genes among *S. enterica* serotypes suggest that during evolution, new combinations of virulence determinants arose through multiple horizontal transfer events (Fig. 2), a process which may have driven the development of host adaptation. In addition, deletion events and sequence divergence by point mutation were likely among the events which contributed to changes in the host ranges of *S. enterica* serotypes. Thus, adaptation to an animal species is a complex phenotype that doubtless involves a large number of gene products. It is not clear which genetic changes account for the adaptation to a particular mammalian or avian species, and it is likely that many of the responsible virulence mechanisms have not been identified to date. For instance, which virulence factors allowed *S. enterica* subspecies I to

breach the defense formed by regional lymph nodes and to spread to the systemic sites of infection? It is also not known which virulence factors are important during infections that remain localized in the intestine and mesenteric lymph node, such as the gastroenteritis caused by most *S. enterica* subspecies I serotypes in humans. Future research into the genetic basis of host adaptation will provide answers to these and other open questions regarding the pathogenesis of *S. enterica*.

ACKNOWLEDGMENTS

We thank Robert A. Kingsley and Stacy M. Townsend for their comments on the manuscript.

Work in A.J.B.'s laboratory is supported by grant AI40124 from the National Institute of Health and a research grant from USDA Formula Animal Health Funds.

REFERENCES

- Ahmer, B. M. M., J. van Reeuwijk, C. D. Timmers, P. J. Valentine, and F. Heffron. 1998. *Salmonella typhimurium* encodes an SdiA homolog, a putative quorum sensor of the LuxR family, that regulates genes on the virulence plasmid. *J. Bacteriol.* **180**:1185–1193.
- Aleksic, S., F. Heinzerling, and J. Bockemühl. 1996. Human infection caused by salmonellae of subspecies II to VI in Germany, 1977–1992. *Zentbl. Bakteriologie* **283**:391–398.
- Alpuche-Aranda, C. M., E. P. Berthiaume, B. Mock, J. A. Swanson, and S. I. Miller. 1995. Spacious phagosome formation within mouse macrophages correlates with *Salmonella* serotype pathogenicity and host susceptibility. *Infect. Immun.* **63**:4456–4462.
- Altmeyer, R. M., J. K. McNern, J. C. Bossio, I. Rosenshine, B. B. Finlay, and J. E. Galán. 1993. Cloning and molecular characterization of a gene involved in *Salmonella* adherence and invasion of cultured epithelial cells. *Mol. Microbiol.* **7**:89–98.
- Anderson, R. M., and R. M. May. 1982. Coevolution of host and parasites. *Parasitology* **85**:411–426.
- Baird, G. D., E. J. Manning, and P. W. Jones. 1985. Evidence for related virulence sequences in plasmids of *Salmonella dublin* and *Salmonella typhimurium*. *J. Gen. Microbiol.* **131**:1815–1823.
- Barnes, D. M., and M. E. Bergland. 1968. *Salmonella typhisuis* infection in Minnesota swine. *J. Am. Vet. Med. Assoc.* **152**:1766–1770.
- Barrow, P. A., M. B. Huggins, and M. A. Lovell. 1994. Host specificity of *Salmonella* infection in chickens and mice is expressed in vivo primarily at the level of the reticuloendothelial system. *Infect. Immun.* **62**:4602–4610.
- Barrow, P. A., and M. A. Lovell. 1988. The association between a large molecular mass plasmid and virulence in a strain of *Salmonella pullorum*. *J. Gen. Microbiol.* **134**:2307–2316.
- Barrow, P. A., J. M. Simpson, M. A. Lovell, and M. M. Binn. 1987. Contribution of *Salmonella gallinarum* large plasmid toward virulence in fowl typhoid. *Infect. Immun.* **55**:388–392.
- Bäumler, A. J. 1997. The record of horizontal gene transfer in *Salmonella*. *Trends Microbiol.* **5**:318–322.
- Bäumler, A. J., A. J. Gilde, R. M. Tsolis, A. W. M. van der Velden, B. M. M. Ahmer, and F. Heffron. 1997. Contribution of horizontal gene transfer and deletion events to the development of distinctive patterns of fimbrial operons during evolution of *Salmonella* serotypes. *J. Bacteriol.* **179**:317–322.
- Bäumler, A. J., R. M. Tsolis, F. Bowe, J. G. Kusters, S. Hoffmann, and F. Heffron. 1996. The *pef* fimbrial operon mediates adhesion to murine small intestine and is necessary for fluid accumulation in infant mice. *Infect. Immun.* **64**:61–68.
- Bäumler, A. J., R. M. Tsolis, and F. Heffron. 1996. Contribution of fimbrial operons to attachment to and invasion of epithelial cell lines by *Salmonella typhimurium*. *Infect. Immun.* **64**:1862–1865.
- Bäumler, A. J., R. M. Tsolis, and F. Heffron. 1997. Fimbrial adhesins of *Salmonella typhimurium*—role in bacterial interactions with epithelial cells, p. 149–158. *In* P. S. Paul, D. H. Francis, and D. Benfield (ed.), *Mechanisms in the pathogenesis of enteric diseases*, vol. 412. Plenum Press, New York, N.Y.
- Bäumler, A. J., R. M. Tsolis, and F. Heffron. 1996. The *lpf* fimbrial operon mediates adhesion to murine Peyer's patches. *Proc. Natl. Acad. Sci. USA* **93**:279–283.
- Bäumler, A. J., R. M. Tsolis, P. J. Valentine, T. A. Ficht, and F. Heffron. 1997. Synergistic effect of mutations in *invA* and *lpfC* on the ability of *Salmonella typhimurium* to cause murine typhoid. *Infect. Immun.* **65**:2254–2259.
- Beninger, P. R., G. Chikami, K. Tanabe, C. Roudier, J. Fierer, and D. G. Guiney. 1988. Physical and genetic mapping of the *Salmonella dublin* virulence plasmid pSDL2. Relationship to plasmids from other *Salmonella* strains. *J. Clin. Investig.* **81**:1341–1347.
- Blaser, M. J., and R. A. Feldman. 1981. *Salmonella* bacteremia: reports from the centers of disease control and prevention. *J. Infect. Dis.* **143**:743–746.
- Bliska, J. B., J. E. Galán, and S. Falkow. 1993. Signal transduction in the mammalian cell during bacterial attachment and entry. *Cell* **73**:903–920.
- Boyd, E. F., and D. L. Hartl. 1997. Recent horizontal transmission of plasmids between natural populations of *Escherichia coli* and *Salmonella enterica*. *J. Bacteriol.* **179**:1622–1627.
- Boyd, E. F., J. Li, H. Ochman, and R. K. Selander. 1997. Comparative genetics of the *inv-spa* invasion gene complex of *Salmonella enterica*. *J. Bacteriol.* **179**:1985–1991.
- Boyd, E. F., F.-S. Wang, P. Beltran, S. A. Plock, K. Nelson, and R. K. Selander. 1993. *Salmonella* reference collection B (SARB): strains of 37 serovars of subspecies I. *J. Gen. Microbiol.* **139**:1125–1132.
- Boyd, E. F., F.-S. Wang, T. S. Whittam, and R. K. Selander. 1996. Molecular genetic relationship of the salmonellae. *Appl. Environ. Microbiol.* **62**:804–808.
- Buisán, M., J. M. Rodríguez-Pena, and R. Rotger. 1994. Restriction map of the *Salmonella enteritidis* virulence plasmid and its homology with the plasmid of *Salmonella typhimurium*. *Microb. Pathog.* **16**:165–169.
- Burnet, M., and D. O. White. 1972. *Natural history of infectious disease*. Cambridge University Press, Cambridge, United Kingdom.
- Buxton, A., and G. Fraser. 1977. *Animal microbiology*, vol. 1. Blackwell Scientific Publications, London, United Kingdom.
- Carter, P. B., and F. M. Collins. 1974. The route of enteric infection in normal mice. *J. Exp. Med.* **139**:1189–1203.
- Cerin, H., and J. Hackett. 1993. The *parVP* region of the *Salmonella typhimurium* virulence plasmid pSLT contains four loci required for incompatibility and partitioning. *Plasmid* **30**:30–38.
- Chart, H., D. Conway, and B. Rowe. 1993. Outer membrane characteristics of *Salmonella enteritidis* phage type-4 growing in chickens. *Epidemiol. Infect.* **111**:449–454.
- Christensen, H., S. Nordentoft, and J. E. Olsen. 1998. Phylogenetic relationships of *Salmonella* based on rRNA sequences. *Int. J. System. Bacteriol.* **48**:605–610.
- Clegg, S., and D. L. Swenson. 1994. *Salmonella* fimbriae, p. 105–113. *In* P. Klemm (ed.), *Fimbriae: adhesion, genetics, biogenesis, and vaccines*. CRC Press, Boca Raton, Fla.
- Crichton, P. B., D. E. Yakubu, D. C. Old, and S. Clegg. 1989. Immunological and genetical relatedness of type 1 and type 2 fimbriae in salmonellas of serotype Gallinarum, Pullorum and Typhimurium. *J. Appl. Bacteriol.* **67**:283–291.
- Danbara, H., R. Moriguchi, S. Suzuki, Y. Tamura, M. Kijima, K. Oishi, H. Matsui, A. Abe, and M. Nakamura. 1992. Effect of 50 kilobase-plasmid, pKDSC50, of *Salmonella choleraesuis* RF-1 strain on pig septicemia. *J. Vet. Med. Sci.* **54**:1175–1178.
- Dimow, I. 1966. Versuche Landschidkröten der Arten *Testudo graeca* und *Testudo hermanni* mit *Salmonella*-bakterien zu infizieren. *Zentbl. Bakteriologie* **199**:181–184.
- Dimow, I. 1966. Versuche zur künstlichen Infektion von Eidechsen mit *Salmonella* und *Arizona*-bakterien. *Zentbl. Vetmed. Reihe B* **13**:587–590.
- Dimow, I., and R. Salwtschew. 1967. Versuche der Experimentalinfizierung von Schlangen *Vipera ammodytes* mit *Salmonella* und *Arizona*-bakterien. *Pathol. Microbiol.* **30**:495–497.
- Duguid, J. P., E. S. Anderson, G. A. Alfredsson, R. Barker, and D. C. Old. 1975. A new biotyping scheme for *Salmonella typhimurium* and its phylogenetic significance. *J. Med. Microbiol.* **8**:149–166.
- Du Pasquier, L. 1982. Antibody diversity in lower vertebrates—why is it so restricted? *Nature* **290**:311–313.
- Du Pasquier, L. 1993. Phylogeny of B-cell development. *Curr. Opin. Immunol.* **5**:185–193.
- Du Pasquier, L. 1989. Evolution of the immune system, p. 139–165. *In* W. E. Paul (ed.), *Fundamental immunology*, 2nd ed. Raven Press, Ltd., New York, N.Y.
- Du Pasquier, L. 1992. Origin and evolution of the vertebrate immune system. *APMIS* **100**:383–392.
- Ernst, R. K., D. M. Dombroski, and J. M. Merrick. 1990. Anaerobiosis, type 1 fimbriae, and growth phase are factors that affect invasion of HEp-2 cells by *Salmonella typhimurium*. *Infect. Immun.* **58**:2014–2016.
- Falkow, S., R. R. Isberg, and D. A. Portnoy. 1992. The interaction of bacteria with mammalian cells. *Annu. Rev. Cell Biol.* **8**:333–363.
- Fields, P. I., R. V. Swanson, C. G. Haidaris, and F. Heffron. 1986. Mutants of *Salmonella typhimurium* that cannot survive within the macrophage are avirulent. *Proc. Natl. Acad. Sci. USA* **83**:5189–5193.
- Fierer, J., M. Krause, R. Tauxe, and D. Guiney. 1992. *Salmonella typhimurium* bacteremia: association with the virulence plasmid. *J. Infect. Dis.* **166**:639–642.
- Friedrich, M. J., N. E. Kinsey, J. Vila, and R. J. Kadner. 1993. Nucleotide sequence of a 13.9 kb segment of the 90 kb virulence plasmid of *Salmonella typhimurium*: the presence of fimbrial biosynthetic genes. *Mol. Microbiol.* **8**:543–558.
- Galán, J. E., and R. Curtiss III. 1989. Cloning and molecular characterization of genes whose products allow *Salmonella typhimurium* to penetrate

- tissue culture cells. Proc. Natl. Acad. Sci. USA **86**:6383–6387.
49. Galyov, E. E., M. W. Wood, R. Rosqvist, P. B. Mullan, P. R. Watson, S. Hedges, and T. S. Wallis. 1997. A secreted effector protein of *Salmonella* dublin is translocated into eukaryotic cells and mediates inflammation and fluid secretion in infected ileal mucosa. Mol. Microbiol. **25**:903–912.
 50. Ginocchio, C., J. Pace, and J. E. Galán. 1992. Identification and characterization of a *Salmonella typhimurium* gene involved in triggering the internalization of salmonellae into cultured epithelial cells. Proc. Natl. Acad. Sci. USA **89**:5976–5980.
 51. Glick, B. 1982. RES structure and function in aves, p. 509–540. In N. Cohen, and M. M. Sigel (ed.), The reticuloendothelial system, vol. 3. Plenum Press, New York, N.Y.
 52. Gulig, P. A., Caldwell, A. L., and Chiodo, V. A. 1992. Identification, genetic analysis and DNA sequence of a 7.8-kb virulence region of the *Salmonella typhimurium* virulence plasmid. Mol. Microbiol. **6**:1395–1411.
 53. Gulig, P. A., and R. Curtiss. 1987. Plasmid-associated virulence of *Salmonella typhimurium*. Infect. Immun. **1987**:2891–2901.
 54. Gulig, P. A., H. Danbara, D. G. Guiney, A. J. Lax, F. Norel, and M. Rhen. 1993. Molecular analysis of *spv* virulence genes of the *Salmonella* virulence plasmids. Mol. Microbiol. **7**:825–830.
 55. Hensel, M., J. E. Shea, A. J. Bäuml, C. Gleeson, F. Blattner, and D. W. Holden. 1997. Analysis of the boundaries of *Salmonella* pathogenicity island 2 and the corresponding chromosomal region of *Escherichia coli* K-12. J. Bacteriol. **179**:1105–1111.
 56. Hoffman, T. A., C. J. Ruiz, G. W. Counts, J. M. Sachs, and J. L. Nitzkin. 1975. Waterborne typhoid fever in Dade County, Florida. Clinical and therapeutic evaluation of 105 bacteremic patients. Am. J. Med. **59**:481–487.
 57. Horiuchi, S., Y. Inagaki, N. Okamura, R. Nakaya, and N. Yamamoto. 1992. Type 1 pili enhance the invasion of *Salmonella baendrup* and *Salmonella typhimurium* to HeLa cells. Microbiol. Immunol. **36**:593–602.
 58. Hurme, R., K. D. Berndt, S. J. Normark, and M. Rhen. 1997. A proteinaceous gene regulatory thermometer in salmonella. Cell **90**:55–64.
 59. Inouye, S., M. G. Sunshine, E. W. Six, and M. Inouye. 1991. Retrophage Φ R73: an *E. coli* phage that contains a retroelement and integrates into a t-RNA gene. Science **252**:969–971.
 60. Ishibashi, Y., and T. Arai. 1996. A possible mechanism for host-specific pathogenesis of *Salmonella* serovars. Microb. Pathog. **21**:435–446.
 61. Ishibashi, Y., and T. Arai. 1990. Roles of the complement receptor type 1 (CR1) and type 3 (CR3) on phagocytosis and subsequent phagosome-lysosome fusion in *Salmonella*-infected murine macrophages. FEMS Microbiol. Lett. **64**:89–96.
 62. Jones, B. D., N. Gori, and S. Falkow. 1994. *Salmonella typhimurium* initiates murine infection by penetrating and destroying the specialized epithelial M cells of the Peyer's patches. J. Exp. Med. **180**:15–23.
 63. Jones, G. W., Rabert, D. K., Svinarich, M., and Whitfield, H. J. 1982. Association of adhesive, invasive, and virulent phenotypes of *Salmonella typhimurium* with autonomous 60-megadalton plasmids. Infect. Immun. **38**:476–86.
 64. Jones, G. W., and L. A. Richardson. 1981. The attachment to and invasion of HeLa cells by *Salmonella typhimurium*: the contribution of mannose-sensitive and mannose-resistant haemagglutinating activities. J. Gen. Microbiol. **127**:361–370.
 65. Jonsson, V. 1985. Comparison and definition of spleen and lymph node: a phylogenetic analysis. J. Theor. Biol. **117**:691–699.
 66. Korpela, K., M. Ranki, S. Sukupolvi, P. H. Mäkelä, and M. Rehn. 1989. Occurrence of *Salmonella typhimurium* virulence plasmid-specific sequences in different serovars of *Salmonella*. FEMS Microbiol. Lett. **58**:49–54.
 67. Lan, R. T., and P. R. Reeves. 1996. Gene transfer is a major factor in bacterial evolution. Mol. Biol. Evol. **13**:47–55.
 68. Lax, A. J., G. D. Pullinger, G. D. Baird, and C. M. Williamson. 1990. The virulence plasmid of *Salmonella dublin*: detailed restriction map and analysis by transposon mutagenesis. J. Gen. Microbiol. **136**:1117–1123.
 69. Le Minor, L., and M. Y. Popoff. 1987. Designation of *Salmonella enterica* sp. nov., nom. rev., as the type and only species of the genus *Salmonella*. Int. J. Syst. Bacteriol. **37**:465–468.
 70. Li, J., H. Ochman, E. A. Groisman, E. F. Boyd, F. Solomon, K. Nelson, and R. K. Selander. 1995. Relationship between evolutionary rate and cellular location among the Inv/Spa invasion proteins of *Salmonella enterica*. Proc. Natl. Acad. Sci. USA **92**:7252–7256.
 71. Li, J., N. H. Smith, K. Nelson, P. B. Crichton, D. C. Old, T. S. Whittam, and R. K. Selander. 1993. Evolutionary origin and radiation of the avian-adapted non-motile salmonellae. J. Med. Microbiol. **38**:129–139.
 72. Libby, S. J., L. G. Adams, T. A. Ficht, C. Allen, H. A. Whitford, N. A. Buchmeier, S. Bossie, and D. G. Guiney. 1997. The *spv* genes of the *Salmonella dublin* virulence plasmid are required for severe enteritis and systemic infection in the natural host. Infect. Immun. **65**:1786–1792.
 73. Lockman, H. A., and R. Curtiss III. 1992. Isolation and characterization of conditional adherent and non-type 1 fimbriated *Salmonella typhimurium* mutants. Mol. Microbiol. **6**:933–945.
 74. Lockman, H. A., and R. Curtiss III. 1992. Virulence of non-type 1-fimbriated and nonfimbriated nonflagellated *Salmonella typhimurium* mutants in murine typhoid fever. Infect. Immun. **60**:491–496.
 75. Mahon, J., and A. J. Lax. 1993. A quantitative polymerase chain reaction method for the detection in avian faeces of *Salmonellas* carrying the *spvR* gene. Epidemiol. Infect. **111**:455–464.
 76. Manning, M. J. 1979. Evolution of the vertebrate immune system. J. R. Soc. Med. **72**:683–688.
 77. McCormick, B. A., S. I. Miller, D. Carnes, and J. L. Madara. 1995. Trans-epithelial signaling to neutrophils by salmonellae: a novel virulence mechanism for gastroenteritis. Infect. Immun. **63**:2302–2309.
 78. Miller, S. L., E. L. Hohmann, and D. A. Pegues. 1995. *Salmonella* (including *Salmonella typhi*), p. 2013–2033. In G. L. Mandell, J. E. Bennett, and R. Dolin (ed.), Principles and practice of infectious diseases, 4th ed., vol. 2. Churchill Livingstone, New York, N.Y.
 79. Mills, D. M., V. Bajaj, and C. A. Lee. 1995. A 40kb chromosomal fragment encoding *Salmonella typhimurium* invasion genes is absent from the corresponding region of the *Escherichia coli* K-12 chromosome. Mol. Microbiol. **15**:749–759.
 80. Montenegro, M. A., G. Morelli, and R. Helmuth. 1991. Heteroduplex analysis of *Salmonella* virulence plasmids and their prevalence in isolates of defined sources. Microb. Pathog. **11**:391–397.
 81. Musher, D. M., and A. D. Rubenstein. 1973. Permanent carriers of nontyphosal *Salmonellae*. Public Health Rep. **132**:869.
 82. Muthukkaruppan, V. R., M. Borysenko, and R. El Ridi. 1982. RES structure and function in reptilia, p. 461–508. In N. Cohen and M. M. Sigel (ed.), The reticuloendothelial system, vol. 3. Plenum Press, New York, N.Y.
 83. Nakamura, M., S. Sato, T. Ohya, S. Suzuki, and S. Ikeda. 1985. Possible relationship of a 36-megadalton *Salmonella enteritidis* plasmid to virulence in mice. Infect. Immun. **47**:831–833.
 84. Ochman, H., and E. A. Groisman. 1996. Distribution of pathogenicity islands in *Salmonella* spp. Infect. Immun. **64**:5410–5412.
 85. Ochman, H., and E. A. Groisman. 1995. The evolution of invasion by enteric bacteria. Can. J. Microbiol. **41**:555–561.
 86. Ochman, H., F. C. Soncini, F. Solomon, and E. A. Groisman. 1996. Identification of a pathogenicity island for *Salmonella* survival in host cells. Proc. Natl. Acad. Sci. USA **93**:7800–7804.
 87. Ochman, H., and A. C. Wilson. 1987. Evolution in bacteria: evidence for a universal substitution rate in cellular genomes. J. Mol. Evol. **26**:74–86.
 88. Old, D. C., and S. B. Payne. 1971. Antigens of the type 2 fimbriae of salmonellae: "cross reacting material" (CRM) of type 1 fimbriae. J. Med. Microbiol. **4**:215–225.
 89. Ou, J. T., M.-Y. Lin, and H.-L. Chao. 1994. Presence of F-like ori-T base-pair sequences on the virulence plasmids of *Salmonella* serovars Gallinarum, Enteritidis, and Typhimurium, but absent in those of Choleraesuis and Dublin. Microbiol. Pathog. **17**:13–21.
 90. Pardon, P., R. Sanchis, J. Marly, F. Lantier, M. Papin, and M. Popoff. 1988. Salmonellosis ovine due a *Salmonella abortusovis*. Ann. Rech. Vet. **19**:221–235.
 91. Pascopella, L., B. Raupach, N. Gori, D. Monack, S. Falkow, and P. L. Small. 1995. Host restriction phenotypes of *Salmonella typhi* and *Salmonella gallinarum*. Infect. Immun. **63**:4329–4335.
 92. Pier, G. B., M. Grout, T. Zaidi, G. Meluleni, S. S. Mueschenborn, G. Banting, R. Ratcliff, M. J. Evans, and W. H. Colledge. 1998. *Salmonella typhi* uses CFTR to enter intestinal epithelial cells. Nature **393**:79–82.
 93. Popoff, M. Y., and L. Le Minor. 1992. Antigenic formulas of the *Salmonella* serovars, 5th ed. WHO Collaborating Center for Reference and Research on *Salmonella*, Institute Pasteur, Paris, France.
 94. Popoff, M. Y., I. Miras, C. Coynault, C. Lasselien, and P. Pardon. 1984. Molecular relationships between virulence plasmids of *salmonella* serotypes typhimurium and dublin and large plasmids of other *Salmonella* serotypes. Ann. Microbiol. **135A**:389–398.
 95. Poppe, C., R. Curtiss III, P. A. Gulig, and C. L. Gyles. 1989. Hybridization with a DNA probe derived from the virulence region of the 60 Mdal plasmid of *Salmonella typhimurium*. Can. J. Vet. Res. **53**:378.
 96. Porter, S. B., and R. Curtiss III. 1997. Effect of *inv* mutations on *Salmonella* virulence and colonization in 1-day-old white leghorn chicks. Avian Dis. **41**:45–57.
 97. Reeves, M. W., G. M. Evins, A. A. Heiba, B. D. Plikaytis, and J. J. Farmer III. 1989. Clonal nature of *Salmonella typhi* and its genetic relatedness to other salmonellae as shown by multilocus enzyme electrophoresis, and proposal of *Salmonella bongori* comb. nov. J. Clin. Microbiol. **27**:313–320.
 98. Rexach, L., F. Dilassier, and P. Fach. 1994. Polymerase chain reaction for *salmonella* virulence-associated plasmid genes: a new tool in salmonella epidemiology. Epidemiol. Infect. **112**:33–43.
 99. Rice, D. H., T. E. Besser, and D. D. Hancock. 1997. Epidemiology and virulence assessment of *Salmonella dublin*. Vet. Microbiol. **56**:111–124.
 100. Rodriguezpena, J. M., I. Alvarez, M. Ibanez, and R. Rotger. 1997. Homologous regions of the salmonella enteritidis virulence plasmid and the chromosome of salmonella typhi encode thiol, disulphide oxidoreductases belonging to the dsba thioredoxin family. Microbiology **143**:1405–1413.
 101. Rodriguezpena, J. M., M. Buisan, M. Ibanez, and R. Rotger. 1997. Genetic map of the virulence plasmid of salmonella enteritidis and nucleotide sequence of its replicons. Gene **188**:53–61.

102. Roggendorf, M., and H. E. Muller. 1976. Enterobacteria from reptiles. Zentbl. Bakteriol. Parasitenkd. Hyg. Abt. I Orig. A **236**:22–35.
103. Roudier, C., M. Krause, J. Fierer, and D. G. Guiney. 1990. Correlation between the presence of sequences homologous to the vir region of *Salmonella dublin* plasmid pSDL2 and the virulence of twenty-two *Salmonella* serotypes in mice. Infect. Immun. **58**:1180–1185.
104. Roy, J. H. B. 1990. Salmonellosis, p. 118–130. In J. H. B. Roy (ed.), The calf, vol. 1 Management of Health, Butterworths, London, United Kingdom.
105. Rychlik, I., M. A. Lovell, and P. A. Barrow. 1998. The presence of genes homologous to the K88 genes *faeH* and *faeI* on the virulence plasmid of *Salmonella gallinarum*. FEMS Microbiol. Lett. **159**:255–260.
106. Schwartz, K. J. 1991. Salmonellosis in swine. Compend. Contin. Educ. Pract. Vet. **13**:139–146.
107. Selander, R. K., P. Beltran, N. H. Smith, R. Helmuth, F. A. Rubin, D. J. Kopecko, K. Ferris, B. D. Tall, A. Cravioto, and J. M. Musser. 1990. Evolutionary genetic relationships of clones of *Salmonella* serovars that cause human typhoid and other enteric fevers. Infect. Immun. **58**:2262–2275.
108. Selander, R. K., J. Li, E. F. Boyd, F.-S. Wang, and K. Nelson. 1994. DNA sequence analysis of the genetic structure of populations of *Salmonella enterica* and *Escherichia coli*, p. 17–49. In F. G. Priest, A. Ramos-Cormenzana, and B. J. Tindall (ed.), Bacterial diversity and systematics. Plenum Press, New York, N.Y.
109. Shea, J. E., M. Hensel, C. Gleeson, and D. W. Holden. 1996. Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium*. Proc. Natl. Acad. Sci. USA **93**:2593–2597.
110. Singh, I. P., V. K. Sharma, and Y. K. Kaura. 1971. Some aspects of the epidemiology of *Salmonella abortus-ovis* infection in equines. Br. Vet. J. **127**:378–383.
111. Stoll, L. 1962. Experimentelle Infektionen mit Keimen der *Salmonella* und *Arizona* Gruppe bei Schlangen. Nord. Vet. Med. **14**:225–232.
112. Stone, B. J., C. M. Garcia, J. L. Badger, T. Hassett, R. I. F. Smith, and V. Miller. 1992. Identification of novel loci affecting entry of *Salmonella enteritidis* into eukaryotic cells. J. Bacteriol. **174**:3945–3952.
113. Tinge, A. S., and R. Curtiss III. 1990. Isolation of the replication and partitioning regions of the *Salmonella typhimurium* virulence plasmid and stabilization of heterologous replicons. J. Bacteriol. **172**:5266–5277.
114. Tinge, S. A., and Curtiss, R. 1990. Conservation of *Salmonella typhimurium* virulence plasmid maintenance regions among *Salmonella* serovars as a basis for plasmid curing. Infect. Immun. **58**:3084–3092.
115. Turner, A. K., A. Lovell, S. D. Hulme, L. Zhang-Barber, and P. A. Barrow. 1998. Identification of *Salmonella typhimurium* genes required for colonization of the chicken alimentary tract and for virulence in newly hatched chicks. Infect. Immun. **66**:2099–2106.
116. van der Velden, A. W. M., A. J. Bäumler, R. M. Tsois, and F. Heffron. 1998. Multiple fimbrial adhesins are required for full virulence of *Salmonella typhimurium* in mice. Infect. Immun. **66**:2803–2808.
117. Vladioianu, I.-R., H. R. Chang, and J.-C. Pechère. 1990. Expression of host resistance to *Salmonella typhi* and *salmonella typhimurium*: bacterial survival within macrophages of murine and human origin. Microb. Pathog. **8**:83–90.
118. Wallis, T. S., S. M. Paulin, J. S. Pledsted, P. R. Watson, and P. W. Jones. 1995. The *Salmonella dublin* virulence plasmid mediates systemic but not enteric phases of salmonellosis in cattle. Infect. Immun. **63**:2755–2761.
119. Watson, P. R., E. E. Galyov, S. M. Paulin, P. W. Jones, and T. S. Wallis. 1998. Mutation of *invH*, but not *stn*, reduces salmonella-induced enteritis in cattle. Infect. Immun. **66**:1432–1438.
120. Watson, P. R., S. M. Paulin, A. P. Bland, P. W. Jones, and T. S. Wallis. 1995. Characterization of intestinal invasion by *Salmonella typhimurium* and *Salmonella dublin* and effect of a mutation in the *invH* gene. Infect. Immun. **63**:2743–54.
121. Wood, M. W., R. Rosqvist, P. B. Mullan, M. H. Edwards, and E. E. Galyov. 1996. SopE, a secreted protein of *Salmonella dublin*, is translocated into the target eukaryotic cell via a *sip*-dependent mechanism and promotes bacterial entry. Mol. Microbiol. **22**:327–338.
122. Woodward, M. J., McLaren, I., and Wray, C. 1989. Distribution of virulence plasmids within salmonellae. J. Gen. Microbiol. **135**:503–511.
123. Zapata, A. G., M. Torroba, A. Vicente, A. Varas, R. Sacedon, and E. Jimenez. 1995. The relevance of cell microenvironments for the appearance of lymphohaemopoietic tissues in primitive vertebrates. Histol. Histopathol. **10**:761–778.

Editor: P. E. Orndorff