



Published in final edited form as:

Nat Chem. 2024 February ; 16(2): 285–293. doi:10.1038/s41557-023-01342-7.

Programmable late-stage functionalization of bridge-substituted bicyclo[1.1.1]pentane bis-boronates

Yangyang Yang^{1,6}, Jet Tsien^{1,6}, Ryan Dykstra², Si-Jie Chen³, James B. Wang¹, Rohan R. Merchant³, Jonathan M. E. Hughes⁴, Byron K. Peters⁴, Osvaldo Gutierrez^{2,5,✉}, Tian Qin^{1,✉}

¹Department of Biochemistry, The University of Texas Southwestern Medical Center, Dallas, TX, USA.

²Department of Chemistry and Biochemistry, University of Maryland, College Park, MD, USA.

³Department of Discovery Chemistry, Merck & Co., Inc., South San Francisco, CA, USA.

⁴Department of Process Research and Development, Merck & Co., Inc., Rahway, NJ, USA.

⁵Department of Chemistry, Texas A&M University, College Station, TX, USA.

⁶These authors contributed equally: Yangyang Yang, Jet Tsien.

Abstract

Modular functionalization enables versatile exploration of chemical space and has been broadly applied in structure–activity relationship (SAR) studies of aromatic scaffolds during drug discovery. Recently, the bicyclo[1.1.1]pentane (BCP) motif has increasingly received attention as a bioisosteric replacement of benzene rings due to its ability to improve the physicochemical properties of prospective drug candidates, but studying the SARs of C₂-substituted BCPs has been heavily restricted by the need for multistep de novo synthesis of each analogue of interest. Here we report a programmable bis-functionalization strategy to enable late-stage sequential derivatization of BCP bis-boronates, opening up opportunities to explore the SARs of drug candidates possessing multisubstituted BCP motifs. Our approach capitalizes on the

Reprints and permissions information is available at www.nature.com/reprints.

✉ Correspondence and requests for materials should be addressed to Osvaldo Gutierrez or Tian Qin., og.labs@tamu.edu; tian.qin@utsouthwestern.edu.

Author contributions

Y.Y. and J.T. performed synthetic experiments; R.D., J.B.W. and O.G. performed DFT theoretical studies. S.-J.C. performed PMI and 3D score calculations. J.M.E.H., B.K.P., R.R.M. and T.Q. designed and supervised the project; Y.Y., J.T., R.D., S.-J.C., J.B.W., J.M.E.H., B.K.P., R.R.M., O.G. and T.Q. contributed to the manuscript writing.

Competing interests

The authors declare the following competing financial interest(s): T.Q., Y.Y. and J.T. from UT Southwestern Medical Center are listed as inventors on US patent application no. 63/146,266, which covers the ‘synthesis of BCP bis-boronates’ in the manuscript, and on the US provisional application no. 63/321,700, which covers ‘C3-functionalization of BCP bis-boronates’ and ‘late-stage C2-functionalization of BCP boronates’ in the manuscript. The remaining authors declare no competing interests.

Additional information

Extended data is available for this paper at <https://doi.org/10.1038/s41557-023-01342-7>.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41557-023-01342-7>.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

inherent chemoselectivity exhibited by BCP bis-boronates, enabling highly selective activation and functionalization of bridgehead (C_3)-boronic pinacol esters (Bpin), leaving the C_2 -Bpin intact and primed for subsequent derivatization. These selective transformations of both BCP bridgehead (C_3) and bridge (C_2) positions enable access to C_1 , C_2 -disubstituted and C_1 , C_2 , C_3 -trisubstituted BCPs that encompass previously unexplored chemical space.

Infinite chemical space provides boundless opportunities for medicinal chemists, within which three-dimensional scaffolds with drug-like properties are particularly sought after^{1–3}. The past decade has witnessed growing interest in the chemical space of bicyclo[1.1.1]pentane (BCP), a class of motifs that are considered medicinally relevant as a three-dimensional bioisosteric replacement of aromatic moieties (Fig. 1a)^{4–11}. Such consideration was driven by its potential to improve physicochemical, pharmacological and toxicological properties of drug candidates^{6,10,11}, with bridgehead (C_1 and C_3)-substituted BCPs widely recognized as saturated bioisosteres for *mono*- and *para*-substituted benzenes in synthetic chemistry^{12–25}. Analogously, it has been hypothesized that C_1 , C_2 -disubstituted BCPs could represent bioisosteres of *ortho*- and *meta*-substituted benzenes^{10,26–28}. However, despite the recent reports from Baran et al.²⁸, Ma et al.^{29,30}, our group³¹ and others^{32–34}, efficient synthesis of bridge-substituted BCPs still proves challenging and the practical applications of such BCPs in medicinal chemistry remain heavily constrained. In contrast to $C(sp^2)$ cross-coupling chemistry, which enables rapid diversification of arenes³⁵, derivatization of the $C(sp^3)$ -rich bioisostere BCPs requires multistep *de novo* synthesis (Fig. 1b)³¹. Such constraints represent the biggest bottleneck to access and evaluate these promising compounds in structure–activity relationship (SAR) studies. Thus, development of a modular and programmable strategy to access C_1 , C_2 -disubstituted and C_1 , C_2 , C_3 -trisubstituted BCPs should enable more comprehensive SAR campaigns and chemical space exploration.

The inert C–H bonds within BCPs not only provide metabolic stability for this three-dimensional scaffold^{36–38}, but also create the synthetic hurdles to generally access bridge (C_2)-substituted derivatives via C–H functionalization²⁷. Despite our earlier success in multisubstituted BCP syntheses via intramolecular coupling strategy³¹, the installation of the bridge-substitution group before cyclization substantially increased the synthetic steps required to access each analogue. To address these issues, our goal was to identify a late-stage BCP intermediate with two functional handles that could be selectively and rapidly modified to pursue divergent SAR exploration. To that end, our group previously reported³¹ the preparation of two C_1 -alkyl BCPs bearing Bpin substituents at both C_2 and C_3 . Inspired by breakthrough reports by Morken^{39–41}, Aggarwal^{42,43} and others^{44,45}, we postulated that these BCP bis-boronates could be selectively and sequentially functionalized (Fig. 1c) and this synthetic approach would provide rapid access to bridge-substituted C_1 , C_2 -di- and C_1 , C_2 , C_3 -tri-substituted BCPs. Here we describe a strategy that leverages the endogenous properties of BCP bis-boronates to enable the modular and efficient synthesis of multisubstituted BCPs via sequential functionalization.

Results and discussion

The striking reactivity difference between C₂- and C₃-Bpin units was initially observed in the boronate ligand exchange reaction with BCP **13** (Fig. 2a). Under treatment of 1,8-diaminonaphthalene in toluene, selective formation of C₃-Bdan-substituted product **15** was identified by NMR analysis and confirmed by single-crystal X-ray diffraction. In comparison, the C₂-cyclopentyl-substituted BCP **14** afforded much lower yield of the corresponding product (**16**, 7%). In a similar vein, this chemoselectivity was also observed in the hydrazone coupling⁴⁶. The bridgehead C₃-Bpin in BCP bis-boronates displayed an enhanced reactivity and underwent coupling with sulfonyl hydrazone **17**, whereas C₂-Bpin in bis-boronates and BCP **14** remained mostly unreacted (Fig. 2a). We initially hypothesized that—as noted by Morken^{39,40}, and others^{42,43,44,45}—selective activation at the C₃ position in this bis-boronate system is probably due to intramolecular coordination (for example, an oxygen lone pair–boron interaction) of the Bpin at the C₂ position; however, careful analysis of B₃–C₃–C₁ angles of these BCP X-ray crystal structures (see Fig. 3 and ‘X-ray crystallographic data for BCP compounds’ section in the Supplementary Information) revealed no obvious initial interaction between the two adjacent Bpin groups in the BCP scaffolds, presumably because the strained bicyclic scaffold restricts the spatial orientations of these groups (Fig. 2a).

Quantum mechanical calculations

We turned to quantum mechanical calculations to gain further insights into the selective reactivity of BCP bis-boronates. First, analysis of the lowest energy conformations of BCP bis-boronate **13** found no apparent Bpin–Bpin non-covalent interactions, consistent with the crystal structures (as determined by natural bond orbital analysis and O–B distances; see Supplementary Information for details). Comparison between the putative anionic intermediates generated revealed an approximately tenfold thermodynamic preference for the C₃ (bridgehead) 3° anionic intermediate ($G_{\text{rel}} \mathbf{13}^{\text{C}2\text{-}/\text{C}3\text{-}} = +1.8 \text{ kcal mol}^{-1}$; Fig. 2b). In contrast, DFT calculations show an energetic preference for the formation of the 2° alkyl radical, which might occur via C₂-Bpin homolysis ($G_{\text{rel}} \mathbf{13}^{\text{C}2\text{·}/\text{C}3\text{·}} = -3.0 \text{ kcal mol}^{-1}$; Fig. 2b). Such results can be rationalized by the higher *s*-orbital-character of the C₃ carbon (behaving as *sp*²) compared with the C₂ carbon (*sp*^{2.5})^{37,38}, leading to a higher electronegativity at the bridgehead position and an increased Lewis acidity of the attached Bpin. Given that exclusive reactivity at the bridgehead position was observed for all transformations (see below), these calculations support an anion-mediated C–B activation pathway (Fig. 2c).

Further study probed into the bond-forming step, in which we modelled the hydrazone coupling reaction⁴⁶ between bis-boronate **13** and a diazo compound **20** presumably generated from the corresponding sulfonyl hydrazone under basic conditions (Fig. 2c). In agreement with the synthetic outcomes (Fig. 2a), these calculations highlighted a strong preference for activation and functionalization of the C₃ carbon versus C₂-Bpin functionalization. Specifically, DFT calculations identified an energetically much lower concerted addition/migration/N₂ extrusion transition state that will lead to the corresponding C–C coupling at the C₃- than at the C₂-carbon (**13**–TS^{C3} versus **13**–TS^{C2}, $G_{\text{rel}} = +6.0$

kcal mol⁻¹). Moreover, the resulting C₃-product is also more thermodynamically favourable than the C₂-coupling product by similar energy (see Supplementary Information). Inspection of the Mulliken charges (MC) of the transition states (Fig. 2c) revealed a greater buildup of negative charge in **13-TS^{C2}** (MC C₂ = -0.181; MC C₃ = -0.045). Further energy decomposition analysis of **13-TS^{C2}** and **13-TS^{C3}** (Fig. 2d) suggested that despite greater non-covalent electrostatic, orbital, and dispersion stabilizing interactions in **13-TS^{C2}**, much more substantial steric interactions were observed in **13-TS^{C2}** ($E_{\text{steric}}^{\text{13-TS}^{\text{C2}}} = +310.3$ kcal mol⁻¹, $E_{\text{steric}}^{\text{13-TS}^{\text{C3}}} = +268.1$ kcal mol⁻¹), overall resulting in a higher barrier via the C₂ transition state ($E_{\text{rel}} = +11.0$ kcal mol⁻¹; Fig. 2c)⁴⁷.

Overall, we attribute the selectivity to (1) greater buildup of negative charge and (2) less destabilizing steric interactions between the BCP scaffold and pinacol ligand in the transition states at the C₃ position (see Supplementary Information for further details). Taking advantage of the higher reactivity of C₃-Bpin, we envisioned a sequential functionalization protocol of BCP bis-boronates, where C₃- and C₂-Bpin can be transformed into multiple functional groups and access divergent C₁, C₂, C₃-tri-substituted BCPs (Fig. 2e).

Synthesis of BCP bis-boronates

The advantages of this strategy to access bridge-substituted BCPs not only draws from its modularity, but also is reinforced by the synthetic efficacy and practicality in the preparation of various BCP bis-boronates. In our previous study³¹, the alkyl substituted BCP bis-boronate (**13**) was synthesized from readily accessible aldehyde via diborylation of its derived sulfonyl hydrazones⁴⁸ and intramolecular coupling³¹ on gram scale and in moderate yield. Bicyclo[1.1.1] pentanes **24** and **25** were prepared via similar routes (Fig. 3a); however, the carboxylate ester-, trifluoromethyl- and aryl-substituted BCP bis-boronates were found unobtainable by the challenging access to geminal bisBpin precursors (yield < 5%) and the diminished yield in intramolecular coupling. These BCPs (**23**, **26**, **27**) were successfully accessed (Fig. 3a) through crucial modifications including the diborylation of geminal dibromo compounds⁴⁹ and intramolecular coupling under dry conditions (pre-preparation of sulfonylhydrazone, dry solvent and base). As an example, synthesis of ester C₁-substituted BCP **23** (Fig. 3b) starts from an affordable cyclobutyl diester **28**. The cyclobutyl sulfonylhydrazone **31** was generated in high yields through a sequence of DIBAL-H reduction, dibromination and diborylation. The following cyclization mediated by anhydrous Cs₂CO₃ robustly afforded **23** on 37-gram scale in a single pass without deterioration of yield. Notably, all the synthesized BCP bis-boronates are stable crystalline solids and can be stored at -20 °C without any noticeable degradation for several months.

C₃-functionalization of BCP bis-boronates

With the knowledge gained from computational results and a variety of BCP bis-boronates prepared, we sought to investigate selective C₃-Bpin deborylative functionalization strategies (Table 1). Consistent with theoretical modelling, BCP bis-boronates underwent selective alkylation at the bridgehead (C₃) Bpin with sulfonylhydrazones³¹ (**18**, **32-35**), leaving the resultant alkyl Bpins intact. It is worth noting that other non-BCP alkyl Bpins were generally incompatible with this hydrazone coupling⁴⁶.

Furthermore, our study continued with C₃-protodeborylation of BCP bis-boronates to afford C₂-Bpin-substituted BCPs, which provides a modular tool to access a library of diverse 1,2-disubstituted BCPs, a class of potential bioisosteres for *ortho*- and *meta*-disubstituted arenes. Inspired by Renaud's studies^{50,51}, BCP bis-boronates (**13**, **25**, **27**) underwent effective protodeborylation with *tert*-butyl catechol (TBC) upon heating under argon atmosphere and C₂-Bpin-disubstituted BCPs were afforded with good yields (**36**, **38–39**). Other BCPs containing electron-withdrawing groups (**23**, **26**) at the C₁ position featured lower reactivities, and their protodeborylation was successfully promoted by air (**37**, **40**); however, this TBC strategy only led to decomposition of starting material in preparation of nitrogen C₃-substituted BCP (**41**), which was only made possible with a photoinduced protodeborylation approach¹⁸.

Enabled by the TBC strategy, the C₃-centred radicals formed in situ can also react with tosyl cyanide (TsCN), PhSO₂SPh and di-*tert*-butyl azodicarboxylate (DBAD) to afford nitrile- (**42**, **43**), thioether- (**44**, **45**) and hydrazide- (**46**, **47**) containing BCPs in moderate yields. C₃-selective Giese alkylation of BCPs (**48–54**) was achieved with Ley's conditions^{52,53} via the bridgehead C₃ radical from photoinduced oxidation of a Bpin-DMAP (4-dimethylaminopyridine) 'ate' complex.

Assessment of nickel-catalysed cross-coupling started with aryl bromides—an abundant class of building blocks—yet all of the previously reported conditions failed to afford the desired product^{54–58}. Extensive screening efforts identified a photosensitized method with 4-CzIPN as the photosensitizer and Zn(OTf)₂ as a key Lewis acid additive, where arylbromides with diverse electronic properties were well tolerated (**55–69**; see Supplementary Information for more details).

Minisci reactivities⁵⁹ could also be achieved with selective introduction of a series of heteroarenes including quinoxaline (**70**, **75**), caffeine (**71**, **76**), pyrazine (**72**, **77**), pyridazine (**73**, **78**), quinoline (**74**) and pyrimidine (**79**), and moderate to good yields of the products were afforded.

Late-stage C₂-functionalization of BCP boronates

With several scalable preparations of BCP C₂-Bpins (**36** and **37**, **54–55**, **65**, **77**), further derivatizations of such BCPs were explored to demonstrate broad applications of a sequential functionalization strategy for accessing C₁, C₂-di- (Fig. 4a) and C₁, C₂, C₃-tri-substituted BCPs (Fig. 4b) at a late stage. Successful transformation of Bpin group in **36** to the more stable trifluoroborate salt (**80**, 73%) and the boronic acid (**81**, 68%) created further functionalization opportunities. In comparison with other non-BCP secondary boronic species, BCP C₂-boronates were found to exhibit inferior reactivity owing to a combination of specific hybridization³⁸ at the C₂ bridge position and steric hinderance³⁶. Fortunately, attempts to leverage BCP **36** in C–O and C–S formation proved successful, as demonstrated in oxidation (**82**) and Renaud's radical trapping⁵¹ (**83**). Nickel-catalysed 'ate' complex coupling (**84**)⁴² with PhLi and Aggarwal's arylation (**85**)⁶⁰ from boronate **36**, Molander's photo-induced cross-coupling (**86**)^{55–58} and Minisci-type heteroarylation (**87**)⁵⁹ from trifluoroborate salt **80** enables the installation of aryl and heteroaryl groups at the

bridge (C_2) position. *tert*-Butyl-catechol-mediated C–N formation affords hydrazine (**88**)⁵¹ in 95% yield and amination with nitroarene via Radosevich's protocol affords anilines (**89**, **90**)⁶¹ in moderate yields. Sulfonyl hydrazone coupling (**91**)⁴⁶ of boronic acid (**81**), Giese-type alkylation (**92**) and cyanation (**93**) via 'ate' complex⁴² using PhLi from boronate **36**, and Matteson homologation (**94**)⁶² afford alkylation, cyanation and CH_2 insertion product on the BCP motif in moderate to high yields.

To further exploit the potential of BCP bis-boronates as a versatile building block, a series of structurally diverse trisubstituted BCPs were accessed through a C_3 – C_2 functionalization sequence. Representatives of bridgehead disubstituted BCP C_2 -Bpin (**42**, **54**, **55**, **65**, **77**) enabled employment of Bpin functionalization strategies, including Giese-type alkylation (**96**, **97**), Minisci-type heteroarylation (**98**, **99**), amination (**100**) arylation (**101**, **102**, **103**, **104**) and sulfuration (**105**).

Alternatively, to first achieve C_2 -Bpin functionalization, C_3 -Bpin was transformed into *N*-methylimidodiacetic boronic acid ester (BMIDA) or tetramethyl *N*-methylimidodiacetic boronic acid ester (BTIDA) (**109**, **110**)^{63,64}, followed by selective functionalization of C_2 -Bpin via oxidation (**111**) or Giese-type alkylation (**112**) with the C_3 -boronates remaining intact (Extended Data Fig. 1).

Furthermore, the medicinal chemistry rationale for pursuing poly-substituted BCPs was showcased via the improvement in physicochemical and ADME properties of biologically relevant compounds. The BCP match-pair **115** of arene **116** a p-38 kinase inhibitor⁶⁵, was prepared via the sequential functionalization strategy from **43**, and physicochemical and ADME properties for both analogues profiled (Extended Data Fig. 2). The data highlights that the theoretical improvement in F_{sp^3} (fraction of sp^3 carbon atoms) values by introducing the BCP also results in a substantial improvement in measured solubility (>10-fold) and reduction of lipophilicity, while retaining excellent permeability.

Bicyclo[1.1.1]pentane derivatives feature characteristic three-dimensional topology and represent a $C(sp^3)$ -rich chemical space. In a comparative study we enumerated libraries of trisubstituted BCP derivatives (1,2,4- and 1,3,4-trisubstituted arenes) with substituents selected from synthetically accessible moieties demonstrated earlier (with protecting groups removed; see Supplementary Information for more details). These compounds were evaluated with principal moments of inertia (PMI) metrics, where shape diversity of a given library can be assessed via computation of principal moments of inertia for each molecule (see Supplementary Information for more details)⁶⁶. In contrast to arene compounds, trisubstituted BCP compounds featured much greater structural diversity as they occupied much broader space (Fig. 5a). More importantly, BCP compounds exhibited higher three-dimensionality as they extend into the sphere-like space, whereas the corresponding arenes are heavily concentrated along the 'rod-disk' axis.

Further study focused on a selection of difunctionalized BCP compounds and arenes sharing the same set of substituents (Fig. 5b; see the Supplementary Information for details). Analysis of these compounds indicated that C_1 , C_2 -difunctionalized BCP compounds are more three-dimensional than 1, 2- and 1,3-difunctionalized arenes and C_1 , C_3 -disubstituted

BCP compounds. Calculation of 3D scores⁶⁷, that is, the sum of PMI (X) and PMI (Y), quantitatively demonstrated the merit of this C₂-functionalization strategy, as the C₁, C₂-difunctionalized BCP compounds scored broader and overall higher (average = 1.17, median = 1.13; Fig. 5b) than C₁, C₂-BCP compounds (average = 1.09, median = 1.05) and arenes (average = 1.04, median = 1.01) in the box-whisker plot. These results together suggested that this BCP C₂-functionalization strategy represent a unique entry to a novel three-dimensional chemical space.

In summary we developed a selective and sequential bis-functionalization strategy of BCP bis-boronates. Computational studies shed insights into the differentiated reactivities between C₂- and C₃-Bpins and such results proved consistent with synthetic outcomes. Combination of the scalable, crystalline BCP bis-boronates and enabling Bpin functionalization strategies allowed the access to an array of novel, diverse di- and tri-substituted BCP compounds. Systematic investigation of large and yet unexplored chemical space of C₂-substitution on BCPs demonstrated their higher shape diversity and three-dimensionality than conventional arene derivatives. Further application of this approach should help assemble molecular libraries with more three-dimensional structures and create a broader avenue to SAR studies^{68–71}.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, Supplementary Information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41557-023-01342-7>.

Methods

General procedure for intramolecular coupling to access BCP bis-boronates

A one-necked (24/40 joint) round-bottomed flask, equipped with a teflon-coated magnetic stir bar, was flame-dried under vacuum and then cooled to 23 °C under an atmosphere of argon. Then the flask was charged with sulfonyl hydrazone (1.0 equiv.) and dried cesium carbonate (3.0 equiv.). (Note: cesium carbonate was dried at 120 °C under vacuum for 18 h.). After being evacuated and backfilled with argon from a balloon three times, dioxane (0.2 M) was added into the flask and the reaction mixture was allowed to stir at 100 °C. After it was confirmed that the starting material, sulfonyl hydrazone, was consumed through TLC analysis, the reaction was cooled to room temperature, filtered through Celite, washed with excess hexanes, and concentrated to remove excess solvents. The crude reaction was purified through flash chromatography (hexanes: ethyl acetate, 20:1 to 10:1) on silica gel to afford the title compound, which was further purified through recrystallization in hexanes at –40 °C, affording the product with >99% purity as white solids³¹.

Recrystallization procedure.

The product after chromatography was dissolved in hexanes (around 1.0 ml g⁻¹) at room temperature and then cooled to –40 °C. After the solution of the product was slowly stirred

at $-40\text{ }^{\circ}\text{C}$ for 1 h, the suspension was filtered and the white solid was washed with cooled hexanes quickly and dried under vacuum for 1 h.

General procedure A for hydrazone coupling of BCP bis-boronates

A screw-capped 13×100 mm Pyrex culture tube was charged with cesium carbonate (3.0 equiv.), 2-mesitylsulfonyl hydrazone (2.0 equiv.) and BCP bis-boronate (1.0 equiv.). Then the tube was evacuated and backfilled with argon for three times, followed by addition of toluene (0.1 M) via a syringe. After stirring for at $70\text{ }^{\circ}\text{C}$ for 18–48 h, the reaction mixture was cooled to room temperature. The suspended solution was then filtered over celite and washed with diethyl ether. The solvent was removed under high vacuum, and the crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **18** and **32–35**.

General procedure B for deborylation of BCP bis-boronates

A screw-capped 13×100 mm Pyrex culture tube or a flame-dried 100 ml Pyrex flask was charged with BCP bisboronate (1.0 equiv.) and *tert*-butyl catechol (2.5 equiv.). The tube or the flask was then evacuated and backfilled with argon or air for three times, followed by addition of toluene (0.1 M) via a syringe. After stirring for at $100\text{ }^{\circ}\text{C}$ for 2–12 h when it is confirmed that the starting material was consumed totally, the reaction mixture was cooled to room temperature. Next, the solvent was removed under high vacuum, and the crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **36–41**.

General procedures C for cyanation of BCP bis-boronates

A flame-dried screw-capped 13×100 mm Pyrex culture tube was charged with BCP bisboronate (1.0 equiv.), *para*-toluenesulfonyl cyanide (2.0 equiv.) and *tert*-butyl catechol (0.25 equiv.). Then the tube or the flask was evacuated and backfilled with argon for three times, followed by addition of toluene (0.2 M) via a syringe. After stirring for at $70\text{ }^{\circ}\text{C}$ for 18–24 hours when it is confirmed that the starting material was consumed totally, the reaction mixture was cooled to room temperature. Next, the solvent was removed under high vacuum, and the crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **42** and **43**.

General procedures D for C–S and C–N formation of BCP bis-boronates

A flame-dried screw-capped 13×100 mm Pyrex culture tube was charged with BCP bisboronate (1.0 equiv.), radical trapping reagent (2.0 equiv., PhSO_2SPh or DBAD) and *tert*-butyl catechol (0.25 equiv.). Then the tube or the flask was evacuated and backfilled with argon for three times, followed by addition of toluene (0.2 M) via a syringe. After stirring for at $70\text{ }^{\circ}\text{C}$ or $100\text{ }^{\circ}\text{C}$ for 18–24 h when it is confirmed that the starting material was consumed totally, the reaction mixture was cooled to room temperature. Next, the solvent was removed under high vacuum, and the crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **44–47**.

General procedure E for Giese-type reaction of BCP bis-boronates

A screw-capped 13 × 100 mm Pyrex culture tube was charged with $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbpy})\text{PF}_6$ (5 mol%), DMAP (30 mol%), and BCP bisboronate (1.0 equiv.). Then the tube or the flask was evacuated and backfilled with argon for three times, followed by addition of Michael acceptor (2.0 equiv.) and methanol/acetone (0.1 M, 1:1) solvent via a syringe. Then the headspace of the tube was purged with a gentle stream of argon for approximately 10 s. After stirring in a 450 nm photoreactor for 24 h when it is confirmed that the starting material was consumed totally, the reaction mixture was concentrated under high vacuum, and the crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **48–54**.

General procedure F for cross-coupling of BCP bis-boronates General procedure F1.

A flame-dried screw-capped 13 × 100 mm Pyrex culture tube was charged with BCP bisboronate (1.0 equiv.), aryl bromide (3.0 equiv.), 4-CzIPN (5 mol%), $\text{Ni}(\text{dtbbpy})\text{Cl}_2$ (10 mol%), $\text{Zn}(\text{OTf})_2$ (2.0 equiv.) and DMAP (4.0 equiv.). Then the tube was evacuated and backfilled with argon three times, followed by addition of DMA (0.2 M) solvent via a syringe. The headspace of the tube was then purged with a gentle stream of argon for approximately 10 s and the reaction was allowed to stir in a 450 nm PennPhD integrated photoreactor (M2) for 24–60 h. After it is confirmed that the starting material was consumed totally, the reaction mixture quenched with water, extracted with ethyl acetate or diethyl ether, washed with brine, dried by Na_2SO_4 and concentrated under high vacuum. The crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **56** and **69**.

General procedure F2.

Preparation of [Ni] catalyst: a flame-dried screw-capped 13 × 100 mm Pyrex culture tube was charged with $\text{Ni}(\text{cod})_2$ (0.2 mmol, 55 mg) and dtbbpy (0.24 mmol, 64.4 mg). Then the tube was evacuated and backfilled with argon three times, followed by addition of DMA (4.0 ml) solvent via a syringe. The tube was then sonicated for 20 min to dissolve the catalyst.

A flame-dried screw-capped 13 × 100 mm Pyrex culture tube was charged with BCP bisboronate (1.0 equiv.), aryl bromide (3.0 equiv.), $\text{Zn}(\text{OTf})_2$ (2.0 equiv.) and DMAP (4.0 equiv.). Then the tube was evacuated and backfilled with argon for three times, followed by addition of 4-CzIPN solution (0.02 equiv., 0.02 M) in DMA and [Ni] catalyst solution (0.2 equiv., 0.05 M) in DMA via a syringe. The headspace of the tube was then purged with a gentle stream of argon for approximately 10 s and the reaction was allowed to stir in a 450 nm PennPhD integrated photoreactor (M2) for 24–60 h. After it is confirmed that the starting material was consumed totally, the reaction mixture quenched with water, extracted with ethyl acetate or diethyl ether, washed with brine, dried by Na_2SO_4 and concentrated under high vacuum. The crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **55**, **57–68**.

General procedure G for Minisci reaction of BCP bis-boronates


A screw-capped 13 × 100 mm Pyrex culture tube was charged with BCP bisboronate (1.0 equiv.), heteroarene (3.0 equiv.) and Mn(OAc)₃ (2.5 equiv.). Then the tube or the flask was evacuated and backfilled with argon for three times, followed by addition of acetic acid/water (0.1 M, 1:1) solvent via a syringe. Trifluoroacetic acid (5.0 equiv.) was then added into the reaction. The headspace of the tube was then purged with a gentle stream of argon for approximately 10 s and the reaction was stirred at 50 °C for 18 h. After it was confirmed that the starting material was completely consumed, the reaction mixture was concentrated under high vacuum to remove excess acetic acid, quenched with Na₂CO₃ solution, extracted with ethyl acetate, dried with Na₂SO₄ and concentrated under high vacuum. The crude residue was purified by chromatography on silica gel.

The procedure was applied in the synthesis of compounds **70–79**.

Data availability


Experimental data as well as characterization data for all new compounds prepared in the course of these studies are provided in the Supplementary Information. Crystallographic data for the structures reported in this Article have been deposited at the Cambridge Crystallographic Data Centre under deposition nos. CCDC 2158998 (**15**), 2159002 (**23**), 2158995 (**24**), 2159016 (**25**), 2159001 (**27**), 2162135 (**33**), 2160325 (**55**), 2160336 (**59**) and 2162136 (**76**), see the ‘X-ray crystallographic data’ section in the Supplementary Information. Copies of the data can be obtained free of charge via <https://www.ccdc.cam.ac.uk/structures/>. Source Data are provided with this paper.

Extended Data



Extended Data Fig. 1 | Selective functionalization of C₂-boronate.

To achieve C₂-functionalization with C₃-boron retained, BCP bis-boronate **23** was firstly transformed into C₃-BMIDA and BTIDA esters (**109**, **110**), followed by oxidation and Giese-type alkylation to afford C₂-functionalized products (**111**, **112**).



Extended Data Fig. 2 |. Synthesis and ADME study of p-38 kinase inhibitor **116** and its BCP analogue **115**.

The BCP analogue **115** of arene **116**, a p-38 kinase inhibitor⁶⁵, was prepared via the selective functionalization sequence: 1) cyanation; 2) hydrolysis and esterification; 3) arylation; 4) hydrolyses and amide couplings. The physicochemical and ADME properties for both compounds were profiled. F_{sp}³, the fraction of sp³ carbon atoms; Log D, distribution coefficient; Solubility, high-throughput equilibrium solubility; MDCKII, Madin-Darby canine kidney cells; Papp, apparent permeability.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support for this work was provided by the National Science Foundation (CAREER CHE-2143925 to T.Q.), National Institutes of Health (R01GM141088 to T.Q. and R35GM137797 to O.G.), Camille and Henry Dreyfus Foundation (to O.G.) and UT Southwestern Eugene McDermott Scholarship (to T.Q.). Preliminary results were made possible by the support of Welch Foundation (I-2010–20190330 to T.Q.) and American Chemistry Society Petroleum Research Fund (62223-DN11 to T. Q.). UT Southwestern Amgen Scholars program supported a fellowship to J.B.W. We thank F. Lin (UTSW) for assistance with NMR spectroscopy; H. Baniyadi (UTSW) for HRMS; and V. Lynch (UT-Austin) for X-ray crystallographic analysis. We gratefully acknowledge the Texas A&M University HPRC resources (<https://hprc.tamu.edu>), UMD Deepthought2, MARCC/BlueCrab HPC clusters and XSEDE (CHE160082 and CHE160053) for computational resources. We thank the Chen, Tambar, Ready, De Brabander, Smith and Falck groups (UTSW) for generous access to equipment, and helpful discussions. We are grateful to S. W. Krska, X. Ma, and D. Leverage (Merck & Co., Inc.) for feedback on this manuscript and assistance with ADME profiling.

References

- Lovering F, Bikker J & Humblet C Escape from flatland: increasing saturation as an approach to improving clinical success. *J. Med. Chem.* 52, 6752–6756 (2009). [PubMed: 19827778]
- Talele TT Opportunities for tapping into three-dimensional chemical space through a quaternary carbon. *J. Med. Chem.* 63, 13291–13315 (2020). [PubMed: 32805118]
- Blakemore DC et al. Organic synthesis provides opportunities to transform drug discovery. *Nat. Chem.* 10, 383–394 (2018). [PubMed: 29568051]

4. Costantino G et al. Synthesis and biological evaluation of 2-(3'-(1H-tetrazol-5-yl)bicyclo[1.1.1]pent-1-yl)glycine (S-TBPG), a novel mGlu1 receptor antagonist. *Bioorg. Med. Chem.* 9, 221–227 (2001). [PubMed: 11249114]
5. Mikhailiuk PK et al. Conformationally rigid trifluoromethyl-substituted α -amino acid designed for peptide structure analysis by solid-state ^{19}F NMR spectroscopy. *Angew. Chem. Int. Ed.* 45, 5659–5661 (2006).
6. Stepan AF et al. Application of the bicyclo[1.1.1]pentane motif as a nonclassical phenyl ring bioisostere in the design of a potent and orally active γ -secretase inhibitor. *J. Med. Chem.* 55, 3414–3424 (2012). [PubMed: 22420884]
7. Westphal MV, Wolfstaedter BT, Plancher J, Gatfield J & Carreira EM Evaluation of tert-butyl isosteres: case studies of physicochemical and pharmacokinetic properties, efficacies, and activities. *ChemMedChem* 10, 461–469 (2015). [PubMed: 25630804]
8. Measom ND et al. Investigation of a bicyclo[1.1.1]pentane as a phenyl replacement within an LpPLA2 Inhibitor. *ACS Med. Chem. Lett.* 8, 43–48 (2017). [PubMed: 28105273]
9. Auberson YP et al. Improving nonspecific binding and solubility: bicycloalkyl groups and cubanes as para-phenyl bioisosteres. *ChemMedChem* 12, 590–598 (2017). [PubMed: 28319646]
10. Mikhaliuk PK Saturated bioisosteres of benzene: where to go next? *Org. Biomol. Chem.* 17, 2839–2849 (2019). [PubMed: 30672560]
11. Subbaiah MAM & Meanwell NA Bioisosteres of the phenyl ring: recent strategic applications in lead optimization and drug design. *J. Med. Chem.* 64, 14046–14128 (2021). [PubMed: 34591488]
12. Gianattasio R et al. Strain-release amination. *Science* 351, 241–246 (2016). [PubMed: 26816372]
13. Caputo DFJ et al. Synthesis and applications of highly functionalized 1-halo-3-substituted bicyclo[1.1.1]pentanes. *Chem. Sci.* 9, 5295–5300 (2018). [PubMed: 29997886]
14. Nugent J et al. A general route to bicyclo[1.1.1]pentanes through photoredox catalysis. *ACS Catal.* 9, 9568–9574 (2019).
15. Zhang X et al. Copper-mediated synthesis of drug-like bicyclopentanes. *Nature* 580, 220–226 (2020). [PubMed: 32066140]
16. Trongsirivat N et al. Reactions of 2-aryl-1,3-dithianes and [1.1.1]propellane. *Angew. Chem. Int. Ed.* 58, 13416–13420 (2019).
17. Yu S, Jing C, Noble A & Aggarwal VK 1,3-Difunctionalizations of [1.1.1]propellane via 1,2-metalate rearrangements of boronate complexes. *Angew. Chem. Int. Ed.* 59, 3917–3921 (2020).
18. Kim JH, Ruffoni A, Al-Faiyz YSS, Sheikh N & Leonori D Divergent strain-release amino-functionalization of [1.1.1]propellane with electrophilic nitrogen-radicals. *Angew. Chem. Int. Ed.* 59, 8225–8231 (2020).
19. Garlets ZJ et al. Enantioselective C–H functionalization of bicyclo[1.1.1]pentanes. *Nat. Catal.* 3, 351–357 (2020).
20. Harmata AS, Spiller TE, Sowden MJ & Stephenson CRJ Photochemical formal (4 + 2)-cycloaddition of imine-substituted bicyclo[1.1.1]pentanes and alkenes. *J. Am. Chem. Soc.* 143, 21223–21228 (2021). [PubMed: 34902245]
21. Yu S, Jing C, Noble A & Aggarwal VK Iridium-catalyzed enantioselective synthesis of α -chiral bicyclo[1.1.1]pentanes by 1,3-difunctionalization of [1.1.1]propellane. *Org. Lett.* 22, 5650–5655 (2020). [PubMed: 32638587]
22. Polites VC, Badir SO, Keess S, Jolit A & Molander GA Nickel-catalyzed decarboxylative cross-coupling of bicyclo[1.1.1]pentyl radicals enabled by electron donor-acceptor complex photoactivation. *Org. Lett.* 23, 4828–4833 (2021). [PubMed: 34100624]
23. Nugent J, Sterling AJ, Frank N, Mousseau JJ & Anderson EA Synthesis of α -quaternary bicyclo[1.1.1]pentanes through synergistic organophotoredox and hydrogen atom transfer catalysis. *Org. Lett.* 23, 8628–8633 (2021). [PubMed: 34699248]
24. Wong MLJ, Sterling AJ, Mousseau JJ, Duarte F & Anderson EA Direct catalytic asymmetric synthesis of α -chiral bicyclo[1.1.1]pentanes. *Nat. Commun.* 12, 1644 (2021). [PubMed: 33712595]
25. Pickford HD et al. Twofold radical-based synthesis of *N,C*-difunctionalized bicyclo[1.1.1]pentanes. *J. Am. Chem. Soc.* 143, 9729–9736 (2021). [PubMed: 34161076]

26. Denisenko A, Garbuz P, Shishkina SV, Voloshchuk NM & Mikhailiuk PK Saturated bioisosteres of *ortho*-substituted benzenes. *Angew. Chem. Int. Ed.* 59, 20515–20521 (2020).
27. Anderson JM, Measom ND, Murphy JA & Poole DL Bridge functionalisation of bicyclo[1.1.1]pentane derivatives. *Angew. Chem. Int. Ed.* 60, 24754–24769 (2021).
28. Zhao J-X et al. 1,2-Difunctionalized bicyclo[1.1.1]pentanes: long-sought-after mimetics for *ortho/meta*-substituted arenes. *Proc. Natl Acad. Sci. USA* 118, e2108881118 (2021). [PubMed: 34244445]
29. Ma X, Han Y & Bennett DJ Selective synthesis of 1-dialkylamino-2-alkylbicyclo-[1.1.1]pentanes. *Org. Lett.* 22, 9133–9138 (2020). [PubMed: 33170018]
30. Ma X, Sloman DL, Han Y & Bennett DJ A selective synthesis of 2,2-difluorobicyclo[1.1.1]pentane analogues: 'BCP-F2'. *Org. Lett.* 21, 7199–7203 (2019). [PubMed: 31294572]
31. Yang Y et al. An intramolecular coupling approach to alkyl bioisosteres for the synthesis of multisubstituted bicycloalkyl boronates. *Nat. Chem.* 13, 950–955 (2021). [PubMed: 34584254]
32. Applequist DE, Renken TL & Wheeler JW Polar substituent effects in 1,3-disubstituted bicyclo[1.1.1]pentanes. *J. Org. Chem.* 47, 4985–4995 (1982).
33. Bychek RM et al. Difluoro-substituted bicyclo[1.1.1]pentanes for medicinal chemistry: design, synthesis, and characterization. *J. Org. Chem.* 84, 15106–15117 (2019). [PubMed: 31553875]
34. Ryan EM, Amber LT & Edward AA Synthesis and applications of polysubstituted bicyclo[1.1.0]butanes. *J. Am. Chem. Soc.* 143, 21246–21251 (2021). [PubMed: 34904841]
35. Buskes MJ & Blanco MJ Impact of cross-coupling reactions in drug discovery and development. *Molecules* 25, 3493–3514 (2020). [PubMed: 32751973]
36. Wiberg KB & Williams VZ Bicyclo[1.1.1]pentane derivatives. *J. Org. Chem.* 35, 369–373 (1970).
37. Levin MD, Kaszynski P & Michl J Bicyclo[1.1.1]pentanes, [n] saffanes, [1.1.1]propellanes, and tricyclo[2.1.0.0_{2,5}]pentanes. *Chem. Rev.* 100, 169–223 (2000). [PubMed: 11749237]
38. Jarret RM & Cusumano L ¹³C–¹³C coupling in [1.1.1]propellane. *Tetrahedron Lett.* 31, 171–174 (1990).
39. Mlynarski SN, Schuster CH & Morken JP Asymmetric synthesis from terminal alkenes by cascades of diboration and cross-coupling. *Nature* 505, 386–390 (2014). [PubMed: 24352229]
40. Blaisdell TP & Morken JP Hydroxyl-directed cross-coupling: a scalable synthesis of debromohamigeran E and other targets of interest. *J. Am. Chem. Soc.* 137, 8712 (2015). [PubMed: 26125083]
41. Liu X, Sun C, Mlynarski S & Morken JP Synthesis and stereochemical assignment of arenolide. *Org. Lett.* 20, 1898 (2018). [PubMed: 29565134]
42. Kaiser D, Noble A, Fasano V & Aggarwal VK 1,2-Boron shifts of β-boryl radicals generated from bis-boronic esters using photoredox catalysis. *J. Am. Chem. Soc.* 141, 14104–14109 (2019). [PubMed: 31461622]
43. Fawcett A et al. Regio- and stereoselective homologation of 1,2-bis(boronic esters): stereocontrolled synthesis of 1,3-diols and Sch725674. *Angew. Chem., Int. Ed.* 55, 14663–14667 (2016).
44. Nóvoa L, Trulli L, Parra A & Tortosa M Stereoselective diboration of spirocyclo-butenes: a platform for the synthesis of spirocycles with orthogonal exit vectors. *Angew. Chem. Int. Ed.* 60, 11763–11768 (2021).
45. Crudden CM et al. Iterative protecting group-free cross-coupling leading to chiral multiply arylated structures. *Nat. Commun.* 7, 11065–11071 (2016). [PubMed: 27040494]
46. Yang Y et al. Practical and modular construction of C(sp³)-rich alkyl boron compounds. *J. Am. Soc. Chem.* 143, 471–480 (2021).
47. Qi X, Kohler DG, Hull KL & Liu P Energy decomposition analyses reveal the origins of catalyst and nucleophile effects on regioselectivity in nucleopalladation of alkenes. *J. Am. Chem. Soc.* 141, 11892–11904 (2019). [PubMed: 31322875]
48. Li H, Wang L, Zhang Y & Wang J Transition-metal-free synthesis of pinacol alkylboronates from tosylhydrazones. *Angew. Chem. Int. Ed.* 51, 2943–2946 (2012).

49. Yang CT, Zhang ZQ, Liu YC & Liu L Copper-catalyzed cross-coupling reaction of organoboron compounds with primary alkyl halides and pseudohalides[†]. *Angew. Chem. Int. Ed.* 50, 3904–3907 (2011).
50. Pozzi D, Scanlan EM & Renaud P A mild radical procedure for the reduction of B-alkylcatecholboranes to alkanes. *J. Am. Chem. Soc.* 127, 14204–14205 (2005). [PubMed: 16218613]
51. André-Joyaux E, Kuzovlev A, Tappin ND & Renaud P A general approach to deboronative radical chain reaction with pinacol alkylboronic esters. *Angew. Chem. Int. Ed.* 59, 13859–13864 (2020).
52. Lima F et al. A Lewis base catalysis approach for the photoredox activation of boronic acids and esters. *Angew. Chem. Int. Ed.* 56, 15136–15140 (2017).
53. Lima F et al. Organic photocatalysis for the radical couplings of boronic acid derivatives in batch and flow. *Chem. Commun.* 54, 5606–5609 (2018).
54. Lima F et al. Visible light activation of boronic esters enables efficient photoredox C(sp²)-C(sp³) cross-couplings in flow. *Angew. Chem. Int. Ed.* 55, 14085–14089 (2016).
55. Tellis JC, Primer DN & Molander GA Single-electron transmetalation in organoboron cross-coupling by photoredox/nickel dual catalysis. *Science* 345, 433–436 (2014). [PubMed: 24903560]
56. Gutierrez O, Tellis JC, Primer DN, Molander GA & Kozlowski MC Nickel-catalyzed cross-coupling of photoredox-generated radicals: uncovering a general manifold for stereoconvergence in nickel-catalyzed cross-couplings. *J. Am. Chem. Soc.* 137, 4896–4899 (2015). [PubMed: 25836634]
57. Primer DN & Molander GA Enabling the cross-coupling of tertiary organoboron nucleophiles through radical-mediated alkyl transfer. *J. Am. Chem. Soc.* 139, 9847–9850 (2017). [PubMed: 28719197]
58. Yuan M, Song Z, Badir SO, Molander GA & Gutierrez O On the nature of C(sp³)-C(sp²) bond formation in nickel-catalyzed tertiary radical cross-couplings: a case study of Ni/photoredox catalytic crosscoupling of alkyl radicals and aryl halides. *J. Am. Chem. Soc.* 142, 7225–7234 (2020). [PubMed: 32195579]
59. Molander GA, Colombel V & Braz VA Direct alkylation of heteroaryls using potassium alkyl- and alkoxyethyltrifluoroborates. *Org. Lett.* 13, 1852–1855 (2011). [PubMed: 21391555]
60. Odachowski M et al. Development of enantiospecific coupling of secondary and tertiary boronic esters with aromatic compounds. *J. Am. Chem. Soc.* 138, 9521–9532 (2016). [PubMed: 27384259]
61. Nykaza TV et al. Intermolecular reductive C–N cross coupling of nitroarenes and boronic acids by P^{III}/P^V=O catalysis. *J. Am. Chem. Soc.* 140, 15200–15205 (2018). [PubMed: 30372615]
62. Sadhu KM & Matteson DS (Chloromethyl)lithium: efficient generation and capture by boronic esters and a simple preparation of diisopropyl (chloromethyl)boronate. *Organometallics* 4, 1687–1689 (1985).
63. Li J, Grillo AS & Burke MD From synthesis to function via iterative assembly of *N*-methyliminodiacetic acid boronate building blocks. *Acc. Chem. Res.* 48, 2297–2307 (2015). [PubMed: 26200460]
64. Blair DJ et al. Automated iterative C_sp³-C bond formation. *Nature* 604, 92–97 (2022). [PubMed: 35134814]
65. Angell RM et al. Biphenyl amide p38 kinase inhibitors 4: DFG-in and DFG-out binding modes. *Bioorg. Med. Chem. Lett.* 18, 4433–4437 (2008). [PubMed: 18602262]
66. Sauer WH & Schwarz MK Molecular shape diversity of combinatorial libraries: a prerequisite for broad bioactivity. *J. Chem. Inf. Comput. Sci.* 43, 987–1003 (2003). [PubMed: 12767158]
67. Prosser KE, Stokes RW & Cohen SM Evaluation of 3-dimensionality in approved and experimental drug space. *ACS Med. Chem. Lett.* 11, 1292–1298 (2020). During manuscript editing process after it is accepted, several publications on synthesis of bridge-substituted BCPs and BCP boronates were reported.
68. Dong W et al. Exploiting the sp² character of bicyclo[1.1.1] pentyl radicals in the transition-metal-free multi-component difunctionalization of [1.1.1]propellane. *Nat. Chem.* 14, 1068–1077 (2022). [PubMed: 35864151]

69. Yu IF et al. Catalytic undirected borylation of tertiary C–H bonds in bicyclo[1.1.1]pentanes and bicyclo[2.1.1]hexanes. *Nat. Chem.* 15, 685–693 (2023). [PubMed: 36973434]
70. Wright BA et al. Skeletal editing approach to bridge-functionalized bicyclo[1.1.1] pentanes from azabicyclo[2.1.1] hexanes. *J. Am. Chem. Soc.* 145, 0960–10966 (2023).
71. Garry OL et al. Rapid access to 2-substituted bicyclo[1.1.1] pentanes. *J. Am. Chem. Soc.* 145, 3092–3100 (2023). [PubMed: 36696089]




Fig. 1 |. Significance, challenges and strategy for accessing multi-substituted BCPs.

a, Significance of bicyclo[1.1.1]pentanes as a 3D bioisostere for arenes in medicinal chemistry. Bicyclo[1.1.1]pentanes has been applied as benzene 3D-surrogates in drug discovery owing to their metabolic stability from their high BDE and kinetically inhibited HAT process. Several BCP derivatives of drug molecules such as BCP-avagacestat and BCP-darapladib have demonstrated their potential to improve physicochemical, pharmacological (ADME) and toxicological (safety) properties. **b**, Synthetic challenges in SAR studies with a BCP core structure. Owing to well-established $C(sp^2)-C(sp^2)$ cross-coupling reactions and abundant available building blocks, it is synthetically accessible to obtain diverse derivatives of arene lead compounds for further SAR studies; however, derivatization of the $C(sp^3)$ -rich bioisostere BCPs has been synthetically challenging due to non-programmable synthetic route, requirement of de novo synthesis and inaccessible chemical space. **c**, Programmable and sequential functionalization of bridge-substituted BCPs. To access

multisubstituted BCPs (**9**), a sequential functionalization strategy from BCP bis-boronate (**10**) was proposed taking advantage of different reactivity of two boronates. Diversification of R¹, chemoselectivity of two boronates and reactivity of boronic esters are main challenges in this strategy.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 2 | Preliminary chemoselectivity of BCP bis-boronates and theoretical explanation.
a, Preliminary results for BCP bis-boronates reactivities. In the ligand-exchange and hydrazone coupling reactions, the bridgehead $\text{C}_3\text{-Bpin}$ in BCP bis-boronates presented an increased reactivity and underwent the two transformations, whereas $\text{C}_2\text{-Bpin}$ in bis-boronates and $\text{C}_2\text{-alkyl}$ BCP boronates remained mostly intact. **b**, Energetic comparison between BCP C_3^- and C_2^- anionic and radical intermediates suggest an anion-mediated transformation at the bridgehead position (C_3^-). **c**, Concerted addition/migration/ N_2 extrusion hydrazone coupling at the C_3^- and C_2^- positions of BCP bis-boronates show substantial preference for reactivity at C_3^- . **d**, Energy decomposition analysis of the concerted transition states in hydrazone coupling reveal that steric interactions greatly impact the selectivity. **e**, Working hypothesis for selective BCP bis-boronates functionalizations. See Supplementary Information for computational methods and details. Energy values in **b**, **c** and **d** are in kcal mol^{-1} . cPent, cyclopentyl.




Fig. 3 |. Scope and synthesis of BCP bis-boronates.

a. Substrate scope of BCP bis-boronates (**13**, **23–27**). Six C_1 -substituted BCP bis boronates (esters, amino, alkyls, CF_3 , aryl) were obtained via intramolecular coupling of sulfonyl hydrazones and boronic esters. **b.** Representative synthesis route towards BCP **23** via (1) DIBAL reduction; (2) dibromination; (3) diborylation/hydrolysis/ hydrazine condensation; and (4) intramolecular coupling. See ‘Synthetic details and optimization of intramolecular coupling’ section in the Supplementary Information for more details.




Fig. 4 | Syntheses of structurally divergent BCP compounds.

a, Functionalization of C_2 -Bpin as a versatile building block towards C_1 , C_2 -disubstituted BCPs. **b**, Late-stage functionalization to access C_1 , C_2 , C_3 -trisubstituted BCPs. Reaction conditions: ^aBCP C_2 -Bpin (1.0 equiv.), PhLi (1.2 equiv.) THF (0.2 M), -78°C to r.t., 1 h; then 4-CzIPN (5 mol%), *tert*-butyl acrylate (2.0 equiv.), THF/MeCN (0.1 M), blue LEDs, 15 h. ^bBCP C_2 - BF_3K (1.0 equiv.), heteroarene (3.0 equiv.), $\text{Mn}(\text{OAc})_3$ (2.5 equiv.), TFA (2.5 equiv.), AcOH/ H_2O (1:1, 0.1 M), 50°C , 18 h. ^cBCP C_2 - $\text{B}(\text{OH})_2$ (1.0 equiv.), ArNO_2 (1.0 equiv.), 1,2,2,3,4,4-hexamethyl-phosphetane 1-oxide (15 mol%), PhSiH_3 (2.0 equiv.),

m-xylene (0.5 M), 120 °C, 8 h. ^dBCP C₂-BF₃K (1.0 equiv.), [Ir] (5 mol%), Ni(dtbbpy) Cl₂ (20 mol%), ArBr (5.0 equiv.), Cs₂CO₃ (6.0 equiv.), dioxane or THF (0.1 M), blue LEDs, 24 h. ^eBCP C₂-Bpin (1.0 equiv.), PhLi (1.2 equiv.) THF (0.2 M), -78 °C to r.t., 1 h; then 4-CzIPN (5 mol%), Ni(dtbbpy)Cl₂ (10 mol%), ArBr (3.0 equiv.), THF/DMA (0.1 M), blue LEDs, 15 h. ^fBCP C₂-Bpin (1.0 equiv.), PhSO₂SPh (4.0 equiv.), MeOBcat (1.0 equiv.), TBC (0.3 equiv.), toluene (0.2 M), 80 °C, 24 h. See the 'Experimental procedures and characterization data of substrates' section in Supplementary Information for details.





Fig. 5 |. PMIs between substituted BCPs and arenes.

a, PMI analysis of trisubstituted BCP and arene chemical space. **b**, PMI and 3D scores comparison between selected disubstituted BCP and arenes. 3D scores were presented as box-whisker plots ranging between 1.00 and 2.00. The lines are the median, the average is marked by crosses, and outliers are shown as dots. See Methods and Source Data Fig. 5 for details (quartiles, boundaries and so on) on the box-whisker plots.

Table 1 |

Selective C₃-Bpin functionalization of BCP bis-boronates



Reaction conditions with BCP BisBpin as the Limiting reagent at 0.05–0.2mmol scale:

^a sulfonyl hydrazones (2.0 equiv.), Cs₂CO₃ (3.0 equiv.), toluene (0.2M), 70°C, 18–48h

^b TBC (2.5 equiv.), toluene (0.2M), argon or air atmosphere, 100°C, 2–12h

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^c 4-CzIPN (5mol%), MeOBeat (30 mol%), acetone/MeOH (1:1, 0.1 M), blue LEDs, r.t., 2 h

^d TsCN (2.0 equiv.), TBC (20mol%), toluene (0.2M), 18h

^e PhSO₂SPh or DBAD (2.0 equiv.), TBC (20mol%), toluene (0.2M), 70°C or 100 °C, 18–24h

^f [Ir] (5mol%), DMAP (30mol%), Michael acceptor (2.0 equiv.), acetone/MeOH (1:1, 0.1M), blue LEDs, r.t., 24h

^g 4-CzIPN (2–5mol%), [Ni] (10–20mol%), ArBr (3.0 equiv.), Zn(OTf)₂ (2.0 equiv.), DMAP (4.0 equiv.), DMA (0.2M), blue LEDs, r.t., 24–60h

^h heteroarene (3.0 equiv.), Mn(OAc)₃·2H₂O (2.5 equiv.), TFA (2.5 equiv.), AcOH/H₂O (1:1, 0.1M), 50°C, 18h.

ⁱ The structure was verified by X-ray experiments

ⁱⁱ d.r. = 1:1

ⁱⁱⁱ 2.0 mmol scale

^{iv} 10mmol scale

^v 1.0 mmol scale. [Ir], (Ir[dF(CF₃)ppy]₂(dtbbpy))PF₆. [Ni], Ni(dtbbpy)Cl₂ or Ni(cod)₂+dtbbpy.

EWG, electron-withdrawing group; Beat, boronic acid catechol ester; ppy, 2-phenylpyridine; dtbbpy, 4,4-di-tert-butyl-2,2-dipyridyl; cod, 1,5-cyclooctadiene; DMA, dimethylacetamide.