

Electrosynthesis

How to cite: *Angew. Chem. Int. Ed.* **2023**, *62*, e202218195

International Edition: doi.org/10.1002/anie.202218195

German Edition: doi.org/10.1002/ange.202218195

Electronically Ambivalent Hydrodefluorination of Aryl-CF₃ groups enabled by Electrochemical Deep-Reduction on a Ni Cathode

John R. Box, Mickaël E. Avanthay, Darren L. Poole, and Alastair J. J. Lennox*

Abstract: The Ar-CF₂H moiety is featured in an increasing number of bioactive compounds due to its unique combination of properties. The hydrodefluorination of Ar-CF₃ compounds is a direct and efficient route toward this motif. As reported methods for this transformation have focused on specific substrate families, herein we describe a general—electronically ambivalent—procedure for the single-step direct mono-hydrodefluorination of a variety of feedstock and functionalized Ar-CF₃ compounds. Exploiting the inherent tunability of electrochemistry and the selectivity enabled by a Ni cathode, the deep reduction garners high selectivity for ArCF₂H products, with good to excellent yields up to gram scale. The protocol has been extended to a single-step *di*-hydrodefluorination yielding benzyl fluorides. The late-stage peripheral editing of a single CF₃ feedstock to construct fluoromethyl (CF₂H, CFH₂) moieties will aid the rapid diversification of lead-compounds and compound libraries.

The difluoromethyl group has been the subject of much recent attention in drug and agrochemical design.^[1–6] In particular, difluoromethylarenes (ArCF₂H) are incorporated as lipophilic^[7,8] hydrogen-bond donors,^[9,10] a unique combination of properties that provides discovery chemists with a robust, non-nucleophilic bioisostere for OH and SH groups.^[2,10–12] While not as prevalent as trifluoromethylarenes (ArCF₃),^[3,13] the frequency of ArCF₂H moieties reported in biologically-relevant systems has exploded over the last decade.^[14] Intramolecular H-bonding in a CF₂H-pyrazole containing fungicide provides structural rigidity and enhances potency,^[9] Figure 1A, as well as intermolecular H-bonding^[11] in biological systems.^[15] The ability to modulate logP^[16] and pK_a^[17] are also enabled by a C–F to C–H switch.

The requirement for rapid syntheses amenable to late-stage intermediates has initiated a shift toward molecular editing strategies.^[18,19] Hence, while the direct difluoromethylation of prefunctionalized substrates is known, albeit with limited scope, or expensive/impractical reagents,^[20,21] editing the more easily installed and traditionally inert CF₃ group has synthetic and strategic advantages, Figure 1B. Pioneering work from Perichon,^[22–24] Savéant^[25,26] and Bordeaux^[27,28] established a route toward their functionalization through single electron reduction that initiates mesolytic C–F bond cleavage.^[29] As the CF₃ group progressively defluorinates, the C–F bond strength decreases^[30–33] and the rate of mesolytic cleavage increases, Figure 1C. This favours further defluorination, as encountered in early electrochemical attempts where total defluorination could not be avoided.^[24,34–36] Nevertheless, the use of excess Mg as reductant was reported for the hydrodefluorination of the readily reducible *bis*(CF₃) arenes under acidic conditions.^[37] The reductant in this 2-electron defluorination/protonation (ET/PT, Figure 1C) strategy must preferentially reduce the ArCF₃ and avoid direct proton reduction, a compromise that naturally limits the scope of amenable ArCF₃ substrates.^[38] Broader scope and functional group tolerance has since been achieved by Jui,^[39] Gouverneur,^[40] and Shang,^[41] who rely on SET from an excited organophotocatalyst and then hydrogen atom transfer (HAT, Figure 1C). Although good hydrodefluorination yields and improved selectivity are reported, each system is specific to a set range of arene electronics, as the reduction potential window is determined by the photocatalyst/H-atom donor system.^[42,43] Hence, a method that is broadly applicable on a range of electronically-variable ArCF₂H compounds remains elusive.





Electrochemistry has the innate ability to safely apply deep-reduction potentials (< –2.0 V vs Fc/Fc⁺). Hence, compounds can be accessed that require reduction potentials outside the range of traditional chemical reductants and excited state photoredox catalysts. As the reduction potential of each intermediate becomes more negative with each defluorination, Figure 1D, by applying only the minimum required potential, the intermediates can be differentiated and over defluorination is easier to avoid. Herein, we detail our successful efforts in the development of a selective electrochemical *mono*- and *di*-hydrodefluorination strategy on a broad range of electronically variable substrates.

The primary challenge in the development of this reaction was to accommodate the deeply reducing potentials required while avoiding over-reduction and competing proton reduction. Investigations initially focused on the electronically neutral Ar-CF₃ substrate **1a** ($E_{\text{red(onset)}}$ –2.8 V

[*] J. R. Box, M. E. Avanthay, A. J. J. Lennox
 School of Chemistry, University of Bristol
 Cantock's Close, Bristol, BS8 1TS (UK)
 E-mail: a.lennox@bristol.ac.uk

D. L. Poole
 Discovery High-Throughput Chemistry, Medicinal Chemistry
 GSK Medicines Research Centre, Stevenage, SG1 2NY (UK)

© 2023 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

[A] Pharmaceutical and agrochemical utility of Ar-CF₂H[B] Strategies and synthesis of Ar-CF₂H groups[C] Reductive defluorination of Ar-CF₃ groups

HAT: photoredox catalysis (Jui,³³ Gouverneur,³⁴ Shang³⁶)

→ Electron-poor + -neutral

ET/PT: Mg (Prakash³¹)

→ Electron-poor

Electrochemistry: **this work**

→ Electron-poor, neutral and electron-rich

[D] Electrochemical reduction to achieve broad scope and differentiation of intermediates: this work




Figure 1. [A–C]: The utility of CF₂H groups, their synthesis and challenges; [D]: Electrochemical defluorination strategy. CV of **1a–d** (averaged forward and return currents). -Ni || +Pt, 5 mM [**1a–d**], degassed DMF, 0.1 V/s, NBu₄PF₆, N₂. See Supporting Information for CVs run in with different electrode materials and in MeCN.

vs Fc/Fc⁺). A divided cell was adopted with a reductively stable electrolyte system (MeCN, $E_{\text{window}} +2.7\text{--}3.4\text{ V vs Fc/}$

Fc⁺ with Et₄NPF₆).^[44] Following extensive optimization studies (see Table 1 and Supporting Information), 76 % of ArCF₂H **1b** was achieved with 25:1 selectivity for the *mono*-hydrodefluorinated **1b** vs *di*-hydrodefluorinated benzyl fluoride **1c** (entry 1). It was found that the inclusion of the fluoride trap, TMSCl, enhanced the yield and selectivity (entry 1 vs 2). In place of sacrificial metal anodes, which can competitively reduce and plate the cathode at deep reduction potentials, Bu₄NBr was used as a suitable organic reductant, as previously discovered in our trifluoromethylketone hydrodefluorination work.^[45] As the oxidation product (Br₃⁻) is anionic, it resists migration to the cathodic chamber and any subsequent short circuiting that would lead to a decline in faradaic efficiency. Application of 1 F gave almost ideal efficiency (max = 50 %) and selectivity (entry 3). Bu₄NCl was also suitable but gave lower selectivity (entry 4). The most important factor to achieving high selectivity on this substrate was the choice of a nickel foil working electrode (entries 5–7).^[46] Ni was found to give higher yields than Pt or graphite and superior selectivity, halting at **1b** after 2 F. Extensive electrode fouling and physical decomposition of graphite electrodes were observed at these deep reduction potentials. Anhydrous MeCN was required to avoid competing reduction of either water (or HCl formed from the hydrolysis of TMSCl). With the exception of TBAClO₄, alternative reductively stable solvents or supporting electrolyte salts did not lead to any desired reaction (entries 8–12); competing K⁺ reduction ($E_{\text{red}} = -2.27\text{ V vs Fc/Fc}^+$)^[47] likely decreases efficiency with KPF₆ (entry 11).


With optimized conditions for **1a**, we tested their amenability to a range of electronically variable ArCF₃ substrates, Figure 2A. Hence, a small collection of electron-neutral, -poor and -rich rings were tested under our conditions, as well as in four published state-of-the-art

Table 1: Hydrodefluorination optimisation.^[a]

Entry	Variation from optimized conditions (above)	Yield [%]	Selectivity CF ₂ H:CFH ₂ (1b : 1c)
1	None	76	25:1
2	No TMSCl	64	7:1
3	1 F not 2 F	47	> 25:1
4	Bu ₄ NCl not Bu ₄ NBr	71	10:1
5	+ Ni foil not Pt	50	20:1
6	- Pt not Ni foil	72	5:1
7	- C not Ni foil	20	5:1
8	Me-THF not MeCN	n.d.	n/a
9	DMF not MeCN	n.d.	n/a
10	Me ₃ PhNPF ₆ not Et ₄ NPF ₆	< 5%	n/a
11	Bu ₄ NClO ₄ not Et ₄ NPF ₆	71	20:1
12	KPF ₆ not Et ₄ NPF ₆	n.d.	n/a

[a] See Supporting Information for full details. ¹⁹F NMR yields. Ratios in brackets correspond to ArCF₂H:ArCFH₂ determined from crude ¹⁹F NMR, n.d. = not-detected.

[A] Exploration of scope



[B] Hammett analysis of scope




Figure 2. [A]: Substrate scope. Isolated yields of ArCF₂H, unless indicated. Ratios in brackets correspond to CF₂H:CFH₂ determined from crude ¹⁹F NMR, n.d. = not-detected. CP = Chronopotentiometry. Benchmarking yields are ¹⁹F NMR yields, except where noted 'b'. ^a¹⁹F NMR yield, not isolated due to volatility. ^b % conversion, ^c CF₂H observed in ¹⁹F NMR of crude mixture, product hydrolysed to corresponding aldehyde on silica, ^d reaction run without TMSCl. [B]: Summed Hammett σ value of successful substrates (see Supporting Information for details).

conditions. Electron neutral substrates **1a–4a** were all well tolerated by our conditions, returning products in good to excellent yields. Yields from the reported methods for this family of substrates were generally comparable to ours. Electron-poor substrate **5a** transferred smoothly to **5b**, without ester reduction, comparing well to the Gouverneur conditions [A] that are specific to electron-poor substrates. However, conditions [B–D] did not provide any product **5b**. Pleasingly our electrochemical conditions could tolerate electron-rich substrates **6a–8a**, including tris(methoxy)phenyl **8a**, whilst reported conditions generally did not convert any of these substrates. CV analysis showed that **6a** and **8a** required a reduction potential ≈ 160 mV and 500 mV deeper, respectively, than **1a**. Finally, pyrazole **7a** was amenable, while all reported methods gave no **7b**.

Encouraged by this, an extensive substrate scope of both commodity and functionalized ArCF_3 was undertaken. Several other electron-rich rings were tested, including aniline derivatives (**10–12b**) and with ether (**13–14b**) or alkyl (**15–21b**) substitution, all of which were well tolerated. More substrates with electron-poor rings (**22–25b**) were tested and found to transform in moderate to good yields, including highly electron-deficient bis-ester **25a**. We tested the conditions on more complex drug-type molecules (**26–33b**). Pleasingly, all these compounds were tolerated and gave products in moderate to good yields, demonstrating the capability of this procedure to previously unreported substrate classes and to late-stage functionalization. Analysis of our scope by the total electronic contribution of the substituents revealed the electronic ambivalence of our method,^[48] (see Supporting Information) Figure 2B.

Several functional handles were well tolerated in the scope, which is essential for diversification and application as building blocks. Such handles include ester, pinacol boronic ester, alkenes, primary alcohols, primary, secondary and tertiary amines, phosphonate and sulfonamides. For nucleophilic groups, such as amines, the exclusion of TMSCl facilitated a more efficient reaction, possibly avoiding competitive reduction of HCl generated from the formation of a N-TMS adduct.^[28] Acetal, ketal and Boc protecting groups were also tolerated, as well as extended ring systems, which can be affected by electrode grafting.^[46,49] Functionality that was not tolerated includes halides (except fluoride), nitro and cyano groups, which can be reductively cleaved (see Supporting Information). We successfully scaled the reaction 10-fold to a gram scale (5 mmol) without a significant decrease in yield or selectivity of product **13b**.

To gain insight into the mechanism and to probe for radical intermediates ($\text{ArCF}_2\cdot$), two radical trap experiments were conducted (Figure 3A). When introducing either an intermolecular (**1a** + amylene) or intramolecular alkene (**17a**) to the system, no products expected from the addition of $\text{ArCF}_2\cdot$ to an alkene were observed, suggesting a rapid second reduction to the ArCF_2^- anion.^[35,36] This contrasts the photochemical HAT methods,^[50] where transient $\text{ArCF}_2\cdot$ radicals trap alkenes, *c.f.* conditions [B] returned no **17b**, Figure 2A. These data provide evidence for a rapid transfer of 2 electrons through an ECEC mechanism (ET/PT, Figure 1C). As the reaction solution is required to be




Figure 3. [A] control experiments to probe for radical or anionic intermediates. [B] ArCFH_2 scope, isolated yields of ArCFH_2 , with ^{19}F NMR yields given in parentheses, ratio is CFH_2 : CF_2H . ^a NMR yield given due to volatility of product.

anhydrous, we were intrigued to probe the origin of the proton. Our previous studies on electrochemical C–F activation found evidence for protonation from the NEt_4^+ cation and not from the solvent.^[45] However, conducting this reaction in anhydrous D_3 -MeCN, **1a** was smoothly transformed to D_1 -**1b** (91% D-incorporation). Though the $\text{p}K_a$ has not been directly determined (due to F^- elimination),^[51,52] indirect evidence suggests the ArCF_2^- anion is sufficiently basic to deprotonate MeCN.^[53]

Motivated by this success, efforts were directed toward the development of direct ArCF_3 di-hydrodefluorination. There are currently no prior reports for this challenging transformation. Nevertheless, we were encouraged when a purified sample of **1b** was transformed under the reaction conditions to **1c** in a good yield (see Supporting Information). As such, a more user friendly one-pot procedure from

ArCF₃ was targeted. After decreasing the substrate concentration, increasing the current density, and passing 4 *F*, a satisfying yield and selectivity was achieved for product **1c**, Figure 3C. A modest scope of substrates was examined, demonstrating moderate to good yields of ArCFH₂ products.^[54] Considering how deep the required reduction potentials are to effect this 4-electron/2-proton transformation, Figure 1D, the substitution tolerated was varied, and complex biologically relevant molecules were converted (**2–34c**). Of additional interest, biaryl **34a** was better tolerated in the *di*- than in the *mono*-hydrodefluorination, which is likely due to resonance stabilization of the intermediate radical anion.^[55,56]

In conclusion, we have developed an electronically ambivalent *mono*-hydrodefluorination reaction of readily accessed ArCF₃ substrates. The reactivity and selectivity are enabled by the electrochemical reduction on a Ni cathode at deeply reducing potentials. A broad range of ArCF₂H compounds are prepared, including pharmaceutically and biologically relevant targets. The methodology is shown to be scalable for med-chem use, and evidence was gained for an ECEC mechanism. The strategy was extended to a *di*-hydrodefluorination reaction, yielding ArCFH₂ compounds. Hence, the rapid construction of fluoromethyl motifs from ArCF₃ groups up-values an inexpensive, readily-available feedstock to more desirable and functionally diverse building blocks, which should aid SAR studies and library synthesis.

Acknowledgements

The authors acknowledge funding from the Royal Society (University Research Fellowship, Research Grant for Research Fellow and Enhancement Awards to A.J.J.L.), EPSRC (EP/S018050/1) and GSK (iCASE award).

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available in the supplementary material of this article.

Keywords: Defluorination • Electrochemistry • Fluorine • Nickel • Reduction

-
- [1] J. Wang, M. Sánchez-Roselló, J. L. Aceña, C. Del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok, H. Liu, *Chem. Rev.* **2014**, *114*, 2432–2506.
 [2] Y. Zafrani, S. Saphier, E. Gershonov, *Future Med. Chem.* **2020**, *12*, 361–365.
 [3] Y. Ogawa, E. Tokunaga, O. Kobayashi, K. Hirai, N. Shibata, *iScience* **2020**, *23*, 101467.

- [4] Y. Zhou, J. Wang, Z. Gu, S. Wang, W. Zhu, J. L. Acenã, V. A. Soloshonok, K. Izawa, H. Liu, *Chem. Rev.* **2016**, *116*, 422–518.
 [5] W. K. Hagmann, *J. Med. Chem.* **2008**, *51*, 4359–4369.
 [6] K. Müller, C. Faeh, F. Diederich, *Science* **2007**, *317*, 1881–1886.
 [7] B. Jeffries, Z. Wang, H. R. Felstead, J. Y. Le Questel, J. S. Scott, E. Chiarparin, J. Graton, B. Linclau, *J. Med. Chem.* **2020**, *63*, 1002–1031.
 [8] Y. Zafrani, D. Yeffet, G. Sod-Moriah, A. Berliner, D. Amir, D. Marciano, E. Gershonov, S. Saphier, *J. Med. Chem.* **2017**, *60*, 797–804.
 [9] J. A. Erickson, J. I. McLoughlin, *J. Org. Chem.* **1995**, *60*, 1626–1631.
 [10] Y. Zafrani, G. Sod-Moriah, D. Yeffet, A. Berliner, D. Amir, D. Marciano, S. Elias, S. Katalan, N. Ashkenazi, M. Madmon, E. Gershonov, S. Saphier, *J. Med. Chem.* **2019**, *62*, 5628–5637.
 [11] C. D. Sessler, M. Rahm, S. Becker, J. M. Goldberg, F. Wang, S. J. Lippard, *J. Am. Chem. Soc.* **2017**, *139*, 9325–9332.
 [12] N. A. Meanwell, *J. Med. Chem.* **2018**, *61*, 5822–5880.
 [13] A. S. Nair, A. K. Singh, A. Kumar, S. Kumar, S. Sukumaran, V. P. Koyiparambath, L. K. Pappachen, T. M. Rangarajan, H. Kim, B. Mathew, *Processes* **2022**, *10*, 2054.
 [14] A survey was conducted on the number of ArCF₂H containing compounds reported in *J. Med. Chem.* (Nov 2022), see Supporting Information for data.
 [15] a) B. Zheng, S. V. D'Andrea, L. Q. Sun, A. X. Wang, Y. Chen, P. Hrcnciar, J. Friborg, P. Falk, D. Hernandez, F. Yu, A. K. Sheaffer, J. O. Knipe, K. Mosure, R. Rajamani, A. C. Good, K. Kish, J. Tredup, H. E. Klei, M. Paruchuri, A. Ng, Q. Gao, R. A. Rampulla, A. Mathur, N. A. Meanwell, F. McPhee, P. M. Scola, *ACS Med. Chem. Lett.* **2018**, *9*, 143–148; b) V. S. Kubyshev, P. K. Mykhailiuk, S. Afonin, A. S. Ulrich, I. V. Komarov, *Org. Lett.* **2012**, *14*, 5254–5257.
 [16] K. Liu, H. Kokubo, *J. Comput.-Aided Mol. Des.* **2019**, *33*, 345–356.
 [17] Q. A. Huchet, B. Kuhn, B. Wagner, N. A. Kratochwil, H. Fischer, M. Kansy, D. Zimmerli, E. M. Carreira, K. Müller, *J. Med. Chem.* **2015**, *58*, 9041–9060.
 [18] J. Jurczyk, J. Woo, S. F. Kim, B. D. Dherange, R. Sarpong, M. D. Levin, *Nat. Synth.* **2022**, *1*, 352–364.
 [19] C. Hui, Z. Wang, S. Wang, C. Xu, *Org. Chem. Front.* **2022**, *9*, 1451–1457.
 [20] J. B. I. Sap, C. F. Meyer, N. J. W. Straathof, N. Iwumene, C. W. Am Ende, A. A. Trabanco, V. Gouverneur, *Chem. Soc. Rev.* **2021**, *50*, 8214–8247.
 [21] D. E. Yerien, S. Barata-Vallejo, A. Postigo, *Chem. Eur. J.* **2017**, *23*, 14676–14701.
 [22] C. Saboureau, M. Troupel, S. Sibille, J. Périchon, *J. Chem. Soc. Chem. Commun.* **1989**, 1138–1139.
 [23] O. Sock, M. Troupel, J. Périchon, *Tetrahedron Lett.* **1985**, *26*, 1509–1512.
 [24] H. Hebri, E. Duñach, J. Périchon, *Synth. Commun.* **1991**, *21*, 2377–2382.
 [25] C. P. Andrieux, A. Le Gorande, J. M. Savéant, *J. Am. Chem. Soc.* **1992**, *114*, 6892–6904.
 [26] C. P. Andrieux, C. Combellas, F. Kanoufi, J. M. Savéant, A. Thiébaud, *J. Am. Chem. Soc.* **1997**, *119*, 9527–9540.
 [27] P. Clavel, M. P. Léger-Lambert, C. Biran, F. Serein-Spirau, M. Bordeau, N. Roques, H. Marzouk, *Synthesis (Stuttg.)* **1999**, 1999, 829–834.
 [28] P. Clavel, G. Lessene, C. Biran, M. Bordeau, N. Roques, S. Trévin, D. de Montauzon, *J. Fluorine Chem.* **2001**, *107*, 301–310.
 [29] J. L. Röckl, E. L. Robertson, H. Lundberg, *Org. Biomol. Chem.* **2022**, *20*, 6707–6720.
 [30] L. Radom, W. J. Hehre, J. A. Pople, *J. Am. Chem. Soc.* **1971**, *93*, 289–300.

- [31] P. Kirsch, *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*, Wiley, Blackwell, **2005**.
- [32] H. A. Bent, *Chem. Rev.* **1961**, *61*, 275–311.
- [33] P. J. Elving, J. T. Leone, *J. Am. Chem. Soc.* **1957**, *79*, 1546–1550.
- [34] H. Lund, I. Fischmeister, E. Stenhagen, G. Andersson, E. Stenhagen, H. Palmstierna, *Acta Chem. Scand.* **1959**, *13*, 192–194.
- [35] J. P. Coleman, H. G. Gilde, J. H. P. Utley, B. C. L. Weedon, *J. Chem. Soc. D* **1970**, 738–739.
- [36] J. P. Coleman, N.-U.-D. Naser-Ud-Din, H. G. Gilde, J. H. P. Utley, B. C. L. Weedon, L. Ebersson, *J. Chem. Soc. Perkin Trans. 2* **1973**, 1903–1908.
- [37] S. B. Munoz, C. Ni, Z. Zhang, F. Wang, N. Shao, T. Mathew, G. A. Olah, G. K. S. Prakash, *Eur. J. Org. Chem.* **2017**, 2322–2326.
- [38] The alternative approach using metal(oid) hydrides readily leads to exhaustive hydrodefluorination, see H. Dang, A. M. Whittaker, G. Lalic, *Chem. Sci.* **2016**, *7*, 505–509.
- [39] D. B. Vogt, C. P. Seath, H. Wang, N. T. Jui, *J. Am. Chem. Soc.* **2019**, *141*, 13203–13211.
- [40] J. B. I. Sap, N. J. W. Straathof, T. Knauber, C. F. Meyer, C. F. Meyer, M. Médebielle, L. Buglioni, C. Genicot, A. A. Trabanco, T. Noël, C. W. Am Ende, V. Gouverneur, *J. Am. Chem. Soc.* **2020**, *142*, 9181–9187.
- [41] C. Liu, K. Li, R. Shang, *ACS Catal.* **2022**, *12*, 4103–4109.
- [42] J. Mattay, *Synthesis (Stuttg.)* **1989**, *1989*, 233–252.
- [43] I. A. MacKenzie, L. Wang, N. P. R. Onuska, O. F. Williams, K. Begam, A. M. Moran, B. D. Dunietz, D. A. Nicewicz, *Nature* **2020**, *580*, 76–80.
- [44] N. Elgrishi, K. J. Rountree, B. D. McCarthy, E. S. Rountree, T. T. Eisenhart, J. L. Dempsey, *J. Chem. Educ.* **2018**, *95*, 197–206.
- [45] J. R. Box, A. P. Atkins, A. J. J. Lennox, *Chem. Sci.* **2021**, *12*, 10252–10258.
- [46] D. M. Heard, A. J. J. Lennox, *Angew. Chem. Int. Ed.* **2020**, *59*, 18866–18884; *Angew. Chem.* **2020**, *132*, 19026–19044.
- [47] L. Deng, Y. Zhang, R. Wang, M. Feng, X. Niu, L. Tan, Y. Zhu, *ACS Appl. Mater. Interfaces* **2019**, *11*, 22449–22456.
- [48] C. Hansch, A. Leo, R. W. Taft, *Chem. Rev.* **1991**, *91*, 165–195.
- [49] S. N. Mailu, T. T. Waryo, P. M. Ndangili, F. R. Ngece, A. A. Baleg, P. G. Baker, E. I. Iwuoha, *Sensors* **2010**, *10*, 9449–9465.
- [50] H. Wang, N. T. Jui, *J. Am. Chem. Soc.* **2018**, *140*, 163–166.
- [51] A. Streitwieser, F. Mares, *J. Am. Chem. Soc.* **1968**, *90*, 2444–2445.
- [52] L. Wang, J. Wei, R. Wu, G. Cheng, X. Li, J. Hu, Y. Hu, R. Sheng, *Org. Chem. Front.* **2017**, *4*, 214–223.
- [53] Szymczak reported the deprotonation of MeCN ($pK_a=31.3$, DMSO) from KCH₂Ph (a) J. B. Geri, E. Y. Aguilera, N. K. Szymczak, *Chem. Commun.* **2019**, *55*, 5119–5122 and b) J. B. Geri, M. M. Wade Wolfe, N. K. Szymczak, *J. Am. Chem. Soc.* **2018**, *140*, 9404–9408.), which has a pKa of 42 in DMSO, see reference 51. This evidence implies Ar-CF₂⁻ should be sufficiently basic to deprotonate MeCN.
- [54] Isolated yields may be slightly diminished due to incompatibility of benzyl fluorides with glass: a) M. M. Nielsen, C. M. Pedersen, *Chem. Sci.* **2022**, *13*, 6181–6196, or instability towards elimination or hydrolysis, see reference 52.
- [55] N. L. Holy, *Chem. Rev.* **1974**, *74*, 243–277.
- [56] H. Senboku, M. Takahashi, T. Fukuhara, S. Hara, *Chem. Lett.* **2007**, *36*, 228–229.

Manuscript received: December 12, 2022

Accepted manuscript online: January 27, 2023

Version of record online: February 10, 2023