

REVIEW ARTICLE

Pooling the evidence: A review of swimming and atopic dermatitis

Cathal O'Connor MD^{1,2,3,4}  | Siobhan McCarthy PhD^{1,3}  | Michelle Murphy MD^{1,3} ¹Dermatology, South Infirmary Victoria University Hospital, Cork, Ireland²Paediatrics and Child Health, Cork University Hospital, Cork, Ireland³Medicine, University College Cork, Cork, Ireland⁴INFANT Research Centre, University College Cork, Cork, Ireland**Correspondence**

Cathal O'Connor, INFANT Research Centre, University College Cork, Cork, Ireland.

Email: cathal.oconnor@ucc.ie**Funding information**

Health Research Board, Grant/Award Number: 223047/Z/21/Z; Wellcome Trust; Health Service Executive National Doctors Training and Planning; Health and Social Care, Research and Development Division, Northern Ireland

Abstract

Swimming is an excellent form of aerobic exercise and is an essential life skill. Many children with atopic dermatitis (AD) are advised not to swim because of concerns about negative impacts on their skin disease, and some children with AD do not swim because they are self-conscious about the appearance of their skin. We aimed to perform a narrative review of the available literature on swimming and AD and scientifically analyze the potential impact of all components of swimming in AD—water, skin barrier, swimming gear, and exercise. Studies examined the impact of swimming on the skin barrier and the relative contraindications to swimming. Constituents of water which may affect AD include hardness, pH, temperature, antiseptics, and other chemicals. Potential interventions to reduce damage included emollient application, special swim gear, and showering post-submersion. The benefits of swimming as a form of exercise in AD included reduced sweating, cardiorespiratory fitness, and maintenance of healthy weight. Drawbacks of swimming as a form of exercise in AD included the limited benefit on bone mineral density. Future research should examine the impact of swimming on flares of AD using noninvasive biomarkers as well as clinical severity assessment and assess the role for different types of emollient as an intervention for optimal eczema control. This review highlights gaps in the scientific literature on swimming and AD and provides evidence-based guidance on interventions to minimize deleterious effects on skincare and maximize opportunities for children with AD to swim.

KEYWORDS

atopic dermatitis, atopy, exercise, quality of life, swimming

1 | INTRODUCTION

Swimming in childhood is known to be associated with enhanced physical and mental health,¹ as well as being an essential life skill to reduce the risk of drowning.² In particular, outdoor swimming has dramatically increased in popularity during the COVID-19 pandemic, with myriad reported health benefits.³ Swimming is a suitable choice of exercise for children with atopic dermatitis (AD), as the water contact

can be cooling for exposed eczematous skin, and excessive sweating is avoided. However, up to a third of children with AD have disrupted access to swimming due to the severity of their disease.⁴ Some children with AD are advised not to swim due to concerns about potential negative effects on their skin, and some children may be reluctant to swim due to self-consciousness about the visual impact of their eczematous skin in front of others. Potential benefits and risks of swimming for children with AD are outlined in Figure 1.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Pediatric Dermatology* published by Wiley Periodicals LLC.




FIGURE 1 Potential benefits and risk of swimming for children with atopic dermatitis.

AD may have a profoundly negative effect on health-related quality of life (QOL).⁵ Key factors in reduced QOL are lifestyle restrictions or limitations imposed because of AD. Prohibiting children with AD from swimming can lead to deterioration in both physical and mental health. Exclusion from peer activities such as swimming can lead to “othering” of children with AD, and put them at risk of teasing, bullying, and further emotional upset.⁶ Subsequent inability to swim can lead to future embarrassment, and increase the risk of drowning, which is the third leading cause of premature death globally, with children accounting for the highest rate (<https://www.who.int/news-room/fact-sheets/detail/drowning>).

This narrative review aims to synthesize the currently available evidence on swimming and AD, and analyze the components of water,

skin barrier protection, swimming gear, and exercise that may be relevant to children with AD. The content of the review is used to formulate guidance for dermatologists and pediatricians on what advice to provide to children with AD regarding swimming.

2 | METHODS

We performed a narrative review by searching PubMed for English language articles published up until March 10, 2023, including the terms “atopic dermatitis” OR “atopic eczema” OR “eczema” AND “swimming.” This resulted in 46 unique articles which were then screened by title and abstract for pertinence, resulting in 20 articles that were read in full by all authors. Further targeted reviews were performed to identify research on basic science related to the skin barrier and swimming.

3 | RESULTS

3.1 | Current evidence on swimming and AD

One of the key concerns relating to swimming exposure and impact on AD development and severity is the effect of swimming on the skin barrier, which is known to be a primary pathogenic factor in the development and persistence of AD.⁷ Transepidermal water loss (TEWL) is a surrogate marker for skin barrier function, and aberrations in TEWL are known to predate the clinical development of AD.⁸ One study in elite teenage swimmers without AD has shown TEWL increases immediately following 2 h of intensive swimming training, but returns to normal after 30 min.⁹ However, most children are unlikely to spend 2 h in the pool, and TEWL rapidly returns to normal levels after submersion, which is reassuring.

Recently, there has been keen interest in skin barrier-focused interventions for the prevention of AD in early life, with mixed results.¹⁰ The effect of infant swimming and emollient application versus swimming alone was examined in a German study in 2009.¹¹ Children with a personal or parental history of AD were excluded. Both groups went swimming for 25–40 min weekly for 4 weeks ($n = 44$), and the intervention limb ($n = 20$) applied an emollient after swimming. The intervention group had stable sebum and pH 1 week following the final swim, while a significant decrease in sebum and pH was noted in the group that was not moisturized after swimming.

A Spanish birth cohort study examined the impact of indoor and outdoor swimming pool attendance and eczema development during the first year of life.¹² When adjusted for parental atopy, type of swimming pool, cumulative duration of pool exposure, and socioeconomic group, there was no difference in AD prevalence in those who were exposed to swimming in the first year of life and those who were not.

Longitudinal studies such as the Avon Longitudinal Study of Parents and Children cohort study¹³ have failed to show an association between early life exposure to swimming and subsequent development of AD. A birth cohort study from Germany showed no increased rates of AD by 6 years in children with early swimming pool attendance.¹⁴ Another German study that asked adults to retrospectively report

childhood exposure to swimming pools showed no correlation between early swimming and lifetime development of AD,¹⁵ with the significant limitations of recall bias and lack of objectively diagnosed AD.

One Spanish study showed a 40% increased risk of AD development in children who swam before the age of 3 years and a higher risk of AD was observed within the highest tercile of years in swimming practice.¹⁶ However, AD was self-reported, and adjustment for parental socioeconomic group and educational status, factors known to be associated with an increased risk of AD in children,¹⁷ was not performed. Another Spanish study reported increased risk of AD in children who swam in pools before 2 years of age,¹⁸ but AD was parent-reported and the possibility of reverse causation cannot be ruled out. Another study looked at current exposure to swimming and AD outcomes,¹⁹ with no association between current swimming and AD prevalence. However, the authors noted their results may be skewed because children with AD may avoid indoor swimming pools because of potential negative effects on their eczema.

Given parental concerns about potential topical corticosteroid side effects and misinformation related to the treatment of eczema,²⁰ parents may seek complementary and alternative medicine (CAM) therapies for AD. One systematic review and meta-analysis of CAM for treatment of AD showed that children randomized to swimming as therapy improved compared to standard clinical care.²¹ However, the methodological quality of these studies was of unclear or high risk of bias in general, and other CAM therapies in the trials included unproven interventions.

One exploratory study, which closely followed up 60 children with AD for environmental exposures, did not show a link between swimming and daily “bother” score, but did show an association with higher “scratch” scores and “step up treatment” scores.²² However, there was no objective assessment of AD severity, and increased “step up treatment” may relate to increased prophylactic emollient usage around the time of swimming. There is no high-quality evidence available related to swimming exposure and AD outcomes during flares of AD, and this should be explored as a future interventional study. Given the potential for aggravation of skin barrier dysfunction during a flare of AD, it is reasonable to recommend delaying swimming during severe flares. Proactive, rather than reactive, treatment of AD should be optimized to treat disease flares and reduce the frequency of exacerbations.²³

As there is a risk of transmission of bacteria, particularly of contagious staphylococcal species, during periods of impetiginization of AD, it may also be reasonable to delay swimming during infective flares until weeping or crusting has resolved. One Italian guideline recommends avoiding swimming until 24 h after antimicrobial therapy has been initiated and crust has lifted.²⁴ However, chlorinated water has antiseptic properties (as discussed below), so swimming during infective flares may have beneficial effects, which must be weighed against the risk of spreading contagion, for example, via shared towels. In addition, studies in remote Australian indigenous communities have shown that the installation of community swimming pools in fact reduces the prevalence of impetigo and other skin infections.²⁵ Molluscum contagiosum (MC), a common pox virus that affects the skin, occurs more frequently both in children with AD and in children without AD who swim regularly.²⁶ The potential risk of developing MC, a benign and self-limiting infection, should not exclude children

with AD from swimming. If children with AD develop MC, swimming is still permissible, but it is advisable to cover lesions with a water-proof plaster and avoid sharing towels.

3.2 | Water constituents

3.2.1 | Hardness

Water hardness is defined as the concentration of divalent metal cations, such as dissolved calcium (Ca^{++}) and magnesium (Mg^{++}) in a water sample. Calcium chloride is added to pools to keep the water hard. This protects the surfaces of the pool from corrosion. However, calcium in hard water may damage the skin barrier, increasing skin dryness and irritation, and both predispose children to AD and provoke flares of established AD.²⁷ Skin contact with hard water has been associated with increased TEWL, particularly in patients with AD and filaggrin (*FLG*) mutations.²⁸ The interaction between hard water, infant swimming, and AD outcomes has also been examined,²⁹ with a linear relationship between water hardness and AD prevalence, and no independent relationship between infant swimming and AD, although the combination of hard water and infant swimming was synergistic in increasing AD prevalence. However, randomized controlled trials comparing water softeners with standard care have not shown a significant difference in objective AD severity with softened water.²⁷

3.2.2 | pH

pH is the most important factor in swimming pool water chemistry since it affects chlorine efficacy as well as overall balance in the water. Pool water must meet stringent quality standards to provide a healthy experience for swimmers, including protection against the chlorine-resistant pathogen *Cryptosporidium* (<https://www.pwtag.org/>). The ideal pH value to be comfortable for eyes and to prevent corrosive or scale-forming conditions is at a slightly alkaline value in the range of pH 7.2–7.4, while skin on most parts of the body has a pH level that normally ranges from 4.1 to 5.8.³⁰ The acid mantle of the skin is responsible for retaining moisture and essential lipids, and providing a barrier against pathogens, irritants, and allergens. Excessive alkalization of the skin may cause the skin to become dry and irritated, potentially leading to AD.

3.2.3 | Temperature

Children with eczema may prefer lower pool temperatures (<https://nationaleczema.org/blog/swimming-eczema/>), and warmer temperatures may increase TEWL and skin pH.³¹

3.2.4 | Antiseptics and other chemicals

Pool water contains multiple chemicals (<https://www.pwtag.org/>) that may dry out or irritate eczematous skin. Chlorine is added to pools for

antiseptic activity, but chlorine exposure can cause irritant contact dermatitis ("pool dermatitis") or allergic contact dermatitis ("pool water dermatitis").³² However, the antiseptic activity of diluted chlorine may reduce microbial colonization and may reduce AD severity, as described with bleach baths.³³ Alternative or additional options for disinfecting swimming pools include salt-water chlorination, bromine, ozone, *polyhexamethylene biguanide*, *algaecides*, and *filter aids, flocculants, and clarifiers*.

3.2.5 | Seawater/saltwater

Anecdotally, many patients with AD report benefit from swimming in seawater or saltwater pools, while some report irritation. Despite salt bathing being a common practice in AD, good-quality evidence is lacking on the effect of salt water on AD outcomes. One small Japanese study showed some benefit from balneotherapy with natural mineral dissolved water.³⁴ Some studies have reported strong associations between marine water contact and staphylococcal skin infections.³⁵

3.3 | Skin barrier protection while swimming

3.3.1 | Emollient barrier

Few studies have examined the benefit of emollient application prior to or after swimming. Garcia Bartels et al. showed persistent enhancement of the cutaneous barrier in infants with emollient application immediately following swimming.¹¹ Future research should examine the impact of application of different emollients before and after swimming on transcutaneous biomarkers in children with AD, as well as clinical severity scoring. Children may tolerate emollients with much higher lipid content during submersion than they would otherwise.

3.4 | Swimming gear and AD

3.4.1 | Residual chlorine/irritants

It has been shown that free residual chlorine in bathing water reduces the water-retaining properties of the stratum corneum in AD.³⁶ If swim wear is not removed quickly following submersion, occlusion, and direct irritation can occur quickly, also predisposing children to folliculitis and bacterial infections.³² It is important to remove all swim wear quickly following exit from the pool, followed by rinsing in fresh unchlorinated water (if available). If swimming at the beach parents could consider bringing a container of fresh water to rinse off after exiting the seawater.

3.4.2 | Contact dermatitis to swim gear

Contact with certain swim gear like goggles, scuba masks, and bathing caps can cause irritation or can sensitize children with AD to various rubber constituents, which may require patch testing for confirmation of allergic contact dermatitis.³²

TABLE 1 Advice from patient information leaflets and websites from major patient organizations and dermatology associations (<https://nationaleczema.org>, <https://eczema.org>, <https://www.aad.org>, <https://www.bad.org.uk>).

Organization	Advice
Eczema Foundation	Apply emollient before swimming ^a Rinse skin post swimming and pat dry ^a Wash at home with shower gel containing copper and zin Apply emollient ± topical corticosteroid ^a
National Eczema Society	Avoid swimming if eczema is flaring badly or infected If swimming indoors, apply emollient (preferably ointment) liberally before entering pool ^a If swimming outdoors, apply sunscreen (preferably containing titanium and/or zinc oxide) Shower immediately after swimming using an emollient and re-apply emollient ^a Do not hang around the pool after as skin is still exposed to chlorinated fumes Consider wearing UV-protection swim suits/clothing for self-conscious children If pool water at one pool is irritating, avoid swimming immediately after chlorine is added to that pool, try other pools or a salt-water pool, or try swimming in fresh water or sea water At a new pool, try a short test period to assess your skin's response to the water
American Association of Dermatology	Apply moisturizer before swimming ^a Rinse the skin with warm water after swimming ^a Immediately pat the skin dry after rinsing and apply moisturizer ^a
British Association of Dermatology	Rinse well after swimming and apply plenty of moisturizer after drying ^a Make sure the shower at the swimming pool contains fresh water and not chlorinated water from the swimming pool

^aAdvice based on clinical or scientific studies.

3.5 | Exercise in AD

There is growing awareness of the associations between AD and obesity and cardiovascular disease.³⁷ It is important for dermatologists to encourage children with AD to foster healthy lifestyles which will enhance their lifelong well-being. Sleep disturbance is a cardinal feature of AD, and exercise such as swimming is a valid nonpharmacologic treatment for sleep disruption.³⁸ Moreover, swimming is a form of exercise that is associated with less perceptible sweating, which is advantageous for children with AD who have significant heat intolerance. Importantly, given the strong link between AD and asthma, there is no strong evidence showing that swimming exposure causes subsequent asthma development or exacerbation. As swimming is a low-contact and nonweight bearing form exercise with



FIGURE 2 Recommendations from the authors for swimming for children with atopic dermatitis.

minimal benefit on bone mineral density,³⁹ it is important to consider other weight-bearing forms of exercise in conjunction with swimming, as patients with AD are known to have an increased risk of osteoporosis.⁴⁰

4 | ADVICE FOR SWIMMING FOR CHILDREN WITH AD

Current guidance from eczema support organizations for swimming for children with AD is highlighted in Table 1. Following the extensive literature review on the relationship between swimming and AD, we have formulated the package of advice seen in Figure 2. The following advice is based on a combination of the currently available scientific evidence, and the opinion of the authors.

5 | CONCLUSION

Swimming is an essential life skill, and children with AD should not miss out on this enjoyable hobby and exercise because of unfounded concerns. Our work identifies significant gaps in the literature on the optimal advice on swimming for children living with AD but provides

suggestions for providers based on current evidence and our opinions when evidence was lacking.

Future research should examine the impact of swimming on flares of AD using clinical severity assessment and noninvasive biomarkers and assess the role of different interventions to optimize eczema control when swimming.

AUTHOR CONTRIBUTIONS

Cathal O'Connor identified the literature gap on the topic, reviewed the literature, wrote the manuscript, and reviewed the manuscript. Siobhan McCarthy and Michelle Murphy reviewed the manuscript and provided feedback. Siobhan McCarthy produced a patient information leaflet based on the literature review.

FUNDING INFORMATION

Dr Cathal O'Connor is funded by the Irish Clinical Academic Training (ICAT) program, supported by the Wellcome Trust and the Health Research Board (grant number 223047/Z/21/Z); the Health Service Executive National Doctors Training and Planning; and the Health and Social Care, Research and Development Division, Northern Ireland.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data derived from public domain resources.

ORCID

Cathal O'Connor <https://orcid.org/0000-0001-7084-5293>

Siobhan McCarthy <https://orcid.org/0000-0002-1098-2963>

Michelle Murphy <https://orcid.org/0000-0003-2431-076X>

REFERENCES

- Thomson H, Kearns A, Petticrew M. Assessing the health impact of local amenities: a qualitative study of contrasting experiences of local swimming pool and leisure provision in two areas of Glasgow. *J Epidemiol Community Health*. 2003;57(9):663-667. doi:10.1136/jech.57.9.6632
- Morrongiello BA, Sandomierski M, Spence JR. Changes over swim lessons in parents' perceptions of children's supervision needs in drowning risk situations: "His swimming has improved so now he can keep himself safe". *Health Psychol*. 2014;33(7):608-615. doi:10.1037/a00338813
- Massey H, Gorczynski P, Harper CM, et al. Perceived impact of outdoor swimming on health: web-based survey. *Interact J Med Res*. 2022;11(1):e25589. doi:10.2196/255894
- Hon KL, Leung TF, Wong KY, Chow CM, Chuh A, Ng PC. Does age or gender influence quality of life in children with atopic dermatitis? *Clin Exp Dermatol*. 2008;33(6):705-709. doi:10.1111/j.1365-2230.2008.02853.x
- Lewis-Jones S. Quality of life and childhood atopic dermatitis: the misery of living with childhood eczema. *Int J Clin Pract*. 2006;60(8):984-992. doi:10.1111/j.1742-1241.2006.01047.x
- Magin P. Appearance-related bullying and skin disorders. *Clin Dermatol*. 2013;31(1):66-71. doi:10.1016/j.clindermatol.2011.11.0097
- Irvine AD, McLean WH. Breaking the (un)sound barrier: filaggrin is a major gene for atopic dermatitis. *J Invest Dermatol*. 2006;126(6):1200-1202. doi:10.1038/sj.jid.5700365

8. Flohr C, England K, Radulovic S, et al. Filaggrin loss-of-function mutations are associated with early-onset eczema, eczema severity and transepidermal water loss at 3 months of age. *Br J Dermatol*. 2010; 163(6):1333-1336. doi:10.1111/j.1365-2133.2010.10068.x
9. Paciência I, Rodolfo A, Leão L, et al. Effects of exercise on the skin epithelial barrier of young elite athletes-swimming comparatively to non-water sports training session. *Int J Environ Res Public Health*. 2021;18(2):653. doi:10.3390/ijerph18020653
10. Kelleher MM, Cro S, Cornelius V, et al. Skin care interventions in infants for preventing eczema and food allergy. *Cochrane Database Syst Rev*. 2021;2(2):CD013534. doi:10.1002/14651858.CD013534.pub2
11. Garcia Bartels N, Rösler S, Martus P, et al. Effect of baby swimming and baby lotion on the skin barrier of infants aged 3-6 months. *J Dtsch Dermatol Ges*. 2011;9(12):1018-1025. doi:10.1111/j.1610-0387.2011.07710.x
12. Font-Ribera L, Villanueva CM, Ballester F, et al. Swimming pool attendance, respiratory symptoms and infections in the first year of life. *Eur J Pediatr*. 2013;172(7):977-985. doi:10.1007/s00431-013-1975-x
13. Font-Ribera L, Villanueva CM, Nieuwenhuijsen MJ, Zock JP, Kogevinas M, Henderson J. Swimming pool attendance, asthma, allergies, and lung function in the Avon longitudinal study of parents and children cohort. *Am J Respir Crit Care Med*. 2011;183(5):582-588. doi:10.1164/rccm.201005-0761OC
14. Schoefer Y, Zutavern A, Brockow I, et al. Health risks of early swimming pool attendance. *Int J Hyg Environ Health*. 2008;211(3-4):367-373. doi:10.1016/j.ijheh.2007.08.001
15. Kohlhammer Y, Döring A, Schäfer T, Wichmann HE, Heinrich J, KORA Study Group. Swimming pool attendance and hay fever rates later in life. *Allergy*. 2006;61(11):1305-1309. doi:10.1111/j.1398-9995.2006.01229.x
16. Cavaleiro Rufo J, Paciência I, Silva D, et al. Swimming pool exposure is associated with autonomic changes and increased airway reactivity to a beta-2 agonist in school aged children: a cross-sectional survey. *PLoS One*. 2018;13(3):e0193848. doi:10.1371/journal.pone.0193848
17. Bajwa H, Baghchechi M, Mujahid M, Kang Dufour MS, Langan SM, Abuabara K. Mixed evidence on the relationship between socioeconomic position and atopic dermatitis: a systematic review. *J Am Acad Dermatol*. 2022;86(2):399-405. doi:10.1016/j.jaad.2021.09.018
18. Font-Ribera L, Kogevinas M, Zock JP, Nieuwenhuijsen MJ, Heederik D, Villanueva CM. Swimming pool attendance and risk of asthma and allergic symptoms in children. *Eur Respir J*. 2009;34(6):1304-1310. doi:10.1183/09031936.00180608
19. Andersson M, Hedman L, Nordberg G, Forsberg B, Eriksson K, Rönmark E. Swimming pool attendance is related to asthma among atopic school children: a population-based study. *Environ Health*. 2015;14:37. doi:10.1186/s12940-015-0023-x
20. O'Connor C, Murphy M. Scratching the surface: a review of online misinformation and conspiracy theories in atopic dermatitis. *Clin Exp Dermatol*. 2021;46(8):1545-1547. doi:10.1111/ced.14679
21. Lu CL, Liu XH, Stub T, et al. Complementary and alternative medicine for treatment of atopic eczema in children under 14 years old: a systematic review and meta-analysis of randomized controlled trials. 2019;19(1):16. *BMC Complement Altern Med*. 2018;18(1):260. doi:10.1186/s12906-018-2306-6
22. Langan SM, Silcocks P, Williams HC. What causes flares of eczema in children? *Br J Dermatol*. 2009;161(3):640-646. doi:10.1111/j.1365-2133.2009.09320.x
23. Brar KK, Nicol NH, Boguniewicz M. Strategies for successful Management of Severe Atopic Dermatitis. *J Allergy Clin Immunol Pract*. 2019; 7(1):1-16. doi:10.1016/j.jaip.2018.10.021
24. Assessorato alla Sanità Segreteria Sanità e Sociale Direzione Prevenzione. Manuale per la prevenzione delle malattie infettive nelle comunità infantili e scolastiche. *Linee Guida Regione Veneto*. 2010;1:12.
25. May PJ, Tong SYC, Steer AC, et al. Treatment, prevention and public health management of impetigo, scabies, crusted scabies and fungal skin infections in endemic populations: a systematic review. *Trop Med Int Health*. 2019;24(3):280-293. doi:10.1111/tmi.13198
26. Olsen JR, Gallacher J, Piguat V, Francis NA. Epidemiology of molluscum contagiosum in children: a systematic review. *Fam Pract*. 2014; 31(2):130-136. doi:10.1093/fampra/cmt075
27. Jabbar-Lopez ZK, Ung CY, Alexander H, et al. The effect of water hardness on atopic eczema, skin barrier function: a systematic review, meta-analysis. *Clin Exp Allergy*. 2021;51(3):430-451. doi:10.1111/cea.13797
28. Danby SG, Brown K, Wigley AM, et al. The effect of water hardness on surfactant deposition after washing and subsequent skin irritation in atopic dermatitis patients and healthy control subjects. *J Invest Dermatol*. 2018;138(1):68-77. doi:10.1016/j.jid.2017.08.037
29. Chaumont A, Voisin C, Sardella A, Bernard A. Interactions between domestic water hardness, infant swimming and atopy in the development of childhood eczema. *Environ Res*. 2012;116:52-57. doi:10.1016/j.envres.2012.04.013
30. Kulthanan K, Nuchkull P, Varothai S. The pH of water from various sources: an overview for recommendation for patients with atopic dermatitis. *Asia Pac Allergy*. 2013;3(3):155-160. doi:10.5415/apallergy.2013.3.3.155
31. Herrero-Fernandez M, Montero-Vilchez T, Diaz-Calvillo P, Romera-Vilchez M, Buendia-Eisman A, Arias-Santiago S. Impact of water exposure and temperature changes on skin barrier function. *J Clin Med*. 2022;11(2):298. doi:10.3390/jcm11020298
32. Tloughan BE, Podjasek JO, Adams BB. Aquatic sports dermatoses: part 1. In the water: freshwater dermatoses. *Int J Dermatol*. 2010;49(8): 874-885. doi:10.1111/j.1365-4632.2010.04536.x
33. Maarouf M, Shi VY. Bleach for atopic dermatitis. *Dermatitis*. 2018; 29(3):120-126. doi:10.1097/DER.0000000000000358
34. Uchiyama A, Fujiwara C, Inoue Y, et al. The effect of balneotherapy with natural mineral dissolved water on dry skin in atopic dermatitis: a phase IIa, nonrandomized, controlled study. *J Cutaneous Immunol Allerg*. 2021;4(6):159-165. doi:10.1002/cia2.12195
35. Tloughan BE, Podjasek JO, Adams BB. Aquatic sports dermatoses. Part 2 - in the water: saltwater dermatoses. *Int J Dermatol*. 2010;49(9): 994-1002. doi:10.1111/j.1365-4632.2010.04476.x
36. Seki T, Morimatsu S, Nagahori H, Morohashi M. Free residual chlorine in bathing water reduces the water-holding capacity of the stratum corneum in atopic skin. *J Dermatol*. 2003;30(3):196-202. doi:10.1111/j.1346-8138.2003.tb00371.x
37. Paller A, Jaworski JC, Simpson EL, et al. Major comorbidities of atopic dermatitis: beyond allergic disorders. *Am J Clin Dermatol*. 2018;19(6): 821-838. doi:10.1007/s40257-018-0383-4
38. Kline CE. The bidirectional relationship between exercise and sleep: implications for exercise adherence and sleep improvement. *Am J Lifestyle Med*. 2014;8(6):375-379. doi:10.1177/1559827614544437
39. Gomez-Bruton A, Montero-Marín J, González-Agüero A, et al. The effect of swimming during childhood and adolescence on bone mineral density: a systematic review and meta-analysis. *Sports Med*. 2016;46(3):365-379. doi:10.1007/s40279-015-0427-3
40. Mukovozov IM, Morra DE, Giustini D, Tadrous M, Cheung AM, Drucker AM. Atopic dermatitis and bone health: a systematic review. *J Eur Acad Dermatol Venerol*. 2021;35(3):615-628. doi:10.1111/jdv.16895

How to cite this article: O'Connor C, McCarthy S, Murphy M. Pooling the evidence: A review of swimming and atopic dermatitis. *Pediatr Dermatol*. 2023;40(3):407-412. doi:10.1111/pde.15325