

Detailed Analysis of *ITPR1* Missense Variants Guides Diagnostics and Therapeutic Design

Jussi Pekka Tolonen, MD, PhD,^{1,2}  Ricardo Parolin Schnekenberg, MD,^{1,3}  Simon McGowan, PhD,⁴ David Sims, PhD,⁴ Meriel McEntagart, FRCP,⁵ Frances Elmslie, MD, FRCP,⁵ Debbie Shears, MBBS, PhD,³ Helen Stewart, MD, FRCP,³ George K. Tofaris, FRCP, PhD,^{1,2}  Tabib Dabir, MD, DCH, MRCP,⁶ Patrick J. Morrison, DSc,⁷ Diana Johnson, BM, BSc, MD,⁸ Marios Hadjivassiliou, MD, PhD,⁹ Sian Ellard, PhD, FRCPATH,¹⁰ Charles Shaw-Smith, MRCP, PhD,¹¹ Anna Znaczkó, MRCP,¹¹ Abhijit Dixit, FRCPEDIN,¹² Mohnish Suri, FRCP,¹² Ajoy Sarkar, MRCPCH, FRCPEDIN,¹² Rachel E. Harrison, BM, BS, PhD,¹² Gabriela Jones, MBChB, BSc,¹² Henry Houlden, MD, PhD,¹³  Giorgia Ceravolo, MD,^{13,14} Joanna Jarvis, BM, MD,¹⁵ Jonathan Williams, FRCPATH, PhD,¹⁶ Morag E. Shanks, PhD,¹⁶ Penny Clouston, FRCPATH, PhD,¹⁶ Julia Rankin, MD,¹⁷ Lubov Blumkin, MD,^{18,19} Tally Lerman-Sagie, MD, PhD,^{18,20} Penina Ponger, MD,^{18,21} Salmo Raskin, MD, PhD,²² Katariina Granath, BM,²³ Johanna Uusimaa, MD, PhD,²³ Hector Conti, MD,²⁴ Emma McCann, MBChB, BSc,²⁵ Shelagh Joss, MBChB, Med,²⁶ Alexander J.M. Blakes, BMCh, MSc,^{27,28}  Kay Metcalfe, FRCP, MD,^{27,28} Helen Kingston, MD, FRCP,²⁸ Marta Bertoli, MD,²⁹ Rachel Kneen, BM, BS,³⁰ Sally Ann Lynch, MD,³¹ Inmaculada Martínez Albaladejo, MD,³² Austen Peter Moore, FRCP,³³ Wendy D. Jones, MRCP, PhD,³⁴ Genomics England Research Consortium,³⁵ Esther B.E. Becker, PhD,^{1,2}  and Andrea H. Németh, MBBS, FRCP, DPhil^{1,3*} 

¹Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK

²Kavli Institute of Nanoscience Discovery, University of Oxford, Oxford, UK

³Oxford Center for Genomic Medicine, Oxford University Hospitals National Health Service Foundation Trust, University of Oxford, Oxford, UK

⁴Centre for Computational Biology, MRC Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, UK

⁵South West Regional Genetics Service, St. George's University Hospitals, London, UK

⁶Northern Ireland Regional Genetics Service, Belfast City Hospital, Belfast, UK

⁷Patrick G. Johnston Centre for Cancer Research and Cell Biology, Queen's University Belfast, Belfast, UK

⁸Sheffield Clinical Genetics Service, Sheffield Children's NHS Foundation Trust, Sheffield, UK

⁹Department of Neurology, Royal Hallamshire Hospital, Sheffield Teaching Hospital NHS Foundation Trust, Sheffield, UK

¹⁰Exeter Genomics Laboratory, Royal Devon University Healthcare NHS Foundation Trust, UK

¹¹Peninsula Clinical Genetics Service, Royal Devon University Hospital, Royal Devon University Healthcare NHS Foundation Trust, Exeter, UK

¹²Department of Clinical Genetics, Nottingham University Hospitals NHS Trust, Nottingham, UK

¹³Department of Neuromuscular Disorders, UCL Queen Square Institute of Neurology, University College London, London, UK

¹⁴Unit of Pediatric Emergency, Department of Adult and Childhood Human Pathology, University Hospital of Messina, Messina, Italy

¹⁵Birmingham Women's and Children's NHS Foundation Trust, Birmingham, UK

¹⁶Oxford Regional Genetics Laboratory, Churchill Hospital, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

¹⁷Department of Clinical Genetics, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK

¹⁸Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel

¹⁹Pediatric Movement Disorders Service, Pediatric Neurology Unit, Edith Wolfson Medical Center, Holon, Israel

²⁰Magen Center for Rare Diseases-Metabolic, Neurogenetic, Wolfson Medical Center, Holon, Israel

²¹Movement Disorders Unit, Department of Neurology, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel

²²Genetika Centro de Aconselhamento e Laboratório, Curitiba, Brazil

²³Research Unit of Clinical Medicine, Medical Research Center, Oulu University Hospital and University of Oulu, Oulu, Finland

²⁴All Wales Medical Genomics Service, Wrexham Maelor Hospital, Wrexham, UK

²⁵Liverpool Women's Hospital Foundation Trust, Liverpool, UK

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

***Correspondence to:** Dr. E. Becker, Nuffield Department of Clinical Neurosciences, Dorothy Crowfoot Hodgkin Building, University of Oxford, South Parks Road, Oxford OX1 3QU, UK; E-mail: esther.becker@ndcn.ox.ac.uk; Dr. A. Németh, Nuffield Department of Clinical Neurosciences, University of Oxford, 6th Floor West Wing, John Radcliffe Hospital, Oxford OX3 9DU, UK; E-mail: andrea.nemeth@ndcn.ox.ac.uk

Esther B. E. Becker and Andrea H. Németh share senior authorship.

Relevant conflicts of interest/financial disclosures: The authors declare that no competing interests exist.

Funding agency: This work was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101023312, Nuffield Department of Clinical Neurosciences (University of Oxford), Ataxia UK, Action Medical Research (GN2063), the Wellcome Trust (161/037, 223521/Z/21/Z), Henry Smith Charity, the John Fell Oxford University Press Research Fund, the Medical Research Council (MR/V007068/1), Orionin Tutkimussäätiö, Sigrid Juséliuksen Säätiö, Stiftelsen Alma och K. A. Snellman Säätiö, Lastentautien Tutkimussäätiö, and the Research Council of Finland (356676, 331436).

Received: 30 July 2023; **Revised:** 16 September 2023; **Accepted:** 16 October 2023

Published online 14 November 2023 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.29651

²⁶West of Scotland Centre for Genomic Medicine, Queen Elizabeth University Hospital, Glasgow, UK

²⁷Division of Evolution, Infection and Genomics, School of Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

²⁸Manchester Centre for Genomic Medicine, University of Manchester, St. Mary's Hospital, Manchester Academic Health Science Centre, Manchester, UK

²⁹Northern Genetics Service, International Centre for Life, Newcastle upon Tyne, UK

³⁰Department of Neurology, Alder Hey Children's NHS Foundation Trust, Liverpool, UK

³¹Department of Clinical Genetics, Children's Health Ireland (CHI) at Crumlin, Dublin, Ireland

³²Neurology in Pediatrics, Hospital Santa Lucia, Murcia, Spain

³³The Walton Centre NHS Foundation Trust, Liverpool, UK

³⁴North East Thames Regional Genetics Service, Great Ormond Street Hospital for Children, Great Ormond Street NHS Foundation Trust, London, UK

³⁵Department of Health & Social Care, London, UK

ABSTRACT: Background: The *ITPR1* gene encodes the inositol 1,4,5-trisphosphate (IP₃) receptor type 1 (IP₃R1), a critical player in cerebellar intracellular calcium signaling. Pathogenic missense variants in *ITPR1* cause congenital spinocerebellar ataxia type 29 (SCA29), Gillespie syndrome (GLSP), and severe pontine/cerebellar hypoplasia. The pathophysiological basis of the different phenotypes is poorly understood.

Objectives: We aimed to identify novel SCA29 and GLSP cases to define core phenotypes, describe the spectrum of missense variation across *ITPR1*, standardize the *ITPR1* variant nomenclature, and investigate disease progression in relation to cerebellar atrophy.

Methods: Cases were identified using next-generation sequencing through the Deciphering Developmental Disorders study, the 100,000 Genomes project, and clinical collaborations. *ITPR1* alternative splicing in the human cerebellum was investigated by quantitative polymerase chain reaction.

Results: We report the largest, multinational case series of 46 patients with 28 unique *ITPR1* missense variants. Variants clustered in functional domains of the protein,

especially in the N-terminal IP₃-binding domain, the carbonic anhydrase 8 (CA8)-binding region, and the C-terminal transmembrane channel domain. Variants outside these domains were of questionable clinical significance. Standardized transcript annotation, based on our *ITPR1* transcript expression data, greatly facilitated analysis. Genotype–phenotype associations were highly variable. Importantly, while cerebellar atrophy was common, cerebellar volume loss did not correlate with symptom progression.

Conclusions: This dataset represents the largest cohort of patients with *ITPR1* missense variants, expanding the clinical spectrum of SCA29 and GLSP. Standardized transcript annotation is essential for future reporting. Our findings will aid in diagnostic interpretation in the clinic and guide selection of variants for preclinical studies. © 2023 The Authors. *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society.

Key Words: *ITPR1*; IP₃R1; spinocerebellar ataxia type 29; Gillespie syndrome; cerebellum; next-generation sequencing

Inositol 1,4,5-triphosphate receptors (IP₃R1–3) are critical players in intracellular calcium signaling, mediating the release of calcium ions from the endoplasmic reticulum (ER) into the cytosol.¹ Activation of the tetrameric channel is induced by simultaneous binding of four molecules of inositol 1,4,5-triphosphate (IP₃), one to each subunit comprising the IP₃R channel.^{2,3} The subunits consist of five key domains: an N-terminal suppressor domain, the loss of which appears to increase IP₃ binding to its binding domain⁴; the IP₃-binding domain; a cytosolic regulatory domain; a transmembrane channel domain; and a cytosolic C-terminus. The domains are organized in a tetrameric “mushroom-like” structure, with the stalk inserted into the ER membrane and the cap exposed to the cytosol.⁵ This organization makes the IP₃ binding core accessible to IP₃, and the regulatory domain

available for many protein interactions (eg, with carbonic anhydrase 8, CAR8^{6,7}) and post-translational modifications that regulate the receptor activity.¹ Binding of IP₃ triggers conformational changes that are transmitted intramolecularly over a large distance to open the C-terminal channel pore.^{3,8,9}

Expression of *ITPR1* (OMIM *147265) encoding the type 1 IP₃R is ubiquitous, and IP₃R1 is the predominant neuronal IP₃ receptor enriched in the Purkinje cells of the cerebellar cortex,^{10–13} where it regulates Purkinje cell development and calcium homeostasis.^{14,15} A knockout of the *Itp1* gene in mice results in very early lethality, severe ataxia, and epileptic seizures without an overt cellular phenotype.¹³ In humans, *ITPR1* is considered a hub gene for cerebellar ataxias.¹⁶ Pathogenic variants in *ITPR1* cause neurodegenerative spinocerebellar ataxia

type 15 (SCA15)¹⁷ (OMIM #606658), and congenital SCA29¹⁸ (OMIM #117360), Gillespie syndrome^{19,20} (OMIM #206700), and severe pontine/cerebellar hypoplasia.²¹ The IP₃R1 channel further contributes to neurodegeneration²² in SCA2^{23,24} (OMIM #183090), SCA3²⁵ (OMIM #109150), Huntington's disease^{26,27} (OMIM #143100), familial Alzheimer's disease^{28,29} (OMIM #607822), and has been implicated in mouse models of *ATM*- and *APTX*-related ataxias,³⁰ making the IP₃R1 channel an attractive druggable target. While the enrichment of *ITPR1* expression in the Purkinje cells may account for the predominantly cerebellar features observed in *ITPR1*-related disorders,^{17-20,31} the pathophysiological basis for these different phenotypes is poorly understood. Furthermore, cerebellar atrophy was recently highlighted as a hallmark of *ITPR1*-related disease,³² but concordance between cerebellar atrophy and symptom progression is unclear.

With increasing interest in disease-specific American College of Medical Genetics and Genomics/American Association of Molecular Pathology (ACMG/AMP) variant interpretation guidelines,³³⁻³⁵ and with prospects for gene-specific therapies being developed, we performed detailed genotype–phenotype analyses of *ITPR1* missense variants. We describe the genetic variants and clinical features for a cohort of 46 patients with early-onset ataxia, highlighting cases where cerebellar atrophy was demonstrated by serial imaging, and correlate the imaging findings with the clinical phenotypes. Genotype–phenotype correlation has previously been hindered by inconsistent transcript annotation in the literature.^{18-20,31,36} We investigated *ITPR1* expression by quantitative polymerase chain reaction (PCR) to determine relative transcript levels and standardized *ITPR1* variant nomenclature. Our data provide valuable information for the clinical interpretation of *ITPR1* missense variants and a focus for future preclinical studies towards new therapeutics.

Methods

The Deciphering Developmental Disorders Cohort

The Deciphering Developmental Disorders (DDD) study investigated children with undiagnosed developmental disorders across the UK and Ireland, utilizing exome sequencing in molecular diagnostics. Written informed consent for all patients was obtained through the DDD study,^{37,38} or directly by their physician.

The DDD diagnostics framework has been described previously.^{39,40} Briefly, fragmented genomic DNA was used for targeted pull-down with a custom Agilent SureSelect 55 MB Exome Plus Enrichment System (Agilent, Santa Clara, CA, USA) and 75-base pair paired-end reads were sequenced on an Illumina HiSeq. Average sequencing

depth (ratio of sequenced bases to targeted bases) was 903 across the whole targeted sequence or 933 across autosomal targets only. Alignment was performed with the Burrows–Wheeler Aligner (v.0.59), and realignment around indels was performed with the Genome Analysis Toolkit (GATK).⁴⁰ Putative de novo variants were identified from exome data with DeNovoGear software.

The DDD study identified 62 patients with variants in *ITPR1*. We filtered these individuals according to: (1) absence of other pathogenic variants; (2) conservation of nucleotides and amino acids; (3) low allele frequency in ExAC and gnomAD; and (4) availability of clinical information. Variants were considered disease-causing if they met the ACMG criteria for likely pathogenic or pathogenic³³ and were associated with an appropriate phenotype. Five cases where a variant of unknown significance (VUS) was considered the most likely cause of the probands' phenotypes were included to increase the likelihood of determining pathogenicity in the future.

Clinical Collaborations

Additional individuals were identified through specialist ataxia or neurogenetics clinics in Brazil, Finland, Ireland, Israel, Italy, Spain, and the UK. Targeted sequencing across a panel of known ataxia genes, or exome sequencing, was performed, and candidate variants were confirmed by Sanger sequencing. Sequencing, sequence analysis, variant calling, and variant annotation were performed according to in-house protocols of the respective National Health Service (NHS) genetics laboratories or external laboratories.

All variants were analyzed by the variant interpretation programme Alamut (<http://www.interactive-biosoftware.com>), and by the standard pathogenicity prediction programmes Polyphen,⁴¹ SIFT,⁴² and CADD.⁴³ Nucleotide conservation was estimated by PhyloP.⁴⁴

The 100,000 Genomes Cohort

The 100,000 Genomes Project, funded by NHS England, investigates patients with undiagnosed rare disease and/or cancer using genome sequencing. Details of the diagnostics pipeline are published.⁴⁵ Briefly, genome sequencing was performed with TruSeq DNA PCR-free sample preparation (Illumina) on a HiSeq 2500 sequencer with mean depth of 32× and a depth greater than 15× for at least 95% of the reference human genome. Genome sequencing reads were aligned to the Genome Reference Consortium human genome build 37 (GRCh37) with Isaac Genome Alignment Software. Family-based variant calling of single-nucleotide variants (SNVs) and indels was performed with Platypus variant caller. Variants were interpreted against the ACMG criteria.

Total RNA Isolation and cDNA Synthesis

Total RNA from human fetal cerebellar tissue (female, 20 post-conception weeks) was acquired

commercially (AMS Biotechnology (Europe) Ltd., Abingdon, UK). Postnatal cerebellar samples were obtained from the Oxford Brain Bank (REC 15/SC/0639). Total RNA was isolated using the RNeasy Mini kit (QIAGEN Ltd., Manchester, UK) according to the manufacturer's instructions and eluted in nuclease-free water. Approximately 30 mg of cortical cerebellar tissue was sonicated for 10 s using a Soniprep 150 Ultrasonic Disintegrator (MSE (UK) Ltd, London, UK) in RLT lysis buffer (QIAGEN). Samples were kept on ice during handling. 1000 ng of total RNA and oligo(dT) primers were used to synthesize cDNA with the SuperScript III First-Strand Synthesis System (Thermo Fisher Scientific, Paisley, UK) according to the manufacturer's instructions. RNA concentrations were measured using a NanoDrop 2000 Spectrophotometer (Thermo Fisher Scientific).

Quantitative PCR

Quantitative PCR (qPCR) was performed using the Fast SYBR Green Master Mix (Thermo Fisher Scientific) with a StepOnePlus Real-Time PCR System (Applied Biosystems, Paisley, UK). Custom qPCR primers were designed to target a region shared by all *ITPR1* transcripts (exon–exon junction between exons 1 and 2) or alternatively spliced isoforms. A duplication event around the S1 site prevented the reliable quantification of the transcripts that lack S1 (ie, S1– transcripts). Thus, the S1+ transcripts are presented as a fraction of total *ITPR1* mRNA. For the S2 and S3 sites, we used one primer pair to quantify transcripts that include the alternatively spliced site (S2+ and S3+) and one primer pair spanning the junction of the flanking sequences (S2– and S3–). The data are presented as a fraction of S+ to S– transcripts. The qPCR primers are listed in Supplementary Table S1. Relative expression levels were normalized to *ACTINB* and *GAPDH* using the standard $2^{-\Delta\Delta CT}$ technique.

Analysis of Single-Nucleus RNA Sequencing Data

The single-nucleus RNA sequencing dataset used in this study was published previously.¹¹ Re-analysis was performed as in the original publication using Seurat⁴⁶ and other necessary packages in R. Data on *ITPR1* in the developing human brain was extracted from the BrainSpan Developmental Transcriptome dataset (RNA-Seq Genome v10).^{47,48}

Results

Novel *ITPR1* Variants and Standardized Variant Nomenclature

To characterize previously unreported cases of spinocerebellar ataxia type 29 (SCA29) and Gillespie

TABLE 1 Summary table of *ITPR1* variants in the present study

Parameter	Percentage % (n)
Variant ^a	
Known	52% (24/46)
Novel	48% (22/46)
Variant origin	
De novo	52% (24/46)
Inherited	9% (4/46)
Unknown	39% (18/46)
Variant classification	
Pathogenic	36% (10/28)
Likely pathogenic	46% (13/28)
Variant of unknown significance (VUS)	18% (5/28)

^aFor each individual; range of individuals with the same variant, 1–7.

syndrome (GLSP), we utilized next-generation sequencing data from the DDD study (Table S2), the 100,000 Genomes Project, and multiple clinical collaborations (see Supplementary Fig. S1 for study flow). Overall, we identified 46 probands who met our filtering criteria with 52% of individuals carrying a known missense variant (Table 1). Some 52% (24/46) of cases were de novo and 82% (23/28) of variants had been classified as disease-causing by the reporting laboratories (Table 1). The molecular characteristics of each variant are shown in Supplementary Tables S3 (DDD, clinical collaborations) and S4 (100,000 Genomes Project).

The 46 probands carried 28 unique *ITPR1* variants, of which 17 had not been reported previously and were distributed in the N-terminus (n = 8), the regulatory domain (n = 4), and the C-terminus (n = 5) (Fig. 1, Supplementary Tables S3 and S4). The first 200 residues in the N-terminus comprise the suppressor domain,⁴ containing a single known pathogenic variant (p.-Arg36Cys) with a gain-of-function effect of increased IP₃ binding.⁴⁹ As part of our study, we identified two cases harboring the same p.Arg36Cys variant (one published recently with limited phenotypic information⁵⁰) and three cases of previously unreported variants p.-Asp34Val and p.Glu106Lys. The aspartate and arginine at positions 34 and 36, respectively, have been shown to regulate the function of the suppressor domain.^{49,51} We therefore confirmed the p.Arg36Cys as a key suppressor domain variant, potentially destabilizing the inhibitory effect this domain has on IP₃ binding.⁴ A review of the literature and our data further highlighted variants p.-Thr267Met, p.Arg269Trp, p.Val1562Met, p.Gly2554Arg, and p.Lys2611del as mutational hotspots (Fig. 1). In our entire dataset, only one case, harboring the p.Val1562Met




FIG. 1. Pathogenic IP_3R1 missense variants cluster in three functional domains. The *ITPR1* gene encodes a protein of 2758 residues with four functional domains (residues 1–223: suppressor [green], residues 226–578: IP_3 -binding [blue], residues 605–2217: regulatory [gray], residues 2227–2758: transmembrane channel [yellow]), and multiple interaction partners. Shown are previously published variants and variants identified as part of the present study, categorized as internal truncating (red square), frameshift (blue circle), missense (yellow circle), and in-frame codon deletion (red circle), and grouped by diagnosis (SCA29/pontocerebellar hypoplasia [PCH], and GLSP). Homozygous variants are shown in red text, whereas novel variants are highlighted on a white background. Variants associated particularly with PCH are shown in blue text. Variants with published experimental validation are further denoted by a blue star. Each variant is listed with the known number of cases per variant indicating multiple mutational hotspots.

variant, was not associated with ataxia but global development delay (Supplementary Table S4). This variant lies within the CAR8-binding region and was the first published missense variant associated with SCA29 of a very mild phenotype.⁵² Finally, in our cohort, the only variants associated with GLSP were p.Gly2554Arg and p-Lys2611del, both previously reported in GLSP.²⁰

Variant calling for the present study revealed significant inconsistencies in *ITPR1* variant nomenclature which confounded variant interpretation. The variable terminology is explained by alternative splicing of the *ITPR1* mRNA at three sites (S1, S2, and S3, corresponding to exons 12, 40–42, and 23, respectively, Supplementary Fig. S2),^{53–55} resulting in at least three different mRNA transcripts being used for variant mapping.^{18–20,31,36} Based on previously published RNA sequencing datasets,^{11,12,47,48} *ITPR1* mRNA is expressed in cerebellar Purkinje cells from early embryonic development to adulthood (Supplementary Fig. S3). Using custom qPCR primers (Supplementary Fig. S2A), we determined that the three splice sites are also expressed from fetal to postnatal

samples (Supplementary Fig. S2B–D). Concurring with previous data in rodents,⁵⁴ the S3 site undergoes a shift towards shorter isoforms whereby the S3– transcripts predominated in the postnatal samples (Supplementary Fig. S2D). Taken together, our data suggest that the three alternatively spliced sites are expressed in the human cerebellum. Consequently, we mapped disease-causing variants, both from published literature (Supplementary Table S5) and our own datasets (Supplementary Tables S3 and S4) to the longest *ITPR1* transcript (NM_001378452.1) (Fig. 1), which contains all three splice sites. This protein isoform is 2758 amino acid residues in length and is listed as the canonical IP_3R1 isoform in UniProt (Q14643-1) and Ensembl (ENST00000649015.2, MANE select transcript).

Core Phenotypes, Neuroradiological Findings, and Atypical Features

Detailed clinical information is summarized in Table 2. Most cases (n = 40, 87%) resemble SCA29

TABLE 2 Summary table of phenotypic findings in the present study

Parameter	Percentage % (n)
Age at onset	
Congenital	48% (13/27)
0–1 y	44% (12/27)
1–5 y	8% (2/27)
Symptom at presentation	
Hypotonia	36% (9/25)
Developmental delay	48% (12/25)
Ophthalmological finding	36% (9/25)
Ataxia	12% (3/25)
Delayed motor milestones	
Independent sitting >1 y	67% (14/21)
Independent walking >2 y	23% (6/26)
Independent walking not attained	50% (13/26)
Developmental delays	
Motor	93% (26/28)
Speech	76% (19/25)
Global	60% (21/35)
Intellectual disability	45% (14/31)
Cerebellar symptoms	
Hypotonia	87% (27/31)
Ataxia	97% (34/35)
Dysmetria	71% (17/24)
Tremor	67% (16/24)
Eye phenotype	
Normal	29% (10/35)
Aniridia ^a	14% (5/35)
Nystagmus	31% (11/35)
Strabismus	9% (3/35)
Ptosis	11% (4/35)

^aAniridia, iris hypoplasia, or a large, non-reactive pupil with irido-lenticular straining.

clinically with five cases of aniridia resulting in the diagnosis of GLSP. Only one individual presented with adult-onset ataxia requiring further evaluation of variant pathogenicity (p.Glu1449Ala, VUS). Some 92% of cases with sufficient data were identified within the first year of life with initial symptoms consisting of hypotonia, delayed developmental milestones, or ophthalmological findings (Table 2). Independent sitting was delayed until after the first year of life in 67% of individuals, and a significant proportion of individuals had

not attained independent walking at last assessment (Table 2). Ataxia was observed in 97% of individuals for whom clinicians had specifically reported presence or absence of ataxia. Intellectual disability was formally diagnosed in 45% of cases. Considering the high proportion of cases with milder cognitive impairments (ie, learning disability, Supplementary Tables S6 and S7), the core phenotype of SCA29 is ataxia with cognitive impairment (Table 2). Nevertheless, our dataset does include individuals with normal cognition (5/34, 13.9%) or limited motor symptoms (4/28, 14.3%), which is in agreement with the range of phenotypes described in published case reports of *ITPR1* missense variants (n = 86, Supplementary Tables S5 and S8). Three patients received a genetic diagnosis of SCA29 as part of this study.

Atypical features and neuroradiological findings are presented in Table 3 and Supplementary Tables S9 and S10. The neuroradiological findings were heterogenous (Table 3, Supplementary Tables S9 and S10). In our dataset, brain imaging of cases was variously reported as unremarkable (28%), cerebellar hypoplasia (24%), or cerebellar atrophy (62%), and occasionally both hypoplasia and atrophy. In those cases in which serial scanning was available (see variants p.Arg269Trp, p.-Leu605Phe, p.Leu1688Pro, p.Ile2439Phe, p.-Gly2554Arg, p.Phe2594Ser, Supplementary Tables S9 and S10), the degree of atrophy did not correlate with the subjective severity of ataxia or intellectual disability, both of which commonly remain stable or may improve. Importantly, in none of the cases with proven cerebellar atrophy was a decline in function or clinical regression reported. Extra-neurological features were reported in several patients without genotype–phenotype association (Table 3, Supplementary Tables S9 and S10). These range from dysmorphic features in the face or extremities to abnormalities in other organ systems such as the heart, where atrioventricular septal defects and pulmonary stenosis are seen in 9% (3/35) and 6% (2/35) of individuals with available data. The individual with adult-onset ataxia (p.Glu1449Ala, VUS) presented with distal sensory neuropathy, which has not previously been reported with *ITPR1* variants.

Pathogenic Missense Variants Cluster in Three Functional Domains

To evaluate genotype–phenotype correlations in the three-dimensional space, we projected published missense variants onto the rat IP₃R1 tetrameric protein structure.⁹ Homozygous truncating variants (p.-His93fs*64, p.Arg728*, and p.Gln1567*) are only reported in GLSP.^{19,56,57} Known missense variants cluster in two major groups: an N-terminal cluster consisting solely of SCA29-associated missense variants (Supplementary Fig. S4A) and a C-terminal cluster in

TABLE 3 Summary of neuroimaging and extra-cerebellar findings in the present study

Parameter	Percentage % (n)
Neuroimaging	
Unremarkable	28% (8/29)
Cerebellar hypoplasia	24% (7/29)
Cerebellar atrophy	62% (18/29)
Cardiovascular abnormalities	
Septal defects	9% (3/35)
Pulmonic stenosis	6% (2/35)
Musculoskeletal abnormalities	
Microcephaly	11% (4/35)
Scoliosis	3% (1/35)
Facial findings	23% (8/35)
Extremities	17% (6/35)
Seizures	9% (3/35)

the channel domain containing both SCA29 and GLSP variants (Supplementary Fig. S4B). However, there were no obvious genotype–phenotype correlations: for example, the C-terminal p.Gly2554Arg, p.Lys2611del, and p.Gly2666Glu variants are mainly associated with GLSP, but there are recent reports of single individuals carrying these variants without aniridia.^{31,58} The N- and C-terminal clusters correlate well with regions of relatively low missense variant density in gnomAD (Supplementary Fig. S5A), indicating that these regions are sensitive to missense variation, whereas the lowest regional missense variant constraint (ie, highest tolerance to missense variation) in ExAC is found in the last third of the regulatory domain (Supplementary Fig. S5B).⁵⁹

Despite the lower missense constraint in the regulatory domain, three *ITPR1* variants have previously been identified in the CAR8-binding region within the regulatory domain,^{31,60} at least two of which impair IP₃R1-CAR8 interaction, thereby releasing IP₃R1 from CAR8-mediated inhibition.⁷ Our dataset includes nine individuals with a variant located in the CAR8-binding region (p.Glu1449Ala (VUS, n = 1), p.Lys1554_Ala1557del (likely pathogenic, n = 1), p.Val1562Met (likely pathogenic, n = 5), p.Leu1688Pro (likely pathogenic, n = 1), and p.Lys1873_Val1874delinsAsnGly (VUS, n = 1)) (Fig. 1), pointing to the CAR8-binding region as an additional cluster of variants. Thus, disease-causing variants in *ITPR1* group in three regions: the N-terminus (49% of variants, 47% of cases), the CAR8-binding region in the regulatory domain (16% of variants, 13% of cases), and the C-terminal

channel domain (35% of variants, 40% of cases) (Fig. 1).

Finally, to ensure that our data were unbiased towards predefined ACMG guidelines, we mapped the *ITPR1* SNVs in the 100,000 Genomes data that were excluded as benign. The 100,000 Genomes dataset contained 225 probands with 193 unique *ITPR1* variants of which 11% were predicted to be deleterious to protein function by Polyphen⁴¹ and Sift⁴²; variants were more frequently classified as deleterious by Polyphen than Sift (29% vs. 13%). Using Human Phenotype Ontology terms, we further narrowed down the number of cases to focus on neurological phenotypes (ataxia [n = 27], global developmental delay [n = 39], and intellectual disability [n = 2]). The suppressor, IP₃-binding, and channel domains contained 2–3 excluded variants each, whereas 30 variants were found in the regulatory domain (Supplementary Fig. S6). When normalized to the number of amino acid residues for each domain, the number of these variants in the regulatory domain is 2–14-fold higher. The data confirm the higher missense tolerance observed for the regulatory domain, and that combined with the clinical phenotypes, variant frequency, and prediction algorithms in each case, these variants are unlikely to be pathogenic (Supplementary Fig. S5B).

Discussion

To our knowledge, this dataset represents the largest cohort of patients with missense variants in *ITPR1*. The data show that these variants cluster around three functional domains of IP₃R1. Two have been previously described: the IP₃-binding domain and the C-terminal channel domain. In addition, we found an enrichment of variants with the core phenotype in the CAR8-binding region. Variants outside these clusters should be interpreted with considerable caution. We have defined the core phenotype of SCA29 and GLSP as ataxia with cognitive impairment and hypothesize that cerebellar atrophy does not correspond with symptom progression. Finally, we have standardized *ITPR1* variant nomenclature to the longest transcript (NM_001378452), enabling easier cross-comparison between datasets. Overall, these results have an important bearing on variant interpretation in clinical practice as well as the pursuit for new therapeutics.

The distribution of variants across different domains of the IP₃R1 protein suggests both loss-of-function (LOF) and gain-of-function (GOF) disease mechanisms (Fig. 2) with important implications for therapeutic development. The majority of SCA29 and GLSP cases harbor variants in the IP₃-binding and channel domains. In the IP₃-binding domain, hotspot variants p.Thr267Met, p.Arg269Trp and others (ie, Tyr567,




FIG. 2. Different functional clusters of IP₃R1 missense variants guide therapeutic design. **(A)** Loss-of-function (LOF) variants such as p.Thr267Met, p.Arg269Trp, and p.Arg568Gly are likely to require potentiators of IP₃R1 channel function to be targeted therapeutically. **(B)** Gain-of-function (GOF) variants p.Arg36Cys, p.Val1562Met, and p.Ser1502Asp interfere with different suppression mechanisms of the IP₃R1 channel potentially requiring IP₃R1 inhibitors as therapeutics. Polyglutamine-expansion disorders (ie, SCA2, SCA3, Huntington's disease, and familial Alzheimer's disease) involving a dysregulated IP₃R1 channel may also respond to IP₃R1 inhibition. The figure shows the protein structure for the rat IP₃R1 (Protein Data Bank: 7LHF) and ataxin-3 (light pink, Protein Data Bank: 3O65).

Arg568) directly impact hydrogen bond formation between IP₃ and its binding domain,^{3,61} significantly reducing IP₃-binding affinities and leading to dominant negative effects. These variants may respond to positive modulators of IP₃R1 channel function. Interestingly, the first IP₃R1 potentiator was recently published with evidence to suggest it may reverse the LOF effect of the p.Thr267Met and p.Arg269Trp variants.⁶² In contrast, variants p.Arg36Cys, p.Val1562Met, and p-Ser1502Asp confer a GOF effect by interfering with different suppression mechanisms of IP₃R1 channel function (ie, destabilizing the suppressor domain and the interaction with CAR8).^{7,49} Although these variants are rare in our dataset, a similar GOF effect is seen in other spinocerebellar ataxias (SCA2, SCA3),

Huntington's disease, familial Alzheimer's disease, and a recent mouse model of IP₃R1 function, again having therapeutic implications.^{23-26,28,29,63} Thus, an IP₃R1 inhibitor may provide a plausible therapeutic option to target a range of disorders involving a dysregulated IP₃R1 channel.

Our SCA29/GLSP cohort reveals variable phenotypic expressivity without correlation with the genotype of the individual, a feature seen in several non-repeat expansion ataxias.⁵⁰ However, our data are the first to evaluate a large cohort of *ITPR1* missense variants alone. This is of relevance, as deletions in *ITPR1* cause SCA15 with a very different phenotype and underlying mechanistic basis from SCA29.⁵⁰ Currently, early-onset *ITPR1*-associated ataxias are diagnosed as GLSP

distinguished by aniridia^{19,20} or SCA29¹⁸ with a few cases of severe pontine/cerebellar hypoplasia.²¹ N-Terminal variants are solely associated with SCA29, while C-terminal variants can cause either GLSP (p-Gly2554Arg and p.Lys2611del) or SCA29. Some evidence points to a downstream transcription initiation site (TSS) 5' to exon 57 of the *ITPR1* gene giving rise to the aniridia in GLSP,⁶⁴ but this TSS does not account for variants that have been associated with both GLSP and SCA29,^{31,58} or homozygous N-terminal truncating variants associated with GLSP.^{19,56,57} Even within the SCA29 diagnosis,^{18,52} the phenotype can range from mild learning disabilities without ataxia to severe, debilitating ataxia and significant intellectual disability. Therefore, the continuum of symptoms is unexplained by current genotype–phenotype evidence and points to the involvement of additional modulators. Nevertheless, the severity of symptoms in SCA29 emphasizes the necessity for early diagnosis and targeted rehabilitation.

Our data highlight two features of SCA29/GLSP that are less characterized: the distinction between cerebellar hypoplasia and atrophy, and extra-neurological features. Available evidence suggests that the natural history of SCA29 and GLSP is non-progressive. However, recent publications have provided evidence that superior vermian and/or hemispheric cerebellar atrophy may represent a hallmark of *ITPR1*-related disorders.^{32,65} We report on at least six individuals in whom cerebellar hypoplasia was excluded by early brain imaging (3 months to 2 years) but who later developed cerebellar atrophy without evidence of clinical regression. At least four individuals were shown to have cerebellar hypoplasia shortly after birth. This distinction between hypoplasia and atrophy is important, as atrophy usually implies neurodegeneration and is unexpected in patients with clinically non-progressive disease. Repeated imaging can distinguish between the two, but we hypothesize that the clinical significance and mechanistic basis of cerebellar atrophy is uncertain. A prospective natural history study of SCA29 utilizing serial scanning and objective disease severity measures is necessary to confirm this finding. Such a study should further aim to provide systematic data on prevalence of *ITPR1*-related diseases, and to add genomic diversity by including individuals from underrepresented populations.

Several individuals in the cohort present with structural malformations (eg, dysmorphic features and congenital heart defects). Previously, *ITPR1* variants have been linked to hemifacial microsomia⁶⁶ and cardiovascular malformations,⁵⁷ mainly in association with GLSP. Interestingly, atrial septal defects and other cardiac abnormalities are seen in at least seven published cases and five individuals in our cohort suggesting a higher-than-expected prevalence considering the size of the cohort and the prevalence of congenital heart

disease in the general population (1.4 and 3.1 per 1000 live births for atrial and ventricular defects, respectively).⁶⁷ IP₃R1 and IP₃R2 have been implicated in the perturbation of cardiogenesis via deficient calcineurin-NFATc signaling,⁶⁸ but causation between dysfunctional IP₃R1 and congenital heart disease remains unclear.

In conclusion, our data provide evidence for a core SCA29/GLSP phenotype resulting from pathogenic IP₃R1 missense variants in specific protein domains. We have standardized *ITPR1* variant nomenclature to enable easier cross-comparison of variant novelty and pathogenicity between datasets. Our data highlight a diverse expression of *ITPR1* transcripts in the cerebellum raising the question whether alternative splicing of *ITPR1* mRNA could help explain why *ITPR1* haploinsufficiency (SCA15) only manifests in adulthood. Finally, we identify key IP₃R1 variants for pre-clinical research to guide therapeutic design for SCA29/GLSP and other disorders involving a dysregulated IP₃R1 channel. ■

Acknowledgments: We thank the patients and their families involved in this study. We acknowledge the Oxford Brain Bank, supported by the Medical Research Council (MRC), Brains for Dementia Research (BDR) (Alzheimer Society and Alzheimer Research UK), Autistica UK, and the NIHR Oxford Biomedical Research Centre. The Deciphering Developmental Disorders (DDD) study presents independent research commissioned by the Health Innovation Challenge Fund (Grant No. HICF-1009-003), a parallel funding partnership between Wellcome and the Department of Health, and the Wellcome Sanger Institute (Grant No. WT098051). The views expressed in this publication are those of the authors and not necessarily those of Wellcome or the Department of Health. The study has UK Research Ethics Committee approval (10/H0305/83, granted by the Cambridge South REC, and GEN/284/12 granted by the Republic of Ireland REC). The research team acknowledges the support of the National Institute for Health Research (NIHR), through the Comprehensive Clinical Research Network. This study makes use of data generated by the DECIPHER community. A full list of centers that contributed to the generation of the data is available from <https://deciphergenomics.org/about/stats> and via email from contact@deciphergenomics.org. Funding for the DECIPHER project was provided by Wellcome (Grant No. WT223718/Z/21/Z). This research was made possible through access to the data and findings generated by the 100,000 Genomes Project. The 100,000 Genomes Project is managed by Genomics England Limited (a wholly owned company of the Department of Health and Social Care). The 100,000 Genomes Project is funded by the NIHR and NHS England. The Wellcome Trust, Cancer Research UK, and the MRC have also funded research infrastructure. The 100,000 Genomes Project uses data provided by patients and collected by the National Health Service as part of their care and support.

Data Availability Statement

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

References

1. Hamada K, Mikoshiba K. IP(3) receptor plasticity underlying diverse functions. *Annu Rev Physiol* 2020;82:151–176.
2. Alzayady KJ, Wang L, Chandrasekhar R, Wagner LE 2nd, Van Petegem F, Yule DL. Defining the stoichiometry of inositol 1,4,5-trisphosphate binding required to initiate Ca²⁺ release. *Ci Signal* 2016;9(422):ra35.

3. Fan G, Baker MR, Terry LE, et al. Conformational motions and ligand-binding underlying gating and regulation in IP(3)R channel. *Nat Commun* 2022;13(1):6942.
4. Yoshikawa F, Morita M, Monkawa T, Michikawa T, Furuichi T, Mikoshiba K. Mutational analysis of the ligand binding site of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* 1996;271(30):18277–18284.
5. Gambardella J, Morelli MB, Wang X, Castellanos V, Mone P, Santulli G. The discovery and development of IP3 receptor modulators: an update. *Expert Opin Drug Discov* 2021;16(6):709–718.
6. Hirota J, Ando H, Hamada K, Mikoshiba K. Carbonic anhydrase-related protein is a novel binding protein for inositol 1,4,5-trisphosphate receptor type 1. *Biochem J* 2003;372(Pt 2):435–441.
7. Ando H, Hirose M, Mikoshiba K. Aberrant IP(3) receptor activities revealed by comprehensive analysis of pathological mutations causing spinocerebellar ataxia 29. *Proc Natl Acad Sci U S A* 2018;115(48):12259–12264.
8. Schmitz EA, Takahashi H, Karakas E. Structural basis for activation and gating of IP(3) receptors. *Nat Commun* 2022;13(1):1408.
9. Baker MR, Fan G, Seryshev AB, Agosto MA, Baker ML, Serysheva II. Cryo-EM structure of type 1 IP(3)R channel in a lipid bilayer. *Commun Biol* 2021;4(1):625.
10. Sharp AH, Nucifora FC Jr, Blondel O, et al. Differential cellular expression of isoforms of inositol 1,4,5-trisphosphate receptors in neurons and glia in brain. *J Comp Neurol* 1999;406(2):207–220.
11. Aldinger KA, Thomson Z, Phelps IG, et al. Spatial and cell type transcriptional landscape of human cerebellar development. *Nat Neurosci* 2021;24(8):1163–1175.
12. Sepp M, Leiss K, Sarropoulos I, et al. Cellular development and evolution of the mammalian cerebellum. *bioRxiv* 2021;2021.12.20.473443.
13. Matsumoto M, Nakagawa T, Inoue T, et al. Ataxia and epileptic seizures in mice lacking type 1 inositol 1,4,5-trisphosphate receptor. *Nature* 1996;379(6561):168–171.
14. Hisatsune C, Kuroda Y, Akagi T, et al. Inositol 1,4,5-trisphosphate receptor type 1 in granule cells, not in Purkinje cells, regulates the dendritic morphology of Purkinje cells through brain-derived neurotrophic factor production. *J Neurosci* 2006;26(42):10916–10924.
15. Sugawara T, Hisatsune C, Le TD, et al. Type 1 inositol trisphosphate receptor regulates cerebellar circuits by maintaining the spine morphology of purkinje cells in adult mice. *J Neurosci* 2013;33(30):12186–12196.
16. Bettencourt C, Rytten M, Forabosco P, et al. Insights from cerebellar transcriptomic analysis into the pathogenesis of ataxia. *JAMA Neurol* 2014;71(7):831–839.
17. van de Leemput J, Chandran J, Knight MA, et al. Deletion at ITPR1 underlies ataxia in mice and spinocerebellar ataxia 15 in humans. *PLoS Genet* 2007;3(6):e108.
18. Synofzik M, Helbig KL, Harmuth F, et al. De novo ITPR1 variants are a recurrent cause of early-onset ataxia, acting via loss of channel function. *Eur J Hum Genet* 2018;26(11):1623–1634.
19. Gerber S, Alzayady KJ, Burglen L, et al. Recessive and dominant de novo ITPR1 mutations cause Gillespie syndrome. *Am J Hum Genet* 2016;98(5):971–980.
20. McEntagart M, Williamson KA, Rainger JK, et al. A restricted repertoire of de novo mutations in ITPR1 cause Gillespie syndrome with evidence for dominant-negative effect. *Am J Hum Genet* 2016;98(5):981–992.
21. van Dijk T, Barth P, Reneman L, Appelhof B, Baas F, Poll-The BT. A de novo missense mutation in the inositol 1,4,5-trisphosphate receptor type 1 gene causing severe pontine and cerebellar hypoplasia: expanding the phenotype of ITPR1-related spinocerebellar ataxia's. *Am J Med Genet A* 2017;173(1):207–212.
22. Egorova PA, Bezprozvanny IB. Inositol 1,4,5-trisphosphate receptors and neurodegenerative disorders. *FEBS J* 2018;285(19):3547–3565.
23. Liu J, Tang TS, Tu H, et al. Deranged calcium signaling and neurodegeneration in spinocerebellar ataxia type 2. *J Neurosci* 2009;29(29):9148–9162.
24. Kasumu AW, Liang X, Egorova P, Vorontsova D, Bezprozvanny I. Chronic suppression of inositol 1,4,5-trisphosphate receptor-mediated calcium signaling in cerebellar purkinje cells alleviates pathological phenotype in spinocerebellar ataxia 2 mice. *J Neurosci* 2012;32(37):12786–12796.
25. Chen X, Tang TS, Tu H, et al. Deranged calcium signaling and neurodegeneration in spinocerebellar ataxia type 3. *J Neurosci* 2008;28(48):12713–12724.
26. Tang TS, Tu H, Chan EY, et al. Huntingtin and huntingtin-associated protein 1 influence neuronal calcium signaling mediated by inositol-(1,4,5) triphosphate receptor type 1. *Neuron* 2003;39(2):227–239.
27. Kaltenbach LS, Romero E, Becklin RR, et al. Huntingtin interacting proteins are genetic modifiers of neurodegeneration. *PLoS Genet* 2007;3(5):e82.
28. Shilling D, Müller M, Takano H, et al. Suppression of InsP3 receptor-mediated Ca²⁺ signaling alleviates mutant presenilin-linked familial Alzheimer's disease pathogenesis. *J Neurosci* 2014;34(20):6910–6923.
29. Cheung KH, Shineman D, Müller M, et al. Mechanism of Ca²⁺ disruption in Alzheimer's disease by presenilin regulation of InsP3 receptor channel gating. *Neuron* 2008;58(6):871–883.
30. Kwak YD, Shaw TI, Downing SM, et al. Chromatin architecture at susceptible gene loci in cerebellar Purkinje cells characterizes DNA damage-induced neurodegeneration. *Sci Adv* 2021;7(51):eabg6363.
31. Zamboni JL, Bellomo A, Ben-Pazi H, et al. Spinocerebellar ataxia type 29 due to mutations in ITPR1: a case series and review of this emerging congenital ataxia. *Orphanet J Rare Dis* 2017;12(1):121.
32. Romaniello R, Pasca L, Panzeri E, et al. Superior cerebellar atrophy: an imaging clue to diagnose ITPR1-related disorders. *Int J Mol Sci* 2022;23(12):6723.
33. Harrison SM, Biesecker LG, Rehm HL. Overview of specifications to the ACMG/AMP variant interpretation guidelines. *Curr Protoc Hum Genet* 2019;103(1):e93.
34. Brnich SE, Abou Tayoun AN, Couch FJ, et al. Recommendations for application of the functional evidence P3/S3 criterion using the ACMG/AMP sequence variant interpretation framework. *Genome Med* 2019;12(1):3.
35. Kelly MA, Caleshu C, Morales A, et al. Adaptation and validation of the ACMG/AMP variant classification framework for MYH7-associated inherited cardiomyopathies: recommendations by ClinGen's inherited cardiomyopathy expert panel. *Genet Med* 2018;20(3):351–359.
36. Hara K, Shiga A, Nozaki H, et al. Total deletion and a missense mutation of ITPR1 in Japanese SCA15 families. *Neurology* 2008;71(8):547–551.
37. Firth HV, Richards SM, Bevan AP, et al. DECIPHER: database of chromosomal imbalance and phenotype in humans using Ensembl resources. *Am J Hum Genet* 2009;84(4):524–533.
38. Deciphering Developmental Disorders Study. Large-scale discovery of novel genetic causes of developmental disorders. *Nature* 2015;519(7542):223–228.
39. Wright CF, Fitzgerald TW, Jones WD, et al. Genetic diagnosis of developmental disorders in the DDD study: a scalable analysis of genome-wide research data. *Lancet* 2015;385(9975):1305–1314.
40. Slevin H, Welsh SJ, Yu J, et al. De novo mutations in EBF3 cause a neurodevelopmental syndrome. *Am J Hum Genet* 2017;100(1):138–150.
41. Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense mutations using PolyPhen-2. *Curr Protoc Hum Genet* 2013;Chapter 7:Unit7.20.
42. Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* 2009;4(7):1073–1081.
43. Rentzsch P, Schubach M, Shendure J, Kircher M. CADD-splice-improving genome-wide variant effect prediction using deep learning-derived splice scores. *Genome Med* 2021;13(1):31.
44. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res* 2010;20(1):110–121.

45. Smedley D, Smith KR, Martin A, et al. 100,000 Genomes pilot on rare-disease diagnosis in health care—preliminary report. *N Engl J Med* 2021;385(20):1868–1880.
46. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data. *Cell* 2021;184(13):3573–3587.e3529.
47. Miller JA, Ding SL, Sunkin SM, et al. Transcriptional landscape of the prenatal human brain. *Nature* 2014;508(7495):199–206.
48. Li M, Santpere G, Imamura Kawasawa Y, et al. Integrative functional genomic analysis of human brain development and neuropsychiatric risks. *Science* 2018;362(6420):eaat7615.
49. Casey JP, Hirouchi T, Hisatsune C, et al. A novel gain-of-function mutation in the *ITPR1* suppressor domain causes spinocerebellar ataxia with altered Ca²⁺ signal patterns. *J Neurol* 2017;264(7):1444–1453.
50. Cunha P, Petit E, Coutelier M, et al. Extreme phenotypic heterogeneity in non-expansion spinocerebellar ataxias. *Am J Hum Genet* 2023;110(7):1098–1109.
51. Bosanac I, Yamazaki H, Matsu-Ura T, Michikawa T, Mikoshiba K, Ikura M. Crystal structure of the ligand binding suppressor domain of type 1 inositol 1,4,5-trisphosphate receptor. *Mol Cell* 2005;17(2):193–203.
52. Dudding TE, Friend K, Schofield PW, Lee S, Wilkinson IA, Richards RI. Autosomal dominant congenital non-progressive ataxia overlaps with the *SCA15* locus. *Neurology* 2004;63(12):2288–2292.
53. Danoff SK, Ferris CD, Donath C, et al. Inositol 1,4,5-trisphosphate receptors: distinct neuronal and nonneuronal forms derived by alternative splicing differ in phosphorylation. *Proc Natl Acad Sci U S A* 1991;88(7):2951–2955.
54. Nucifora FC Jr, Li SH, Danoff S, Ullrich A, Ross CA. Molecular cloning of a cDNA for the human inositol 1,4,5-trisphosphate receptor type 1, and the identification of a third alternatively spliced variant. *Brain Res Mol Brain Res* 1995;32(2):291–296.
55. Nakagawa T, Okano H, Furuichi T, Aruga J, Mikoshiba K. The subtypes of the mouse inositol 1,4,5-trisphosphate receptor are expressed in a tissue-specific and developmentally specific manner. *Proc Natl Acad Sci U S A* 1991;88(14):6244–6248.
56. Paganini L, Pesenti C, Milani D, et al. A novel splice site variant in *ITPR1* gene underlying recessive Gillespie syndrome. *Am J Med Genet A* 2018;176(6):1427–1431.
57. Carvalho DR, Medeiros JEG, Ribeiro DSM, Martins B, Sobreira NLM. Additional features of Gillespie syndrome in two Brazilian siblings with a novel *ITPR1* homozygous pathogenic variant. *Eur J Med Genet* 2018;61(3):134–138.
58. Raslan IR, França MC Jr, Oliveira JB, et al. Quadrupedal gait and cerebellar hypoplasia, the Uner tan syndrome, caused by *ITPR1* gene mutation. *Parkinsonism Relat Disord* 2021;92:33–35.
59. Samocha KE, Kosmicki JA, Karczewski KJ, et al. Regional missense constraint improves variant deleteriousness prediction. *bioRxiv* 2017;148353.
60. Parolin Schneckenberg R, Perkins EM, Miller JW, et al. De novo point mutations in patients diagnosed with ataxic cerebral palsy. *Brain* 2015;138(Pt 7):1817–1832.
61. Ida Y, Kidera A. The conserved Arg241-Glu439 salt bridge determines flexibility of the inositol 1,4,5-trisphosphate receptor binding core in the ligand-free state. *Proteins* 2013;81(10):1699–1708.
62. Genovese M, Guidone D, Buccirosi M, et al. Pharmacological potentiators of the calcium signaling cascade identified by high-throughput screening. *PNAS Nexus* 2023;2:pgac288.
63. Yao J, Ni M, Tian S, et al. A gain-of-function mutation in the gating domain of *ITPR1* impairs motor movement and increases thermal and mechanical sensitivity. *Neuroscience* 2023;522:11–22.
64. Kinoshita A, Ohyama K, Tanimura S, et al. *Itp1* regulates the formation of anterior eye segment tissues derived from neural crest cells. *Development* 2021;148(16):dev188755.
65. Sakamoto M, Iwama K, Sasaki M, et al. Genetic and clinical landscape of childhood cerebellar hypoplasia and atrophy. *Genet Med* 2022;24(12):2453–2463.
66. Liu Z, Sun H, Dai J, Xue X, Sun J, Wang X. *ITPR1* mutation contributes to hemifacial microsomia Spectrum. *Front Genet* 2021;12:616329.
67. Liu Y, Chen S, Zühlke L, et al. Global birth prevalence of congenital heart defects 1970–2017: updated systematic review and meta-analysis of 260 studies. *Int J Epidemiol* 2019;48(2):455–463.
68. Uchida K, Aramaki M, Nakazawa M, et al. Gene knock-outs of inositol 1,4,5-trisphosphate receptors types 1 and 2 result in perturbation of cardiogenesis. *PLoS One* 2010;5(9):e12500.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.