



HHS Public Access

Author manuscript

Eur Neuropsychopharmacol. Author manuscript; available in PMC 2024 March 28.

Published in final edited form as:

Eur Neuropsychopharmacol. 2020 January ; 30: 44–55. doi:10.1016/j.euroneuro.2017.11.012.

“RBF0X1, encoding a splicing regulator, is a candidate gene for aggressive behavior”

Noèlia Fernández-Castillo^{1,2,3,4}, Gabriela Gan⁵, Marjolein M.J. van Donkelaar⁶, Mariliis Vaht⁷, Heike Weber^{8,9}, Wolfgang Retz¹⁰, Andreas Meyer-Lindenberg⁵, Barbara Franke^{6,11}, Jaanus Harro⁷, Andreas Reif⁸, Stephen V. Faraone^{12,13}, Bru Cormand^{1,2,3,4}

¹Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona, Barcelona, Catalonia, Spain.

²Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Instituto de Salud Carlos III, Spain.

³Institut de Biomedicina de la Universitat de Barcelona (IBUB), Barcelona, Catalonia, Spain.

⁴Institut de Recerca Sant Joan de Déu (IR-SJD), Esplugues de Llobregat, Catalonia, Spain

⁵Systems Neuroscience in Psychiatry, Department of Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany.

⁶Radboud university medical center, Donders Institute for Brain, Cognition and Behaviour, Department of Human Genetics, Nijmegen, The Netherlands

⁷Division of Neuropsychopharmacology, Department of Psychology, University of Tartu, Tartu, Estonia.

⁸Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt - Goethe University, Frankfurt am Main, Germany.

⁹Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital of Würzburg, Würzburg, Germany.

¹⁰Department of Psychiatry and Psychotherapy, University Medical Center Mainz, Mainz, Germany.

¹¹Radboud university medical center, Donders Institute for Brain, Cognition and Behaviour, Department of Psychiatry, Nijmegen, The Netherlands

Corresponding authors: Noèlia Fernández Castillo. Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona, Catalonia, Spain, Avinguda Diagonal 643, edifici annex, planta 3, 08028, Barcelona, Catalonia, Spain. Tel.: (+34) 93 4037082. Fax: (+34) 93 4034420; noefernandez@ub.edu, Bru Cormand. Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona, Catalonia, Spain, Avinguda Diagonal 643, edifici annex, planta 3, 08028, Barcelona, Catalonia, Spain. Tel.: (+34) 93 4021013. Fax: (+34) 93 4034420; bcormand@ub.edu.

Contributors

N. Fernández-Castillo and B. Cormand identified the gene as a possible candidate and led the work and together with S.V. Faraone coordinated the study. N. Fernández-Castillo performed the literature review and wrote the first draft of the manuscript. B. Cormand and S.V. Faraone gathered new data. G. Gan, M.J. van Donkelaar, B. Franke and A. Meyer-Lindenberg performed neuroimaging genetic analyses. G. Gan contributed to the writing of the neuroimaging section. H. Weber, A. Reif and W. Retz performed the association study in the German sample of prisoners, and M. Vaht and J. Harro in the Estonian sample. All authors contributed to the final version of the manuscript.

Conflict of interest

None of the authors declare any conflict of interest.

¹²Departments of Psychiatry and of Neuroscience and Physiology, SUNY Upstate Medical University, Syracuse NY, USA.

¹³K.G. Jepsen Centre for Research on Neuropsychiatric Disorders, University of Bergen, Bergen, Norway.

Abstract

The *RBFOX1* gene (or *A2BPI*) encodes a splicing factor important for neuronal development that has been related to autism spectrum disorder and other neurodevelopmental phenotypes. Evidence from complementary sources suggests that this gene contributes to aggressive behavior. Suggestive associations with *RBFOX1* have been identified in genome-wide association studies (GWAS) of anger, conduct disorder, and aggressive behavior. Nominal association signals in *RBFOX1* were also found in an epigenome-wide association study (EWAS) of aggressive behavior. Also, variants in this gene affect temporal lobe volume, a brain area that is altered in several aggression-related phenotypes. In animals, this gene has been shown to modulate aggressive behavior in *Drosophila*. *RBFOX1* has also been associated with canine aggression and is upregulated in mice that show increased aggression after frustration of an expected reward. Associated common genetic variants as well as rare duplications and deletions affecting *RBFOX1* have been identified in several psychiatric and neurodevelopmental disorders that often are comorbid with aggressive behaviors. In this paper, we comprehensively review the cumulative evidence linking *RBFOX1* to aggression behavior and provide new results implicating *RBFOX1* in this phenotype. Most of these studies (genetic and epigenetic analyses in humans, neuroimaging genetics, gene expression and animal models) are hypothesis-free, which strengthens the validity of the findings, although all the evidence is nominal and should therefore be taken with caution. Further studies are required to clarify in detail the role of this gene in this complex phenotype.

Keywords

RBFOX1 ; *A2BPI* ; aggression; genetics

Introduction

Aggressive behavior and violence are major causes of mortality and morbidity in humans. These traits are observed in several psychiatric and neurodevelopmental disorders. Aggressive behavior is an evolutionary conserved trait of high importance for species survival. For this reason, it has been subject to selection throughout evolution and has a substantial genetic underpinning, while staying responsive to environmental cues. Accordingly, the heritability of aggressive behaviors has been estimated to be around 50% (reviewed by Veroude et al., 2016). Several genes and pathways contributing to aggression have been identified, such as those involved in the serotonergic and dopaminergic neurotransmission and in hormone regulation, although most of the genetic contribution to aggression is still unexplained (Fernández-Castillo and Cormand, 2016).

In a recent review on the genetics of human aggressive behavior, we searched for genes and pathways involved in this phenotype using data from genome-wide association studies

(GWAS), which so far have not produced genome-wide significant genes/loci (Fernández-Castillo and Cormand, 2016). Among others, we highlighted the *RBFOX1* gene (also known as *A2BPI*), which showed suggestive associations in three different GWAS (Anney et al., 2008; Mick et al., 2014; Sonuga-Barke et al., 2008). Interestingly, other complementary sources of evidence provide additional support for the contribution of this gene to the susceptibility to aggressive behavior and to several psychiatric disorders as well.

RBFOX1 encodes the RNA Binding Protein, Fox-1 Homolog 1, also known as Ataxin-2-binding protein. It is expressed mainly in the nervous system, heart, and muscle (Jin et al., 2003; Underwood et al., 2005). The gene encodes a splicing factor that plays an important role in the regulation of the alternative splicing of large neuronal gene networks important for brain development (Bill et al., 2013; Conboy, 2017; Li et al., 2015). Cytoplasmic and nuclear *RBFOX1* isoforms seem to play different roles, with the first one contributing to mRNA stability and promoting translation and the second one acting as a splicing regulator (Hamada et al., 2016; Lee et al., 2016). The nuclear isoform is involved in neuron migration and synapse network formation during corticogenesis (Hamada et al., 2016) and is important for the control of neuronal excitation in the mammalian brain (Gehman et al., 2011). Alterations in the *RBFOX1* gene have been associated with several neurodevelopmental pathologies, especially autism spectrum disorder (reviewed by Bill et al., 2013).

Here, we reviewed the cumulative evidence supporting a contribution of *RBFOX1* to aggressive behaviors and to other psychiatric and neurodevelopmental disorders that often display aggressive behavior. We also present hitherto unpublished data derived from genetic association and neuroimaging genetics studies supporting our hypothesis.

***RBFOX1* and aggressive behavior in humans**

Association studies—In a previous review of the genetic basis of aggressive behavior in humans (Fernández-Castillo and Cormand, 2016), *RBFOX1* showed suggestive associations ($p < 5e-05$, Table 1) with aggressive traits or diagnostic categories in three GWAS (Anney et al., 2008; Merjonen et al., 2011; Sonuga-Barke et al., 2008). A common variant located within the first intron of the *RBFOX1* gene, rs6500744, was identified as one of the top association signals in a GWAS assessing gene by environment interactions (GxE) (Sonuga-Barke et al., 2008). The C allele of this single nucleotide polymorphism (SNP) was associated with conduct disorder (CD) symptoms in interaction with mothers' warmth (Table 1, Figure 1). Another SNP, also located in intron 1 of *RBFOX1*, rs8062784, was associated with anger in a GWAS assessing hostility in adolescents and adults (Merjonen et al., 2011), and two variants located in intron 3 of the gene, rs10153149 and rs12921846, were associated with CD in a sample of ADHD trios (Anney et al., 2008) (see Table 1 and Figure 1).

A meta-analysis of nine population-based GWASs including around 19,000 children provided more evidence for the contribution of *RBFOX1* to aggressive behavior (Pappa et al., 2016). Four SNPs in this gene (rs809682, rs12922093, rs12373031, and rs10521042, all located in intron 2) showed suggestive associations with children's aggressive behavior in a meta-analysis ($p < 5e-04$, Table 1 and Figure 1, data kindly provided by the authors).

Overall, eight *RBFOX1* variants from four GWASs showed a nominal association with aggression. The variants associated in each GWAS were not in linkage disequilibrium (LD) with those from the other studies. The LD between the four variants found associated in the meta-analysis (Pappa et al., 2016) was moderate to high. In an attempt to investigate the association of *RBFOX1* with aggressive behavior further, we genotyped four SNPs (the most significant one from each GWAS) in a sample of German male prisoners, not included in any of the GWASs (n = 188, Supplementary Material). A significant association was found for rs809682, when comparing aggressive prisoners to controls (p = 0.03, additive model) as well as aggressive prisoners to non-aggressive prisoners (p = 0.02), but not when comparing non-aggressive prisoners and controls (p = 0.76) (Supplementary Table 1). However, the direction of the effect was opposite as that of the original study, with the A allele linked to aggression (Supplementary Table 1 and Table 1). The same variant was also genotyped in a population-representative sample of 1176 individuals from the Estonian Children Personality Behaviour and Health Study, ECPBHS (Harro et al., 2001; Vaht et al., 2016) (see Supplementary Material for details). Several traits related to aggressiveness were assessed in this population-based cohort, in particular those related to personality and anxiety. In the ECPBHS, five-factor personality data at young adulthood were available from the Estonian version of the NEO-PI-R questionnaire (Kallasmaa et al., 2000) and anxiety information from Spielberger's Trait Anxiety questionnaire (Spielberger, 1983). Aggressiveness is associated with certain facets in basic personality traits, in particular with low agreeableness and neuroticism (Caprara et al., 1996; Sharpe and Desai, 2001), and is also associated with anxiety (Clement and Chapouthier, 1998; Siddaway et al., 2017; Van Praag, 1998), higher negative emotionality (Caprara et al., 1996; Kodžopelji et al., 2014; Tremblay and Ewart, 2005), and lower extraversion levels (Sharpe and Desai, 2001; Tremblay and Ewart, 2005). In the Estonian sample, rs809682 was nominally associated with extraversion (p = 0.024), the T/T homozygotes showing the lowest levels of extraversion (Supplementary Table 2). We recently showed that sex can influence the contribution of genes to aggression (van Donkelaar et al., under review). We therefore also investigated genotype by sex interactions. *RBFOX1* genotype and sex indeed had a significant interaction effect on agreeableness (p = 6e-03); male T/T homozygotes had the lowest scores of agreeableness. While NEO-PI neuroticism was not significantly associated with the genotype, data on Spielberger's Trait Anxiety showed a nominal association (p = 0.043), with the T-allele carriers having higher levels of anxiety. Findings in the Estonian sample are in the direction expected for a gene associated with aggression, and suggest that higher expression of aggressive behavior in the T-allele carriers of the *RBFOX1* rs809682 polymorphism may be related to basic personality traits and anxiety.

As indicated above, aggressive behavior is influenced by both genes and environment. For this reason, looking at epigenetic profiles linked to aggression also may provide insight into *RBFOX1*'s role. A recent epigenome-wide association study (EWAS) of aggressive behavior investigated DNA methylation levels associated with this phenotype in a sample of 2,029 individuals (van Dongen et al., 2015). This study did not produce any epigenome-wide significant findings, but two nominal associations between aggressive behavior and the methylation levels of sites located in *RBFOX1* were identified at cg12310850 (GRCh37/hg19 position; chr16: 6,533,700, p = 8.7e-03) and cg00499781 (chr16: 7,568,364, p = 0.040)

(data kindly provided by the authors). The authors also investigated the methylation levels of monozygotic twins highly discordant for aggression, identifying five nominal associations in sites located in *RBFOX1* at cg03934713 (chr16: 6,069,198, $p = 0.047$), cg16396980 (chr16: 6,633,344, $p = 0.041$), cg06705265 (chr16: 6,696,222, $p = 0.024$), cg00514665 (chr16: 7,703,812, $p = 0.049$), and cg03986562 (chr16: 7,703,893, $p = 0.037$).

Neuroimaging genetics studies and gene expression—Human neuroimaging genetics studies suggest that the *RBFOX1* gene contributes to brain function and structure. Thus, *RBFOX1* has been shown to influence temporal lobe volume with genome-wide significance in the Alzheimer’s Disease Neuroimaging Initiative (ADNI) cohort including patients with Alzheimer’s disease, patients with mild cognitive impairment, and cognitively healthy elderly controls (Kohannim et al., 2012b). In a related genome-wide study on the ADNI cohort, Kohannim et al. (Kohannim et al., 2012a) observed that a variant in the *RBFOX1* gene was highly predictive of temporal brain volume. The association between a specific variant in *RBFOX1* and temporal lobe structure has been confirmed in voxel-wise whole-brain analyses in the ADNI cohort (Kohannim et al., 2012a, 2012b). In line with these findings, the *RBFOX1* gene has been identified as specifically contributing to grey matter loss in the temporal lobe in patients with mild cognitive impairment, in a sample of Alzheimer’s Disease (Vounou et al., 2012). While the aforementioned studies consistently show a link between *RBFOX1* and temporal lobe volume at genome-wide significance, caution in drawing conclusions for healthy younger adults is required as all findings are based on the ADNI cohort studying associations between temporal lobe volume in older adults with and without cognitive impairment and manifest Alzheimer’s disease. Thus, associations between *RBFOX1* and temporal lobe volume might be explained by the decline in temporal lobe volume observed in Alzheimer’s disease and mild cognitive impairment (e.g., Vounou et al., 2012). Still, the link between *RBFOX1* and temporal lobe volume is intriguing because Alzheimer’s disease often presents with aggressive behavior (Zhao et al., 2016). Moreover, animal and human studies have underlined the role of the temporal lobe for aggression, especially with the amygdala and hippocampus being located in its medial part (Gregg and Siegel, 2001; Potegal, 2012; Siever, 2008). Abnormalities in temporal lobe function and structure have been observed in aggressive populations including individuals with conduct disorder (CD) (Cappadocia et al., 2009; Kruesi et al., 2004), psychopathy (Raine et al., 2004), and aggressive schizophrenia (Hoptman et al., 2011; Soyka, 2011). Structural and functional changes in the temporal lobe in aggressive populations may be related to impaired emotion regulation (Bufkin, 2005) and/or to the lack of empathy in CD and antisocial behavior (Cappadocia et al., 2009). Still, although *RBFOX1* seems to contribute to temporal lobe functioning, that does not directly support its contribution to aggressive behavior and further studies should investigate the effect of these variants.

In a recent study, we found the first evidence that the risk variant rs6500744 in the *RBFOX1* gene, which had been associated with aggression in the GWAS of Sonuga-Barke et al. (2008), might be relevant to brain activity during neurocognitive processes such as inhibitory control and emotional reactivity (Gan et al., in preparation). In a sample of 331 healthy human participants, carriers of the risk allele C (C/C and C/T) showed an increased brain response in the dorsal anterior cingulate cortex (ACC) during emotion processing, and

reduced brain responses in the left inferior/middle frontal gyrus during inhibitory control measured with a combined Flanker/Go-Nogo task. Moreover, we observed a sex by SNP interaction, in which female C/C-carriers showed increased and male C/C-carriers showed decreased responding in the fusiform face area and the hypothalamus/ventral striatum compared to T-allele carriers during a Hariri emotional face recognition task (for fMRI tasks, compare (Meyer-Lindenberg et al., 2006)). Importantly, these findings converge with neuroimaging phenotypes including impaired brain functioning in prefrontal-limbic networks during inhibitory control and emotional reactivity in carriers of the MAOA-L genotype, the most widely studied risk genotype for aggression (Alia-Klein et al., 2011; Buckholtz and Meyer-Lindenberg, 2008; Fan et al., 2003; Meyer-Lindenberg et al., 2006; Passamonti et al., 2006).

As expected, given its role in neurodevelopment, the expression of *RBFOX1* in humans is mainly restricted to the brain, although it is also expressed in skeletal muscle and heart (Figure 2). Interestingly, the prefrontal cortex (PFC) including the anterior cingulate cortex (ACC) shows the highest *RBFOX1* expression levels in humans (Figure 2). The PFC has been shown to play a role in impulsive aggression (Blair, 2016; Fan et al., 2003; Passamonti et al., 2006), most likely through its involvement in inhibitory control and behavioral self-regulation (Buckholtz and Meyer-Lindenberg, 2008; Davidson et al., 2000; Heatherton and Wagner, 2011). Also, ACC impaired functioning and structural abnormalities of this area has been repeatedly linked to a propensity for impulsive/reactive aggression (Buckholtz and Meyer-Lindenberg, 2008; Meyer-Lindenberg et al., 2006; Sterzer et al., 2005). Moreover, high expression levels of *RBFOX1* are also found in the basal ganglia including the nucleus accumbens, putamen and caudate, brain areas previously linked to impulsive aggression across species (Buckholtz et al., 2010; Couppis and Kennedy, 2008; Ferrari et al., 2003; Gan et al., 2015; Krämer et al., 2007). A similar but less pronounced *RBFOX1* expression level emerges for the amygdala, a crucial brain area for impulsive reactive aggression due to its involvement in emotional reactivity to threatening cues (Buckholtz et al., 2008; Coccaro et al., 2007; Meyer-Lindenberg et al., 2006). To conclude, although the observation of high *RBFOX1* expression in brain areas relevant to aggression is interesting, this fact does not necessarily link this gene with aggression, as many other genes may show similar expression patterns and also many brain regions are involved in aggressive behavior. Future imaging genetic studies may shed light on the impact of *RBFOX1* genotypes and pathways on brain function and structure within these areas.

***RBFOX1* and aggressive behavior in animals**

RBFOX1, also known as *A2BPI*, has been related to aggressive behavior in animal models of aggression. In *Drosophila*, *A2bp1* was one of the genes identified through a genome-wide analysis aimed at mapping loci involved in aggressive behavior (Shorter et al., 2015). The study used the sequenced inbred lines of the *Drosophila melanogaster* Genetic Reference Panel (DGRP) to investigate natural genetic variation related to aggression. Male-to-male aggression was quantified in 200 DGRP lines to identify the underlying genetic variation. Extreme quantitative trait locus (QTL) genome-wide association analysis was also performed in a population derived from DGRP lines with extremely high and extremely low aggression scores. From both analyses, the authors obtained a genetic interaction network

with 741 genes, in which *A2bp1* was present as an important node. The functional validation of the finding for the *A2bp1* gene, using a mutant line with a *Mi{ET1}* element insertional mutation, yielded significant evidence that homozygous mutant flies for this gene show decreased aggression ($p < 0.001$).

In a genome-wide mapping study of aggression in dogs, several nominal associations with *A2bp1* were identified in two different cohorts (Zapata et al., 2016). Some of these signals in *A2bp1* were associated with the four measures of aggression that were assessed: stranger-directed aggression (towards unfamiliar humans), dog-directed aggression (towards unfamiliar dogs), owner-directed aggression (towards familiar humans) and dog rivalry (towards familiar dogs). Four *loci* in *A2bp1* were nominally associated with all four measures in one cohort (chr6: 35,449,934, chr6: 35,459,495, chr6: 35,557,330 and chr6: 35,557,812) and two in another cohort (chr6: 35,641,555 and chr6: 35,672,733).

Finally, in a murine frustration model that showed increased aggressive behavior when the access to an expected reward was denied (Burokas et al., 2012), *A2bp1* expression was upregulated (fold change = 1.32, $p = 1.29e-03$, FDR < 5%) in ventral striatum of frustrated mice (Martín-García et al., 2015).

Mutant models of *Rbfox1* exist, but they have not been evaluated for aggressive behavior: a knockdown of *rbfox1* in zebrafish was investigated for heart phenotypes and was shown to produce cardiac dysfunction and heart failure (Frese et al., 2015); knockout mice for *Rbfox1* ($-/-$), central nervous system-specific, have been reported to present seizures and increased neuronal excitability (Gehman et al., 2011). Electrophysiological recordings of rat cortical neurons with increased expression of this gene also showed increased neuronal activity (Wen et al., 2015). A very recent study shows that RBFOX1 plays an important role in coordinating the synaptic downscaling of excitatory synapses (Rajman et al., 2017). Previous studies have reported that both excitatory and inhibitory neurotransmission are altered in aggression and that changes in neuronal excitation or inhibition modify aggressive behavior (Ende et al., 2015; Lin et al., 2011; Luque et al., 2009; Takahashi et al., 2015; Vekovischeva et al., 2004). Also, RBFOX1 is highly expressed in GABAergic neurons of the developing forebrain in mice (Hammock and Levitt, 2011). Thus, since changes in RBFOX1 seem to affect neuronal excitability, we could hypothesize that it may contribute to aggressive behavior by triggering alterations in inhibition/excitability balance, although no direct evidence of this has been observed so far. Further studies assessing aggressive behavior in knockout and knockin animals for *Rbfox1* are required to confirm this hypothesis and the role of this gene in this complex phenotype.

RBFOX1 and psychiatric neurodevelopmental and neurodegenerative disorders

Genetic variation in *RBFOX1* – both common and rare - has been associated with anxiety disorder, substance use disorders, schizophrenia, bipolar disorder, attention-deficit/hyperactivity disorder (ADHD), autism spectrum disorder (ASD) as well as Alzheimer disease. All these disorders present with aggressive behaviors (Brady et al., 1998; Bubier and Drabick, 2009; Fitzpatrick et al., 2016; Granic, 2014; Hoaken and Stewart, 2003; King and Waschbusch, 2010; Látalová, 2009; Volavka, 2013; Zhao et al., 2016).

Findings for associations with common variants in the gene originate mainly from GWAS. In anxiety disorder, a variant in intron 1 of the *RBFOX1* gene (rs13334105) showed a genome-wide significant association with sensitivity to anxiety. Several other SNPs within the same intron also showed suggestive associations ($p < 1e-07$) (Davies et al., 2015). The authors also inspected this region in a GWAS meta-analysis of anxiety disorder and observed several nominal associations. *RBFOX1* variants have been found associated with substance dependence and the ability to quit smoking in 13 different datasets (reviewed by Zhong et al., 2015). Also, several genetic markers in a genomic region containing *RBFOX1* displayed suggestive linkage to substance dependence-related phenotypes (Zhong et al., 2015). For schizophrenia, suggestive associations were found with *RBFOX1* in Ashkenazi Jews (Goes et al., 2015). Also, suggestive signals in this gene were found associated with both Schizophrenia and bipolar disorder (Wang et al., 2010). In Alzheimer disease, genome-wide significant signals have been identified in the gene (Herold et al., 2016). Alzheimer Disease presents a high occurrence of aggression, estimated to occur in around 40% of individuals with this disorder (Zhao et al., 2016).

Using the Ricopili web tool (<https://data.broadinstitute.org/mpg/ricopili/>) we identified additional common genetic variants in *RBFOX1* that show suggestive association with several psychiatric phenotypes: bipolar disorder and schizophrenia (cross-disorder meta-analysis; rs12444931, $p = 5e-06$), schizophrenia (rs12447542, $p = 1.1e-06$), smoking behavior (rs3112740, $p = 6e-06$) and number of cigarettes per day (rs8055842, $p = 3.4e-05$).

Evidence for rare *RBFOX1* variants being linked to brain disease come from several study designs. Copy number variants (CNVs) spanning the *RBFOX1* gene have been reported in individuals with several psychiatric disorders. A partial duplication of *RBFOX1* was associated with risk for schizophrenia, with an estimated eight-fold increased risk for males, but not for females (Melhem et al., 2011). Another partial duplication was identified in another patient with schizophrenia (Xu et al., 2008). A gain CNV spanning the first two exons of *RBFOX1* and a loss in an intronic region have been reported for bipolar disorder (Noor et al., 2014). Also, hemizygous intronic *RBFOX1* deletions and a duplication were identified in ADHD patients (Elia et al., 2010).

Another study identified CNVs in *RBFOX1* in patients with neuropsychiatric and neurodevelopmental disorders such as ASD and global developmental delay, many of them presenting also with epilepsy (Zhao, 2013). Interestingly, in this study, one patient with intellectual disability, marked aggressive behavior, and epilepsy was found to bear a deletion in the *RBFOX1* gene. Other studies identified deletions in *RBFOX1* in autistic patients (Davis et al., 2012; Griswold et al., 2012; Martin et al., 2007; Sebat et al., 2007). One of these studies characterized a deletion in a proband with autism, global developmental delay, and epilepsy. The deletion was located at the boundary between the first exon and intron and reduced *RBFOX1* mRNA expression in lymphocytes from the subject (Martin et al., 2007). More evidence connects *RBFOX1* with ASD. A *de novo* truncating mutation in this gene and a duplication were identified in autistic patients (Griswold et al., 2015; Kanduri et al., 2016). A transcriptomic analysis of post-mortem autistic brains identified a module of co-expressed genes in which *RBFOX1* was an important node (Voineagu et al., 2011). This gene module was enriched for associated genetic variants in an autism GWAS dataset.

Furthermore, the authors performed RNAseq to compare brain samples of autistic patients with a decreased expression of *RBFOX1* (FC = -5.9) and controls with average expression of this gene, and observed a broad dysregulation of alternative splicing in the brain of autistic patients that is dependent on *RBFOX1*.

Several genetic alterations in *RBFOX1* have been identified in individuals with epilepsy, a neurological disorder that has also been related to aggressive behavior (Brodie et al., 2016). In line with this, as mentioned above, the knockout mice for *Rbfox1* show susceptibility to seizures and increased neuronal excitability (Gehman et al., 2011). Another study observed upregulation of *RBFOX1* in patients with malformed cortex and epilepsy, and observed that this upregulation also showed an increased neuronal activity of rat cortical neurons (Wen et al., 2015).

The fact that *RBFOX1* has been related to many psychiatric disorders make it an appealing candidate gene for aggression, although the increased incidence of aggression in many of these disorders does not connect *RBFOX1* with aggressive behavior. In this regard, exploration of large cohorts of aggression is needed to investigate the effect of genetic variants in *RBFOX1* that have been reported as risk factors for other psychiatric disorders. Also, we need to get more insight into the contribution to aggression of CNVs spanning *RBFOX1*.

Discussion and future perspectives

In this review, we bring together different lines of evidence that implicate *RBFOX1* in the etiology of aggressive behavior, including genetic association and mutation studies, neuroimaging genetics data, transcriptomic analyses, and animal models. Although the convergence of evidence is impressive, most of this supportive data is of nominal significance in the individual studies and should therefore be interpreted with caution.

Four independent GWASs (one of them including nine datasets) associated *RBFOX1* with aggression. However, it should be noted that this gene spans a wide region in the genome (1.7 Mbp, more than 30 times the average length of a human gene) and carries many common variants, increasing the probability for identifying spurious associations. Speaking against such false positive effects, according to the GWAS catalog (<https://www.ebi.ac.uk/gwas/>), so far only very few disease or trait associations have been identified for *RBFOX1* with p-values < 1e-05, (including obesity and body mass index, heart rate, periodontitis, and eye related diseases). In any case, we cannot discard that gene length may influence the appearance of false positive associations. Finally, given the high degree of phenotypic heterogeneity of aggressive behaviors, association studies in larger and more homogeneous samples are required to confirm the involvement of *RBFOX1*.

All the *RBFOX1* variants identified by the different GWASs of aggression lie in the first three introns of the gene. This localization also applies to the SNPs and CNVs associated with other psychiatric disorders. This clustering of risk variants may be due to the presence of regulatory elements of *RBFOX1* in this region, as suggested by others (Martin et al., 2007). To our knowledge, CNVs in *RBFOX1* have not been investigated in relation

to aggressive behavior directly. Such studies may be informative as they have been for schizophrenia, bipolar disorder, ADHD or autism.

The neuroimaging genetics studies reported here highlight structural and functional brain alterations correlating with *RBFOX1* variation, both in population-based cohorts and in clinical samples. These results point to the temporal lobe and to alterations in neurocognitive performance. However, to our knowledge, no investigations have been performed in clinical samples with pathological levels of aggression (e.g., conduct disorder, antisocial personality disorder, intermittent explosive disorder). Such studies may shed light on the functional significance of *RBFOX1* with regard to brain regions previously linked to aggressive behavior, such as the temporal and frontal lobes, the basal ganglia and the amygdala, where the gene is highly expressed.

Animal models can provide more insights into the role of this gene in aggression. The data from *Drosophila* reviewed above are convincing. In this regard (and given the *Drosophila* phenotype as well as the findings of duplications in the gene in human brain disorders), it might be interesting to assess aggressive behavior not only in knockout animals for *RBFOX1*, but also in knockin animals, where the gene is upregulated.

Taken together, evidence from complementary study designs point to *RBFOX1* as a strong candidate for susceptibility to aggressive behavior and to several psychiatric disorders. Still, further association and neuroimaging genetics studies in larger samples, as well as studies in transgenic animals for *RBFOX1*, are needed to confirm the contribution of this gene to aggressive behavior. If confirmation is obtained, *RBFOX1* could be a promising pharmacological target for the treatment of aggression, given its broad role in the development and functioning of brain processes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

We would like to thank the EAGLE consortium, Dorret Boomsma and Jenny van Dongen for kindly providing us with the results from their human GWAS and EWAS studies, and to Carlos Álvarez and Isain Zapata for the results of their study on canine aggression. We would like to thank Jeffrey Glennon, Heike Tost and Anais Harneit for providing useful comments and literature.

Role of the funding source

The research leading to these results has received funding from the European Union Seventh Framework Programme [FP7/2007-2013] under grant agreements n° 602805, 602450 and 278948 and from the European Union H2020 Programme [H2020/2014-2020] under grant agreements n° 643051 and 667302. Additionally, N. Fernández-Castillo is supported by a contract of the 'Centro de Investigación Biomédica en Red de Enfermedades Raras' (CIBERER) and B. Cormand by the Spanish 'Ministerio de Economía y Competitividad' (SAF2015-68341-R) and AGAUR, 'Generalitat de Catalunya' (2014SGR932). Further funding was received from the European College for Neuropsychopharmacology (ECNP) by a grant for the ECNP Network 'ADHD across the lifespan'. G. Gan was supported by the German academic exchange service (DAAD), and the Olympia-Morata Program of the University of Heidelberg, Germany. J. Harro received funding from EMES Grant IUT-20-40. Prof. Faraone is supported by the K.G. Jebsen Centre for Research on Neuropsychiatric Disorders, University of Bergen, Norway, and NIMH grants 5R01MH101519 and U01 MH109536-01. Prof. B. Franke and M. van Donkelaar are supported by funding from a personal Vici grant of the Netherlands Organisation for Scientific Research (NWO); grant

016-130-669, to BF). None of the funding institutions had any role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Bibliography

- Alia-Klein N, Parvaz MA, Woicik PA, Konova AB, Maloney T, Shumay E, Wang R, Telang F, Biegon A, Wang G-J, Fowler JS, Tomasi D, Volkow ND, Goldstein RZ, 2011. Gene x disease interaction on orbitofrontal gray matter in cocaine addiction. *Arch. Gen. Psychiatry* 68, 283–294. [PubMed: 21383264]
- Anney RJL, Lasky-Su J, Ó'Dúshláine C, Kenny E, Neale BM, Mulligan A, Franke B, Zhou K, Chen W, Christiansen H, Arias-Vásquez A, Banaschewski T, Buitelaar J, Ebstein R, Miranda A, Mulas F, Oades RD, Roeyers H, Rothenberger A, Sergeant J, Sonuga-Barke E, Steinhausen H, Asherson P, Faraone SV, Gil M, 2008. Conduct disorder and ADHD: Evaluation of conduct problems as a categorical and quantitative trait in the international multicentre ADHD genetics study. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 147, 1369–1378.
- Bill BR, Lowe JK, DyBuncio CT, Fogel BL, 2013. Orchestration of neurodevelopmental programs by RBFOX1: Implications for autism spectrum disorder. *Int. Rev. Neurobiol.* 113, 251–267. [PubMed: 24290388]
- Blair RJR, 2016. The Neurobiology of Impulsive Aggression. *J. Child Adolesc. Psychopharmacol.* 26, 4–9. [PubMed: 26465707]
- Brady KT, Myrick H, McElroy SL, 1998. The relationship between substance use disorders, impulse control disorders, and pathological aggression. *Am. J. Addict.* 7, 221–230. [PubMed: 9702290]
- Brodie MJ, Besag F, Ettinger AB, Mula M, Gobbi G, Comai S, Aldenkamp AP, Steinhoff BJ, 2016. Epilepsy, Antiepileptic Drugs, and Aggression: An Evidence-Based Review. *Pharmacol. Rev.* 68, 563–602. [PubMed: 27255267]
- Bubier JL, Drabick DAG, 2009. Co-occurring anxiety and disruptive behavior disorders: The roles of anxious symptoms, reactive aggression, and shared risk processes. *Clin. Psychol. Rev.* 29, 658–669. [PubMed: 19729235]
- Buckholtz JW, Callicott JH, Kolachana B, Hariri AR, Goldberg TE, Genderson M, Egan MF, Mattay VS, Weinberger DR, Meyer-Lindenberg A, 2008. Genetic variation in MAOA modulates ventromedial prefrontal circuitry mediating individual differences in human personality. *Mol. Psychiatry* 13, 313–324. [PubMed: 17519928]
- Buckholtz JW, Meyer-Lindenberg A, 2008. MAOA and the neurogenetic architecture of human aggression. *Trends Neurosci.* 31, 120–129. [PubMed: 18258310]
- Buckholtz JW, Treadway MT, Cowan RL, Woodward ND, Benning SD, Li R, Ansari MS, Baldwin RM, Schwartzman AN, Shelby ES, Smith CE, Cole D, Kessler RM, Zald DH, 2010. Mesolimbic dopamine reward system hypersensitivity in individuals with psychopathic traits. *Nat. Neurosci.* 13, 419–421. [PubMed: 20228805]
- Bufkin JL, 2005. Neuroimaging Studies of Aggressive and Violent Behavior: Current Findings and Implications for Criminology and Criminal Justice. *Trauma, Violence, Abus.* 6, 176–191.
- Burokas A, Gutiérrez-Cuesta J, Martín-García E, Maldonado R, 2012. Operant model of frustrated expected reward in mice. *Addict. Biol.* 17, 770–782. [PubMed: 22264360]
- Cappadocia MC, Desrocher M, Pepler D, Schroeder JH, 2009. Contextualizing the neurobiology of conduct disorder in an emotion dysregulation framework. *Clin. Psychol. Rev.* 29, 506–518. [PubMed: 19573964]
- Caprara GV, Barbaranelli C, Zimbardo PG, 1996. Understanding the Complexity of Human Aggression: Affective, Cognitive, and Social Dimensions of Individual Differences in Propensity Toward Aggression. *Eur. J. Pers.* 10, 133–155.
- Clement Y, Chapouthier G, 1998. Biological bases of anxiety. *Neurosci. Biobehav. Rev.* 22, 623–633. [PubMed: 9662724]
- Coccaro EF, McCloskey MS, Fitzgerald DA, Phan KL, 2007. Amygdala and Orbitofrontal Reactivity to Social Threat in Individuals with Impulsive Aggression. *Biol. Psychiatry* 62, 168–178. [PubMed: 17210136]

- Conboy JG, 2017. Developmental regulation of RNA processing by Rbfox proteins. *Wiley Interdiscip. Rev. RNA* 8, e1398.
- Couppis MH, Kennedy CH, 2008. The rewarding effect of aggression is reduced by nucleus accumbens dopamine receptor antagonism in mice. *Psychopharmacology (Berl)*. 197, 449–456. [PubMed: 18193405]
- Davidson RJ, Putnam KM, Larson CL, 2000. Dysfunction in the Neural Circuitry of Emotion Regulation--A Possible Prelude to Violence. *Science (80-.)*. 289, 591–594.
- Davies MN, Verdi S, Burri A, Trzaskowski M, Lee M, Hettema JM, Jansen R, Boomsma DI, Spector TD, 2015. Generalised Anxiety Disorder - A twin study of genetic architecture, genome-wide association and differential gene expression. *PLoS One* 10, e0134865.
- Davis LK, Maltman N, Mosconi MW, Macmillan C, Schmitt L, Moore K, Francis SM, Jacob S, Sweeney JA, Cook EH, 2012. Rare inherited A2BP1 deletion in a proband with autism and developmental hemiparesis. *Am. J. Med. Genet. Part A* 158 A, 1654–1661. [PubMed: 22678932]
- Elia J, Gai X, Xie HM, Perin JC, Geiger E, Glessner JT, D'arcy M, DeBerardinis R, Frackelton E, Kim C, Lantieri F, Muganga BM, Wang L, Takeda T, Rappaport EF, Grant SFA, Berrettini W, Devoto M, Shaikh TH, Hakonarson H, White PS, 2010. Rare structural variants found in attention-deficit hyperactivity disorder are preferentially associated with neurodevelopmental genes. *Mol. Psychiatry* 15, 637–46. [PubMed: 19546859]
- Ende G, Cackowski S, VanEijk J, Sack M, Demirakca T, Kleindienst N, Bohus M, Sobanski E, Krause-Utz A, Schmahl C, 2015. Impulsivity and aggression in female BPD and ADHD patients: association with ACC Glutamate and GABA concentrations. *Neuropsychopharmacology* 41, 410–418. [PubMed: 26040503]
- Fan J, Fossella J, Sommer T, Wu Y, Posner MI, 2003. Mapping the genetic variation of executive attention onto brain activity. *Proc. Natl. Acad. Sci. U. S. A.* 100, 7406–7411. [PubMed: 12773616]
- Fernández-Castillo N, Cormand B, 2016. Aggressive behavior in humans: Genes and pathways identified through association studies. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 171, 676–696.
- Ferrari PF, Van Erp AMM, Tornatzky W, Miczek KA, 2003. Accumbal dopamine and serotonin in anticipation of the next aggressive episode in rats. *Eur. J. Neurosci.* 17, 371–378. [PubMed: 12542674]
- Fitzpatrick SE, Srivorakiat L, Wink LK, Pedapati EV, Erickson CA, 2016. Aggression in autism spectrum disorder: Presentation and treatment options. *Neuropsychiatr. Dis. Treat.* 12, 1525–1538. [PubMed: 27382295]
- Frese KS, Meder B, Keller A, Just S, Haas J, Vogel B, Fischer S, Backes C, Matzas M, Köhler D, Benes V, Katus HA, Rottbauer W, 2015. RNA splicing regulated by RBFOX1 is essential for cardiac function in zebrafish. *J. Cell Sci.* 128, 3030–3040. [PubMed: 26116573]
- Gan G, Sterzer P, Marxen M, Zimmermann US, Smolka MN, 2015. Neural and Behavioral Correlates of Alcohol-Induced Aggression Under Provocation. *Neuropsychopharmacology* 40, 2886–2896. [PubMed: 25971590]
- Gehman LT, Stoilov P, Maguire J, Damianov A, Lin C-H, Shiue L, Ares M, Mody I, Black DL, 2011. The splicing regulator Rbfox1 (A2BP1) controls neuronal excitation in the mammalian brain. *Nat. Genet.* 43, 706–711. [PubMed: 21623373]
- Goes FS, Mcgrath J, Avramopoulos D, Wolyniec P, Pirooznia M, Ruczinski I, Nestadt G, Kenny EE, Vacic V, Peters I, Lencz T, Darvasi A, Mulle JG, Warren ST, Pulver AE, 2015. Genome-wide association study of schizophrenia in Ashkenazi Jews. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 168, 649–659.
- Granic I, 2014. The role of anxiety in the development, maintenance, and treatment of childhood aggression. *Dev. Psychopathol.* 26, 1515–1530. [PubMed: 25422976]
- Gregg TR, Siegel A, 2001. Brain structures and neurotransmitters regulating aggression in cats: Implications for human aggression. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* 25, 91–140.
- Griswold AJ, Dueker ND, Van Booven D, Rantus JA, Jaworski JM, Slifer SH, Schmidt MA, Hulme W, Konidari I, Whitehead PL, Cuccaro ML, Martin ER, Haines JL, Gilbert JR, Hussman JP, Pericak-Vance MA, 2015. Targeted massively parallel sequencing of autism spectrum disorder-associated

genes in a case control cohort reveals rare loss-of-function risk variants. *Mol. Autism* 6, 43. [PubMed: 26185613]

- Griswold AJ, Ma D, Cukier HN, Nations LD, Schmidt MA, Chung RH, Jaworski JM, Salyakina D, Konidari I, Whitehead PL, Wright HH, Abramson RK, Williams SM, Menon R, Martin ER, Haines JL, Gilbert JR, Cuccaro ML, Pericak-Vance MA, 2012. Evaluation of copy number variations reveals novel candidate genes in autism spectrum disorder-associated pathways. *Hum. Mol. Genet.* 21, 3513–3523. [PubMed: 22543975]
- Hamada N, Ito H, Nishijo T, Iwamoto I, Morishita R, Tabata H, Momiyama T, Nagata K-I, Shibata H, Huynh DP, Pulst SM, Jin Y, Underwood JG, Boutz PL, Dougherty JD, Stoilov P, Black DL, Zhou HL, Baraniak AP, Lou H, Lee JA, Tang ZZ, Black DL, Gehman LT, Zhang C, Philippe A, Bucan M, Abrahams BS, Wang K, Glessner JT, Gai X, Pinto D, Sanders SJ, Buxbaum JD, Lucarelli P, Voineagu I, Weyn-Vanhentenryck SM, Bhalla K, Elia J, Hamshere ML, Xu B, Hamada N, Nakahata S, Kawamoto S, Kim KK, Kim YC, Adelstein RS, Kawamoto S, Konno D, Koresawa Y, Kawabata I, Umeda T, Yamamoto K, Okabe S, Hamada N, Hanai N, Nagata K-I, Ito H, Iwamoto I, Morishita R, Asano T, Shinoda T, Ito H, Tabata H, Nakajima K, Uchino S, Nishimura YV, Shinoda T, Inaguma Y, Ito H, Nagata K-I, Gehman LT, Lal D, Friocourt G, Tabata H, Nakajima K, Noctor SC, Martínez-Cerdeño V, Ivic L, Kriegstein AR, Solecki DJ, Model L, Gaetz J, Kapoor TM, Hatten ME, Tanaka T, Nadarajah B, Brunstrom JE, Grutzendler J, Wong RO, Pearlman AL, Craig AM, Banker G, Tepper JM, Sharpe NA, Koós TZ, Trent F, Yoshikawa G, Sasaki J, Raj B, Blencowe BJ, Carninci P, Ihara M, Zhao L, Longo-Guess C, Harris BS, Lee J-W, Ackerman SL, Inaguma Y, Lee J-A, Reiner O, Sapir T, Tsai L-H, Gleeson JG, Fogel BL, Zhang J, O'Brien JE, 2016. Essential role of the nuclear isoform of RBFOX1, a candidate gene for autism spectrum disorders, in the brain development. *Sci. Rep.* 6, 30805. [PubMed: 27481563]
- Hammock EAD, Levitt P, 2011. Developmental expression mapping of a gene implicated in multiple neurodevelopmental disorders, A2bp1 (Fox1). *Dev. Neurosci.* 33, 64–74. [PubMed: 21346316]
- Harro M, Eensoo D, Kiive E, Merenák L, Alep J, Oreland L, Harro J, 2001. Platelet monoamine oxidase in healthy 9- and 15-years old children: The effect of gender, smoking and puberty. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* 25, 1497–1511.
- Heatherington TF, Wagner DD, 2011. Cognitive neuroscience of self-regulation failure. *Trends Cogn. Sci.* 15, 132–139. [PubMed: 21273114]
- Herold C, Hooli BV, Mullin K, Liu T, Roehr JT, Mattheisen M, Parrado AR, Bertram L, Lange C, Tanzi RE, 2016. Family-based association analyses of imputed genotypes reveal genome-wide significant association of Alzheimer's disease with OSBPL6, PTPRG, and PDCL3. *Mol. Psychiatry* 21, 1608–1612. [PubMed: 26830138]
- Hoaken PNS, Stewart SH, 2003. Drugs of abuse and the elicitation of human aggressive behavior. *Addict. Behav.* 28, 1533–1554. [PubMed: 14656544]
- Hoptman MJ, Antonius D, Kline N, 2011. Neuroimaging correlates of aggression in schizophrenia: an update. *Curr. Opin. Psychiatry* 24, 100–106. [PubMed: 21178624]
- Jin Y, Suzuki H, Maegawa S, Endo H, Sugano S, Hashimoto K, Yasuda K, Inoue K, 2003. A vertebrate RNA-binding protein Fox-1 regulates tissue-specific splicing via the pentanucleotide GCAUG. *EMBO J.* 22, 905–912. [PubMed: 12574126]
- Kallasmaa T, Allik J, Realo A, McCrae RR, 2000. The Estonian version of the NEO-PI-R: An examination of universal and culture-specific aspects of the five-factor model. *Eur. J. Pers.* 14, 265–278.
- Kanduri C, Kantojärvi K, Salo PM, Vanhala R, Buck G, Blancher C, Lähdesmäki H, Järvelä I, 2016. The landscape of copy number variations in Finnish families with autism spectrum disorders. *Autism Res.* 9, 9–16. [PubMed: 26052927]
- King S, Waschbusch DA, 2010. Aggression in children with attention-deficit/hyperactivity disorder. *Expert Rev. Neurother.* 10, 1581–1594. [PubMed: 20925473]
- Kodžopelji J, Smederevac S, Mitrovi D, Dini B, olovi P, 2014. School bullying in adolescence and personality traits: a person-centered approach. *J. Interpers. Violence* 29, 736–757. [PubMed: 24255068]
- Kohannim O, Hibar DP, Jahanshad N, Stein JL, Hua X, Toga AW, Jack CR, Weiner MW, Thompson PM, the Alzheimer's Disease Neuroimaging Initiative, P.M., Initiative, the A.D.N., 2012a.

Predicting temporal lobe volume on MRI from genotypes using L(1)-L(2) regularized regression. *Proceedings. IEEE Int. Symp. Biomed. Imaging* 1160–1163.

- Kohannim O, Hibar DP, Stein JL, Jahansha N, Hua X, Rajagopalan P, Toga AW, Jack CR, Weiner MW, de Zubicaray GI, McMahon KL, Hansell NK, Martin NG, Wright MJ, Thompson PM, 2012b. Discovery and replication of gene influences on brain structure using LASSO regression. *Front. Neurosci.* 6, 115. [PubMed: 22888310]
- Krämer UM, Jansma H, Tempelmann C, Münte TF, 2007. Tit-for-tat: The neural basis of reactive aggression. *Neuroimage* 38, 203–211. [PubMed: 17765572]
- Kruesi MJP, Casanova MF, Mannheim G, Johnson-Bilder A, 2004. Reduced temporal lobe volume in early onset conduct disorder. *Psychiatry Res.* 132, 1–11. [PubMed: 15546698]
- Látalová K, 2009. Bipolar disorder and aggression. *Int. J. Clin. Pract.* 63, 889–899. [PubMed: 19490199]
- Lee JA, Damianov A, Lin CH, Fontes M, Parikshak NN, Anderson ES, Geschwind DH, Black DL, Martin KC, 2016. Cytoplasmic Rbfox1 Regulates the Expression of Synaptic and Autism-Related Genes. *Neuron* 89, 113–128. [PubMed: 26687839]
- Li YI, Sanchez-Pulido L, Haerty W, Ponting CP, 2015. RBFOX and PTBP1 proteins regulate the alternative splicing of micro-exons in human brain transcripts. *Genome Res.* 25, 1–13. [PubMed: 25524026]
- Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, Anderson DJ, 2011. Functional identification of an aggression locus in the mouse hypothalamus. *Nature* 470, 221–226. [PubMed: 21307935]
- Luque MJ, Martin-Lopez M, Navarro JF, 2009. Effects of LY379268, A Selective Agonist of mGlu2/3 Receptors, on Isolation-Induced Aggression in Male Mice. *Open Pharmacol. J.* 3, 17–20.
- Martin CL, Duvall JA, Ilkin Y, Simon JS, Arreaza MG, Wilkes K, Alvarez-Retuerto A, Whichello A, Powell CM, Rao K, Cook E, Geschwind DH, 2007. Cytogenetic and molecular characterization of A2BP1/FOX1 as a candidate gene for autism. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 144, 869–876.
- Martín-García E, Fernández-Castillo N, Burokas A, Gutiérrez-Cuesta J, Sánchez-Mora C, Casas M, Ribasés M, Cormand B, Maldonado R, 2015. Frustrated expected reward induces differential transcriptional changes in the mouse brain. *Addict. Biol.* 20, 22–27. [PubMed: 25288320]
- Melhem N, Middleton F, McFadden K, Klei L, Faraone SV, Vinogradov S, Tiobech J, Yano V, Kuarstei S, Roeder K, Byerley W, Devlin B, Myles-Worsley M, 2011. Copy number variants for schizophrenia and related psychotic disorders in Oceanic Palau: Risk and transmission in extended pedigrees. *Biol. Psychiatry* 70, 1115–1121. [PubMed: 21982423]
- Merjonen P, Keltikangas-Järvinen L, Jokela M, Seppälä I, Lyytikäinen L-P, Pulkki-Råback L, Kivimäki M, Elovainio M, Kettunen J, Ripatti S, Kähönen M, Viikari J, Palotie A, Peltonen L, Raitakari OT, Lehtimäki T, 2011. Hostility in adolescents and adults: a genome-wide association study of the Young Finns. *Transl. Psychiatry* 1, e11. [PubMed: 22832427]
- Meyer-Lindenberg A, Buckholtz JW, Kolachana B, R Hariri A, Pezawas L, Blasi G, Wabnitz A, Honea R, Verchinski B, Callicott JH, Egan M, Mattay V, Weinberger DR, 2006. Neural mechanisms of genetic risk for impulsivity and violence in humans. *Proc. Natl. Acad. Sci. U. S. A.* 103, 6269–6274. [PubMed: 16569698]
- Mick E, McGough J, Deutsch CK, Frazier JA, Kennedy D, Goldberg RJ, 2014. Genome-wide association study of proneness to anger. *PLoS One* 9, e87257.
- Noor A, Lionel AC, Cohen-Woods S, Moghimi N, Rucker J, Fennell A, Thiruvahindrapuram B, Kaufman L, Degagne B, Wei J, Parikh SV, Muglia P, Forte J, Scherer SW, Kennedy JL, Xu W, McGuffin P, Farmer A, Strauss J, Vincent JB, 2014. Copy number variant study of bipolar disorder in Canadian and UK populations implicates synaptic genes. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 165, 303–313.
- Pappa I, St Pourcain B, Benke K, Cavadino A, Hakulinen C, Nivard MG, Nolte IM, Tiesler CMT, Bakermans-Kranenburg MJ, Davies GE, Evans DM, Geoffroy MC, Grallert H, Groen-Blokhuis MM, Hudziak JJ, Kemp JP, Keltikangas-Järvinen L, McMahon G, Mileva-Seitz VR, Motazed E, Power C, Raitakari OT, Ring SM, Rivadeneira F, Rodriguez A, Scheet PA, Seppälä I, Snieder H, Standl M, Thiering E, Timpson NJ, Veenstra R, Velders FP, Whitehouse AJO, Smith GD, Heinrich J, Hyppönen E, Lehtimäki T, Middeldorp CM, Oldehinkel AJ, Pennell CE, Boomsma DI, Tiemeier

- H, 2016. A genome-wide approach to children's aggressive behavior: The EAGLE consortium. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 171, 562–572.
- Passamonti L, Fera F, Magariello A, Cerasa A, Gioia MC, Muglia M, Nicoletti G, Gallo O, Provinciali L, Quattrone A, 2006. Monoamine oxidase-A genetic variations influence brain activity associated with inhibitory control: New insight into the neural correlates of impulsivity. *Biol. Psychiatry* 59, 334–340. [PubMed: 16202396]
- Potegal M, 2012. Temporal and frontal lobe initiation and regulation of the top-down escalation of anger and aggression. *Behav. Brain Res.* 231, 386–395. [PubMed: 22085875]
- Raine A, Ishikawa SS, Arce E, Lencz T, Knuth KH, Bihrlé S, LaCasse L, Colletti P, 2004. Hippocampal structural asymmetry in unsuccessful psychopaths. *Biol. Psychiatry* 55, 185–191. [PubMed: 14732599]
- Rajman M, Metge F, Fiore R, Khudayberdiev S, Aksoy-Aksel A, Bicker S, Reschke CR, Raoof R, Brennan GP, Delanty N, Farrell MA, O'Brien DF, Bauer S, Norwood B, Veno MT, Krüger M, Braun T, Kjems J, Rosenow F, Henshall DC, Dieterich C, Schratt G, 2017. A microRNA-129-5p/Rbfox crosstalk coordinates homeostatic downscaling of excitatory synapses. *EMBO J.* ahead of print.
- Sebat J, Lakshmi B, Malhotra D, Troge J, Lese-martin C, Walsh T, Yamrom B, Yoon S, Krasnitz A, Kendall J, Leotta A, Pai D, Zhang R, Lee Y, Hicks J, Spence SJ, Lee AT, Puura K, Lehtimäki T, Ledbetter D, Gregersen PK, Bregman J, Sutcliffe JS, Jobanputra V, Chung W, Warburton D, 2007. Strong association of de novo copy number mutations with Autism. *Science.* 316, 445–449. [PubMed: 17363630]
- Sharpe JP, Desai S, 2001. The Revised Neo Personality Inventory and the MMPI-2 Psychopathology Five in the prediction of aggression. *Pers. Individ. Dif.* 31, 505–518.
- Shorter J, Couch C, Huang W, Carbone MA, Peiffer J, Anholt RRH, Mackay TFC, 2015. Genetic architecture of natural variation in *Drosophila melanogaster* aggressive behavior. *Proc. Natl. Acad. Sci.* 112, E3555–E3563. [PubMed: 26100892]
- Siddaway AP, Taylor PJ, Wood AM, Siddaway AP, Taylor PJ, Wood AM, 2017. Reconceptualizing Anxiety as a Continuum That Ranges From High Calmness to High Anxiety: The Joint Importance of Reducing Distress and Increasing Well-Being. *J. Pers. Soc. Psychol.* in press.
- Siever LJ, 2008. Neurobiology of Aggression and Violence. *Am. J. Psychiatry* 165, 429–442. [PubMed: 18346997]
- Sonuga-Barke EJS, Lasky-Su J, Neale BM, Oades R, Chen W, Franke B, Buitelaar J, Banaschewski T, Ebstein R, Gill M, Anney R, Miranda A, Mulas F, Roeyers H, Rothenberger A, Sergeant J, Steinhausen HC, Thompson M, Asherson P, Faraone SV, 2008. Does parental expressed emotion moderate genetic effects in ADHD? An exploration using a genome wide association scan. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 147, 1359–1368.
- Soyka M, 2011. Neurobiology of aggression and violence in schizophrenia. *Schizophr. Bull.* 37, 913–920. [PubMed: 21860037]
- Spielberger CD, 1983. *Manual for the State-Trait Anxiety Inventory (STAI)*. Palo Alto: Consulting Psychologists Press.
- Sterzer P, Stadler C, Krebs A, Kleinschmidt A, Poustka F, 2005. Abnormal neural responses to emotional visual stimuli in adolescents with conduct disorder. *Biol. Psychiatry* 57, 7–15. [PubMed: 15607294]
- Takahashi A, Lee RX, Iwasato T, Itohara S, Arima H, Bettler B, Miczek KA, Koide T, 2015. Glutamate input in the dorsal raphe nucleus as a determinant of escalated aggression in male mice. *J. Neurosci.* 35, 6452–6463. [PubMed: 25904796]
- Tremblay PF, Ewart LA, 2005. The Buss and Perry Aggression Questionnaire and its relations to values, the Big Five, provoking hypothetical situations, alcohol consumption patterns, and alcohol expectancies. *Pers. Individ. Dif.* 38, 337–346.
- Underwood JG, Boutz PL, Dougherty JD, Stoilov P, Black DL, 2005. Homologues of the *Caenorhabditis elegans* Fox-1 protein are neuronal splicing regulators in mammals. *Mol. Cell. Biol.* 25, 10005–10016. [PubMed: 16260614]
- Vaht M, Kiive E, Veidebaum T, Harro J, 2016. A functional vesicular monoamine transporter 1 (VMAT1) gene variant is associated with affect and the prevalence of anxiety, affective,

- and alcohol use disorders in a longitudinal population-representative birth cohort study. *Int. J. Neuropsychopharmacol.* 19, 1–9.
- van Dongen J, Nivard MG, Baselmans BML, Zilhão NR, Ligthart L, Heijmans BT, Bartels M, Boomsma DI, 2015. Epigenome-wide association study of aggressive behavior. *Twin Res. Hum. Genet.* 18, 686–698.
- Van Praag HM, 1998. Anxiety and increased aggression as pacemakers of depression. *Acta Psychiatr. Scand.* 98, 81–88. [PubMed: 9696519]
- Vekovischeva OY, Aitta-Aho T, Echenko O, Kankaanpää A, Seppälä T, Honkanen A, Sprengel R, Korpi ER, 2004. Reduced aggression in AMPA-type glutamate receptor GluR-A subunit-deficient mice. *Genes, Brain Behav.* 3, 253–265. [PubMed: 15344919]
- Veroude K, Zhang-James Y, Fernández-Castillo N, Bakker MJ, Cormand B, Faraone SV, 2016. Genetics of aggressive behavior: An overview. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 171, 3–43.
- Voineagu I, Wang X, Johnston P, Lowe JK, Tian Y, Horvath S, Mill J, Cantor RM, Blencowe BJ, Geschwind DH, 2011. Transcriptomic analysis of autistic brain reveals convergent molecular pathology. *Nature* 474, 380–384. [PubMed: 21614001]
- Volavka J, 2013. Violence in schizophrenia and bipolar disorder. *Psychiatr. Danub.* 25, 24–33. [PubMed: 23470603]
- Vounou M, Janousova E, Wolz R, Stein JL, Thompson PM, Rueckert D, Montana G, 2012. Sparse reduced-rank regression detects genetic associations with voxel-wise longitudinal phenotypes in Alzheimer’s disease. *Neuroimage* 60, 700–716. [PubMed: 22209813]
- Wang KS, Liu XF, Aragam N, 2010. A genome-wide meta-analysis identifies novel loci associated with schizophrenia and bipolar disorder. *Schizophr. Res.* 124, 192–199. [PubMed: 20889312]
- Wen M, Yan Y, Yan N, Chen XS, Liu SY, Feng ZH, 2015. Upregulation of RBFOX1 in the malformed cortex of patients with intractable epilepsy and in cultured rat neurons. *Int. J. Mol. Med.* 35, 597–606. [PubMed: 25571999]
- Xu B, Roos JL, Levy S, van Rensburg EJ, Gogos JA, Karayiorgou M, 2008. Strong association of de novo copy number mutations with sporadic schizophrenia. *Nat. Genet.* 40, 880–885. [PubMed: 18511947]
- Zapata I, Serpell JA, Alvarez CE, 2016. Genetic mapping of canine fear and aggression. *BMC Genomics* 17, 572. [PubMed: 27503363]
- Zhao Q-F, Tan L, Wang H-F, Jiang T, Tan M-S, Tan L, Xu W, Li J-Q, Wang J, Lai T-J, Yu J-T, 2016. The prevalence of neuropsychiatric symptoms in Alzheimer’s disease: Systematic review and meta-analysis. *J. Affect. Disord.* 190, 264–271. [PubMed: 26540080]
- Zhao W-W, 2013. Intragenic deletion of RBFOX1 associated with neurodevelopmental/neuropsychiatric disorders and possibly other clinical presentations. *Mol. Cytogenet.* 6, 26. [PubMed: 23822903]
- Zhong X, Drgonova J, Li CY, Uhl GR, 2015. Human cell adhesion molecules: Annotated functional subtypes and overrepresentation of addiction-associated genes. *Ann. N. Y. Acad. Sci.* 1349, 83–95. [PubMed: 25988664]




Figure 1.
 A) Diagram of the human *RBFOX1* gene (NM_018723, GRCh38 Chr16: 6,019,131–7,713,338). Solid boxes indicate exons, with the coding regions in black and the non-coding ones in grey. Exon numbering is shown on top of the figure. Eight SNPs found nominally associated with aggressive phenotypes in several genome-wide association studies (GWAS) are shown below, with their locations indicated by discontinuous lines. B) Linkage disequilibrium (LD) plot of the SNPs considering r^2 , according to Haploview 4.2. Black boxes indicate complete LD and grey is for moderate LD.

Table 1.

Suggestive associations in the *RBFox1* gene identified in GWAS of aggressive behaviors.

SNP	Reference (GWAS)	Chr	bp (GRCh38)	Alleles ^{&}	MAF*	Phenotype associated	Sample	P-value	Risk allele
rs6500744	Sonuga-Barke et al., 2008	16	6063660	C>T	0.46	Conduct disorder symptoms interacting with mother's warmth	938 trios	3.65 e-05	C
rs8062784	Merjonen et al., 2011	16	6096633	A>T	0.085	Anger (mean of 4 measurement phases)	2,242 adults	2.34 e-05	A
rs809682	Pappa et al., 2016	16	6346372	T>A	0.26	Children's aggressive behavior	18,988 children	2.62 e-04	T
rs12922093	Pappa et al., 2016	16	6346947	T>C	0.26	Children's aggressive behavior	18,988 children	3.21 e-04	T
rs12373031	Pappa et al., 2016	16	6347436	T>C	0.14	Children's aggressive behavior	18,988 children	4.93 e-04	T
rs10521042	Pappa et al., 2016	16	6349836	C>T	0.14	Children's aggressive behavior	18,988 children	4.35 e-04	C
rs12921846	Anney et al., 2008	16	6860384	A>T	0.16	Conduct disorder	938 trios	1.39 e-05	A
rs10153149	Anney et al., 2008	16	6875239	A>C	0.16	Conduct disorder	938 trios	3.64 e-05	A

[&] According to the forward strand.

* Minimum allele frequency in the European population from 1000 Genomes (<http://www.internationalgenome.org/>).