



## EDITORIAL

## Peptide drugs: a new direction in cancer immunotherapy

Xinghua Sui, Xiaoshuang Niu, Xiuman Zhou, Yanfeng Gao

School of Pharmaceutical Sciences (Shenzhen), Shenzhen Campus of Sun Yat-sen University, Shenzhen 518107, China

Cancer immunotherapy has emerged as a promising approach in cancer treatment and is considered a major advancement after surgical interventions, radiotherapy, chemotherapy, and targeted therapy. The clinical use of immunotherapeutic drugs, particularly antibody-based drugs that target immune checkpoints, has notably increased<sup>1</sup>. Unlike traditional antitumor drugs, which exclusively targeting tumor cells, these drugs have a unique mechanism of action, as they inhibit protein-protein interactions across multiple cell types, such as PD-1/PD-L1 blockade may function to interfere among T cell, tumor cell, macrophage, and dendritic cell (Figure 1). Nevertheless, timely withdrawal of antibody therapeutics in the presence of immune-mediated adverse reactions poses a substantial challenge, because of their high molecular weight and long half-life. The cellular uptake of most small molecule chemical drugs is generally feasible but may be accompanied by off-target effects. Peptide therapeutics, in contrast, occupy an intermediate position between monoclonal antibody therapeutics and small molecule chemical drugs in the spectrum of therapeutic agents. Peptides have notable advantages, including remarkable selectivity, particularly to drug targets on the cell surface, robust infiltration into solid tumors, and facile synthesis; therefore, they are critical contenders for immune checkpoint inhibition.

However, the clinical utility of peptide therapeutics is constrained by 2 primary obstacles: enzymatic degradation within the physiological milieu and suboptimal oral bioavailability. Various approaches have been implemented to avoid enzymatic degradation, including the use of non-natural amino acids, cyclization modification, and mirror-image phage

display technology to develop peptides consisting of amino acids in all-D configuration<sup>2,3</sup>. The strategic development of suitable formulations can improve the oral characteristics and bioavailability of peptides. The primary objective of this review is to provide a brief overview of advancements in peptide immune checkpoint inhibitors. In addition, the potential applications and future prospects of these peptides in cancer immunotherapy and diagnosis are examined.

### Peptide inhibitors that selectively target T cell immune checkpoints

Through high-throughput screening or rational design guided by the crystal structure of the PD-1/PD-L1 complex, peptide inhibitors have been designed to target PD-1/PD-L1<sup>4-6</sup>. Nevertheless, these peptides, comprising L-amino acids, have shown suboptimal stability and have encountered challenges in achieving robust target blockade. Peptide cyclization has been demonstrated to enhance the enzymatic degradation stability and conformational stability of peptides<sup>7</sup>. BMS-986189 (Figure 2), a macrocyclic peptide developed by Bristol-Myers Squibb (BMS), has been designed to target PD-L1. This promising therapeutic candidate has progressed to a phase I clinical trial<sup>8</sup>. In an alternative approach, mirror-image phage display technology enables the screening of peptides with an all-D configuration. This innovative method effectively addresses concerns regarding enzymatic degradation stability and offers a comprehensive solution. We have successfully generated a novel class of anti-proteolytic peptides denoted <sup>D</sup>PPA-1 (nyskptdrqyhf), which target PD-L1<sup>9</sup> and comprise exclusively all-D amino acids. Moreover, another D-peptide targeting PD-L1, OPBP-1 (Gqsehhrvysf), has been obtained and loaded with trimethyl chitosan hydrogel for oral administration. This peptide shows notably enhanced oral bioavailability and half-life in rats. Therefore, OPBP-1 is the first peptide for oral administration for immune checkpoint blockade<sup>10</sup>.

Correspondence to: Yanfeng Gao

E-mail: gaoyf29@mail.sysu.edu.cn


ORCID ID: <https://orcid.org/0000-0001-5533-7100>

Received August 7, 2023; accepted September 28, 2023;


published online December 7, 2023.

Available at [www.cancerbiomed.org](http://www.cancerbiomed.org)

©2024 The Authors. Creative Commons Attribution-NonCommercial 4.0 International License



**Figure 1** Immune checkpoints as drug targets for cancer immunotherapy. The interaction of immune checkpoint receptors and ligands expressed on the surfaces of MDSCs, macrophages, T cells, and tumor cells are shown.



**Figure 2** Possible structure of the PD-L1 macrocyclic peptide inhibitor BMS-986189, which is undergoing a phase I clinical trial<sup>8</sup>.

Beyond the PD-1/PD-L1 axis, other pairs of immune checkpoints have major roles in the context of tumor-induced immunosuppression; these pairs include the T cell immunoreceptor with immunoglobulin and ITIM domains (TIGIT) and poliovirus receptor (PVR); lymphocyte activation gene 3 (LAG-3) with major histocompatibility complex class II (MHC-II); and V-domain Ig suppressor of T cell activation (VISTA) with P-selectin glycoprotein ligand 1 (PSGL-1). Concurrently inhibiting these targets is anticipated to increase the efficacy of PD-1/PD-L1 blockade.

Using mirror-image phage display, our team has successfully acquired <sup>D</sup>TBP-3 (GGYtfhwrlnp), the first D peptide inhibitor that blocks the TIGIT/PVR interaction. The remarkable efficacy of <sup>D</sup>TBP-3 has been observed in 2 distinct murine cancer models:

the anti-PD-1 antibody-responsive CT26 colorectal cancer model and the anti-PD-1 antibody-resistant 4T1 breast cancer model<sup>11</sup>. By using phage display, we have discovered the initial cyclic peptide inhibitor C25 (CVPMTYRAC), which blocks the LAG-3/MHC-II interaction. The administration of C25 peptide notably increases both the quantity and efficacy of CD8<sup>+</sup> T cells, but not CD4<sup>+</sup> T cells, localized at tumor sites. Additionally, the relative abundance of Treg cells within tumor tissue decreases<sup>12</sup>. Given the presence of an acidic microenvironment within tumor tissues, we have identified peptide inhibitors that selectively target immune checkpoints within this acidic milieu. Pal-DVS3 (dpGWSFGKLHWPGS-Pal), a peptide inhibitor with incorporation of D-amino acids to enhance enzymatic stability and fatty acid modifications to prolong the *in vivo* half-life, has been specifically designed to target VISTA/PSGL-1. This peptide exhibits enduring anti-tumor effects by mitigating the immunosuppressive effects of myeloid-derived suppressor cells (MDSCs) on CD8<sup>+</sup> T cells<sup>13</sup>.

## Peptide inhibitors that selectively target immune checkpoints on other immune cell subsets

The intriguing phenomenon of “cold tumors”, distinguished by a lack of T cell infiltration within specific tumor tissues, is a formidable obstacle in immunotherapy. Nevertheless, most solid tumors have substantial presence of macrophages—immune cells with crucial roles in the tumor microenvironment. Interestingly, certain immune checkpoints, such as CD47/SIRPα and CD24/Siglec10, and even

PD-1/PD-L1, have been identified as key players in regulating the “don’t eat me” signal. These checkpoints effectively hinder macrophage phagocytosis, thereby impeding the elimination of tumor cells by macrophages.

Our group has used phage display technology to develop peptide inhibitors targeting CD47/SIRP $\alpha$ . Notably, we have successfully identified pep-20, along with its terminal D-amino acid substituted analogue, pep-20-D12 (awsATWSNYwrh), as potential inhibitors. Our findings suggest that peptides without Fc-mediated ADCC effects may be superior alternatives to CD47 antibodies because these peptides can specifically activate macrophages in the tumor microenvironment rather than inducing the phagocytosis of peripheral red blood cells, thus avoiding anemia<sup>14</sup>. Recently, using phage display and subsequent retro-inverse strategies, we have successfully developed a D-peptide CSBP (ldvflyse) that targets both CD24/Siglec10 and PD-1/PD-L1. These peptides increase the phagocytic activity of both macrophages and M-MDSCs, thus fortifying the innate immune response. Moreover, they increase the functionality of CD8<sup>+</sup> T cells and consequently foster adaptive anti-tumor immune responses. Particularly when used in conjunction with radiotherapy, these peptides have robust synergistic anti-tumor effects, owing to substantial infiltration of macrophages and MDSCs into tumor tissues induced by radiotherapy<sup>15</sup>.

Other immune cells also express immune checkpoints, such as TIPE2<sup>16</sup> and KIR2DL4<sup>17</sup> on NK cells; TIM-1<sup>18</sup> on B cells; and gp49B<sup>19</sup> on neutrophils, NK cells, monocytes, etc. Although these immune checkpoints and various immune cells play a crucial immune modulatory role in the anti-tumor immune response, studies on their peptide inhibitors remain limited. Future research will greatly benefit from the development of peptides targeting these immune checkpoints.


## Dual-function peptides uniquely exhibiting distinct biological functions

Dual-function peptides, akin to bispecific antibodies, are peptide entities that have been conjugated to effectively target 2 functionally related or complementary entities. This unique characteristic enables them to concurrently regulate distinct signaling pathways, thereby exhibiting multifaceted modulatory effects (**Figure 3**). In contrast to bispecific antibodies, dual-function peptides have relatively streamlined, less complex design and synthesis processes.

In our recent study, we designed a novel peptide inhibitor known as Pal-DMPOP (Pal-PEG<sub>4</sub>-WSMTWWNYWrvysf), by combining the minimal active fragment of the PD-1/PD-L1 blocking peptide OPBP-1 with the minimal variant of the CD47/SIRP $\alpha$  peptide inhibitor pep-20. Pal-DMPOP exerts its immunomodulatory effects by selectively engaging and modulating the signaling pathways associated with both T cell and macrophage checkpoints, thus leading to the activation and enhancement of anti-tumor immune responses<sup>20</sup>. In another study, we have conjugated the minimal active fragment of OPBP-1 with the anti-angiogenesis peptide <sup>D</sup>A7R, and subsequently coupled this bispecific peptide with the albumin-binding peptide <sup>D</sup>SP, thus yielding a long-acting dual-functional peptide termed <sup>D</sup>SPOGS. The <sup>D</sup>A7R peptide has notable anti-tumor effects through impeding neovascularization within tumor tissues. Unfortunately, these anti-angiogenic effects impede the infiltration of immune cells into tumor tissue. However, <sup>D</sup>SPOGS effectively addresses this concern and has demonstrated synergistic anti-tumor effects<sup>21</sup>.

## Peptide-drug conjugates (PDCs) and self-assembly nanoparticles

In contrast to antibody-drug conjugates (ADCs), peptides can be more easily conjugated with small molecule drugs, thereby facilitating straightforward synthesis. Amphiphilic PDCs have been engineered by conjugating the small molecule radiosensitizers with the PD-1/PD-L1 blocking peptide <sup>D</sup>PPA-1. The self-assembly of these molecules results in the formation of nanoparticles, which effectively encapsulate TLR7/8 agonists. This intricate process has remarkable synergistic outcomes, wherein radiotherapy sensitization, activation of innate immunity, and enhancement of adaptive immune responses are achieved concurrently<sup>22</sup>. In another study, the <sup>D</sup>PPA-1 peptide has been conjugated with the chemotherapeutic agent doxorubicin by using a substrate sequence specifically designed for matrix metalloproteinase 2 (MMP2). This strategic approach has led to the development of nanoparticles that are responsive to the enzymatic activity of MMP2. The nanoparticles have a dual mechanism of action, thus effectively inducing both the cytotoxic effects of chemotherapeutics and T cell-mediated immune killing functions<sup>23</sup>. PDCs comprising peptides targeting non-endocytic receptors and non-toxic drugs targeting immunotherapeutic targets hold immense future promise, because their mechanisms of action can deviate from those



**Figure 3** Strategies for the design and modification of peptides. The stability of peptides can be enhanced by cyclization, stapling, or the use of non-natural amino acids. FcRn or albumin targeting designs or lipid acid modifications can extend the *in vivo* half-life of peptides. Rational conjugation designs can endow peptides with dual functions. Nanomaterials can be used for oral peptide delivery. Peptides can be self-assembled, coupled to create PDCs, or conjugated with radiolabels for tumor diagnosis or treatment.

of conventional ADCs, thereby avoiding unforeseen toxicity associated with toxin molecules in ADCs.

## Radiolabeled peptides for tumor diagnosis and therapy

The efficacy of immune checkpoint inhibitors is intricately associated with the expression levels of immune checkpoint molecules. The dynamic nature of immune checkpoint molecule expression is evident throughout tumor progression and treatment. Hence, non-invasive *in vivo* detection of immune checkpoint expression with high sensitivity is imperative. The tissue penetration and long half-life of radiolabeled immune

checkpoint antibodies poses several limitations, thus potentially resulting in extended clearance from the body and subsequent unnecessary toxicity. Radiolabeled peptides, serving as diagnostic probes, have remarkable attributes, including robust tumor infiltration, negligible toxicity, and favorable half-life characteristics.

The <sup>64</sup>Cu-labeled <sup>D</sup>PPA-1 peptide has notable efficacy in rapid and precise visualization of PD-L1 expression within tumor tissues. This peptide has potential as a radiotherapeutic agent<sup>24</sup>. Furthermore, <sup>68</sup>Ga-GP12 (<sup>D</sup>TBP-3), as a positron emission tomography (PET) tracer, has been reported to specifically target TIGIT in patients diagnosed with advanced non-small cell lung cancer. The imaging findings obtained with <sup>68</sup>Ga-GP12 PET/CT are similar to those obtained with

the widely used  $^{18}\text{F}$ -FDG PET/CT, but confer particular advantages in the identification of lymph node metastases<sup>25</sup>.

## Discussion

In cancer immunotherapy, in addition to the immune checkpoints present on T cells and macrophages, newly identified cell subsets and targets in the tumor microenvironment have been continually explored. These discoveries have been instrumental in advancing peptide-based therapeutics. In long-acting peptide design, cutting-edge methods, such as “click chemistry”, and intricate peptide modification techniques, such as stapled peptides, alongside approaches aimed at prolonging *in vivo* half-life with albumin-binding properties, will undoubtedly have critical roles<sup>2</sup>. As PDCs continue to be extensively investigated, the fundamental strategy remains rooted in the paradigm of antibody-drug conjugates. This approach entails the transport of toxin molecules into the intracellular milieu of tumor cells *via* endocytosis, which cannot fully harness the potential benefits of peptides. Unlike the component of ADCs, the peptide mass is similar to that of the small molecule payload in PDCs. Therefore, novel methods and strategies must be explored to increase therapeutic efficacy and avoid unwanted adverse effects. The advent of  $^{177}\text{Lu}$ -octreotate has propelled scientific investigation. Furthermore, given their *in vivo* stability and half-life of in tumor diagnosis, peptides have substantial research potential that might surpass that of therapeutic agents.

The oral route of administration has consistently posed a major hurdle in peptide therapeutics. The investigation of orally administered drugs for cancer immunotherapy has significance in the field, by enabling the flexible adaptation of dosing frequency in accordance with a patient’s immune status. The gastrointestinal tract is the primary site of absorption for peptide drugs administered orally. Importantly, the intestinal mucosal immune system, which encompasses the gut-associated lymphoid tissue, is the largest lymphoid organ in the human body<sup>26</sup>. Hence, deeper investigation of the mechanisms underlying the specific targeting of intestinal cells is of considerable scientific interest, to enhance the oral bioavailability of peptide-based therapeutics and stimulate these cells to elicit robust anti-tumor immune responses.

Beyond serving as immune checkpoint inhibitors, peptides may be used as vaccines or vaccine delivery carriers, thus playing critical roles in cancer immunotherapy. Hundreds of peptide vaccines, including long peptide vaccines, short peptide

vaccines, and other forms, have entered clinical research. The combination of immune checkpoint blockade and vaccines has been widely recognized to achieve a synergistic antitumor response.

## Grant support

This work was supported by grants from the National Natural Science Foundation of China (Grant No. U20A20369) and GuangDong Basic and Applied Basic Research Foundation (Grant No. 2022B1515120085).

## Conflict of interest statement

No potential conflicts of interest are disclosed.

## Author contributions

Conceived and designed the analysis: Xinghua Sui, Xiaoshuang Niu, Xiuman Zhou, Yanfeng Gao.

Wrote the paper: Xinghua Sui, Yanfeng Gao.

## References

1. Kiyotani K, Toyoshima Y, Nakamura Y. Personalized immunotherapy in cancer precision medicine. *Cancer Biol Med.* 2021; 18: 955-65.
2. Muttenthaler M, King GF, Adams DJ, Alewood PF. Trends in peptide drug discovery. *Nat Rev Drug Discov.* 2021; 20: 309-25.
3. Lyu P, Kwok HF. High-throughput strategy accelerates the progress of marine anticancer peptide drug development. *Recent Pat Anticancer Drug Discov.* 2019; 14: 2-4.
4. Liu H, Zhao Z, Zhang L, Li Y, Jain A, Barve A, et al. Discovery of low-molecular weight anti-PD-L1 peptides for cancer immunotherapy. *J Immunother Cancer.* 2019; 7: 270.
5. Li C, Zhang N, Zhou J, Ding C, Jin Y, Cui X, et al. Peptide blocking of PD-1/PD-L1 interaction for cancer immunotherapy. *Cancer Immunol Res.* 2018; 6: 178-88.
6. Yin H, Zhou X, Huang YH, King GJ, Collins BM, Gao Y, et al. Rational design of potent peptide inhibitors of the PD-1:PD-L1 interaction for cancer immunotherapy. *J Am Chem Soc.* 2021; 143: 18536-47.
7. Zhai W, Zhou X, Zhai M, Li W, Ran Y, Sun Y, et al. Blocking of the PD-1/PD-L1 interaction by a novel cyclic peptide inhibitor for cancer immunotherapy. *Sci China Life Sci.* 2021; 64: 548-62.
8. Pan C, Yang H, Lu Y, Hu S, Wu Y, He Q, et al. Recent advance of peptide-based molecules and nonpeptidic small-molecules modulating PD-1/PD-L1 protein-protein interaction or targeting PD-L1 protein degradation. *Eur J Med Chem.* 2021; 213: 113170.

9. Chang HN, Liu BY, Qi YK, Zhou Y, Chen YP, Pan KM, et al. Blocking of the PD-1/PD-L1 interaction by a D-peptide antagonist for cancer immunotherapy. *Angew Chem Int Ed Engl.* 2015; 54: 11760-4.
  10. Li W, Zhu X, Zhou X, Wang X, Zhai W, Li B, et al. An orally available PD-1/PD-L1 blocking peptide OPBP-1-loaded trimethyl chitosan hydrogel for cancer immunotherapy. *J Control Release.* 2021; 334: 376-88.
  11. Zhou X, Zuo C, Li W, Shi W, Zhou X, Wang H, et al. A novel D-peptide identified by mirror-image phage display blocks TIGIT/PVR for cancer immunotherapy. *Angew Chem Int Ed Engl.* 2020; 59: 15114-8.
  12. Zhai W, Zhou X, Wang H, Li W, Chen G, Sui X, et al. A novel cyclic peptide targeting LAG-3 for cancer immunotherapy by activating antigen-specific CD8(+) T cell responses. *Acta Pharm Sin B.* 2020; 10: 1047-60.
  13. Niu X, Wu M, Li G, Zhou X, Cao W, Zhai W, et al. Identification and optimization of peptide inhibitors to block VISTA/PSGL-1 interaction for cancer immunotherapy. *Acta Pharm Sin B.* 2023; 13: 4341-666.
  14. Wang H, Sun Y, Zhou X, Chen C, Jiao L, Li W, et al. CD47/Sirp $\alpha$  blocking peptide identification and synergistic effect with irradiation for cancer immunotherapy. *J Immunother Cancer.* 2020; 8: e000905.
  15. Shen W, Shi P, Dong Q, Zhou X, Chen C, Sui X, et al. Discovery of a novel dual-targeting D-peptide to block CD24/Siglec-10 and PD-1/PD-L1 interaction and synergize with radiotherapy for cancer immunotherapy. *J Immunother Cancer.* 2023; 11: e007068.
  16. Bi J, Cheng C, Zheng C, Huang C, Zheng X, Wan X, et al. TIPE<sub>2</sub> is a checkpoint of natural killer cell maturation and antitumor immunity. *Sci Adv.* 2021; 7: eabi6515.
  17. Zheng G, Jia L, Yang AG. Roles of HLA-G/KIR<sub>2</sub>DL<sub>4</sub> in breast cancer immune microenvironment. *Front Immunol.* 2022; 13: 791975.
  18. Bod L, Kye YC, Shi J, Torlai Triglia E, Schnell A, Fessler J, et al. B-cell-specific checkpoint molecules that regulate anti-tumour immunity. *Nature.* 2023; 619: 348-56.
  19. Lee HN, Manangeeswaran M, Lewkowicz AB, Engel K, Chowdhury M, Garige M, et al. NK cells require immune checkpoint receptor LILRB4/gp49B to control neurotropic Zika virus infections in mice. *JCI Insight.* 2022; 7: e151420.
  20. Hu Z, Li W, Chen S, Chen D, Xu R, Zheng D, et al. Design of a novel chimeric peptide via dual blockade of CD47/Sirp $\alpha$  and PD-1/PD-L1 for cancer immunotherapy. *Sci China Life Sci.* 2023; 66: 2310-28.
  21. Jiao L, Dong Q, Zhai W, Zhao W, Shi P, Wu Y, et al. A PD-L1 and VEGFR2 dual targeted peptide and its combination with irradiation for cancer immunotherapy. *Pharmacol Res.* 2022; 182: 106343.
  22. Zhu X, Wang X, Li B, Zhang Y, Chen Y, Zhang W, et al. A three-in-one assembled nanoparticle containing peptide-radio-sensitizer conjugate and TLR7/8 agonist can initiate the cancer-immunity cycle to trigger antitumor immune response. *Small.* 2022; 18: e2107001.
  23. Zhu X, Li C, Lu Y, Liu Y, Wan D, Zhu D, et al. Tumor microenvironment-activated therapeutic peptide-conjugated prodrug nanoparticles for enhanced tumor penetration and local T cell activation in the tumor microenvironment. *Acta Biomater.* 2021; 119: 337-48.
  24. Hu K, Wu W, Xie L, Geng H, Zhang Y, Hanyu M, et al. Whole-body pet tracking of a D-dodecapeptide and its radiotheranostic potential for PD-L1 overexpressing tumors. *Acta Pharm Sin B.* 2022; 12: 1363-76.
  25. Wang X, Zhou M, Chen B, Liu H, Fang J, Xiang S, et al. Preclinical and exploratory human studies of novel <sup>68</sup>Ga-labeled D-peptide antagonist for PET imaging of TIGIT expression in cancers. *Eur J Nucl Med Mol Imaging.* 2022; 49: 2584-94.
  26. Chen G, Kang W, Li W, Chen S, Gao Y. Oral delivery of protein and peptide drugs: From non-specific formulation approaches to intestinal cell targeting strategies. *Theranostics.* 2022; 12: 1419-39.
- Cite this article as:** Sui X, Niu X, Zhou X, Gao Y. Peptide drugs: a new direction in cancer immunotherapy. *Cancer Biol Med.* 2024; 21: 198-203. doi: 10.20892/j.issn.2095-3941.2023.0297