



REVIEW

Bronchoalveolar lavage fluid assessment facilitates precision medicine for lung cancer

Hantao Zhang^{1,2*}, Dan Deng^{2*}, Shujun Li², Jing Ren³, Wei Huang², Dan Liu³, Weiya Wang¹

¹Department of Pathology, West China Hospital, Sichuan University, Chengdu 610000, China; ²West China Biobank, West China Hospital, Sichuan University, Chengdu 610000, China; ³Department of Respiratory and Critical Care Medicine, Clinical Research Center for Respiratory Disease, West China Hospital, Sichuan University, Chengdu 610000, China

ABSTRACT

Lung cancer is the most common and fatal malignant disease worldwide and has the highest mortality rate among tumor-related causes of death. Early diagnosis and precision medicine can significantly improve the survival rate and prognosis of lung cancer patients. At present, the clinical diagnosis of lung cancer is challenging due to a lack of effective non-invasive detection methods and biomarkers, and treatment is primarily hindered by drug resistance and high tumor heterogeneity. Liquid biopsy is a method for detecting circulating biomarkers in the blood and other body fluids containing genetic information from primary tumor tissues. Bronchoalveolar lavage fluid (BALF) is a potential liquid biopsy medium that is rich in a variety of bioactive substances and cell components. BALF contains information on the key characteristics of tumors, including the tumor subtype, gene mutation type, and tumor environment, thus BALF may be used as a diagnostic supplement to lung biopsy. In this review, the current research on BALF in the diagnosis, treatment, and prognosis of lung cancer is summarized. The advantages and disadvantages of different components of BALF, including cells, cell-free DNA, extracellular vesicles, and microRNA are introduced. In particular, the great potential of extracellular vesicles in precision diagnosis and detection of drug-resistant for lung cancer is highlighted. In addition, the performance of liquid biopsies with different body fluid sources in lung cancer detection are compared to facilitate more selective studies involving BALF, thereby promoting the application of BALF for precision medicine in lung cancer patients in the future.

KEYWORDS

Lung cancer; bronchoalveolar lavage fluid; extracellular vesicles; liquid biopsy; precision medicine

Introduction

Lung cancer is one of the most common malignancies and accounts for approximately 21% of cancer-related deaths worldwide¹. The incidence of lung cancer ranks first and second among all cancers in China among adult males and females, respectively, representing substantial medical and economic burdens². Lung cancer is also considered one of the most fatal malignancies, with a 5-year survival rate of only 23%, which is largely attributable to late diagnosis. It has

been reported that when lung cancer is detected in a localized stage, the 5-year survival rate can reach 60%, while the 5-year survival rate drops to 6% when detected in a distant stage¹. Therefore, an effective screening method to prolong the life of lung cancer patients by early diagnosis has always been highly sought. Currently, low-dose computerized tomography (LDCT) screening is prioritized for the early detection of lung cancer, and its implementation is associated with reduced mortality due to improvements in nodule characterization³. However, LDCT alone lacks specificity for distinguishing between benign and malignant nodules, with a false-positive rate of up to 24%, which may lead to overtreatment and cause unnecessary anxiety^{3,4}. Thus, it is necessary to establish improved detection techniques with high sensitivity and specificity in the early stage of lung cancer.

Lung cancer is predominantly categorized into two subtypes [non-small cell lung cancer (NSCLC) and small cell lung cancer (SCLC)]. Over the past two decades, molecular targeted therapies and immunotherapies have achieved encouraging results in treating NSCLC, with unprecedented improvement

*These authors contributed equally to this work.

Correspondence to: Weiya Wang

E-mail: wangweiya@scu.edu.cn

ORCID ID: <https://orcid.org/0000-0003-3927-6218>

Received September 27, 2023; accepted November 24, 2023;

published online December 29, 2023.

Available at www.cancerbiomed.org

©2024 The Authors. Creative Commons Attribution-NonCommercial

4.0 International License

in patient survival. Unfortunately, not all patients respond to targeted therapies, and for those who do, resistance typically emerges during subsequent treatment⁵. The main obstacle for immunotherapy is how to select appropriate responders⁶. SCLC is characterized by a high proliferative capacity, genomic instability, and early metastasis⁷. Precision medicine for SCLC involves multiple dimensions, such as genetics, epigenetics, proteomics, and the tumor microenvironment. Drug resistance and high tumor heterogeneity are the primary challenges to effective and accurate treatment⁸. Therefore, to improve the survival rate and prognosis of lung cancer patients, both early diagnosis and the identification of the key tumor characteristics, including the tumor subtype, gene mutation type, and tumor microenvironment, are important. Dynamic detection of relevant indicators during treatment is conducive to monitoring efficacy, optimizing treatment strategies, and improving drug resistance. Tissue biopsy remains the preferred method for detecting these indicators, but tissue biopsy is impeded by the scarcity of tissue samples, sampling frequency, and incomplete representation of the entire tumor bulk⁹. It is important to note that liquid biopsy may be able to compensate for these defects.

Liquid biopsy, a method for detecting circulating biomarkers in body fluids with genetic information from primary tumor tissues, is a minimally invasive, repeatable, and comprehensive option that can provide personalized and dynamic diagnostic and prognostic information for patient care¹⁰. Circulating tumor cells (CTCs), cell-free DNA (cfDNA), extracellular vesicles (EVs), cfRNA, and microRNA (miRNA) are the most commonly used biologic materials in liquid biopsy. Among these biologic materials, several assays for detecting CTCs and cfDNA have been approved by the FDA⁹. The clinical use of peripheral blood-derived cfDNA as a supplementary specimen for assessing epidermal growth factor receptor (EGFR) mutation status and detecting the T790M mutation when tumor tissue is not readily available has been approved for NSCLC¹¹. Currently, the concept of liquid biopsy has been expanded from blood to bronchoalveolar lavage fluid (BALF), pleural effusions, and other body fluids.

BALF is obtained from the lung through flexible bronchoscopy. BALF is an important bioactive material that includes cellular and non-cellular components which provides information on biological processes occurring in the alveoli, such as inflammation and infection. Over the past few years, BALF has mainly been used to diagnose interstitial and infectious lung diseases¹². At the same time, there have been corollary studies

focusing on the role of BALF in lung cancer. A review has summarized studies of BALF for lung cancer in analyzing the tumor immune microenvironment and assessing immune-related adverse events of immunotherapy¹³. Another review described the cell metabolism changes in BALF at various stages of lung cancer¹⁴. With the advent of liquid biopsy technology, liquid biopsy using BALF as the medium is expected to become a diagnostic supplement to lung biopsy, but a comprehensive summary on BALF in liquid biopsy for lung cancer is lacking. To fill this gap, we have provided a comprehensive overview of the current research findings based on BALF in lung cancer according to the different liquid components and addressed the advantages and disadvantages of BALF compared to other liquid biopsy techniques using different body fluids. Finally, we discussed the challenges and highlight future directions for implementing these techniques in lung cancer clinical trials.

Overview of the development of BALF

Bronchoalveolar lavage technology has a long history of development worldwide. Initially, bronchoalveolar lavage was used for clinical testing *via* a rigid bronchoscope, followed by major breakthroughs in the invention of the flexible bronchoscope, which is safer and better tolerated by patients. In 1974, bronchoalveolar lavage was first described as a diagnostic method¹⁵. Since then this technique has been progressively utilized to diagnose abnormal lung diseases, including interstitial, infectious, and even malignant diseases¹². There are three steps involved in BALF collection. First, the flexible bronchoscope is “wedged” into the target bronchial segment after local anesthesia. Second, 100–200 mL (\leq 300 mL) of normal saline is injected into the alveolar cavity through the bronchoscope, followed by negative pressure suction to recover perfusion. The recollected fluid is termed BALF, and different centers may have personalized modifications for the collection protocol. Third, cells and supernatant are obtained from BALF through pretreatment and centrifugation, which can then be used for further detection (**Figure 1**).

Under normal circumstances, “healthy” BALF is sterile; extra cells, bacteria, or acellular material are aberrant and can be utilized as a diagnostic tool for various lung disorders¹². Cellular components of “healthy” BALF include $<$ 0.5% eosinophils, $<$ 5% neutrophils, $<$ 20% lymphocytes, and $>$ 80% macrophages¹³; however, this pattern is not static. For




Figure 1 Overall workflow of BALF. Normal saline is injected into the alveolar cavity through the flexible bronchoscope, followed by negative pressure suction to recover the perfusate. The recollected fluid is BALF. After pretreatment and centrifugation, BALF is divided into two parts (cells and supernatant). Cells are used for cell smears, liquid-based cytology, and flow cytometry analysis. The supernatant can be subjected to ELISA, gene sequencing, or multi-omics analysis, such as proteomics, metabolomics, and microbiome analysis, allowing in-depth analysis of proteins, nucleic acids, metabolites, and the microbiome. (Created with BioRender.com).

example, compared to non-smokers, smokers have a significant increase in the absolute cell count, leading to a three-fold higher number of macrophages. Furthermore, smokers have a lower percentage of lymphocytes and the ratio of different subtypes of lymphocytes is also changed.

Application of BALF in the diagnosis and prognosis of lung cancer

The application of BALF in lung cancer diagnosis and prognosis is still in the research stage. The BALF supernatant obtained by centrifugation is rich in various non-cellular components, including proteins, nucleic acids, and EVs (Figure 2). Because it is possible to distinguish cellular and non-cellular content

from proximal airways and lung alveoli, BALF is the ideal “lung liquid biopsy” material for the acquisition of biomarkers pertaining to lung cancer^{1,3}.

Cytologic examination

Cancer cells

Cell pellets can be obtained from BALF through centrifugation, allowing for cytologic examination. Conventional methods, such as cell smears, have poor diagnostic performance due to the limitations of smear thickness, overlap, and immune cells. With the advent of liquid-based cytology, the diagnostic rate of lung cancer has improved, but the sensitivity of cytologic examinations based on BALF fluctuates widely




Figure 2 Application of BALF in lung cancer. BALF contains many different components. The cell components can be divided into cancer and immune cells. The presence of tumor cells reflects the possibility of malignancy. Different proportions of immune cells reflect the local immune microenvironment to some extent. The studies of protein focus on serum tumor markers and some soluble factors. Molecular analysis includes microsatellite alterations, mutations in oncogenes, or tumor suppressors and gene methylation. Multiple miRNAs are specifically expressed in lung cancer. DNA, miRNA, mRNA, and proteins in BALF-EVs also have the potential to serve as biomarkers for lung cancer. Some metabolites or microbiota are also differentially expressed in BALF of lung cancer patients. (Created with BioRender.com).

in different studies, ranging from 28%–87%^{14,16–20}. The possible reason for such a large fluctuation is that various clinical factors, such as pneumonia, viral infections, and inflammatory changes, lead to severely dysplastic cells, which are difficult to distinguish from malignant cells. Some artificial factors also affect the accuracy of cytological examinations. The effects of different bronchoscopic specimens combined with liquid-based cytology on the early diagnosis of lung cancer have been compared. The sensitivity and specificity of BALF are comparable to bronchoscopic wash cultures and transbronchial lung biopsies; the performance of endobronchial ultrasound-guided transbronchial needle aspiration has the best sensitivity and specificity¹⁷. Other studies have

shown that the sensitivity of liquid-based cytology of BALF is higher in visible lesions, but can be significantly improved by combining BALF with bronchial brushing when the lesion is not visible^{16,19}. Another study evaluated the performance of peripheral blood-derived CTCs and BALF-derived tumor cells, and found that the sensitivity and specificity of the two samples were similar in distinguishing benign lesions from lung cancer²⁰.

Immune cells and the tumor microenvironment

BALF contains a variety of immune cells that can be analyzed by flow cytometry. Most studies have demonstrated that compared to healthy subjects, lung cancer patients exhibit increased

levels of neutrophils and T cells in BALF^{13,21}. Further analysis of T-cell subtypes in BALF revealed an increase in CD8⁺ T cells, while the CD4:CD8 ratio was decreased¹³. In addition, multiple studies have indicated that BALF from lung cancer patients has an immunosuppressive microenvironment that is characterized by an elevated proportion of Treg cells and CTLA-4⁺ Treg cells, as well as polarization of macrophages toward the M2 population^{13,21-23}. Compared to NSCLC patients, the proportion of IL10⁺CD206⁺CD14⁺ M2-like macrophages are elevated in BALF from SCLC patients, positively correlated with tumor stage, and negatively correlated with survival time²⁴.

With the development of immunotherapy, immune checkpoint inhibitors (ICIs) have been used to treat various advanced tumors, including NSCLC. PD-1-PD-L1 pathway blockers have been approved as first-line therapy for advanced NSCLC, with approximately one-half of patients achieving a response to ICIs and experiencing longer survival rates, ranging from 20%–30%¹³. Therefore, it is essential to analyze the tumor microenvironment to identify which patients will benefit from immunotherapy. The tumor microenvironment includes tumor, stromal, and immune cells, as well as related cytokines. Tumor biopsies available for tumor microenvironment analysis are very limited, so BALF may serve as a superior source of information. Compared to peripheral blood, BALF is more proximal or “vicinal” to the tumor cells in the lung, so BALF can provide more direct information on the local immune status in lung tissue¹³. The detection of PD-L1 expression by cells derived from BALF is more consistent with paired tissue biopsies, which in turn is better than pleural effusion cells²⁵. A tumor microenvironment analysis based on BALF showed that the expression of CXCL9 in BALF of NSCLC patients who responded to nivolumab is higher than non-responders²⁶. BALF can also be used to evaluate immune-related adverse events of immunotherapy¹³, which manifests in the lung as pneumonia, and is referred to as checkpoint inhibitor pneumonitis²⁷. The number of central memory T cells is increased in BALF of patients with checkpoint inhibitor pneumonitis and the expression of CTLA-4 and PD-1 is decreased in BALF Tregs compared to patients without this complication²⁸. Therefore, dysregulation of immune cell subsets in BALF may serve as a potential target for the treatment of immune-related adverse events.

There are significant differences in the immune cell profile between BALF and peripheral blood. The same cells account for a high proportion in BALF and may be very low in peripheral blood^{21,23,29}. The consistency between the immune cell

profile based on BALF and the tumor-infiltrating lymphocytes detected in tumor resection tissues also differs. A study has shown that a higher proportion of Treg cells is present in the BALF of adenocarcinoma patients²², which is consistent with the results directly obtained from tumor tissues³⁰. However, another study comparing the immune cell subtypes of lymphocytes, dendritic cells, myeloid suppressor cells, and macrophages in BALF and tumor tissues showed that except for myeloid-derived suppressor cells, the proportion of other cells in the two samples had no significant correlation³¹. The above disparities may be because BALF reflects the overall immune environment of the lung, and various contributing factors, including smoking and chronic obstructive pulmonary disease (COPD), ultimately affect the proportion of immune cells, while tumors principally cause a local immune response.

Tumor-related proteins

Tumor markers

There are some common serum tumor markers that have been used in the clinical diagnosis of lung cancer. Carcinoembryonic antigen (CEA) is a glycoprotein complex expressed on the surface of tumor cells. Compared with a control group, CEA expression in the BALF of lung cancer patients is significantly upregulated³²⁻³⁴. Numerous studies have reported that the level of CEA expression in the BALF of lung cancer patients is higher than serum, so the sensitivity of detecting CEA in BALF is higher^{35,36}. By comparing the expression of CEA in BALF and serum among patients with lung cancer and pneumonia, and healthy individuals, it was found that CEA expression in serum could not discriminate between lung cancer and pneumonia patients, while these levels were significantly higher in BALF from lung cancer patients than BALF from pneumonia patients³⁷. Another study showed that CEA expression in BALF was significantly increased in non-smokers with tumors, while no significant difference was observed in smokers with tumors³⁸. Another study demonstrated non-significant variations in the CEA concentration in BALF from patients with various pathologic types³⁴. Neuron-specific enolase (NSE) is a neuronal form of the glycolytic enzyme, enolase. NSE mainly exists in neural and neuroendocrine tissues, as well as tumors derived from these tissues³⁹. SCLC is classified as a neuroendocrine tumor and serum levels of NSE are elevated in 66%–81% of patients with SCLC⁴⁰. The NSE content in BALF samples from tumor groups has been shown to be significantly higher than BALF from control groups^{32,34,41,42}. However, unlike NSE

in serum, most studies (with the exception of one study³⁴) have indicated no difference in the expression of BALF-derived NSE in SCLC and NSCLC^{32,41,42}. The expression of NSE between BALF and serum was not shown to be significantly different, but simultaneous detection of NSE from two sources could improve the sensitivity⁴². Cytokeratin 19 fragment antigen 21-1 (CYFRA21-1) is a fragment of cytokeratin 19 and serves as the most sensitive serum marker of NSCLC, especially in patients with lung squamous cell carcinoma (LSCC)⁴³; however, there has been ongoing debate regarding the diagnostic value of CYFRA21-1 in BALF. An early study concluded that CHFRA21-1 had poor diagnostic value⁴⁴, while subsequent studies reported improved diagnostic performance^{32,34}. In addition to the tumor markers mentioned above, research has been conducted on other markers, such as squamous cell carcinoma antigen and precursor gastrin-releasing peptide^{33,35}. Based on existing reports, however, the detection of tumor markers in BALF is insufficient for the diagnosis of lung cancer, but combined detection can significantly improve the sensitivity of diagnosis^{32,34}.

Soluble factors

In addition to common tumor markers, BALF also contains several soluble factors that can be detected by ELISA. Among the soluble factors, the vascular endothelial growth factor (VEGF) family has been extensively studied. The content of VEGF-A is notably elevated in the BALF of lung cancer patients and higher than plasma^{45,46}. VEGF-C is another member of the VEGF family, the expression of which in the peripheral blood of lung cancer patients is significantly increased⁴⁷. In the BALF of LSCC patients, VEGF-C expression has been shown to be significantly increased⁴⁸. VEGF-D expression is also upregulated in the BALF of patients with LSCC⁴⁹. Indeed, there are several other soluble factors that warrant consideration. Among these findings, transforming growth factor beta (TGF- β) expression has been shown to be upregulated in the BALF of patients with primary lung cancer⁵⁰. In a subsequent study of NSCLC, researchers also showed that the level of TGF- β in the BALF of cancer patients was higher than healthy controls and positively correlated with the overall survival of patients with stage IIIB NSCLC⁵¹. In the same study high mobility group box 1 in BALF was associated with metastasis and overall survival of patients with stage IIIB NSCLC⁵¹. A series of studies have indicated that the expression of various soluble factors in BALF, such as B-cell-attracting chemokine 1 (BCA-1), hepatocyte growth factor (HGF), interleukin 18 (IL-18), IL-22,

IL-27, and IL-29, are abnormal in NSCLC patients, some of which are related to disease progression and response⁵²⁻⁵⁵. NSCLC patients with higher sRANKL⁵² or BCA-1⁵³ levels in BALF were associated with poorer survival. Another study also demonstrated a significant rise in the level of IL-27 expression in the BALF of early NSCLC patients⁵⁶. Researchers have only demonstrated relevant evidence for the above indicators but have not conducted in-depth studies, which is insufficient for clinical practice. In a recent study involving patients with lung nodules < 8 mm in size that were suspicious for cancer, the BALF was assessed for combination indices (VEGF, TGF- β , and HGF) and compared to the histopathologic findings. The accuracy of detecting malignant and benign nodules was 71.4% and 74.5%, respectively⁵⁷. This finding may be used for preliminary screening of small nodules for which histopathologic specimens could not be obtained.

Proteomics

Proteomics technology has also been applied to BALF research. The relevant clinical studies using this method are summarized in **Table 1**. Matrix-assisted laser desorption/ionization time of flight (MALDI-TOF) mass spectrometry was first reported for detecting BALF in NSCLC patients, suggesting that BALF is a suitable biofluid for screening biomarkers of NSCLC⁵⁸. Subsequently, bidimensional polyacrylamide gel electrophoresis (2D-PAGE) and MALDI-TOF/TOF were utilized to analyze the BALF of NSCLC patients; seven specific proteins were identified (CTSD, ALDO A, FBP1, ERZ, AKR1B10, TKT, and SELENBP1). Further validation *via* western blotting of paired BALF proteins showed that only the AKR1B10 protein was upregulated in the NSCLC group⁵⁹. The proteome of BALF was analyzed by 2D-PAGE and mass spectrometry (MS). In combination with bead-based immunoassays, a panel of proteins (APOA1, CO4A, CRP, GSTP1, and SAMP) was screened out. The sensitivity and specificity of this panel were 95% and 81%, respectively, indicating an effective assay for lung cancer diagnosis. This study also identified two proteins, STMN1 and GSTP1, which can be used to distinguish SCLC from NSCLC with a sensitivity and specificity of 90% and 57%, respectively⁶¹. Proteomics quantification of BALF in patients with primary lung adenocarcinoma was performed by liquid chromatography-mass spectrometry (LC-MS). Thirty-three proteins were identified to be consistently overexpressed in all samples, some of which were shown to be associated with lung cancer progression⁶². Proteomic analysis was conducted for the first time using sequential windowed acquisition of all

Table 1 Proteomic analysis based on BALF for lung cancer

Study subjects	Proteomic analysis	Validation subjects	Validation method	Specific proteins (numbers/name)	Comments	Ref.
NSCLC (<i>n</i> = 6) and control (<i>n</i> = 16)	MALDI-TOF/TOF MS	NA	NA	2/HTN3 and S100A12	HTN3 and S100A12 were significantly upregulated in the BALF of NSCLC patients.	58
COPD (<i>n</i> = 15), NSCLC (<i>n</i> = 15), NSCLC with COPD (<i>n</i> = 15), and control group with neither NSCLC nor COPD (<i>n</i> = 15)	2D-PAGE and MALDI-TOF/TOF MS	Paired BALF proteins	Western blot	1/AKR1B10	Compared with the control group, the expression of AKR1B10 was increased in the NSCLC group, while no difference was detected in the COPD group.	59
Lung adenocarcinoma (<i>n</i> = 4) and benign lung diseases (<i>n</i> = 4)	iTRAQ labeling and LC-MS/MS	Lung adenocarcinoma (<i>n</i> = 18), lung SQCC (<i>n</i> = 9), SCLC (<i>n</i> = 6), and benign lung diseases (<i>n</i> = 6)	ELISA in BALF samples	1/Napsin A	The protein level of Napsin A was significantly increased in BALF of adenocarcinoma.	60
Lung cancer (<i>n</i> = 139) and benign lung diseases (<i>n</i> = 49)	2D-PAGE and MS	Lung cancer (<i>n</i> = 204) and benign lung diseases (<i>n</i> = 48)	Bead-based immunoassays in BALF samples.	5/APOA1, CO4A, CRP, GSTP1, SAMP	The levels of these five proteins led to a lung cancer diagnostic panel with 95% Sn and 81% Sp.	61
Lung adenocarcinoma (<i>n</i> = 8) and non-tumor patients (<i>n</i> = 8)	LC-MS/MS	NA	NA	2/STMN1 and GSTP1	The quantification of these two proteins differentiated NSCLC and SCLC patients with 90% Sn and 57% Sp.	62
Lung adenocarcinoma (<i>n</i> = 12) and non-tumor patients (<i>n</i> = 10)	SWATH DJA MS [†]	NA	NA	33/S100-A8, TYMP, annexin A1, annexin A2, and TG2	These five proteins of the DEPs were associated with lung cancer progression in previous studies and may be potential biomarkers for lung adenocarcinoma.	63
NSCLC (<i>n</i> = 26) and non-tumor patients (<i>n</i> = 16)	Label-free MS	NSCLC (<i>n</i> = 46) and non-tumor patients (<i>n</i> = 26)	ELISA in plasma	44/HPT, CO4A, GTSP1	Three of the DEPs that we suggest as potential biomarkers were consistent with previous studies using BALF samples.	64
Lung adenocarcinoma (<i>n</i> = 5), lung SQCC (<i>n</i> = 4), SCLC (<i>n</i> = 4), and benign lung diseases (<i>n</i> = 3)	MS-based quantitative N-glycoproteomic	Lung adenocarcinoma (<i>n</i> = 18), lung SQCC (<i>n</i> = 9), SCLC (<i>n</i> = 6), and benign lung diseases (<i>n</i> = 6)	ELISA in BALF	3/Cystatin-C, Lipocalin-2, TIMP-1	These three proteins were upregulated in BALF of NSCLC patients, and the results were confirmed in plasma.	65
SCLC (<i>n</i> =5): tumor-bearing lungs and non-tumor lungs	TMT-based quantitative MS	Paired tumor tissues	IHC in SCLC tissues	4/DSTN, RNPEP, CNDP2, and CA1	The expression of periostin was increased in BALF from all three tumor types, which is in agreement with the observation of elevated serum levels of this protein in a previous study.	66

Table 1 Continued

Study subjects	Proteomic analysis	Validation subjects	Validation method	Specific proteins (numbers/name)	Comments	Ref.
		NA	SCLC-CellMiner resource [†]	2/CNDP2 and RNPEP	1. CNDP2 and RNPEP appeared to be potential indicators for ASCL1-high and NEUROD1-high subtypes of SCLC. 2. CNDP2 was found to be positively correlated with responses to etoposide, carboplatin, and irinotecan.	

[†]Indicates the analysis of BALF proteins by LC-MS/MS combining a simple sample pre-treatment and a DIA approach.

[‡]Indicates a bioinformatic resource for SCLC cell line genomics and pharmacology based on genomic signatures (<https://discover.nci.nih.gov/rsconnect/ScicellMinerCDB/>).

theoretical fragment ion mass spectra (SWATH) MS on the BALF of patients with lung adenocarcinoma. A total of 44 specific and highly expressed proteins were screened out, among which HPT, CO4A, and GTSP1 were consistent with previous reports⁶³. Label-free MS was performed to analyze the proteome of BALF in NSCLC patients. Compared to patients without tumors, significant changes in protein abundance and molecular function were found in NSCLC patients. The protein expression profile was also different between adenocarcinoma and LSCC patients. According to proteomic analysis, four proteins (cystatin C, TIMP-1, lipocalin-2, and HSP70/HSPA1A) were significantly elevated in NSCLC patients, and the expression of the first three proteins was further verified in plasma samples⁶⁴. An in-depth proteomics method for BALF analysis was further developed that effectively detected both high- and low-abundance proteins, enabling the identification of the largest number of BALF proteins⁶⁷. The control samples for the above studies were all from patients without tumors. In one study, however, BALF was collected from the tumor-bearing and healthy lungs of the same patient. According to tandem mass tag (TMT)-based quantitative MS analysis, four proteins (DSTN, RNPEP, CNDP2, and CA1) were shown to be highly expressed in tumor-bearing lungs, and these proteins were validated by tissue immunohistochemistry⁶⁶. According to the expression of prominent transcription factors, SCLC can be categorized into four subtypes. ASCL1 and NEUROD1 are associated with neuroendocrine-high (NE-high) markers. All five patients in the study were NE-high, and further bioinformatics analysis revealed that CNDP2 and RNPEP are potential markers for ASCL1-high and NEUROD1-high subtypes of SCLC⁶⁶.

Given that the majority of the secreted proteins in BALF are glycoproteins, N-glycoproteins were detected through the use of isobaric tags for relative and absolute quantitation (iTRAQ) labeling and LC-MS. Eight glycoproteins were highly expressed in the BALF of lung adenocarcinoma. Among these glycoproteins, Napsin A has been reported to be highly expressed in lung adenocarcinoma tissues⁶⁸. Therefore, the expression of Napsin A was further validated in BALF samples by ELISA in another independent cohort, and a significant increase in Napsin A was found in the BALF of lung adenocarcinoma patients⁶⁰. The MS-based quantitative N-glycoproteomic technique was also used to analyze BALF samples, revealing periostin in all samples from lung cancer patients. Validation of the screened periostin with ELISA in another 39 BALF samples also yielded the same conclusion⁶⁵.

The above studies collectively indicate that proteomics-based analysis of BALF can be used to screen potential diagnostic markers for lung cancer, especially when combined with other experimental methods for validation to further narrow the range of target proteins. However, owing to the diversity of proteomic techniques, validation populations and sample types, the different selection of control groups, and the limited sample size of the studies, the clinical translation of proteomics technology on BALF analysis has not been established.

Molecular analysis

DNA

Common genetic alterations of tumor-related molecules include microsatellite alterations, mutations in oncogenes or tumor suppressors, and gene hypermethylation. Initially, researchers collected BALF cells for molecular analysis. With the advances in liquid biopsy technology, research on BALF supernatants has gradually increased. The clinical studies pertaining to BALF genetic analysis-based lung cancer detection are listed in **Table 2**.

In 1999 researchers used the molecular analysis results of tumor tissues as a reference to detect the genetic alterations in the corresponding BALF cell-derived DNA and found that the detection rates of microsatellite alterations, *P53* and *KRAS* mutations, and *P16* methylation were 13.6% (3/22), 39.3% (11/28), 33.3% (4/12), and 63.2% (12/19), respectively⁷⁷. The detection of *EGFR* mutations was better than detection of *P53* or *KRAS* mutations^{71,78,79}. The most common mutation in NSCLC involves the *EGFR* gene. It has been shown that BALF has higher sensitivity than pleural fluids for detecting *EGFR* mutations in lung adenocarcinoma patients based on next-generation sequencing (NGS)⁶⁹. The sensitivity of BALF-derived cfDNA for detecting *EGFR* mutations was 91.7% in lung adenocarcinoma patients, which was better than paired bronchial washing samples (18.2%)⁷⁰. In 13 lung adenocarcinoma patients with confirmed *EGFR* mutations, BALF identified 2 additional mutations compared with tissue biopsy and had higher sensitivity (92.3%) than peripheral blood samples (38.5%) for detecting *EGFR* mutations⁷¹. These results suggest that *EGFR* mutation detection in BALF may become a convenient and sensitive method for the diagnosis of NSCLC.

In addition to determining the status of *EGFR* mutations before treatment, monitoring changes in drug resistance gene mutations during treatment is also critical. After *EGFR*-tyrosine kinase inhibitor (TKI) treatment, most patients with

EGFR mutations will develop secondary mutations 9–14 months later, among which the T790M mutation is most common^{78–80}. At present studies involving detection of the T790M mutation in cfDNA mainly focus on serum/plasma^{81,82}, but there are several studies showing that the T790M mutation can also be identified in BALF. A patient with lung adenocarcinoma developed a T790M mutation 3 months after initiation of gefitinib. The detection results of BALF-derived cfDNA were consistent with tumor tissues, but the mutation could not be detected in bronchial washing samples⁷⁰. Two additional mutations were detected in BALF samples compared to the corresponding lung adenocarcinoma tissue samples, one of which was the T790M mutation⁷⁹.

In addition to *EGFR* mutations, *TP53*, *KRAS*, and *BRAF* were also found in BALF cfDNA from lung cancer patients, and the genomic profile had a sensitivity of 67.7%⁷². Analysis of tumor-associated gene mutation profiles using a tumor-naïve or -informed approach showed that BALF cfDNA exhibited higher sensitivity than plasma cfDNA for detecting lung cancer-derived mutations in stage I-II disease. A diagnostic classifier for lung cancer-derived mutations was then developed with a sensitivity of 64.7%, which was superior to BALF cytology (11.8%)⁷³. The above results indicate that BALF, as a liquid biopsy medium, has good sensitivity in the detection of gene mutations from lung cancer and may be used as a diagnostic supplement for tissue biopsy.

Epigenetic studies have revealed a close association between DNA methylation and the occurrence and progression of lung cancer. Aberrant methylation of gene promoters has been used as a biomarker for diagnosing lung cancer⁸³. Methylation typically occurs in the early stage of lung cancer and has been extensively studied in liquid biopsy⁸⁴. Methylation of 8 gene promoters (*CDH1*, *APC*, *MGMT*, *RASSF1A*, *GSTP1*, *P16*, *RAR-β2*, and *ARF*) in BALF cell-derived DNA from 31 patients with primary lung cancer was examined and abnormal methylation of these genes was shown in 67.7% of the samples⁷⁴. Aberrant methylation of *SHOX2* and *RASSF1A* genes has also been widely reported in various studies on lung cancer. The clinical specimens include tumor tissues, plasma, pleural effusions, and bronchial aspirates^{83,85–87}. In recent years studies based on BALF samples have also begun to increase. The combined detection of *SHOX2* and *RASSF1A* methylation improves the detection rate of lung cancer compared with individual assessment or cytologic examination of BALF. Interestingly, *SHOX2* methylation has a histologic subtype preference, with optimal results in SCLC, followed by LSCC and adenocarcinoma⁷⁵.

This finding was consistent with previous studies carried out on plasma and bronchial aspirates^{83,85}. Another study yielded similar results. At all stages of lung cancer, the sensitivity of the BALF-derived *SHOX2* and *RASSF1A* methylation assay was higher than cytology and serum CEA, especially in stage I, with a sensitivity of 85.7%. Therefore, BALF-based methylation analysis of *SHOX2* and *RASSF1A* can be used as an effective complement to cytological examination, especially in the early stage of lung cancer⁸⁸. The results of this study also showed that the *SHOX2* and *RASSF1A* methylation assay has a preference for histological subtypes, and the conclusion is consistent with previous findings that the sensitivity is highest in SCLC⁷⁶. Another study investigated the correlation between gene methylation in BALF and patient prognosis and found that the methylation of *RASSF1A* was associated with shorter progression-free survival in NSCLC patients treated with pemetrexed⁸⁸. The above methylation detection was based on BALF cell-derived DNA. Another study performed methylation profiling of cfDNA from BALF. The findings indicated that the methylation analysis of BALF-cfDNA possessed high sensitivity in distinguishing small malignant tumors (≤ 2 cm) from benign nodules⁷².

In summary, BALF-based genetic analysis exhibits good sensitivity with specific methods for the identification of lung cancer. For precision treatment, BALF also has an auxiliary role in the detection of secondary drug-resistant mutations.

RNA

MicroRNAs, small non-coding RNA composed of 20–24 nucleotides, are an important part of liquid biopsy. In fact, miRNAs have been widely reported in blood and other body fluids but are less studied in BALF. In one study researchers analyzed the expression of a specific miRNA panel (miR-21, miR-143, miR-155, miR-210, and miR-372) in BALF from NSCLC patients and found that the diagnostic sensitivity was 85.7%, which was higher than sputum (67.8%)⁸⁹. In another study, miR-205-5p and miR-944 were used as a prediction model to distinguish LSCC and adenocarcinoma. The accuracy of miR-205-5p and miR-944 in BALF samples was 95.7%, which approximated frozen tissue (96.6%) and paraffin-embedded formalin-fixed tissue (96.4%) and better than cytology (68.1%)⁹⁰. The above results indicate that this prediction model may have clinical value in differentiating LSCC from adenocarcinoma in tumor tissues and BALF liquid biopsies. Specific miRNAs (miRNA-34a, miRNA-141, miRNA-143, and miRNA-217) that regulate tumor suppressor genes (*RAR β*

and *FHIT*) were subsequently detected in the BALF of patients with LSCC, but the results showed that the expression of these miRNAs may not have diagnostic value for LSCC⁹¹. Based on the above findings, the detection of BALF-derived miRNAs may be a potential means for lung cancer diagnosis in some cases; however, few studies have explored this possibility. Among the studies that have studied BALF-derived miRNAs for lung cancer diagnosis, the miRNAs screened have been different and may require a larger sample size, further screening, and verification.

EVs

EVs are vesicles with lipid bilayer membrane structures that are released by cells into the extracellular matrix and can be divided into exosomes, microvesicles and apoptotic bodies according to the origin and size. A variety of cells secrete EVs into BALF, including bronchial epithelial cells, alveolar macrophages, and endothelial cells. EVs are rich in various components, including DNA, RNA, and proteins. Tumor cells also secrete a large number of EVs. The cargos in EVs reflect information from parental tumor cells, so EVs can be used as biomarkers to assist in the diagnosis of some tumors⁹². To avoid interference from viruses, proteins, or other aggregates, high-purity EVs are a prerequisite for further analysis. Common methods for isolating EVs include differential centrifugation, density gradient centrifugation, size-exclusive chromatography, and commercial kits. BALF contains many soluble proteins, and the methods for EV isolation from BALF are still in the early stage of development⁹³. The current studies of lung cancer based on BALF-EVs are shown in **Table 3**.

BALF-derived EVs are a reliable source for identifying gene mutations. Researchers compared BALF-EV DNA and tissue DNA based on targeted NGS and found that although the yield and coverage depth of BALF-EV DNA were not as good as tissue DNA, the mutation concordance between the two samples could reach 81%¹⁰². Compared with tissue biopsy, *EGFR* genotyping based on BALF-EVs may be more sensitive. BALF-EV DNA was able to detect not only all of the *EGFR* mutations detected in tissue samples but also six additional cases⁹⁵. BALF-EV DNA can also be used to detect the T790M mutation in patients with secondary resistance to *EGFR*-TKIs, even more efficiently and conveniently than tissue re-biopsy⁹⁴. In addition, the sensitivity of *EGFR* genotyping based on BALF-EVs was increased with TMN grading and even reached 100%⁹⁵. This is contrary to the detection results

Table 2 Genetic analysis detection based on BALF for lung cancer

Study population of BALF	Genetic mutations	Source of origin	Methods	Sn/Sp (%)	Comparison (Sn/Sp) (%)	Comments	Ref.
Lung adenocarcinoma (n = 33)	EGFR 33	Cell pellet	Sanger sequencing	9.09/NA	Pleural effusion: 13.3/NA (unpaired)	BALF cell NGS had higher sensitivity than Sanger sequencing, and therefore could be a diagnostic supplement for better characterization of lung cancer; BALF had higher sensitivity than pleural effusion.	69
Lung adenocarcinoma (n = 20)	EGFR 12 vs. WT 8	cfDNA	PNA-PCR and PANAMutyper [†]	91.7/100	Bronchial washing samples: 18.2/100 (paired)	BALF cfDNA was useful in detecting EGFR mutation and identifying T790M mutations; BALF was more useful than bronchial washing samples.	70
Lung adenocarcinoma (n = 26)	EGFR 13 vs. WT 13	Mixture [†]	PCR	92.3/100	Peripheral blood: 38.5/100 (paired)	BALF had much higher diagnostic accuracy than blood for EGFR mutation detection; two additional mutations were identified in BALF compared with tissue biopsy.	71
Malignant (n = 31) and benign (n = 9)	Mutation profiling	cfDNA	NGS	67.7/77.8	Bronchoscopy biopsy: 23.1/100 (unpaired)	BALF cfDNA methylation analysis had better sensitivity in distinguishing small malignant tumors (≤ 2 cm) from benign nodules than mutation analysis; both were better than conventional methods.	72
Malignant (n = 27) and benign (n = 21)	Methylation profiling		Bisulfite sequencing and RT-PCR	81.5/81.0	Bronchial brushing cytology: 9.7/100 (unpaired) Lavage cytology: 3.1/100 (unpaired)		
Lung cancer stage I-II (n = 18)	Mutation profiling	cfDNA	Tumor naïve approach	66.7/NA	Plasma cfDNA 11.1/NA (paired)	BALF cfDNA analysis was more sensitive than plasma cfDNA for detecting lung cancer-derived mutations in stage I-II disease.	73
Lung cancer stage I-II (n = 22)			Tumor-informed analysis	77.3/NA	Plasma cfDNA: 45.5/NA (paired)		
Lung cancer (n = 17) and non-cancer (n = 16)			Genomic classifier	64.7/100	BAL cytology: 11.8/100 (paired)	BALF cfDNA analysis was useful for the diagnosis of lung cancer.	
Lung cancer (n = 31)	Methylation of CDHL, APC, RASSF1A, MGMT, P16, GSTP1, ARF, and RAR- β 2	Cell pellet	RT-PCR and RT-MSP	67.7/NA	NA	Aberrant DNA methylation was detected in the majority of lung cancer patients.	74
Lung cancer (n = 123); benign lung diseases (n = 112); malignancies in other systems (n = 18)	Methylation of SHOX2 and/or RASSF1A	Cell pellet	RT-PCR and Sanger sequencing	SHOX2: 64.2/92.3; RASSF1A: 50.4/96.2; SHOX2 and RASSF1A: 71.5/90.0	BALF cytology: 5.7/99.2	BALF cell methylation analysis (SHOX2 and RASSF1A) improved diagnostic accuracy for lung cancer, and was superior to cytology diagnosis.	75

Table 2 Continued

Study population of BALF	Genetic mutations	Source of origin	Methods	Sn/Sp (%)	Comparison (Sn/Sp) (%)	Comments	Ref.
Lung cancer (n = 284); benign lung diseases (n = 35); malignancies in other systems (n = 3)	Methylation of <i>SHOX2</i> and <i>RASSF1A</i>	Cell pellet	RT-PCR and Sanger sequencing	Total: 81/97.4 Stage I: 85.7/NA	Total Cytology: 68.3/97.4; Serum CEA: 30.6/100 Stage I Cytology: 46.4/NA Serum CEA: 10.7/NA	BALF cell methylation analysis had relatively satisfactory diagnostic accuracy for lung cancer, especially at early stage, and was complementary to cytology diagnosis.	76

Sn, sensitivity; Sp, specificity; NA, not available; NGS, next-generation sequencing; WT, wild type; PCR, polymerase chain reaction; PNA-PCR, peptide nucleic acid-mediated PCR; RT-PCR, real-time PCR; RT-MSP, real-time methylation-specific polymerase chain reaction.

[†]Indicates PANAMutyper™ R *EGFR* kit with PNA clamping-assisted fluorescence melting curve analysis was used for mutation detection and genotyping.

[‡]Indicates non-centrifuged BALF.

of BALF cfDNA, which is more sensitive in the early stage of lung cancer. For liquid biopsy, DNA derived from EVs is better than cfDNA. *EGFR* genotyping of EV-DNA and cfDNA from BALF and plasma samples was conducted for NSCLC diagnosis. The results showed that the sensitivity of EV-DNA was better than cfDNA from both sources, which may be due to the protective effect of the EV bilayer membrane on internal DNA⁹⁴. Furthermore, BALF samples exhibited significantly higher sensitivities for cfDNA and EV DNA than plasma samples⁹⁴. Another study of patients with stage III-IV NSCLC also showed that *EGFR* genotyping based on BALF-EVs had higher sensitivity and specificity than *EGFR* genotyping based on plasma cfDNA⁹⁶. The above results indicate that *EGFR* genotyping with BALF-EVs has satisfactory accuracy and is more convenient than solid tissue biopsy and cfDNA- or plasma-based liquid biopsy. Moreover, *EGFR* genotyping based on BALF-EVs has higher sensitivity in advanced tumors and can detect the T790M mutation. Therefore, *EGFR* genotyping based on BALF-EVs can be utilized as a substitute for tissue biopsy to guide the prompt treatment of *EGFR*-TKIs, especially for patients with an advanced disease stage or from whom sample collection is difficult.

For RNA in EVs, current studies primarily focus on detecting the expression of tumor-related gene mRNA and miRNA. Exosomes are one type of EV with a diameter of 30–100 nm. Compared to plasma, BALF contains fewer exosomes. Moreover, the miRNA content in BALF exosomes showed no difference between patients with and without tumors, while the miRNA content in plasma exosomes of tumor patients was higher than patients without tumors. This finding suggested that more tumor-derived miRNAs are released into plasma than into BALF⁹⁷. MiR 126 and Let-7a concentrations are elevated in the BALF of lung adenocarcinoma patients compared with to control subjects. However, subsequent validation utilizing an independent set of matched adenocarcinoma and normal tissue samples (n = 4) demonstrated that only miR-126 expression is increased in cancer tissues compared to healthy tissues⁹⁸. In a study related to drug resistance, plasma and BALF samples were collected from NSCLC patients before and after icotinib/gefitinib treatment. Exosomes were isolated and mRNA expression of 10 metastasis-related genes was detected. It was found that MET expression in exosomes was associated with lung cancer metastasis, but there was no significant difference between BALF and plasma⁹⁹. These results indicated that BALF may not offer benefits in identifying RNA related to lung cancer.

Table 3 BALF-derived EVs for lung cancer

Study population	Type	Methods of isolation	Cargos	Targets	Sn/Sp (%)	Comparison (Sn/Sp) (%)	Comments	Ref.
NSCLC (<i>n</i> = 23): <i>EGFR</i> mutation 14 vs. WT 9	EVs	Differential centrifugation [†] 1. 1000 g × 15 min 2. 200,000 g × 1 h	DNA	<i>EGFR</i> mutation	100/ 100	BALF cfDNA: 71.4/100. Plasma EV DNA: 55.0/NA (unpaired) Plasma cfDNA: 30.0/NA (unpaired)	1. Liquid biopsy with EVs DNA was more sensitive compared to using cfDNA. 2. Sensitivities of BALF samples were significantly higher compared to plasma samples for both cfDNA and EVs DNA.	94
Patients resistant to <i>EGFR</i> -TKIs (<i>n</i> = 9)				T790M mutation	55.5/NA	BALF cfDNA: 33.3/NA Re-biopsy tissue: 22.2/NA (paired)	BALF-EVs DNA had higher efficiency for detecting T790M mutation compared to tissue re-biopsy.	
Lung cancer (<i>n</i> = 137): <i>EGFR</i> mutation 54 vs. WT 83	EVs	Differential centrifugation 1. 1000 g × 10 min 2. 200,000 g × 1 h	DNA	<i>EGFR</i> mutation	75.9/86.7	T1: 40/NA; T4: 100/NA N0: 63.3/NA; N3: 100/NA M0: 43.5/NA; M1: 100/NA	BALF EVs-based <i>EGFR</i> genotyping had increased sensitivity with escalating TMN stage and even more useful when biopsy was not feasible.	95
NSCLC III-IV (<i>n</i> = 224): <i>EGFR</i> mutation 93 vs. WT 131	EVs	Differential centrifugation 1. 1000 g × 10 min 2. 200,000 g × 1 h	DNA	<i>EGFR</i> mutation	97.8/96.9	Plasma cfDNA: 48.5/86.3 (unpaired)	BALF-derived EVs testing in advanced NSCLC was rapid with high accuracy, serving as an alternative for tissue biopsy for guiding prompt <i>EGFR</i> -TKI treatment.	96
NSCLC (<i>n</i> = 30) and non-tumor patients (<i>n</i> = 75)	Exosomes	Differential centrifugation 1. 500 g × 10 min 2. 17,000 g × 20 min 3. 120,000 g × 90 min	miRNA	miRNA profiles	NA	NA	1. Exosome levels were significantly higher in plasma than in BALF samples in patients with and without cancer. 2. The concentrations of plasma miRNAs was higher in tumor patients when comparing non-tumors but were similar for BALF miRNA.	97
Lung adenocarcinoma (<i>n</i> = 13) and benign lung diseases (<i>n</i> = 15)	Exosomes	Commercial kit (ExoQuick Exosome Precipitation Solution)	miRNA	miR-126/ Let-7a	NA	NA	Exosomal miRNA-126 and Let-7a levels were higher in the BALF of adenocarcinoma than benign BALF.	98
NSCLC patients with <i>EGFR</i> 19del mutation (<i>n</i> = 10)	Exosomes	Differential centrifugation 1. 300 g × 10 min 2. 16,500 g × 20 min 3. 120,000 g × 70 min	mRNA	MET	NA	NA	MET was detected in plasma- and BALF-derived exosomes in metastasis without a significant difference between the two body fluids.	99
NSCLC (<i>n</i> = 22) and healthy volunteers (<i>n</i> = 12)	EVs	Differential centrifugation 1. 500 g × 10 min 2. 2,500 g × 12 min 3. 17,000 g × 160 min	Protein	SOC33	NA	NA	The protein level of SOC33 in the BALF was significantly reduced, independent of histopathologic type and smoking status.	100

Table 3 Continued

Study population	Type	Methods of isolation	Cargos	Targets	Sn/Sp (%)	Comparison (Sn/Sp) (%)	Comments	Ref.
Lung cancer (n = 12) and non-lung cancer (n = 12)	EVs	Differential centrifugation 1. 3000 g × 20 min 2. 12,000 g × 1 h 3. 100,000 g × 2 h	Proteome	DNMT3B	NA	NA	1. Proteome complexity of BALF-EVs correlated with lung cancer stage IV and mortality within 2 years of follow-up. 2. DNMT3B complex was significantly up-regulated in tumor tissue and BALF-EVs.	101

Sn, sensitivity; Sp, specificity; NSCLC, non-small-cell lung cancer; WT, wild type; EVs, extracellular vesicles; NA, not available; TKIs, tyrosine kinase inhibitors.

*Indicates a separation technique separating and removing components other than EVs from BALF in a stepwise manner, and all the centrifugation steps are performed at 4°C.

†Plasma and BALF were collected at the time of primary diagnosis and after the treatment with icotinib/gefitinib within a follow-up period of 3–6 months.

The miRNAs screened in the previous study need to be fully validated before clinical use.

In addition to nucleic acids, EVs contain numerous protein components. Exosome proteins make up 35% of the differentially expressed proteins in the BALF of lung cancer patients¹⁰³. Protein detection in BALF-EVs can be used to further study the therapeutic mechanism underlying lung cancer. Suppressors of cytokine signaling 3 (SOCS3) vesicles from alveolar macrophages are secreted into BALF. The level of SOCS3, as detected by ELISA, is significantly downregulated in NSCLC patients, independent of adenocarcinoma, LSCC, and smoking status¹⁰⁰. In another study, proteomic analysis of BALF-EVs was performed with LC-MS. The results showed that the DNMT3B complex is greatly augmented in tumor tissues and BALF-EVs, and protein markers associated with innate immune cells and fibroblasts in BALF-EVs are associated with a poorer prognosis (4 of 5 patients died during the 2-year follow-up)¹⁰¹.

Others

Metabolomics

In addition to the above common components, some studies have analyzed other components in BALF. Metabolomics is a systems biology method that emerged after genomics and proteomics. The non-targeted qualitative and quantitative analysis of metabolites is expected to provide a new perspective for the study of pathologic processes and biomarkers for diseases¹⁰⁴. There have been numerous studies on metabolomics analysis of serum, plasma, urine, and sputum¹⁰⁵⁻¹⁰⁸ of lung cancer patients, but there are few metabolomics studies involving BALF. Two complementary metabolomics techniques were performed to detect lung cancer-related metabolites in BALF samples (24 cases of lung cancer and 31 cases of non-tumor lung disease) for the first time and identified 42 different metabolites. Glutamic acid and glutamine metabolic pathways are the main changed pathways. Glycerol and phosphate have promising potential as biomarkers that are both sensitive and specific for diagnosing and predicting lung cancer outcomes¹⁰⁹. Subsequently, a combined metabolomics analysis was performed on serum, urine, and BALF from patients with lung cancer. Compared to non-tumor lung disease, researchers identified 16 metabolites that are altered in the BALF of lung cancer patients, 6 of which were detected in serum and urine. Among these metabolites, phosphoric acid was the most sensitive and specific¹¹⁰. The aforementioned

studies indicate that phosphoric acid/phosphate may have a significant involvement in the pathologic process underlying lung cancer.

Microbiome

The pulmonary microbiome has a vital role in respiratory diseases, but the significance in lung cancer has not been established. Some studies have reported that dysbiosis of the microbiota during the disease process mainly includes the following: enrichment of potential pathogenic bacteria; loss of probiotics; and change in microbial composition¹¹¹. The BALF microbiota of patients with lung cancer (20 cases) and benign lung masses (8 cases) were assessed *via* 16S rRNA-based NGS. The findings demonstrated that two phyla (*Firmicutes* and *TM7*) were increased and two genera (*Veillonella* and *Megasphaera*) were enriched in lung cancer¹¹², among which *Veillonella* has also been reported to be enriched in the saliva of patients with lung cancer¹¹³. Another study showed that *Veillonella* is dominant in BALF of NSCLC patients in high-PD-L1 and immunotherapy responder groups¹¹⁴. Our group collected BALF samples for the purpose of investigating the microbiome of the lower respiratory tract (LRT) in individuals with lung cancer. Compared to healthy subjects (30 cases), LRT microbiome richness was decreased in lung cancer patients (91 cases). However, there was no significant distinction between the lung cancer and non-malignant lung disease groups (29 cases). We further established a diagnostic model of lung cancer (including age, smoking history, and 11 types of bacteria), with areas under the curve (AUCs) of 0.882 and 0.796 in the training and validation sets (57 lung cancer and 28 non-malignant lung disease patients), respectively, confirming that biomarkers based on the BALF microbiome may be effective in identifying lung cancer¹¹⁵. Similarly, another study discovered that there was no significant distinction in the main microbial composition in BALF between NSCLC ($n = 32$) and non-tumor patients ($n = 15$)¹¹⁶. The BALF microbiome of patients with stage I NSCLC ($n = 48$) before resection was analyzed. It was shown that the BALF microbiome differs between patients with and without a tumor recurrence after surgery during the 32-month follow-up¹¹⁷. A distinctive microbiome signature was also identified, which was associated with shorter recurrence-free survival¹¹⁷. Despite several differentially expressed genera being identified in the above studies, the particular function of these genera in the pathogenesis of lung cancer needs further investigation due to the dynamics and diversity of the microbial composition and the complexity of patient diets, regions, and

lifestyles. In addition, a study analyzing the microbiomes of the saliva samples, BALF samples, and non-malignant, peritumoral, and tumor tissue samples from 18 NSCLC patients revealed that BALF has a unique microbial diversity compared to the other three types of lung tissues¹¹⁸.

BALF and liquid biopsy

Tissue biopsy is the gold standard for the clinical diagnosis of lung cancer. Percutaneous lung aspiration and transbronchial lung biopsies are invasive diagnostic procedures with restrictions on sampling tumors that are undetectable by bronchoscopy. Liquid biopsy, in contrast, offers many advantages, including simple sampling, low invasiveness, and relative safety. In addition, traditional surgical biopsy tends to be a single-site sample that cannot be used to obtain comprehensive information on the disease. Liquid biopsy contains more representative information and provides a more accurate assessment of the disease because liquid biopsy is used to detect molecules shed/released from different parts of the tumor. Furthermore, liquid biopsy can be used to conduct real-time detection during treatment process and provide objective evidence for clinicians to adjust treatment plans and appraise prognosis.

In the liquid biopsy of lung cancer, distinct characteristics of various body fluids are summarized (**Figure 3**). Conventional serum markers are metabolic byproducts frequently generated and released by tumor cells. However, false-positives and -negatives are common since these markers can also be present in some benign lesions¹¹⁹. With respect to molecular analysis, the components in peripheral blood are relatively complex and cfDNA may come from any part of the body, so cfDNA lacks specificity. Other body fluids also have certain limitations. Sputum contains few cells, making sputum difficult to meet detection requirements. Moreover, it is also rich in pathogens from the upper respiratory tract, which easily causes interference. Pleural effusions, which are usually produced after lung cancer or lung infection, are invasive and challenging to sample early. In addition, the environment of these body fluids is intricate and affects the integrity of DNA sequences to some extent. Compared with these body fluids, BALF has advantages. BALF is in proximity to the tumor site, contains a larger amount of lung cancer cells, and is less likely to be contaminated with cancer cells from other organs, thus has a relatively higher sensitivity and specificity in early diagnosis. Furthermore, the low invasiveness of BALF is appropriate for




Figure 3 Characteristics of different body fluids. (Created with BioRender.com).

advanced patients who are difficult to sample and holds the possibility for serial sampling during treatment.

Conclusions and future directions

Precision medicine involves accurate diagnosis and classification of disease, accurate administration of drugs, accurate evaluation of curative effects, and accurate prediction of prognosis. Precision medicine is not only a public requirement, but also a clinical necessity for advancement. At present, BALF has been utilized in the accurate diagnosis of several diseases. For example, a lymphocytic-predominant BALF is sufficient for diagnosing lung sarcoidosis or upper respiratory infection with matching medical symptoms and imaging findings¹²⁰. In patients with intense alveolar phacoemulsification on chest imaging, the presence of significant eosinophils in BALF accurately predicts eosinophilic

pneumonia¹²¹. For pulmonary infectious diseases, assessment of BALF can be used to effectively identify etiologic pathogens, including bacteria, fungi, parasites, and viruses, to provide focused antimicrobial therapy, reduce the use of unnecessary antibiotics, and slow the rise of antimicrobial drug resistance. With respect to pulmonary alveolar proteinosis (PAP), researchers showed that serum levels of α - and β -defensins cannot differentiate PAP from COPD and sarcoidosis patients, while the BALF of patients with PAP has extremely high levels of these defensins¹²². For critically ill patients with COVID-19 pneumonia, analysis of cellular composition in BALF as biomarkers is reliable for disease management and prediction¹²³.

The high heterogeneity of lung cancer determines the necessity for precision medicine. *EGFR* genotyping based on BALF has a satisfactory detection rate, which is expected to achieve accurate diagnosis of lung cancer^{69-71,94-96}. The prediction

model established by proteomics or microbiomics exhibits good sensitivity and specificity and may serve as an effective indicator for the precise prediction of lung cancer^{61,115}. Some special proteins or miRNAs have been identified from BALF that can be used to distinguish SCLC from NSCLC or LSCC from adenocarcinoma, which may make precise typing of lung cancer possible^{61,90}. The analysis of BALF components is also helpful to determine whether a patient is a candidate for immunotherapy of NSCLC with nivolumab²⁶. Additionally, some soluble factors⁵¹⁻⁵³, methylation of genes⁸⁸, and specific cell subsets²⁴ from BALF may also have been shown to be associated with the prognosis of lung cancer. Thus, the assessment of BALF exhibits promising potential in advancing precision medicine for lung cancer.

Although ongoing research into the clinical utility of the various components of BALF is still evolving and is far from being as comprehensive as similar studies conducted on blood, there is no doubt that BALF has great potential for lung cancer detection, especially using the EV DNA in BALF. Given the protective nature of the EV bilayer membrane, EV DNA exhibits heightened stability and is less susceptible to degradation by external factors compared with cfDNA⁹⁶. In the detection of oncogene mutation, it is noteworthy that BALF-EV DNA exhibits superior sensitivity and specificity compared to both BALF cfDNA and plasma-derived EV DNA. Remarkably, the mutation detection rate of BALF-EV DNA even exceeds that of tissue biopsy, indicating that gene mutation detection based on BALF-EV holds significant promise as the optimal choice for lung cancer prediction. In addition, the sensitivity of BALF-EV DNA detection escalates with the TMN stage, and it is also a valuable tool for detecting the T790M mutation in patients with acquired drug resistance. These features enhance the advantages of BALF, especially in advanced lung cancer patients or in patients in whom sample collection is difficult, and can replace re-biopsy technique in guiding prompt treatment. It is worth noting that two clinical trials (NCT05469022 and NCT03228277) based on BALF-EVs in the context of lung cancer have advanced to phase II, further underlining the reliability and efficacy of BALF-EVs as a potential detection tool.

Nevertheless, the translation of BALF-EV detection into clinical applications faces several substantial challenges. Interference from viruses, proteins, and other aggregates, coupled with the inherently viscous nature of BALF, limits the availability of valuable specimens. In addition, the relatively low concentration of EVs in BALF along with the

presence of a myriad of EVs from various cellular sources may often result in limited acquisition of the desired tumor-related EVs. Consequently, there is an urgent need to develop a standardized, high-yield and effective protocol for BALF-EV isolation. Such efforts are crucial to ensure the comparability and reproducibility of the research findings across different laboratories^{124,125} as well as to develop specialized and commercialized kits for BALF EV.

Currently, the assessment of BALF from lung cancer patients is evolving in various new directions, especially involving the immune response to lung cancer and immune checkpoint inhibitors for precision treatment. Elucidating the clinical utility of BALF as a complementary liquid biopsy for the examination of tumor-related markers will be of paramount importance in advancing lung cancer diagnostics and personalized treatment strategies. Additionally, there is a need to promptly commence large-scale, multicenter trials in pulmonology units with the involvement of oncologists to further advance the research and application of BALF in clinical practice for lung cancer.

Grant support

This work was supported by grants from the National Natural Science Foundation of China (Grant No. 82173182), the Sichuan Science and Technology Program (Grant No. 2021YJ0117 to Weiya Wang; Grant No. 2023NSFSC1939 to Dan Liu), and the 1·3·5 project for Disciplines of Excellence—Clinical Research Incubation Project, West China Hospital, Sichuan University (Grant Nos. 2019HXFH034 and ZYJC21074).

Conflict of interest statement

No potential conflicts of interest are disclosed.

Author contributions

Conceived and designed the analysis: Weiya Wang, Dan Liu, Wei Huang.

Collected the data: Hantao Zhang, Dan Deng, Shujun Li, Jing Ren.

Contributed data or analysis tools: Hantao Zhang, Weiya Wang.

Performed the analysis: Hantao Zhang, Wei Huang.

Wrote the paper: Hantao Zhang, Dan Deng.

References

- Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin.* 2023; 73: 17-48.
- Maomao C, He L, Dianqin S, Siyi H, Xinxin Y, Fan Y, et al. Current cancer burden in China: epidemiology, etiology, and prevention. *Cancer Biol Med.* 2022; 19: 1121-38.
- Oudkerk M, Liu S, Heuvelmans MA, Walter JE, Field JK. Lung cancer LDCT screening and mortality reduction - evidence, pitfalls and future perspectives. *Nat Rev Clin Oncol.* 2021; 18: 135-51.
- Li C, Wang H, Jiang Y, Fu W, Liu X, Zhong R, et al. Advances in lung cancer screening and early detection. *Cancer Biol Med.* 2022; 19: 591-608.
- Boumahdi S, de Sauvage FJ. The great escape: tumour cell plasticity in resistance to targeted therapy. *Nat Rev Drug Discov.* 2020; 19: 39-56.
- Wang M, Herbst RS, Boshoff C. Toward personalized treatment approaches for non-small-cell lung cancer. *Nat Med.* 2021; 27: 1345-56.
- Rudin CM, Brambilla E, Faivre-Finn C, Sage J. Small-cell lung cancer. *Nat Rev Dis Primers.* 2021; 7: 3.
- Zhang C, Wang H. Accurate treatment of small cell lung cancer: current progress, new challenges and expectations. *Biochim Biophys Acta Rev Cancer.* 2022; 1877: 188798.
- Heitzer E, Haque IS, Roberts CES, Speicher MR. Current and future perspectives of liquid biopsies in genomics-driven oncology. *Nat Rev Genet.* 2019; 20: 71-88.
- Ignatiadis M, Sledge GW, Jeffrey SS. Liquid biopsy enters the clinic - implementation issues and future challenges. *Nat Rev Clin Oncol.* 2021; 18: 297-312.
- Rolfo C, Mack P, Scagliotti GV, Aggarwal C, Arcila ME, Barlesi F, et al. Liquid biopsy for advanced NSCLC: a consensus statement from the international association for the study of lung cancer. *J Thorac Oncol.* 2021; 16: 1647-62.
- Davidson KR, Ha DM, Schwarz MI, Chan ED. Bronchoalveolar lavage as a diagnostic procedure: a review of known cellular and molecular findings in various lung diseases. *J Thorac Dis.* 2020; 12: 4991-5019.
- Domagala-Kulawik J. The relevance of bronchoalveolar lavage fluid analysis for lung cancer patients. *Expert Rev Respir Med.* 2020; 14: 329-37.
- Kalkanis A, Papadopoulos D, Testelmans D, Kopitopoulou A, Boeykens E, Wauters E. Bronchoalveolar lavage fluid-isolated biomarkers for the diagnostic and prognostic assessment of lung cancer. *Diagnostics (Basel).* 2022; 12: 2949.
- Reynolds HY, Newball HH. Analysis of proteins and respiratory cells obtained from human lungs by bronchial lavage. *J Lab Clin Med.* 1974; 84: 559-73.
- Cao C, Yu X, Zhu T, Jiang Q, Li Y, Li X. Diagnostic role of liquid-based cytology of bronchial lavage fluid in addition to bronchial brushing specimens in lung cancer. *Tumori.* 2021; 107: 325-28.
- Han S, Yang W, Li H. A study of the application of fiberoptic bronchoscopy combined with liquid-based cytology test in the early diagnosis of lung cancer. *Oncol Lett.* 2018; 16: 5807-12.
- Debeljak A, Mermolja M, Sorli J, Zupancic M, Zorman M, Remskar J. Bronchoalveolar lavage in the diagnosis of peripheral primary and secondary malignant lung tumors. *Respiration.* 1994; 61: 226-30.
- Ma S, Yu X, Jin X, Qiu F, Chen X, Wang R, et al. The usefulness of liquid-based cytology of bronchoalveolar lavage fluid combined with bronchial brush specimens in lung cancer diagnosis. *J Int Med Res.* 2022; 50: 3000605221132708.
- Zhong CH, Tong D, Zhou ZQ, Su ZQ, Luo YL, Xing J, et al. Performance evaluation of detecting circulating tumor cells and tumor cells in bronchoalveolar lavage fluid in diagnosis of peripheral lung cancer. *J Thorac Dis.* 2018; 10: S830-7.
- Domagala-Kulawik J, Raniszewska A. How to evaluate the immune status of lung cancer patients before immunotherapy. *Breathe (Sheff).* 2017; 13: 291-6.
- Osińska I, Stelmaszczyk-Emmel A, Polubiec-Kownacka M, Dziedzic D, Domagała-Kulawik J. CD4+/CD25(high)/FoxP3+/CD127-regulatory T cells in bronchoalveolar lavage fluid of lung cancer patients. *Hum Immunol.* 2016; 77: 912-5.
- Kwiecien I, Stelmaszczyk-Emmel A, Polubiec-Kownacka M, Dziedzic D, Domagała-Kulawik J. Elevated regulatory T cells, surface and intracellular CTLA-4 expression and interleukin-17 in the lung cancer microenvironment in humans. *Cancer Immunol Immunother.* 2017; 66: 161-70.
- Hu X, Gu Y, Zhao S, Hua S, Jiang Y. Increased IL-10+CD206+CD14+M2-like macrophages in alveolar lavage fluid of patients with small cell lung cancer. *Cancer Immunol Immunother.* 2020; 69: 2547-60.
- Mansour MSI, Hejny K, Johansson F, Mufti J, Vidis A, Mager U, et al. Factors influencing concordance of PD-L1 expression between biopsies and cytological specimens in non-small cell lung cancer. *Diagnostics (Basel).* 2021; 11: 1927.
- Masuiro K, Tamiya M, Fujimoto K, Koyama S, Naito Y, Osa A, et al. Bronchoalveolar lavage fluid reveals factors contributing to the efficacy of PD-1 blockade in lung cancer. *JCI Insight.* 2022; 7: e157915.
- Suresh K, Naidoo J, Lin CT, Danoff S. Immune checkpoint immunotherapy for non-small cell lung cancer: benefits and pulmonary toxicities. *Chest.* 2018; 154: 1416-23.
- Suresh K, Naidoo J, Zhong Q, Xiong Y, Mammen J, de Flores MV, et al. The alveolar immune cell landscape is dysregulated in checkpoint inhibitor pneumonitis. *J Clin Invest.* 2019; 129: 4305-15.
- Kwiecien I, Skirecki T, Polubiec-Kownacka M, Raniszewska A, Domagała-Kulawik J. Immunophenotype of T cells expressing programmed death-1 and cytotoxic T cell antigen-4 in early lung cancer: local vs. systemic immune response. *Cancers (Basel).* 2019; 11: 567.
- Schneider T, Kimpfler S, Warth A, Schnabel PA, Dienemann H, Schadendorf D, et al. Foxp3(+) regulatory T cells and natural killer

- cells distinctly infiltrate primary tumors and draining lymph nodes in pulmonary adenocarcinoma. *J Thorac Oncol.* 2011; 6: 432-8.
31. Brcic L, Stanzer S, Krenbek D, Gruber-Moesenbacher U, Absenger G, Quehenberger F, et al. Immune cell landscape in therapy-naïve squamous cell and adenocarcinomas of the lung. *Virchows Arch.* 2018; 472: 589-98.
 32. Li J, Chen P, Mao CM, Tang XP, Zhu LR. Evaluation of diagnostic value of four tumor markers in bronchoalveolar lavage fluid of peripheral lung cancer. *Asia Pac J Clin Oncol.* 2014; 10: 141-8.
 33. Tian K, Li Z, Qin L. Detection of CEA and ProGRP levels in BALF of patients with peripheral lung cancer and their relationship with CT signs. *Biomed Res Int.* 2022; 2022: 4119912.
 34. Wang H, Zhang X, Liu X, Liu K, Li Y, Xu H. Diagnostic value of bronchoalveolar lavage fluid and serum tumor markers for lung cancer. *J Cancer Res Ther.* 2016; 12: 355-8.
 35. Nikliński J, Chyczewska E, Furman M, Kowal E, Kozłowski M. Value of CEA and SCC-Ag in bronchoalveolar lavage (BAL) and serum of patients with lung carcinoma. *Neoplasma.* 1992; 39: 283-5.
 36. Pina TC, Zapata IT, Hernández FC, López JB, Paricio PP, Hernández PM. Tumour markers in serum, bronchoalveolar lavage and biopsy cytosol in lung carcinoma: what environment lends the optimum diagnostic yield? *Clin Chim Acta.* 2001; 305: 27-34.
 37. de Diego A, Compte L, Sanchis J, Enguidanos MJ, Marco V. Usefulness of carcinoembryonic antigen determination in bronchoalveolar lavage fluid. A comparative study among patients with peripheral lung cancer, pneumonia, and healthy individuals. *Chest.* 1991; 100: 1060-3.
 38. Pothal S, Patil KP, Manjhi R, Dutta P. Diagnostic efficacy of broncho-alveolar lavage carcino-embryonic antigen in carcinoma of lung. *J Family Med Prim Care.* 2019; 8: 1725-9.
 39. Duffy MJ, O'Byrne K. Tissue and blood biomarkers in lung cancer: a review. *Adv Clin Chem.* 2018; 86: 1-21.
 40. Ledermann JA. Serum neurone-specific enolase and other neuroendocrine markers in lung cancer. *Eur J Cancer.* 1994; 30a: 574-6.
 41. Prados MC, Alvarez-Sala R, Blasco R, Chivato T, García Satué JL, García Río FJ, et al. The clinical value of neuron-specific enolase as a tumor marker in bronchoalveolar lavage. *Cancer.* 1994; 74: 1552-5.
 42. Dowlati A, Bury T, Corhay JL, Weber T, Mendes P, Radermecker M. High neuron specific enolase levels in bronchoalveolar lavage fluid of patients with lung carcinoma: diagnostic value, relation to serum neuron specific enolase, and staging. *Cancer.* 1996; 77: 2039-43.
 43. Wieskopf B, Demangeat C, Purohit A, Stenger R, Gries P, Kreisman H, et al. Cyfra 21-1 as a biologic marker of non-small cell lung cancer. Evaluation of sensitivity, specificity, and prognostic role. *Chest.* 1995; 108: 163-9.
 44. Cremades MJ, Menéndez R, Pastor A, Llopis R, Aznar J. Diagnostic value of cytokeratin fragment 19 (CYFRA 21-1) in bronchoalveolar lavage fluid in lung cancer. *Respir Med.* 1998; 92: 766-71.
 45. Cao C, Sun SF, Lv D, Chen ZB, Ding QL, Deng ZC. Utility of VEGF and sVEGFR-1 in bronchoalveolar lavage fluid for differential diagnosis of primary lung cancer. *Asian Pac J Cancer Prev.* 2013; 14: 2443-6.
 46. Ohta Y, Ohta N, Tamura M, Wu J, Tsunozuka Y, Oda M, et al. Vascular endothelial growth factor expression in airways of patients with lung cancer: a possible diagnostic tool of responsive angiogenic status on the host side. *Chest.* 2002; 121: 1624-7.
 47. Zhao X, Sun X, Li XL. Expression and clinical significance of STAT3, P-STAT3, and VEGF-C in small cell lung cancer. *Asian Pac J Cancer Prev.* 2012; 13: 2873-7.
 48. Cao C, Chen ZB, Sun SF, Yu YM, Ding QL, Deng ZC. Evaluation of VEGF-C and tumor markers in bronchoalveolar lavage fluid for lung cancer diagnosis. *Sci Rep.* 2013; 3: 3473.
 49. Lv D, Tan L, Zhang Q, Ma H, Zhang Y, Zhang Q, et al. Vascular endothelial growth factor-D (VEGF-D) is elevated in bronchoalveolar lavage fluid of patients with lung squamous carcinoma. *Clin Lab.* 2019; 65: 125-30.
 50. Domagała-Kulawik J, Hoser G, Safianowska A, Grubek-Jaworska H, Chazan R. Elevated TGF-beta1 concentration in bronchoalveolar lavage fluid from patients with primary lung cancer. *Arch Immunol Ther Exp (Warsz).* 2006; 54: 143-7.
 51. Jakubowska K, Naumnik W, Niklińska W, Chyczewska E. Clinical significance of HMGB-1 and TGF-β level in serum and BALF of advanced non-small cell lung cancer. *Adv Exp Med Biol.* 2015; 852: 49-58.
 52. Naumnik W, Płońska I, Ossolińska M, Nikliński J, Naumnik B. Prognostic value of osteoprotegerin and sRANKL in bronchoalveolar lavage fluid of patients with advanced non-small cell lung cancer. *Adv Exp Med Biol.* 2018; 1047: 1-6.
 53. Naumnik W, Panek B, Ossolińska M, Naumnik B. B cell-attracting chemokine-1 and progranulin in bronchoalveolar lavage fluid of patients with advanced non-small cell lung cancer: new prognostic factors. *Adv Exp Med Biol.* 2019; 1150: 11-6.
 54. Naumnik W, Naumnik B, Niewiarowska K, Ossolinska M, Chyczewska E. Novel cytokines: IL-27, IL-29, IL-31 and IL-33. Can they be useful in clinical practice at the time diagnosis of lung cancer? *Exp Oncol.* 2012; 34: 348-53.
 55. Naumnik W, Naumnik B, Niewiarowska K, Ossolinska M, Chyczewska E. Angiogenic axis angiopoietin-1 and angiopoietin-2/Tie-2 in non-small cell lung cancer: a bronchoalveolar lavage and serum study. *Adv Exp Med Biol.* 2013; 788: 341-8.
 56. Kopiński P, Wandtke T, Dyczek A, Wędrowska E, Roży A, Senderek T, et al. Increased levels of interleukin 27 in patients with early clinical stages of non-small cell lung cancer. *Pol Arch Intern Med.* 2018; 128: 105-14.
 57. Xiong W, Ding W, Xu M, Pudasaini B, Sun J, Zhao Y. The screening role of a biomarker panel in BALF among patients with cancer-suspected pulmonary nodules less than 8 mm. *Clin Respir J.* 2020; 14: 829-38.
 58. Oumeraci T, Schmidt B, Wolf T, Zapatka M, Pich A, Brors B, et al. Bronchoalveolar lavage fluid of lung cancer patients: mapping the uncharted waters using proteomics technology. *Lung Cancer.* 2011; 72: 136-8.
 59. Pastor MD, Nogal A, Molina-Pinelo S, Meléndez R, Salinas A, González De la Peña M, et al. Identification of proteomic signatures associated with lung cancer and COPD. *J Proteomics.* 2013; 89: 227-37.

60. Li QK, Shah B, Li Y, Aiyetan PO, Chen J, Yung R, et al. Glycoproteomic analysis of bronchoalveolar lavage (BAL) fluid identifies tumor-associated glycoproteins from lung adenocarcinoma. *J Proteome Res.* 2013; 12: 3689-96.
61. Uribarri M, Hormaeche I, Zalacain R, Lopez-Vivanco G, Martinez A, Nagore D, et al. A new biomarker panel in bronchoalveolar lavage for an improved lung cancer diagnosis. *J Thorac Oncol.* 2014; 9: 1504-12.
62. Almatroodi SA, McDonald CF, Collins AL, Darby IA, Pouniotis DS. Quantitative proteomics of bronchoalveolar lavage fluid in lung adenocarcinoma. *Cancer Genomics Proteomics.* 2015; 12: 39-48.
63. Ortea I, Rodríguez-Ariza A, Chicano-Gálvez E, Arenas Vacas MS, Jurado Gámez B. Discovery of potential protein biomarkers of lung adenocarcinoma in bronchoalveolar lavage fluid by SWATH MS data-independent acquisition and targeted data extraction. *J Proteomics.* 2016; 138: 106-14.
64. Hmmer A, O'Brien ME, Lynch V, Clynes M, Morgan R, Dowling P. Proteomic analysis of bronchoalveolar lavage fluid (BALF) from lung cancer patients using label-free mass spectrometry. *BBA Clin.* 2017; 7: 97-104.
65. Zhou Y, Yang W, Ao M, Höti N, Gabrielson E, Chan DW, et al. Proteomic analysis of the air-way fluid in lung cancer. Detection of periostin in bronchoalveolar lavage (BAL). *Front Oncol.* 2020; 10: 1072.
66. Vu HM, Mohammad HB, Nguyen TNC, Lee JH, Do Y, Sung JY, et al. Quantitative proteomic analysis of bronchoalveolar lavage fluids from patients with small cell lung cancers. *Proteomics Clin Appl.* 2023; 17: e2300011.
67. Sim SY, Choi YR, Lee JH, Lim JM, Lee SE, Kim KP, et al. In-depth proteomic analysis of human bronchoalveolar lavage fluid toward the biomarker iscovery for lung cancers. *Proteomics Clin Appl.* 2019; 13: e1900028.
68. Travis WD, Brambilla E, Noguchi M, Nicholson AG, Geisinger KR, Yatabe Y, et al. International Association for the Study of Lung Cancer/American Thoracic Society/European Respiratory Society International Multidisciplinary Classification of Lung Adenocarcinoma. *J Thorac Oncol.* 2011; 6: 244-85.
69. Buttitta F, Felicioni L, Del Grammastro M, Filice G, Di Lorito A, Malatesta S, et al. Effective assessment of EGFR mutation status in bronchoalveolar lavage and pleural fluids by next-generation sequencing. *Clin Cancer Res.* 2013; 19: 691-8.
70. Park S, Hur JY, Lee KY, Lee JC, Rho JK, Shin SH, et al. Assessment of EGFR mutation status using cell-free DNA from bronchoalveolar lavage fluid. *Clin Chem Lab Med.* 2017; 55: 1489-95.
71. Yanev N, Mekov E, Valev D, Yankov G, Milanov V, Bichev S, et al. EGFR mutation status yield from bronchoalveolar lavage in patients with primary pulmonary adenocarcinoma compared to a venous blood sample and tissue biopsy. *PeerJ.* 2021; 9: e11448.
72. Zeng D, Wang C, Mu C, Su M, Mao J, Huang J, et al. Cell-free DNA from bronchoalveolar lavage fluid (BALF): a new liquid biopsy medium for identifying lung cancer. *Ann Transl Med.* 2021; 9: 1080.
73. Nair VS, Hui AB, Chabon JJ, Esfahani MS, Stehr H, Nabet BY, et al. Genomic profiling of bronchoalveolar lavage fluid in lung cancer. *Cancer Res.* 2022; 82: 2838-47.
74. Topaloglu O, Hoque MO, Tokumaru Y, Lee J, Ratovitski E, Sidransky D, et al. Detection of promoter hypermethylation of multiple genes in the tumor and bronchoalveolar lavage of patients with lung cancer. *Clin Cancer Res.* 2004; 10: 2284-8.
75. Ren M, Wang C, Sheng D, Shi Y, Jin M, Xu S. Methylation analysis of SHOX2 and RASSF1A in bronchoalveolar lavage fluid for early lung cancer diagnosis. *Ann Diagn Pathol.* 2017; 27: 57-61.
76. Zhang C, Yu W, Wang L, Zhao M, Guo Q, Lv S, et al. DNA methylation analysis of the SHOX2 and RASSF1A panel in bronchoalveolar lavage fluid for lung cancer diagnosis. *J Cancer.* 2017; 8: 3585-91.
77. Ahrendt SA, Chow JT, Xu LH, Yang SC, Eisenberger CF, Esteller M, et al. Molecular detection of tumor cells in bronchoalveolar lavage fluid from patients with early stage lung cancer. *J Natl Cancer Inst.* 1999; 91: 332-9.
78. Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med.* 2009; 361: 947-57.
79. Zhou C, Wu YL, Chen G, Feng J, Liu XQ, Wang C, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. *Lancet Oncol.* 2011; 12: 735-42.
80. Arcila ME, Oxnard GR, Nafa K, Riely GJ, Solomon SB, Zakowski MF, et al. Rebiopsy of lung cancer patients with acquired resistance to EGFR inhibitors and enhanced detection of the T790M mutation using a locked nucleic acid-based assay. *Clin Cancer Res.* 2011; 17: 1169-80.
81. Pérez-Barrios C, Sánchez-Herrero E, García-Simón N, Barquín M, Clemente MB, Provencio M, et al. ctDNA from body fluids is an adequate source for EGFR biomarker testing in advanced lung adenocarcinoma. *Clin Chem Lab Med.* 2021; 59: 1221-9.
82. Macías M, Alegre E, Alkorta-Aranburu G, Patiño-García A, Mateos B, Andueza MP, et al. The dynamic use of EGFR mutation analysis in cell-free DNA as a follow-up biomarker during different treatment lines in non-small-cell lung cancer patients. *Dis Markers.* 2019; 2019: 7954921.
83. Kneip C, Schmidt B, Seegebarth A, Weickmann S, Fleischhacker M, Liebenberg V, et al. SHOX2 DNA methylation is a biomarker for the diagnosis of lung cancer in plasma. *J Thorac Oncol.* 2011; 6: 1632-8.
84. Li P, Liu S, Du L, Mohseni G, Zhang Y, Wang C. Liquid biopsies based on DNA methylation as biomarkers for the detection and prognosis of lung cancer. *Clin Epigenetics.* 2022; 14: 118.
85. Dietrich D, Kneip C, Raji O, Liloglou T, Seegebarth A, Schlegel T, et al. Performance evaluation of the DNA methylation biomarker SHOX2 for the aid in diagnosis of lung cancer based on the analysis of bronchial aspirates. *Int J Oncol.* 2012; 40: 825-32.
86. Schneider KU, Dietrich D, Fleischhacker M, Leschber G, Merk J, Schäper F, et al. Correlation of SHOX2 gene amplification and DNA methylation in lung cancer tumors. *BMC Cancer.* 2011; 11: 102.
87. Ilse P, Biesterfeld S, Pomjanski N, Fink C, Schramm M. SHOX2 DNA methylation is a tumour marker in pleural effusions. *Cancer Genomics Proteomics.* 2013; 10: 217-23.

88. Deng Q, Su B, Ji X, Fang Q, Zhou S, Zhou C. Predictive value of unmethylated RASSF1A on disease progression in non-small cell lung cancer patients receiving pemetrexed-based chemotherapy. *Cancer Biomark.* 2020; 27: 313-23.
89. Kim JO, Gazala S, Razzak R, Guo L, Ghosh S, Roa WH, et al. Non-small cell lung cancer detection using microRNA expression profiling of bronchoalveolar lavage fluid and sputum. *Anticancer Res.* 2015; 35: 1873-80.
90. Li H, Jiang Z, Leng Q, Bai F, Wang J, Ding X, et al. A prediction model for distinguishing lung squamous cell carcinoma from adenocarcinoma. *Oncotarget.* 2017; 8: 50704-14.
91. Dutkowska A, Antczak A, Domańska-Senderowska D, Brzezińska-Lasota E. Expression of selected miRNA, RAR β and FHIT genes in BALf of squamous cell lung cancer (squamous-cell carcinoma, SCC) patients: a pilot study. *Mol Biol Rep.* 2019; 46: 6593-7.
92. Rahbarghazi R, Jabbari N, Sani NA, Asghari R, Salimi L, Kalashani SA, et al. Tumor-derived extracellular vesicles: reliable tools for Cancer diagnosis and clinical applications. *Cell Commun Signal.* 2019; 17: 73.
93. Carnino JM, Lee H, Jin Y. Isolation and characterization of extracellular vesicles from Broncho-alveolar lavage fluid: a review and comparison of different methods. *Respir Res.* 2019; 20: 240.
94. Hur JY, Kim HJ, Lee JS, Choi CM, Lee JC, Jung MK, et al. Extracellular vesicle-derived DNA for performing EGFR genotyping of NSCLC patients. *Mol Cancer.* 2018; 17: 15.
95. Hur JY, Lee JS, Kim IA, Kim HJ, Kim WS, Lee KY. Extracellular vesicle-based EGFR genotyping in bronchoalveolar lavage fluid from treatment-naive non-small cell lung cancer patients. *Transl Lung Cancer Res.* 2019; 8: 1051-60.
96. Kim IA, Hur JY, Kim HJ, Kim WS, Lee KY. Extracellular vesicle-based bronchoalveolar lavage fluid liquid biopsy for EGFR mutation testing in advanced non-squamous NSCLC. *Cancers (Basel).* 2022; 14: 2744.
97. Rodríguez M, Silva J, López-Alfonso A, López-Muñiz MB, Peña C, Domínguez G, et al. Different exosome cargo from plasma/bronchoalveolar lavage in non-small-cell lung cancer. *Genes Chromosomes Cancer.* 2014; 53: 713-24.
98. Kim JE, Eom JS, Kim WY, Jo EJ, Mok J, Lee K, et al. Diagnostic value of microRNAs derived from exosomes in bronchoalveolar lavage fluid of early-stage lung adenocarcinoma: a pilot study. *Thorac Cancer.* 2018; 9: 911-5.
99. Yu Y, Abudula M, Li C, Chen Z, Zhang Y, Chen Y. Icotinib-resistant HCC827 cells produce exosomes with mRNA MET oncogenes and mediate the migration and invasion of NSCLC. *Respir Res.* 2019; 20: 217.
100. Speth JM, Penke LR, Bazzill JD, Park KS, de Rubio RG, Schneider DJ, et al. Alveolar macrophage secretion of vesicular SOCS3 represents a platform for lung cancer therapeutics. *JCI Insight.* 2019; 4: e131340.
101. Carvalho AS, Moraes MCS, Hyun Na C, Fierro-Monti I, Henriques A, Zahedi S, et al. Is the proteome of bronchoalveolar lavage extracellular vesicles a marker of advanced lung cancer? *Cancers (Basel).* 2020; 12: 3450.
102. Lee SE, Park HY, Hur JY, Kim HJ, Kim IA, Kim WS, et al. Genomic profiling of extracellular vesicle-derived DNA from bronchoalveolar lavage fluid of patients with lung adenocarcinoma. *Transl Lung Cancer Res.* 2021; 10: 104-16.
103. Carvalho AS, Cuco CM, Lavareda C, Miguel F, Ventura M, Almeida S, et al. Bronchoalveolar lavage proteomics in patients with suspected lung cancer. *Sci Rep.* 2017; 7: 42190.
104. Lindon JC, Holmes E, Nicholson JK. Metabonomics and its role in drug development and disease diagnosis. *Expert Rev Mol Diagn.* 2004; 4: 189-99.
105. Chen Y, Ma Z, Li A, Li H, Wang B, Zhong J, et al. Metabolomic profiling of human serum in lung cancer patients using liquid chromatography/hybrid quadrupole time-of-flight mass spectrometry and gas chromatography/mass spectrometry. *J Cancer Res Clin Oncol.* 2015; 141: 705-18.
106. Maeda J, Higashiyama M, Imaizumi A, Nakayama T, Yamamoto H, Daimon T, et al. Possibility of multivariate function composed of plasma amino acid profiles as a novel screening index for non-small cell lung cancer: a case control study. *BMC Cancer.* 2010; 10: 690.
107. Carrola J, Rocha CM, Barros AS, Gil AM, Goodfellow BJ, Carreira IM, et al. Metabolic signatures of lung cancer in biofluids: NMR-based metabonomics of urine. *J Proteome Res.* 2011; 10: 221-30.
108. O'Shea K, Cameron SJ, Lewis KE, Lu C, Mur LA. Metabolomic-based biomarker discovery for non-invasive lung cancer screening: a case study. *Biochim Biophys Acta.* 2016; 1860: 2682-7.
109. Callejón-Leblic B, García-Barrera T, Grávalos-Guzmán J, Pereira-Vega A, Gómez-Ariza JL. Metabolic profiling of potential lung cancer biomarkers using bronchoalveolar lavage fluid and the integrated direct infusion/gas chromatography mass spectrometry platform. *J Proteomics.* 2016; 145: 197-206.
110. Callejón-Leblic B, García-Barrera T, Pereira-Vega A, Gómez-Ariza JL. Metabolomic study of serum, urine and bronchoalveolar lavage fluid based on gas chromatography mass spectrometry to delve into the pathology of lung cancer. *J Pharm Biomed Anal.* 2019; 163: 122-9.
111. Brüßow H. Problems with the concept of gut microbiota dysbiosis. *Microb Biotechnol.* 2020; 13: 423-34.
112. Lee SH, Sung JY, Yong D, Chun J, Kim SY, Song JH, et al. Characterization of microbiome in bronchoalveolar lavage fluid of patients with lung cancer comparing with benign mass like lesions. *Lung Cancer.* 2016; 102: 89-95.
113. Yan X, Yang M, Liu J, Gao R, Hu J, Li J, et al. Discovery and validation of potential bacterial biomarkers for lung cancer. *Am J Cancer Res.* 2015; 5: 3111-22.
114. Jang HJ, Choi JY, Kim K, Yong SH, Kim YW, Kim SY, et al. Relationship of the lung microbiome with PD-L1 expression and immunotherapy response in lung cancer. *Respir Res.* 2021; 22: 322.
115. Jin J, Gan Y, Liu H, Wang Z, Yuan J, Deng T, et al. Diminishing microbiome richness and distinction in the lower respiratory tract of lung cancer patients: a multiple comparative study design with independent validation. *Lung Cancer.* 2019; 136: 129-35.

116. Zheng L, Sun R, Zhu Y, Li Z, She X, Jian X, et al. Lung microbiome alterations in NSCLC patients. *Sci Rep.* 2021; 11: 11736.
 117. Patnaik SK, Cortes EG, Kannisto ED, Punnanitnont A, Dhillon SS, Liu S, et al. Lower airway bacterial microbiome may influence recurrence after resection of early-stage non-small cell lung cancer. *J Thorac Cardiovasc Surg.* 2021; 161: 419-29.e16.
 118. Bingula R, Filaire E, Molnar I, Delmas E, Berthon JY, Vasson MP, et al. Characterisation of microbiota in saliva, bronchoalveolar lavage fluid, non-malignant, peritumoural and tumour tissue in non-small cell lung cancer patients: a cross-sectional clinical trial. *Respir Res.* 2020; 21: 129.
 119. Lee JH, Chang JH. Diagnostic utility of serum and pleural fluid carcinoembryonic antigen, neuron-specific enolase, and cytokeratin 19 fragments in patients with effusions from primary lung cancer. *Chest.* 2005; 128: 2298-303.
 120. Darlington P, Kullberg S, Eklund A, Grunewald J. Lung CD4+ V α 2.3+ T-cells in sarcoidosis cohorts with Löfgren's syndrome. *Respir Res.* 2020; 21: 61.
 121. Sohn JW. Acute eosinophilic pneumonia. *Tuberc Respir Dis (Seoul).* 2013; 74: 51-5.
 122. Mukae H, Ishimoto H, Yanagi S, Ishii H, Nakayama S, Ashitani J, et al. Elevated BALF concentrations of alpha- and beta-defensins in patients with pulmonary alveolar proteinosis. *Respir Med.* 2007; 101: 715-21.
 123. Singh G, Martin Rumende C, Sharma SK, Rengganis I, Amin Z, Loho T, et al. Low BALF CD4 T cells count is associated with extubation failure and mortality in critically ill covid-19 pneumonia. *Ann Med.* 2022; 54: 1894-905.
 124. Zareba L, Szymanski J, Homoncik Z, Czystowska-Kuzmicz M. EVs from BALF-mediators of inflammation and potential biomarkers in lung diseases. *Int J Mol Sci.* 2021; 22: 3651.
 125. Yu W, Hurley J, Roberts D, Chakraborty SK, Enderle D, Noerholm M, et al. Exosome-based liquid biopsies in cancer: opportunities and challenges. *Ann Oncol.* 2021; 32: 466-77.
- Cite this article as:** Zhang H, Deng D, Li S, Ren J, Huang W, Liu D, et al. Bronchoalveolar lavage fluid assessment facilitates precision medicine for lung cancer. *Cancer Biol Med.* 2024; 21: 230-251. doi: 10.20892/j.issn.2095-3941.2023.0381