

Pest categorisation of *Eulecanium giganteum*

EFSA Panel on Plant Health (PLH) | Claude Bragard | Paula Baptista | Elisavet Chatzivassiliou | Francesco Di Serio | Paolo Gonthier | Josep Anton Jaques Miret | Annemarie Fejer Justesen | Christer Sven Magnusson | Panagiotis Milonas | Juan A. Navas-Cortes | Stephen Parnell | Roel Potting | Philippe Lucien Reignault | Emilio Stefani | Hans-Hermann Thulke | Wopke Van der Werf | Antonio Vicent Civera | Jonathan Yuen | Lucia Zappalà | Jean-Claude Grégoire | Chris Malumphy | Antigoni Akrivou | Virag Kertesz | Dimitrios Papachristos | Oresteia Sfyra | Alan MacLeod

Correspondence: plants@efsa.europa.eu

Abstract

The EFSA Panel on Plant Health performed a pest categorisation of *Eulecanium giganteum* (Hemiptera: Coccidae), the giant eulecanium scale, for the territory of the European Union, following the commodity risk assessment of *Acer palmatum* plants from China, in which *E. giganteum* came to attention as a pest of possible concern. The pest is only known to be present in Asia, where it has been reported from China, India, Iran, Japan and eastern Russia (Primorsky Krai). The pest has not been reported within the EU. It is not listed in Annex II of Commission Implementing Regulation (EU) 2019/2072. It is polyphagous, feeding on broad-leaf trees and shrubs assigned to 41 genera in 22 plant families. Host plant species commonly found in the EU include apricot (*Prunus armeniaca*), elm (*Ulmus* spp.), grapevine (*Vitis vinifera*), maple (*Acer* spp.), oak (*Quercus* spp.), oriental plane (*Platanus orientalis*), pomegranate (*Punica granatum*), quince (*Cydonia oblonga*), silkworm mulberry (*Morus alba*), walnut (*Juglans regia*), and several ornamentals. Climatic conditions and availability of host plants in southern EU countries would most probably allow this species to successfully establish and spread. However, EU native natural enemies are anticipated to provide biological control and therefore reduce potential impacts. Phytosanitary measures are available to reduce the likelihood of entry and spread. *E. giganteum* satisfies all the criteria that are within the remit of EFSA to assess for it to be regarded as a potential Union quarantine pest, other than the criterion on impact which is a key uncertainty.

KEYWORDS

Coccidae, Giant eulecanium scale, Hemiptera, pest risk, plant health, plant pest, quarantine

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs](https://creativecommons.org/licenses/by-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.

© 2024 European Food Safety Authority. *EFSA Journal* published by Wiley-VCH GmbH on behalf of European Food Safety Authority.

CONTENTS

Abstract.....	1
1. Introduction	4
1.1. Background and Terms of Reference as provided by the requestor.....	4
1.1.1. Background	4
1.1.2. Terms of Reference.....	4
1.2. Interpretation of the Terms of Reference	4
1.3. Additional information	4
2. Data and methodologies.....	5
2.1. Data.....	5
2.1.1. Literature search	5
2.1.2. Database search.....	5
2.2. Methodologies.....	5
3. Pest categorisation	6
3.1. Identity and biology of the pest.....	6
3.1.1. Identity and taxonomy	6
3.1.2. Biology of the pest.....	6
3.1.3. Host range/species affected.....	7
3.1.4. Intraspecific diversity	7
3.1.5. Detection and identification of the pest	7
3.2. Pest distribution	8
3.2.1. Pest distribution outside the EU	8
3.2.2. Pest distribution in the EU.....	8
3.3. Regulatory status	9
3.3.1. Commission implementing regulation 2019/2072.....	9
3.3.2. Hosts or species affected that are prohibited from entering the union from third countries.....	9
3.4. Entry, establishment and spread in the EU.....	10
3.4.1. Entry	10
3.4.2. Establishment	10
3.4.2.1. EU distribution of main host plants.....	10
3.4.2.2. Climatic conditions affecting establishment	11
3.4.3. Spread.....	11
3.5. Impacts.....	12
3.6. Available measures and their limitations	12
3.6.1. Identification of potential additional measures.....	12
3.6.1.1. Additional potential risk reduction options	12
3.6.1.2. Additional supporting measures.....	13
3.6.1.3. Biological or technical factors limiting the effectiveness of measures	14
3.7. Uncertainty.....	14
4. Conclusions.....	15
Abbreviations	15
Glossary	15
Conflict of interest	16
Requestor.....	16
Question number.....	16
Copyright for non-EFSA content.....	16
Panel members	16
Map disclaimer	16
References.....	16

Appendix A..... 18

Appendix B.....20

Appendix C.....21

Appendix D.....23

1 | INTRODUCTION

1.1 | Background and Terms of Reference as provided by the requestor

1.1.1 | Background

The new Plant Health Regulation (EU) 2016/2031, on the protective measures against pests of plants, is applying from 14 December 2019. Conditions are laid down in this legislation in order for pests to qualify for listing as Union quarantine pests, protected zone quarantine pests or Union regulated non-quarantine pests. The lists of the EU regulated pests together with the associated import or internal movement requirements of commodities are included in Commission Implementing Regulation (EU) 2019/2072. Additionally, as stipulated in the Commission Implementing Regulation 2018/2019, certain commodities are provisionally prohibited to enter in the EU (high-risk plants, HRP). EFSA is performing the risk assessment of the dossiers submitted by exporting to the EU countries of the HRP commodities, as stipulated in Commission Implementing Regulation 2018/2018. Furthermore, EFSA has evaluated a number of requests from exporting to the EU countries for derogations from specific EU import requirements.

In line with the principles of the new plant health law, the European Commission with the Member States are discussing monthly the reports of the interceptions and the outbreaks of pests notified by the Member States. Notifications of an imminent danger from pests that may fulfil the conditions for inclusion in the list of the Union quarantine pest are included. Furthermore, EFSA has been performing horizon scanning of media and literature.

As a follow-up of the above-mentioned activities (reporting of interceptions and outbreaks, HRP, derogation requests and horizon scanning), a number of pests of concern have been identified. EFSA is requested to provide scientific opinions for these pests, in view of their potential inclusion by the risk manager in the lists of Commission Implementing Regulation (EU) 2019/2072 and the inclusion of specific import requirements for relevant host commodities, when deemed necessary by the risk manager.

1.1.2 | Terms of Reference

EFSA is requested, pursuant to Article 29(1) of Regulation (EC) No 178/2002, to provide scientific opinions in the field of plant health.

EFSA is requested to deliver 53 pest categorisations for the pests listed in Annex 1A, 1B, 1D and 1E (for more details see mandate M-2021-00027 on the [Open.EFSA portal](#)). Additionally, EFSA is requested to perform pest categorisations for the pests so far not regulated in the EU, identified as pests potentially associated with a commodity in the commodity risk assessments of the HRP dossiers (Annex 1C; for more details see mandate M-2021-00027 on the [Open.EFSA portal](#)). Such pest categorisations are needed in the case where there are not available risk assessments for the EU.

When the pests of Annex 1A are qualifying as potential Union quarantine pests, EFSA should proceed to phase 2 risk assessment. The opinions should address entry pathways, spread, establishment, impact and include a risk reduction options analysis.

Additionally, EFSA is requested to develop further the quantitative methodology currently followed for risk assessment, in order to have the possibility to deliver an express risk assessment methodology. Such methodological development should take into account the EFSA Plant Health Panel Guidance on quantitative pest risk assessment and the experience obtained during its implementation for the Union candidate priority pests and for the likelihood of pest freedom at entry for the commodity risk assessment of High Risk Plants.

1.2 | Interpretation of the Terms of Reference

Eulecanium giganteum is one of a number of pests relevant to Annex 1C of the terms of reference (ToR) to be subject to pest categorisation to determine whether it fulfils the criteria of a potential Union quarantine pest for the area of the EU excluding Ceuta, Melilla and the outermost regions of Member States referred to in Article 355(1) of the Treaty on the Functioning of the European Union (TFEU), other than Madeira and the Azores, and so inform EU decision-making as to its appropriateness for potential inclusion in the lists of pests of Commission Implementing Regulation (EU) 2019/ 2072. If a pest fulfils the criteria to be potentially listed as a Union quarantine pest, risk reduction options will be identified.

1.3 | Additional information

This pest categorisation was initiated following the commodity risk assessment of *Acer palmatum* plants from China performed by EFSA (EFSA PLH Panel, 2022), in which *E. giganteum* was identified as a relevant non-regulated EU pest of possible concern, which could potentially enter the EU on *A. palmatum*.

2 | DATA AND METHODOLOGIES

2.1 | Data

2.1.1 | Literature search

A literature search on *E. giganteum* was conducted at the beginning of the categorisation in the ISI Web of Science bibliographic database, using the scientific name of the pest as search term. Papers relevant for the pest categorisation were reviewed, and further references and information were obtained from experts, as well as from citations within the references and grey literature.

2.1.2 | Database search

Data about the import of commodity types that could potentially provide a pathway for the pest to enter the EU and about the area of hosts grown in the EU were obtained from EUROSTAT (Statistical Office of the European Communities).

The Europhyt and TRACES databases were consulted for pest-specific notifications on interceptions and outbreaks. Europhyt is a web-based network run by the Directorate General for Health and Food Safety (DG SANTÉ) of the European Commission as a subproject of PHYSAN (Phyto-Sanitary Controls) specifically concerned with plant health information. TRACES is the European Commission's multilingual online platform for sanitary and phytosanitary certification required for the importation of animals, animal products, food and feed of non-animal origin and plants into the European Union, and the intra-EU trade and EU exports of animals and certain animal products. Up until May 2020, the Europhyt database managed notifications of interceptions of plants or plant products that do not comply with EU legislation, as well as notifications of plant pests detected in the territory of the Member States and the phytosanitary measures taken to eradicate or avoid their spread. The recording of interceptions switched from Europhyt to TRACES in May 2020.

GenBank was searched to determine whether it contained any nucleotide sequences for *E. giganteum* which could be used as reference material for molecular diagnosis. GenBank® (www.ncbi.nlm.nih.gov/genbank/) is a comprehensive publicly available database that as of August 2019 (release version 227) contained over 6.25 trillion base pairs from over 1.6 billion nucleotide sequences for 450,000 formally described species (Sayers et al., 2020).

2.2 | Methodologies

The Panel performed the pest categorisation for *E. giganteum*, following guiding principles and steps presented in the EFSA guidance on quantitative pest risk assessment (EFSA PLH Panel, 2018), the EFSA guidance on the use of the weight of evidence approach in scientific assessments (EFSA Scientific Committee, 2017) and the International Standards for Phytosanitary Measures No. 11 (FAO, 2013).

The criteria to be considered when categorising a pest as a potential Union quarantine pest (QP) is given in Regulation (EU) 2016/2031 Article 3 and Annex I, Section 1 of the Regulation. Table 1 presents the Regulation (EU) 2016/2031 pest categorisation criteria on which the Panel bases its conclusions. In judging whether a criterion is met, the Panel uses its best professional judgement (EFSA Scientific Committee, 2017) by integrating a range of evidence from a variety of sources (as presented above in Section 2.1) to reach an informed conclusion as to whether or not a criterion is satisfied.

The Panel's conclusions are formulated respecting its remit and particularly with regard to the principle of separation between risk assessment and risk management (EFSA founding regulation (EU) No 178/2002); therefore, instead of determining whether the pest is likely to have an unacceptable impact, deemed to be a risk management decision, the Panel will present a summary of the observed impacts in the areas where the pest occurs, and make a judgement about potential likely impacts in the EU. While the Panel may quote impacts reported from areas where the pest occurs in monetary terms, the Panel will seek to express potential EU impacts in terms of yield and quality losses and not in monetary terms, in agreement with the EFSA guidance on quantitative pest risk assessment (EFSA PLH Panel, 2018). Article 3 (d) of Regulation (EU) 2016/2031 refers to unacceptable social impact as a criterion for quarantine pest status. Assessing social impact is outside the remit of the Panel.

TABLE 1 Pest categorisation criteria under evaluation, as derived from Regulation (EU) 2016/2031 on protective measures against pests of plants (the number of the relevant sections of the pest categorisation is shown in brackets in the first column).

Criterion of pest categorisation	Criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest (article 3)
Identity of the pest (Section 3.1)	Is the identity of the pest clearly defined, or has it been shown to produce consistent symptoms and to be transmissible?
Absence/presence of the pest in the EU territory (Section 3.2)	Is the pest present in the EU territory? If present, is the pest in a limited part of the EU or is it scarce, irregular, isolated or present infrequently? If so, the pest is considered to be not widely distributed
Pest potential for entry, establishment and spread in the EU territory (Section 3.4)	Is the pest able to enter into, become established in, and spread within, the EU territory? If yes, briefly list the pathways for entry and spread

(Continues)

TABLE 1 (Continued)

Criterion of pest categorisation	Criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest (article 3)
Potential for consequences in the EU territory (Section 3.5)	Would the pests' introduction have an economic or environmental impact on the EU territory?
Available measures (Section 3.6)	Are there measures available to prevent pest entry, establishment, spread or impacts?
Conclusion of pest categorisation (Section 4)	A statement as to whether (1) all criteria assessed by EFSA above for consideration as a potential quarantine pest were met and (2) if not, which one(s) were not met

3 | PEST CATEGORISATION

3.1 | Identity and biology of the pest

3.1.1 | Identity and taxonomy

Is the identity of the pest clearly defined, or has it been shown to produce consistent symptoms and/or to be transmissible?

Yes, the identity of the pest is established and *Eulecanium giganteum* (Shinji) is the accepted name.

Eulecanium giganteum (Shinji, 1935) is an insect within the order Hemiptera, family Coccidae, and is commonly known as the giant eulecanium scale (García Morales et al., 2016; Kondo & Watson, 2022). It was originally described as *Lecanium gigantea* by Shinji (1935) from specimens collected in Morioka, Japan, on *Magnolia kobus* (northern Japanese magnolia). Later, Borchsenius (1955) redescribed and illustrated specimens of the same species collected in Primorsky Krai, Far East Region of Russia, on *Quercus* spp. as *Eulecanium diminutum* (García Morales et al., 2016). Wang (1980) changed the combination of genus and species to *Eulecanium gigantea* (Ben-Dov, 1993; García Morales et al., 2016).

The EPPO code¹ (EPPO, 2019; Griessinger & Roy, 2015) for this species is: EULCGI (EPPO, online).

3.1.2 | Biology of the pest

E. giganteum reproduces sexually and has one generation per year (García Morales et al., 2016; Kondo & Watson, 2022). Females have three development stages: egg, nymph (two instars) and adult, while males have two additional non-feeding nymphal instars, the prepupa and pupa (Zhao & Xie, 2004). Most of the development stages are found on branches that are 1–3 years old. Females live about 20–34 days while males live only 1–2 days and die after mating (Wang, 2000; Xie, 1985; Yue et al., 2011). Fecundity is high, as an adult female can lay more than 6000 eggs on average during its life span (Kondo & Watson, 2022). The pest develops one annual generation in northern China (Gansu province) (Xie et al., 1995) Shanxi province (Kondo & Watson, 2022) and in Guanzhong region in Shaanxi province (Wang, 2000). It overwinters as a second-instar nymph on twigs (Deng et al., 2016), and the sex ratio of overwintered female to male nymphs is 1:2 (Kondo & Watson, 2022). On average, each female can produce 677 nymphs that successfully reach the leaves (Kondo & Watson, 2022). Xie et al. (1995) have shown that in Taiyuan, Shanxi Province in China, the high level of urban air pollution (sulfur dioxide and lead) caused an increase in population densities of this species (García Morales et al., 2016). The reasons could be: (a) effect of pollutants improved nutrients for the host plant *Styphnolobium japonicum*, so scales became larger and more fertile, and (b) pollutants had negative effect on natural enemies of *E. giganteum* and the scale was no longer controlled (Xue et al., 1999) (Table 2).

¹An EPPO code, formerly known as a Bayer code, is a unique identifier linked to the name of a plant or plant pest important in agriculture and plant protection. Codes are based on genus and species names. However, if a scientific name is changed the EPPO code remains the same. This provides a harmonised system to facilitate the management of plant and pest names in computerised databases, as well as data exchange between IT systems (Griessinger & Roy, 2015; EPPO, 2019).

TABLE 2 Important features of the life history strategy of *Eulecanium giganteum*.

Life stage	Phenology and relation to host	Other relevant information
Egg	In northern China, oviposition occurs from late April to early May. Eggs hatch in late May (Kondo & Watson, 2022)	The eggs take about 25 days to develop (Kondo & Watson, 2022)
Nymph	In spring, overwintering second-instar nymphs complete development and new adults appear in May and start reproducing. The number of the hatched crawlers increases rapidly until June (Deng et al., 2016). From June to September, the crawlers feed on leaves, and then in September–October, the second-instar nymphs move to the branches to overwinter (EFSA PLH Panel, 2022; Kondo & Watson, 2022)	The first-instar nymphs are mobile (crawlers) while the second instars are sedentary (EFSA PLH Panel, 2022; Tao et al., 2002). The crawlers can be dispersed by the wind, insects or birds (EFSA PLH Panel, 2022; Zhao & Xie, 2004)
Prepupa-Pupa (males)	Males have four development stages. Prepupa-pupa stage takes places after the second-instar male nymph (EFSA PLH Panel, 2022)	
Adult	Adults of both sexes emerge and mate from late April to early May (EFSA PLH Panel, 2022)	

3.1.3 | Host range/species affected

E. giganteum is a polyphagous insect, feeding on plants assigned to more than 41 genera in 22 plant families (Appendix A provides a full list of hosts). *E. giganteum* has been recorded on broad-leaf trees and shrubs such as apricot (*Prunus armeniaca*), elm (*Ulmus* spp.), grapevine (*Vitis vinifera*), maple (*Acer* spp.), oak (*Quercus* spp.), oriental plane (*Platanus orientalis*), pomegranate (*Punica granatum*), quince (*Cydonia oblonga*), silkworm mulberry (*Morus alba*), walnut (*Juglans regia*) and several ornamentals (García Morales et al., 2016; Suganthi et al., 2022).

3.1.4 | Intraspecific diversity

Chinese literature between 1989 and 2016 considered what is now recognised as *E. giganteum* and *E. kuwanai* Kanda, 1934 as the same species with wide intraspecific diversity. Shi and Lü (1989) determined that *E. giganteum* and *E. kuwanai* were two different ecological types of the same species, with the different phenotypes resulting from varying population densities. This was because both species are morphologically similar, sympatrically distributed in China, share many of the same hosts, and often appear together on the same plants, and even on the same twigs. Deng et al. (2016), however, demonstrated that they were indeed distinct species using molecular techniques.

There are no reports for intraspecific diversity since *E. giganteum* has been separated from *E. kuwanai*.

3.1.5 | Detection and identification of the pest

Are detection and identification methods available for the pest?

Yes, visual detection is possible, and morphological and molecular identification methods are available.

Detection

Visual examination of plants is an effective way for the detection of *E. giganteum* due to the large size of adult female scales (Kondo & Watson, 2022). Accumulation of honeydew, sooty mould and honeydew-seeking ants are general signs of phloem-feeding insect infestations; they can be used to pinpoint the areas where plants may be inspected for the presence of soft scales (Camacho & Chong, 2015; Deng et al., 2016). Sticky bands around branches can be used to detect crawlers (Bethke & Wilen, 2010).

Symptoms

According to Wang et al. (2012), EFSA PLH Panel (2022) and Kondo and Watson (2022), the main symptoms of *E. giganteum* infestation are:

- honeydew egested by the scales;
- black sooty mould growing on the honeydew;
- partial necrosis and wilting of twigs and leaves, and;
- yellowing, defoliation, reduced plant growth, dieback of the branches or of the entire plant caused by heavy infestations.

These symptoms are similar to those caused by many other phloem-feeding insects and should not be considered as diagnostic.

Identification

The identification of *E. giganteum* requires microscopic examination of slide-mounted adults and verification of the presence of key morphological characteristics. Detailed morphological descriptions, illustrations and keys of adult *E. giganteum* can be found in Danzig (1980), and Zhao and Xie (2004), while egg and nymphal stages are described by Xie (1985).

Molecular identification based on the nucleotide sequence of e.g. the mitochondrial cytochrome c oxidase subunit I (COI) gene can be used for species identification (Deng et al., 2016). GenBank contains gene nucleotide sequences for *E. giganteum* (<https://www.ncbi.nlm.nih.gov/nuccore/?term=Eulecanium+giganteum>).

Description

Young adult females are almost hemispherical, reddish brown to purple-brown with dark irregular lines. The dorsum is covered by thin grey-white powdery wax. At maturity, the body is nearly 19 mm long, 18 mm wide and 14 mm high, making it the largest species in the genus. In mature adult females, there is no visible wax, the scale is dark and often with reddish-brown patches on the dorsum (Kondo & Watson, 2022). Males are winged and have robust legs (Zhao & Xie, 2004).

In addition to its large size, *E. giganteum* can be differentiated from other species in the genus by the following combination of characteristics: (a) marginal setae of one type, conical, present in a single row; (b) the stigmatic spines are not differentiated from the marginal setae; (c) the dorsal tubercles are absent; (d) small dorsal tubular ducts are present; and (e) anal ring with eight setae (Kondo & Watson, 2022).

3.2 | Pest distribution

3.2.1 | Pest distribution outside the EU

E. giganteum is an Asiatic species first described in Morioka, Japan (García Morales et al., 2016). Its present known distribution includes most of northern China, India, Iran, Japan and eastern Russia (Primorsky Krai) (García Morales et al., 2016; Deng et al., 2016; Kondo & Watson, 2022; Suganthi et al., 2022; see Figure 1).



FIGURE 1 Global distribution of *Eulecanium giganteum* (data source: Deng et al., 2016; García Morales et al., 2016; Kondo & Watson, 2022; Suganthi et al., 2022). The polygons with highlighted orange colour indicate the administrative areas where *E. giganteum* is present.

3.2.2 | Pest distribution in the EU

Is the pest present in the EU territory? If present, is the pest in a limited part of the EU or is it scarce, irregular, isolated or present infrequently? If so, the pest is considered to be not widely distributed.

No. *E. giganteum* has not been recorded in the EU territory.

3.3 | Regulatory status

3.3.1 | Commission implementing regulation 2019/2072

E. giganteum is not listed in Annex II of Commission Implementing Regulation (EU) 2019/2072, an implementing act of Regulation (EU) 2016/2031, or in any emergency plant health legislation.

3.3.2 | Hosts or species affected that are prohibited from entering the union from third countries

According to the Commission Implementing Regulation (EU) 2019/2072, Annex VI, introduction of several *E. giganteum* hosts in the Union from certain third countries is prohibited (Table 3).

Plants for planting of *Acer* L., *Corylus* L., *Fraxinus* L., *Juglans* L., *Quercus* L., *Robinia* L., *Rosa* L., *Prunus* L., *Salix* L. and *Ulmus* L., which are hosts of *E. giganteum* (Appendix A), are considered high-risk plants for the EU and their import is prohibited pending risk assessment (EU 2018/2019).

TABLE 3 List of plants, plant products and other objects that are *Eulecanium giganteum* hosts whose introduction into the Union from certain third countries is prohibited (Source: Commission Implementing Regulation (EU) 2019/2072, Annex VI).

List of plants, plant products and other objects whose introduction into the union from certain third countries is prohibited			
	Description	CN code	Third country, group of third countries or specific area of third country
2.	Plants of [...] <i>Quercus</i> L., with leaves, other than fruit and seeds	ex 0602 10 90 ex 0602 20 20 ex 0602 20 80 ex 0602 90 41 ex 0602 90 45 ex 0602 90 46 ex 0602 90 48 ex 0602 90 50 ex 0602 90 70 ex 0602 90 99 ex 0604 20 90 ex 1404 90 00	Third countries other than Albania, Andorra, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canary Islands, Faeroe Islands, Georgia, Iceland, Liechtenstein, Moldova, Monaco, Montenegro, North Macedonia, Norway, Russia (only the following parts: Central Federal District [Tsentralny federalny okrug], Northwestern Federal District [Severo- Zapadny federalny okrug], Southern Federal District (Yuzhny federalny okrug), North Caucasian Federal District [Severo-Kavkazsky federalny okrug] and Volga Federal District [Privolzhsky federalny okrug]), San Marino, Serbia, Switzerland, Türkiye, Ukraine and the United Kingdom
5.	Isolated bark of <i>Quercus</i> L., other than <i>Quercus suber</i> L.	ex 1404 90 00 ex 4401 40 90	Mexico
8.	Plants for planting of [...] <i>Cydonia</i> Mill., [...] <i>Prunus</i> L., [...] and <i>Rosa</i> L., other than dormant plants free from leaves, flowers and fruits	ex 0602 10 90 ex 0602 20 20 ex 0602 20 80 ex 0602 40 00 ex 0602 90 41 ex 0602 90 45 ex 0602 90 46 ex 0602 90 47 ex 0602 90 48 ex 0602 90 50 ex 0602 90 70 ex 0602 90 91 ex 0602 90 99	Third countries other than Albania, Andorra, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canary Islands, Faeroe Islands, Georgia, Iceland, Liechtenstein, Moldova, Monaco, Montenegro, North Macedonia, Norway, Russia (only the following parts: Central Federal District (Tsentralny federalny okrug), Northwestern Federal District (Severo- Zapadny federalny okrug), Southern Federal District (Yuzhny federalny okrug), North Caucasian Federal District (Severo-Kavkazsky federalny okrug) and Volga Federal District (Privolzhsky federalny okrug)), San Marino, Serbia, Switzerland, Türkiye, Ukraine and the United Kingdom
9.	Plants for planting of <i>Cydonia</i> Mill., [...] <i>Prunus</i> L. and [...] and their hybrids, and [...] other than seeds	ex 0602 10 90 ex 0602 20 20 ex 0602 90 30 ex 0602 90 41 ex 0602 90 45 ex 0602 90 46 ex 0602 90 48 ex 0602 90 50 ex 0602 90 70 ex 0602 90 91 ex 0602 90 99	Third countries other than Albania, Algeria, Andorra, Armenia, Australia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canada, Canary Islands, Egypt, Faeroe Islands, Georgia, Iceland, Israel, Jordan, Lebanon, Libya, Liechtenstein, Moldova, Monaco, Montenegro, Morocco, New Zealand, North Macedonia, Norway, Russia (only the following parts: Central Federal District (Tsentralny federalny okrug), Northwestern Federal District (Severo-Zapadny federalny okrug), Southern Federal District (Yuzhny federalny okrug), North Caucasian Federal District (Severo-Kavkazsky federalny okrug) and Volga Federal District (Privolzhsky federalny okrug)), San Marino, Serbia, Switzerland, Syria, Tunisia, Türkiye, Ukraine, the United Kingdom and United States other than Hawaii
10.	Plants of <i>Vitis</i> L., other than fruits	0602 10 10 0602 20 10 ex 0604 20 90 ex 1404 90 00	Third countries other than Switzerland

3.4 | Entry, establishment and spread in the EU

3.4.1 | Entry

Is the pest able to enter into the EU territory? If yes, identify and list the pathways.

Yes, *E. giganteum* could enter the EU territory. Possible pathways of entry are plants for planting (except seeds, bulbs, and tubers), fruits and cut flowers, and isolated bark.

Comment on plants for planting as a pathway.

Plants for planting are the main pathway for *E. giganteum* to enter the EU (Table 4).

Plants for planting, fruits and cut flowers are the main potential pathways for entry of *E. giganteum* (Table 4).

TABLE 4 Potential pathways for *Eulecanium giganteum* into the EU.

Pathways (e.g. host/intended use/source)	Life stage	Relevant mitigations (e.g. prohibitions [Annex VI], special requirements [Annex VII] or phytosanitary certificates [Annex XI] within Implementing Regulation 2019/2072)
Plants for planting	All life stages	Plants for planting that are hosts of <i>E. giganteum</i> and are prohibited to import from third countries (Regulation 2019/2072, Annex VI), are listed in Table 3 Plants for planting from third countries require a phytosanitary certificate (Regulation 2019/2072, Annex XI, Part A) Some hosts are considered high-risk plants (EU 2018/2019) for the EU and their import is prohibited subject to risk assessment
Fruits and cut flowers	All life stages	Fruits and cut flowers from third countries require a phytosanitary certificate to be imported into the EU (2019/2072, Annex XI, Part A). However, no requirements are specified for <i>E. giganteum</i>
Host isolated bark	Eggs	Annex VI prohibitions apply to the bark of some hosts i.e. <i>Quercus</i> sp., Table 3, point 5, but for countries where <i>E. giganteum</i> is not known to occur

Annual imports of *E. giganteum* hosts from countries where the pest is known to occur are provided in Appendix C.

Notifications of interceptions of harmful organisms began to be compiled in Europhyt in May 1994 and in TRACES in May 2020. As of November 2023, there were no records of interception of *E. giganteum* in the Europhyt and TRACES databases.

3.4.2 | Establishment

Is the pest able to become established in the EU territory?

Yes, the climate in EU countries is suitable and there are many available hosts that can support establishment.

Climatic mapping is the principal method for identifying areas that could provide suitable conditions for the establishment of a pest taking key abiotic factors into account (Baker, 2002; Baker et al., 2000). Availability of hosts is considered in Section 3.4.2.1. Climatic factors are considered in Section 3.4.2.2.

3.4.2.1 | EU distribution of main host plants

Many genera of *E. giganteum* host plants are present or are grown widely across the EU. Among others, *Acer*, *Cydonia*, *Ficus*, *Juglans*, *Morus*, *Prunus*, *Quercus*, *Ulmus*, *Vitis* and some ornamental plants. The main hosts of the scale insect cultivated in the EU between 2017 and 2022 are shown in Table 5.

TABLE 5 Crop area of *Eulecanium giganteum* key hosts in the EU in 1000 ha (Eurostat accessed on 30 November 2023).

Crop	2017	2018	2019	2020	2021	2022
Apricots	72.23	72.57	73.22	76.13	73.48	74.90
Berries (excluding strawberries)*	146.27	150.42	154.44	154.27	157.07	–

TABLE 5 (Continued)

Crop	2017	2018	2019	2020	2021	2022
Figs	24.63	24.99	25.59	27.63	25.79	26.42
Walnuts	74.15	80.60	87.62	99.21	97.00	104.74
Grapes	3133.32	3135.50	3155.20	3146.24	3120.22	3132.12

*Only a proportion of these berries are host species, specifically *Morus alba* (mulberries).

3.4.2.2 | Climatic conditions affecting establishment

E. giganteum occurs in continental, temperate and dry areas in Asia. The biology of this pest is little studied and no temperature thresholds for development have been reported. Consequently, there is some uncertainty regarding the climatic requirements of the insect. Figure 2 shows the world distribution of Köppen–Geiger climate types (Kottek et al., 2006) that occur in the EU, and which occur in countries where *E. giganteum* has been reported. In Russia, it was found to occur in the most southerly tip of the Primorsky Krai (Far East) Territory which experiences a humid continental climate (Dfb). Winters are cold and dry with daily mean temperatures between December and March below zero, and the average low temperature in January -13.9°C (EFSA PLH Panel, 2021). Based on locations where *E. giganteum* is reported in literature such as Anhui, Henan and Hunan in China (Cfa climate); Honshu in Japan (Cfa climate); and Kermanshah in Iran (Csa climate), southern EU countries may provide suitable climatic conditions for establishment. Distribution of the pest in its native range might be broader. The comparison of the current distribution of the pest with the suitability of the environment in the EU indicates that Southern Scandinavia and Southern Europe are climatically suitable but not central Europe. However, the Panel considers that climates between these areas would also enable survival of the pest (a map including Cfb is included in Appendix D).




FIGURE 2 World distribution of Köppen–Geiger climate types that occur in the EU and which occur in countries where *Eulecanium giganteum* has been reported.

3.4.3 | Spread

Describe how the pest would be able to spread within the EU territory following establishment?

Natural spread by first instar nymphs crawling or being carried by wind, or by hitchhiking on other animals, humans or machinery, will occur locally and relatively slowly. All stages may be moved over long distances in trade of infested plant materials, specifically plants for planting, fruits, and cut flowers.

Comment on plants for planting as a mechanism of spread.

Plants for planting provide a main spread mechanism for *E. giganteum*.

First-instar nymphs (crawlers) may be carried to neighbouring plants by their own movement, wind or by hitchhiking on clothing, equipment or animals (EFSA PLH Panel, 2020).

Plants for planting, scion and rough wood are the main pathways of spread of *E. giganteum*, especially over long distances (EFSA PLH Panel, 2022).

3.5 | Impacts

Would the pests' introduction have an economic or environmental impact on the EU territory?

This is a key uncertainty. Evidence from Asia indicates *E. giganteum* is a pest. However, the closely related species *E. excrucians*, also a pest in Asia, has been established in England since the 1990s but is not an economically important pest due in part to natural enemies providing control.

E. giganteum feeds on the phloem and egests sugary honeydew, which serves as a medium for the growth of sooty moulds. The mould reduces photosynthesis and gas exchange, causing a loss of vigour and yield (Kondo & Watson, 2022). Infestations of *E. giganteum* may completely cover the lower surfaces of the foliage, forming a dense mat of waxy secretions. Fruits from infested plants and infested ornamental plants are unmarketable (Kondo & Watson, 2022).

In China, *E. giganteum* is reported to cause serious damage to garden trees (EFSA PLH Panel, 2022). The production of jujube (*Ziziphus jujuba*) in Xinjiang, China has been reported to be severely threatened by *E. giganteum* (Deng et al., 2016, Li and Xu, 2013). Xie (1985) reported that in Lanzhou, China, *E. giganteum* causes serious damage between April and May in black locust (*Robinia pseudoacacia*), and Japanese pagoda trees (*S. japonicum*) while in the Kunming area, China, it mainly damages trident maple (*Acer buergerianum*), and triangle maple (*Celtis tetrandra*) (Tao et al., 2002). In Taiyuan, China, the population of *E. giganteum* on its host *S. japonicum* was reported to be positively correlated with air pollutants (Kondo & Watson, 2022; Xie et al., 1995).

The closely related *Eulecanium excrucians* (Ferris) is native to Asia, highly polyphagous, and is recorded feeding on many deciduous orchard and ornamental trees. In China, it is reported to be a pest of apple, pear and peach trees, although in the UK and USA, where it has been introduced (Malumphy, 2005), it does not cause economic damage. In the UK, this is due to high levels of parasitism as native parasitoids were rapidly recruited to attack the new resource (CP Malumphy, DEFRA, oral communication at the working group meeting on 9 February, 2024). Several *E. giganteum* parasitoid species are reported in the EU (Noyes, 2019). In addition, there are predators of *E. giganteum* that also occur in the EU (Kondo & Watson, 2022). Therefore, should *E. giganteum* be introduced into the EU, we would expect these natural enemies to accept this new host and provide biological control.

3.6 | Available measures and their limitations

Are there measures available to prevent pest entry, establishment, spread or impacts such that the risk becomes mitigated?

Yes. Although the existing phytosanitary measures identified in Section 3.3.2 do not specifically target *E. giganteum*, they mitigate the likelihood of its entry into, establishment, and spread within the EU (see also Section 3.6.1).

3.6.1 | Identification of potential additional measures

Phytosanitary measures (prohibitions) are currently applied to some host plants for planting (see Section 3.3.2).

Additional potential risk reduction options and supporting measures are shown in Sections 3.6.1.1 and 3.6.1.2.

3.6.1.1 | Additional potential risk reduction options

Potential additional control measures are listed in Table 6.

TABLE 6 Selected control measures (a full list is available in EFSA PLH Panel, 2018) for pest entry/establishment/spread/impact in relation to currently unregulated hosts and pathways. Control measures are measures that have a direct effect on pest abundance.

Control measure/risk reduction option (Blue underline = Zenodo doc, Blue = WIP)	RRO summary	Risk element targeted (entry/establishment/spread/impact)
Require pest freedom	Pest-free place of production (e.g. place of production and its immediate vicinity is free from pest over an appropriate time period, e.g. since the beginning of the last complete cycle of vegetation, or past 2 or 3 cycles). Pest-free production site	Entry/Establishment/Spread
Growing plants in isolation	Place of production is insect proof originate in a place of production with complete physical isolation	Entry (reduce infestation)/Establishment/Spread
Managed growing conditions	Used to mitigate likelihood of infestation at origin. Plants collected directly from natural habitats, have been grown, held and trained for at least two consecutive years prior to dispatch in officially registered nurseries, which are subject to an officially supervised control regime	Entry (reduce infestation)/Establishment/Spread
Roguing and pruning	Roguing is defined as the removal of infested plants and/or uninfested host plants in a delimited area, whereas pruning is defined as the removal of infested plant parts only without affecting the viability of the plant	Entry/Spread/Impact
Biological control and behavioural manipulation	Zhang and Huang (2001) reported <i>Oriencyrtus liaoi</i> sp. nov. (Hymenoptera: Encyrtidae) as a parasitoid of <i>E. giganteum</i> on willow (<i>Salix</i> spp.) in Zhongwei, China. In northern China, the parasitoid wasp <i>Encyrtus eulecaniumiae</i> sp. nov. (Hymenoptera: Encyrtidae) was reported on <i>E. giganteum</i> (Wang et al., 2016). In Iran (Ghazanchi, Kermanshah) larvae of <i>Dicrodiplosis manihoti</i> (Diptera: Cecidomyiidae) were observed feeding on egg masses and crawlers of <i>E. giganteum</i> on Canadian phlox (<i>Phlox divaricate</i>) (Jalilvand et al., 2013). In Xinjiang, <i>Eunotus aequalivena</i> (Hymenoptera: Pteromalidae) was reported to be a highly parasitic species to <i>E. giganteum</i> (Zhang et al., 2016). Tao et al. (2002) reported that, in Kunming, there are six natural enemies of <i>E. giganteum</i> such as <i>Blastothrix sericea</i> , <i>Metaphycus pulvinariae</i> , <i>Cocophagus hawaiiensis</i> , <i>Microterys ericeri</i> and <i>Cocophagus</i> sp. The first two parasitoid wasps have about 88% parasitism rate under natural conditions (Tao et al., 2002). Some of the parasitoid species that have been recorded to parasitise on <i>E. giganteum</i> in its distribution range, such as <i>Blastothrix sericea</i> , <i>Metaphycus pulvinariae</i> and <i>Cocophagus</i> spp., are also recorded in the EU territory	Impact
Chemical treatments on crops including reproductive material	Used to mitigate likelihood of infestation of pests susceptible to chemical treatments. The effectiveness of insecticide applications against soft scales may be reduced by the waxy coating of the adult. The efficacy of insecticides was tested on different nymphal stages of <i>E. giganteum</i> . Only the control of nymphs at the end of first instar and the beginning of second instar was effective, with mortality rate over 94% (Xie, 1985)	Entry/Establishment/Spread/Impact
Cleaning and disinfection of facilities, tools and machinery	The physical and chemical cleaning and disinfection of facilities, tools, machinery, facilities and other accessories (e.g. boxes, pots, hand tools)	Entry/Spread
Heat and cold treatments	Controlled temperature treatments aimed to kill or inactivate pests without causing any unacceptable prejudice to the treated material itself. Treatments relevant for this risk mitigation measure are: autoclaving; steam; hot water; hot air; cold treatment	Entry/Spread
Controlled atmosphere	Treatment of plants by storage in a modified atmosphere (including modified humidity, O ₂ , CO ₂ , temperature, pressure)	Entry/Spread (via commodity)

3.6.1.2 | Additional supporting measures

Potential additional supporting measures are listed in [Table 7](#).

TABLE 7 Selected supporting measures (a full list is available in EFSA PLH Panel, 2018) in relation to currently unregulated hosts and pathways. Supporting measures are organisational measures or procedures supporting the choice of appropriate risk reduction options that do not directly affect pest abundance.

Supporting measure (Blue underline = Zenodo doc, Blue = WIP)	Summary	Risk element targeted (entry/ establishment/spread/ impact)
Inspection and trapping	ISPM 5 (FAO, 2023) defines inspection as the official visual examination of plants, plant products or other regulated articles to determine if pests are present or to determine compliance with phytosanitary regulations. The effectiveness of sampling and subsequent inspection to detect pests may be enhanced by including trapping and luring techniques	Entry/Establishment/Spread/ Impact
Laboratory testing	Examination, other than visual, to determine if pests are present using official diagnostic protocols. Diagnostic protocols describe the minimum requirements for reliable diagnosis of regulated pests	Entry/Establishment/Spread
Sampling	According to ISPM 31 (FAO, 2008), it is usually not feasible to inspect entire consignments, so phytosanitary inspection is performed mainly on samples obtained from a consignment. It is noted that the sampling concepts presented in this standard may also apply to other phytosanitary procedures, notably selection of units for testing. For inspection, testing and/or surveillance purposes, the sample may be taken according to a statistically based or a non-statistical sampling methodology	Entry/Establishment
Phytosanitary certificate and plant passport	According to ISPM 5 (FAO, 2023), a phytosanitary certificate and a plant passport are official paper documents or their official electronic equivalents, consistent with the model certificates of the IPPC, attesting that a consignment meets phytosanitary import requirements: a) export certificate (import) b) plant passport (EU internal trade)	Entry/Establishment/Spread
Certified and approved premises	Mandatory/voluntary certification/approval of premises is a process including a set of procedures and of actions implemented by producers, conditioners and traders contributing to ensure the phytosanitary compliance of consignments. It can be a part of a larger system maintained by the NPPO in order to guarantee the fulfilment of plant health requirements of plants and plant products intended for trade. Key property of certified or approved premises is the traceability of activities and tasks (and their components) inherent the pursued phytosanitary objective. Traceability aims to provide access to all trustful pieces of information that may help to prove the compliance of consignments with phytosanitary requirements of importing countries	Entry/Spread
Certification of reproductive material (voluntary/ official)	Plants come from within an approved propagation scheme and are certified pest free (level of infestation) following testing; used to mitigate against pests that are included in a certification scheme	Entry/Spread
Delimitation of Buffer zones	ISPM 5 (FAO, 2023) defines a buffer zone as 'an area surrounding or adjacent to an area officially delimited for phytosanitary purposes in order to minimise the probability of spread of the target pest into or out of the delimited area, and subject to phytosanitary or other control measures, if appropriate'. The objectives for delimiting a buffer zone can be to prevent spread from the outbreak area and to maintain a pest-free production place (PFPP), site (PFPS) or area (PFA)	Spread
Surveillance	Surveillance to guarantee that plants and produce originate from a pest-free area could be an option	Establishment/Spread

3.6.1.3 | Biological or technical factors limiting the effectiveness of measures

- *E. giganteum* is polyphagous, making the inspections of all consignments containing hosts from countries where the pest occurs difficult.
- Egg masses may be difficult to detect on large trees.
- Limited effectiveness of insecticides due to the presence of protective cover over the scales.
- Limited biological data on developmental threshold temperatures.

3.7 | Uncertainty

Noting that the related species *E. excrescens* is reported as a pest of apple, pear and peach trees in China, but following its establishment in USA and UK (Malumphy, 2005), has failed to cause any economic and environmental impacts, there is a key uncertainty as to whether *E. giganteum* will cause economic or environmental impact if it were to establish in the EU.

4 | CONCLUSIONS

Eulecanium giganteum satisfies all the criteria that are within the remit of EFSA to assess for it to be regarded as a potential Union quarantine pest, other than the criterion on impact which is a key uncertainty (Table 8).

TABLE 8 The Panel's conclusions on the pest categorisation criteria defined in Regulation (EU) 2016/2031 on protective measures against pests of plants (the number of the relevant sections of the pest categorisation is shown in brackets in the first column).

Criterion of pest categorisation	Panel's conclusions against criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest	Key uncertainties
Identity of the pest (Section 3.1)	The identity of <i>E. giganteum</i> is established. Taxonomic keys based on morphology of adults exist. There are also molecular techniques for species identification	None
Absence/presence of the pest in the EU (Section 3.2)	No, <i>E. giganteum</i> is not known to occur in the EU	None
Pest potential for entry, establishment and spread in the EU (Section 3.4)	<i>E. giganteum</i> is able to enter, become established and spread within the EU territory especially in the southern EU MS. The main pathways are plants for planting, cut flowers, and fruits	None
Potential for consequences in the EU (Section 3.5)	The introduction of the pest could cause yield and quality losses on several crops and reduce the value of ornamental plants	There is uncertainty whether <i>E. giganteum</i> will cause economic or environmental impact if it were to establish in the EU
Available measures (Section 3.6)	There are measures available to prevent entry, establishment and spread of <i>E. giganteum</i> in the EU. Risk reduction options include inspections, chemical and physical treatments on consignments of fresh plant material from infested countries and the production of plants for import in the EU in pest free areas. Natural biological control could prevent impact	None
Conclusion (Section 4)	<i>E. giganteum</i> satisfies all the criteria that are within the remit of EFSA to assess for it to be regarded as a potential Union quarantine pest, other than the criterion on impact which is a key uncertainty	There is uncertainty whether <i>E. giganteum</i> will cause economic or environmental impact if it were to establish in the EU
Aspects of assessment to focus on/ scenarios to address in future if appropriate:		

ABBREVIATIONS

EPPO	European and Mediterranean Plant Protection Organization
FAO	Food and Agriculture Organization
IPPC	International Plant Protection Convention
ISPM	International Standards for Phytosanitary Measures
MS	Member State
PLH	EFSA Panel on Plant Health
PZ	Protected Zone
TFEU	Treaty on the Functioning of the European Union
ToR	Terms of Reference

GLOSSARY

Containment (of a pest)	Application of phytosanitary measures in and around an infested area to prevent spread of a pest (FAO, 2023).
Control (of a pest)	Suppression, containment or eradication of a pest population (FAO, 2023).
Entry (of a pest)	Movement of a pest into an area where it is not yet present, or present but not widely distributed and being officially controlled (FAO, 2023).
Eradication (of a pest)	Application of phytosanitary measures to eliminate a pest from an area (FAO, 2023).
Establishment (of a pest)	Perpetuation, for the foreseeable future, of a pest within an area after entry (FAO, 2023).
Greenhouse	A walk-in, static, closed place of crop production with a usually translucent outer shell, which allows controlled exchange of material and energy with the surroundings and prevents release of plant protection products (PPPs) into the environment.
Hitchhiker	An organism sheltering or transported accidentally via inanimate pathways including with machinery, shipping containers and vehicles; such organisms are also known as contaminating pests or stowaways (Toy & Newfield, 2010).
Impact (of a pest)	The impact of the pest on the crop output and quality and on the environment in the occupied spatial units.

Introduction (of a pest)	The entry of a pest resulting in its establishment (FAO, 2023).
Pathway	Any means that allows the entry or spread of a pest (FAO, 2023).
Phytosanitary measures	Any legislation, regulation or official procedure having the purpose to prevent the introduction or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests (FAO, 2023).
Quarantine pest	A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled (FAO, 2023).
Risk reduction option (RRO)	A measure acting on pest introduction and/or pest spread and/or the magnitude of the biological impact of the pest should the pest be present. A RRO may become a phytosanitary measure, action or procedure according to the decision of the risk manager.
Spread (of a pest)	Expansion of the geographical distribution of a pest within an area (FAO, 2023).

CONFLICT OF INTEREST

If you wish to access the declaration of interests of any expert contributing to an EFSA scientific assessment, please contact interestmanagement@efsa.europa.eu.

REQUESTOR

European Commission

QUESTION NUMBER

EFSA-Q-2023-00324

COPYRIGHT FOR NON-EFSA CONTENT

EFSA may include images or other content for which it does not hold copyright. In such cases, EFSA indicates the copyright holder and users should seek permission to reproduce the content from the original source.

PANEL MEMBERS

Claude Bragard, Paula Baptista, Elisavet Chatzivassiliou, Francesco Di Serio, Paolo Gonthier, Josep Anton Jaques Miret, Annemarie Fejer Justesen, Alan MacLeod, Christer Sven Magnusson, Panagiotis Milonas, Juan A. Navas-Cortes, Stephen Parnell, Roel Potting, Philippe L. Reignault, Emilio Stefani, Hans-Hermann Thulke, Wopke Van der Werf, Antonio Vicent Civera, Jonathan Yuen, and Lucia Zappalà.

MAP DISCLAIMER

The designations employed and the presentation of material on any maps included in this scientific output do not imply the expression of any opinion whatsoever on the part of the European Food Safety Authority concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

REFERENCES

- Baker, R. H. A. (2002). Predicting the limits to the potential distribution of alien crop pests. In G. J. Hallman & C. P. Schwalbe (Eds.), *Invasive arthropods in agriculture: Problems and solutions* (pp. 207–241). Science Publishers Inc.
- Baker, R. H. A., Sansford, C. E., Jarvis, C. H., Cannon, R. J. C., MacLeod, A., & Walters, K. F. A. (2000). The role of climatic mapping in predicting the potential geographical distribution of non-indigenous pests under current and future climates. *Agriculture, Ecosystems & Environment*, 82(1-3), 57–71.
- Ben-Dov, Y. (1993). *A systematic catalogue of the soft scale insects of the world (Homoptera: Coccoidea: Coccidae)* (p. 536). Sandhill Crane Press Gainesville.
- Bethke, J. A., & Wilen, C. A. (2010). UC IPM pest management guidelines: floriculture and ornamental nurseries, UCANR Publication 3392. <https://ipm.ucanr.edu/agriculture/floriculture-and-ornamental-nurseries/soft-scales/>
- Camacho, E. R., & Chong, J. H. (2015). General biology and current management approaches of soft scale pests (Hemiptera: Coccidae). *Journal of Integrated Pest Management*, 6, 1–22. <https://doi.org/10.1093/jipm/pmv016>
- Danzig, E. M. (1980). Coccoids of the Far East USSR (Homoptera, Coccinea) with phylogenetic analysis of scale insects of the world. Nauka, Leningrad. 367 pp. [in Russian, translated into English and published in 1986 by Amerind publishing Co., New Delhi, India. 450 pp.].
- Deng, J., Li, H.-B., Wang, X.-B., Yu, F., Zhang, Y.-Z., & Wu, S.-A. (2016). Molecular identification of two morphologically similar *Eulecanium* species: *E. Giganteum* and *E. Kuwanai* (Hemiptera: Coccidae). *The Canadian Entomologist*, 148(1), 1–7. <https://doi.org/10.4039/tce.2015.42>
- EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Gonthier, P. (2022). Scientific opinion on the commodity risk assessment of *Acer palmatum* plants grafted on *Acer davidii* from China. *EFSA Journal*, 20(5), 7298. <https://doi.org/10.2903/j.efsa.2022.7298>
- EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Dehnen-Schmutz, K., Di Serio, F., Gonthier, P., Jacques, M. A., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Reignault, P. L., Thulke, H.-H., Van der Werf, W., Civera, A. V., Yuen, J., Zappalà, L., ... Potting, R. (2020). Scientific opinion on the commodity risk assessment of *Jasminum polyanthum* plants from Israel. *EFSA Journal*, 18(8), 6928. <https://doi.org/10.2903/j.efsa.2020.6225>
- EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Thulke, H. H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., Grégoire, J.-C., Malumphy, C., ... MacLeod, A. (2021). Scientific opinion on the pest categorisation of *Crisicoccus pini*. *EFSA Journal*, 19(11), 6928. <https://doi.org/10.2903/j.efsa.2021.6928>
- EFSA PLH Panel (EFSA Panel on Plant Health), Jeger, M., Bragard, C., Caffier, D., Candresse, T., Chatzivassiliou, E., Dehnen-Schmutz, K., Gregoire, J.-C., Jaques Miret, J. A., MacLeod, A., Navajas Navarro, M., Niere, B., Parnell, S., Potting, R., Rafoss, T., Rossi, V., Urek, G., Van Bruggen, A., Van Der Werf, W., ... Gilioli, G. (2018). Guidance on quantitative pest risk assessment. *EFSA Journal*, 16(8), 5350. <https://doi.org/10.2903/j.efsa.2018.5350>

- EFSA Scientific Committee, Hardy, A., Benford, D., Halldorsson, T., Jeger, M. J., Knutsen, H. K., More, S., Naegeli, H., Noteborn, H., Ockleford, C., Ricci, A., Rychen, G., Schlatter, J. R., Silano, V., Solecki, R., Turck, D., Benfenati, E., Chaudhry, Q. M., Craig, P., ... Younes, M. (2017). Scientific opinion on the guidance on the use of the weight of evidence approach in scientific assessments. *EFSA Journal*, 15(8), 4971. <https://doi.org/10.2903/j.efsa.2017.4971>
- EPPO (European and Mediterranean Plant Protection Organization). (2019). EPPO codes. https://www.eppo.int/RESOURCES/eppo_databases/eppo_codes
- EPPO (European and Mediterranean Plant Protection Organization). (online). EPPO Global Database. <https://gd.eppo.int> [Accessed on: 20 December 2023].
- FAO (Food and Agriculture Organization of the United Nations). (2008). ISPM (international standards for Phytosanitary measures) No 31. Methodologies for sampling of consignments. FAO, Rome, 19 pp. https://www.ippc.int/static/media/files/publication/en/2016/11/ISPM_31_2008_Sampling_of_consignments_EN.pdf
- FAO (Food and Agriculture Organization of the United Nations). (2013). ISPM (international standards for Phytosanitary measures) No 11. Pest risk analysis for quarantine pests. FAO, Rome, 36 pp. https://www.ippc.int/sites/default/files/documents/20140512/ispm_11_2013_en_2014-04-30_20140512523-494.65%20KB.pdf
- FAO (Food and Agriculture Organization of the United Nations). (2023). ISPM (international standards for Phytosanitary measures) No 5. Glossary of phytosanitary terms. FAO, Rome, 40 pp. https://assets.ippc.int/static/media/files/publication/en/2023/07/ISPM_05_2023_En_Glossary_PostCPM-17_2023-07-12_Fixed.pdf
- García Morales, M., Denno, B. D., Miller, D. R., Miller, G. L., Ben-Dov, Y., & Hardy, N. B. (2016). ScaleNet: A literature-based model of scale insect biology and systematics. *Database: The Journal of Biological Databases and Curation*, 2016, 1–5. <https://doi.org/10.1093/database/bav118> [Accessed on: 20 December 2023].
- Griessinger, D., & Roy, A.-S. (2015). EPPO codes: a brief description. https://www.eppo.int/media/uploaded_images/RESOURCES/eppo_databases/A4_EPPO_Codes_2018.pdf
- Jalilvand, K., Shirazi, M., Vahedi, H., Fallahzadeh, M., Naghadeh, N., & Samih, M. (2013). A preliminary study on natural enemies of coccioidea (Hemiptera, Sternorrhyncha) in Kermanshah Province, Western Iran. *Acta Phytopathologica et Entomologica Hungarica*, 48(2), 299–308. <https://doi.org/10.1556/aphyt.48.2013.2.11>
- Kondo, T., & Watson, G. W. (Eds.). (2022). *Encyclopedia of scale insect pests* (p. 608). CABI International.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., & Rubel, F. (2006). World map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift*, 15, 259–263. <https://doi.org/10.1127/0941-2948/2006/0130>
- Li, J., & Xu, Q. (2013). *Eulecanium gigantea* occurrence and control measures. *Shaanxi Forest Science Technology*, 2, 81–83.
- Malumphy, C. P. (2005). *Eulecanium excrescens* (Ferris) (Hemiptera: Coccidae), an Asian pest of woody ornamentals and fruit trees, new to Britain. *British Journal of Entomology and Natural History*, 18, 45–49. <https://www.biodiversitylibrary.org/partpdf/263854>
- Malumphy, C. P. (2024). Personal communication at the working group meeting. 9 February, 2024.
- Noyes, J. S. (2019). Universal Chalcidoidea database. World Wide Web Electronic Publication. <https://www.nhm.ac.uk/chalcidooids>
- Sayers, E. W., Cavanaugh, M., Clark, K., Ostell, J., Pruitt, K. D., & Karsch-Mizrachi, I. (2020). Genbank. *Nucleic Acids Research*, 48(Database issue), D84–D86. <https://doi.org/10.1093/nar/gkz956>
- Shi, Y. L., & Lü, J. P. (1989). Studies on morphology and biology of *Eulecanium kuwanai* (Kanda). *Journal of Shandong Agricultural University*, 1, 12–19.
- Suganthi, M., Logeshwaran, R., Abirami, G., Rupa Shree, B., Anandaraj, P., & Senthilkumar, P. (2022). Molecular identification of scale insect (*Eulecanium giganteum*) in *Hibiscus rosa-sinensis*. *Journal of Experimental Biology and Agricultural Sciences*, 10(4), 797–804. [https://doi.org/10.18006/2022.10\(4\).797.804](https://doi.org/10.18006/2022.10(4).797.804)
- Tao, M., Guohua, C., & Benli, H. (2002). Research of parasitoid of *E. Giganteum* (Shinji) in Kunming. *Journal of Yunnan Agricultural University*, 17(3), 225–227 [In Chinese].
- Toy, S. J., & Newfield, M. J. (2010). The accidental introduction of invasive animals as hitchhikers through inanimate pathways: A New Zealand perspective. *Revue Scientifique et Technique (International Office of Epizootics)*, 29(1), 123–133.
- Wang, T. C. (1980). *Handbook for the determination of the common coccoids*. Chinese Academy of Sciences, p. 252.
- Wang, H. (2000). A Study on *Eulecanium Gigantea*. *Acta Agriculturae Boreali-Occidentalis Sinica*, 9, 83–86.
- Wang, Y., Zhou, Q. S., Qiao, H. J., Yu, F., Ai-B, Z., Wang, X. B., Zhu, C. D., & Zhang, Y. Z. (2016). Formal nomenclature and description of cryptic species of the *Encyrtus sasakii* complex (hymenoptera: Encyrtidae). *Scientific Reports*, 6, 34372. <https://doi.org/10.1038/srep34372>
- Wang, Y. I., Ai-H, L., Zhang, J. W., Zhao, B. J., Yue, Z. Y., Zhang, X. P., & Tang, L. (2012). Coincidence level in spatial distribution of *Eulecanium gigantea* and *Blastothrix sericae*[J]. *Journal of Zhejiang A&F University*, 29(5), 799–802. <https://doi.org/10.11833/j.issn.2095-0756.2012.05.026>
- Xie, X. X. (1985). A preliminary study on *Eulecanium gigantea*. *Scientia Silvae Sinicae (Linye Kexue)*, 21, 44–52. [In Chinese].
- Xie, Y. P., Liu, X., Li, J., & Tang, M. (1995). The effect of urban air pollution on populations of *Eulecanium gigantea* (Shinji) (Coccidae) in Taiyuan City, China. *Israel Journal of Entomology*, 29, 165–168.
- Xue, J., Xie, Y., Liu, H., Liu, J., & Li, Y. (1999). The effect of air pollution on *Sophora japonica* (Leguminosae) and *Eulecanium giganteum* (Shinji) (Hemiptera: Coccoidea: Coccidae) in urban areas in China. *Entomologica-Bari*, 33, 383–388.
- Yue, C., Zhang, J., & Zhang, X. (2011). Damage Degularity and control techniques of *Eulecanium giganteum* around Tarim Basin in Xinjiang. *Procedia Engineering*, 18, 133–138. <https://doi.org/10.1016/j.proeng.2011.11.021>
- Zhang, P., Xiao, H., Li, Q., Ma, J., & Xue, G. (2016). *Eunotus aequalivena*, an important parasitoid of *Eulecanium gigantea*. *Chinese Journal of Biological Control*, 32(1), 125–128. <https://doi.org/10.16409/j.cnki.2095-039x.2016.01.019>
- Zhang, Y. Z., & Huang, D. W. (2001). Two new Encyrtid parasites (hymenoptera: Chalcidoidea) from China. *Oriental Insects*, 35(1), 311–319. <https://doi.org/10.1080/00305316.2001.10417310>
- Zhao, X., & Xie, Y. (2004). Morphological characteristics of the different developmental stages of the male scale insect, *Eulecanium giganteum*. *Entomological Knowledge*, 41, 60–64.
- Zhu, X. S., Wang, D. Y., & Yu, J. N. (2014). Genetic variation of *Eulecanium giganteum* (Shinji) geographical populations and its correlations with ecological factors. *Chinese Journal of Ecology*, 33, 1267–1273.

How to cite this article: EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L. ... MacLeod, A. (2024). Pest categorisation of *Eulecanium giganteum*. *EFSA Journal*, 22(4), e8666. <https://doi.org/10.2903/j.efsa.2024.8666>

APPENDIX A

***Eulecanium giganteum* host plants/species affected**

Source: García Morales et al. (2016) (ScaleNet, online), and literature.

Plant family	Host name	Common name	Reference ^A
Anacardiaceae	<i>Spondias pinnata</i>	Andaman mombin, Indian hog plum, Indian mombin, wild mango	García Morales et al. (2016)
Asteraceae	<i>Taraxacum mongolicum</i>	Dandelion	García Morales et al. (2016)
Betulaceae	<i>Corylus heterophylla</i>	Japanese hazel, Siberian filbert, Siberian hazel	García Morales et al. (2016)
Betulaceae	<i>Corylus sieboldiana</i>	Japanese filbert	García Morales et al. (2016)
Cannabaceae	<i>Celtis tetrandra</i>	Nilgiri elm	García Morales et al. (2016)
Elaeagnaceae	<i>Elaeagnus angustifolia</i>	Oleaster, Russian olive, Trebizond date, wild olive	García Morales et al. (2016)
Fabaceae	<i>Albizia julibrissin</i>	Persian acacia, pink siris, silk tree, varay cotton	García Morales et al. (2016)
Fabaceae	<i>Amorpha fruticosa</i>	Bastard indigo, desert false indigo, false indigo, indigo bush	García Morales et al. (2016)
Fabaceae	<i>Caragana sinica</i>	Chinese Peashrub	García Morales et al. (2016)
Fabaceae	<i>Gleditsia sinensis</i>	Chinese honey locust	García Morales et al. (2016)
Fabaceae	<i>Glycyrrhiza uralensis</i>	Chinese liquorice	García Morales et al. (2016)
Fabaceae	<i>Halimodendron halodendron</i>	Russian salt tree, Siberian salt tree	García Morales et al. (2016)
Fabaceae	<i>Maackia amurensis</i>	Amur maackia	García Morales et al. (2016)
Fabaceae	<i>Robinia hispida</i>	Bristly locust, moss locust	García Morales et al. (2016)
Fabaceae	<i>Robinia pseudoacacia</i>	Black locust, false acacia, locust, locust tree, robinia	García Morales et al. (2016)
Fabaceae	<i>Styphnolobium japonicum</i> *	Japanese pagoda tree, pagoda tree, Chinese scholar tree	Deng et al. (2016), García Morales et al. (2016)
Fabaceae	<i>Wisteria sinensis</i>	Chinese wisteria, purple wisteria	Deng et al. (2016), García Morales et al. (2016)
Fagaceae	<i>Quercus acutissima</i>	Japanese chestnut oak, sawtooth oak	García Morales et al. (2016)
Fagaceae	<i>Quercus mongolica</i>	Mongolian oak	García Morales et al. (2016)
Juglandaceae	<i>Juglans mandshurica</i>	Manchurian walnut	García Morales et al. (2016)
Juglandaceae	<i>Juglans regia</i>	Common walnut, Persian walnut, walnut	Deng et al. (2016), García Morales et al. (2016)
Lythraceae	<i>Lagerstroemia indica</i>	Cannonball, carrion tree, crepe myrtle, Indian crape myrtle, June rose, lilac of the south	García Morales et al. (2016)
Lythraceae	<i>Punica granatum</i>	Carthaginian apple, pomegranate	García Morales et al. (2016)
Magnoliaceae	<i>Magnolia denudata</i>	Magnolia yulan, yulan	García Morales et al. (2016)
Magnoliaceae	<i>Magnolia kobus</i>	Northern Japanese magnolia	García Morales et al. (2016)
Malvaceae	<i>Hibiscus rosa-sinensis</i>	China rose, Chinese hibiscus, Chinese rose, Hawaiian hibiscus, rose mallow, rose of China, shoe-black plant, shoe-flower	Suganthi et al. (2022)
Moraceae	<i>Broussonetia papyrifera</i>	Common paper mulberry, paper mulberry, tapa-cloth tree	Deng et al. (2016), García Morales et al. (2016)
Moraceae	<i>Ficus carica</i>	Common fig, fig	García Morales et al. (2016)
Moraceae	<i>Morus alba</i>	Silkworm mulberry, white mulberry	García Morales et al. (2016)
Oleaceae	<i>Fraxinus bungeana</i>	Bunge ash	García Morales et al. (2016)
Oleaceae	<i>Fraxinus chinensis</i>	Chinese ash	García Morales et al. (2016)
Oleaceae	<i>Ligustrum quihoui</i>	Waxyleaf privet	García Morales et al. (2016)
Platanaceae	<i>Platanus orientalis</i>	Chenar tree, oriental plane	García Morales et al. (2016)
Poaceae	<i>Stipa splendens</i>	Chee grass	García Morales et al. (2016)
Rhamnaceae	<i>Ziziphus jujuba</i>	Chinese date, Chinese jujube, common jujube, Indian jujube, Indian plum, jujube tree	Deng et al. (2016), García Morales et al. (2016), Zhu et al. (2014)
Rosaceae	<i>Cydonia oblonga</i>	Quince	García Morales et al. (2016)
Rosaceae	<i>Prunus armeniaca</i> **	Apricot	Deng et al. (2016), García Morales et al. (2016)

Plant family	Host name	Common name	Reference ^A
Rosaceae	<i>Prunus cerasifera</i> var. <i>divaricata</i>		Jalilvand et al. (2013)
Rosaceae	<i>Rosa</i>	Rose	Deng et al. (2016), García Morales et al. (2016)
Rosaceae	<i>Sorbaria kirilowii</i>	Chinese sorbaria	García Morales et al. (2016)
Rutaceae	<i>Zanthoxylum bungeanum</i>	Chinese pepper tree, Chinese prickly ash, flat-spine prickly ash	García Morales et al. (2016)
Salicaceae	<i>Populus tomentosa</i>	Chinese white poplar	Deng et al. (2016), García Morales et al. (2016)
Salicaceae	<i>Salix</i>		García Morales et al. (2016)
Salicaceae	<i>Salix babylonica</i>	Chinese willow, mourning willow, Peking willow, weeping willow	Deng et al. (2016), García Morales et al. (2016)
Sapindaceae	<i>Acer buergerianum</i>	Trident maple	García Morales et al. (2016)
Sapindaceae	<i>Acer elegantulum</i>	Elegant maple	Deng et al. (2016), García Morales et al. (2016)
Sapindaceae	<i>Acer negundo</i>	Ash-leaf maple, ash-leaved maple, box elder, Manitoba maple	García Morales et al. (2016)
Sapindaceae	<i>Acer oliverianum</i> ssp. <i>formosanum</i>	Oliver's maple	García Morales et al. (2016)
Sapindaceae	<i>Acer pictum</i>	Yellow-paint maple	García Morales et al. (2016)
Sapindaceae	<i>Koelreuteria paniculata</i>	Chinese varnish tree, golden rain, pride of India	Deng et al. (2016), García Morales et al. (2016)
Sapindaceae	<i>Xanthoceras sorbifolium</i> ^{***}	Chinese flowering chestnut, goldenhorn, shiny-leaved yellowhorn, yellowhorn	García Morales et al. (2016)
Ulmaceae	<i>Ulmus</i>		García Morales et al. (2016)
Ulmaceae	<i>Ulmus macrocarpa</i>	Large-fruited elm	Zhang et al. (2016), García Morales et al. (2016)
Ulmaceae	<i>Ulmus pumila</i>	Dwarf Asiatic elm, Siberian elm	Deng et al. (2016), García Morales et al. (2016)
Vitaceae	<i>Vitis vinifera</i>	Common grapevine	García Morales et al. (2016)

^AReported as *Sophora japonica*.

**Reported as *Armeniaca vulgaris*.

***Reported as *Xanthoceras sorbifolia*.

APPENDIX B

Distribution of *Eulecanium giganteum*

Distribution records based on García Morales et al. (ScaleNet, online) and literature (Deng et al., 2016; Jalilvand et al., 2013; Suganthi et al., 2022; Zhu et al., 2014).

Region	Country	Sub-national (e.g. state)	Status
Asia	China	Anhui	Present, no details
		Beijing	Present, no details
		Lanzhou, Gansu	Present, no details
		Hebei	Present, no details
		Henan	Present, no details
		Hunan	Present, no details
		Liaoning	Present, no details
		Nei Mongol	Present, no details
		Ningxia	Present, no details
		Qinghai	Present, no details
		Shaanxi	Present, no details
		Shandong	Present, no details
		Taiyuan, Shanxi	Present, no details
		Hami, Xinjiang	Present, no details
		Bazhou, Xinjiang	Present, no details
		Aksu, Xinjiang	Present, no details
		Kashgar, Xinjiang	Present, no details
		Khotan, Xinjiang	Present, no details
		Heilongjiang	Present, no details
	Jilin	Present, no details	
	India	Thyagaraja Nagar, Chennai	Present, no details
	Iran	Kermanshah	Present, no details
	Japan	Honshu	Present, no details
		Morioka*	Present, no details
	Russia (non-European)	Primorsky Krai*	Present, no details

* Deng et al. (2016) declare that in the last decades, the pest had not been reported to be present in Morioka, Japan and Primorsky Krai, non-European Russia. However, this reference is almost 10 years old and there is no more recent information.

APPENDIX C

Import data

TABLE C.1 Edible fruit or nut trees, shrubs and bushes, whether or not grafted) imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China	404.63	642.61	305.32	31.80	1.68
India		0.22		0.03	
Iran, Islamic Republic of		8.17			
Japan	0.95	41.26	0.55	0.40	139.38
Russian Federation (Russia)	2.08	189.39		1777.42	112.21

TABLE C.2 Conifer and evergreen outdoor trees, shrubs and bushes, incl. their roots (excl. with bare roots, cuttings, slips, young plants and fruit, nut and forest trees) imported in 100 kg into the EU (27) from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China	68.30		14.50	36.58	
Japan	735.11	705.20	437.92	303.18	202.38
Russian Federation (Russia)				295.54	

TABLE C.3 Grapes, fresh or dried imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 07 November 2023).

Country	2018	2019	2020	2021	2022
China	87,690.22	191,986.55	156,789.04	80,255.76	23,689.94
India	741,303.06	970,130.19	767,803.65	852,065.11	896,702.51
Iran, Islamic Republic of	101,488.05	165,329.68	201,689.92	298,066.29	114,871.64
Japan	1.52	1.19	21.09	34.49	4.99
Russian Federation (Russia)	1.00	0.71	16.90	7.85	25.88

TABLE C.4 Roses, whether or not grafted imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China	2510.23	623.75	3.01	623.10	0.00
India	17.18	17.67	17.80	24.68	0.05
Japan	0.01	0.15	0.85	0.02	
Russian Federation (Russia)	2.50			7.25	

TABLE C.5 Fresh or dried figs imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China	340.30	192.97	55.21	141.58	250.59
India	15.49	20.64	8.03	1.63	0.14
Iran, Islamic Republic of	780.01	540.56	1055.88	718.26	542.18
Japan		0.00			0.03
Russian Federation (Russia)				0.01	0.00

TABLE C.6 Fresh quinces imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China			178.68	0.01	
India					
Iran, Islamic Republic of			21.75		
Japan					
Russian Federation (Russia)					

TABLE C.7 Apricots, cherries, peaches incl. nectarines, plums and sloes, fresh imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China	0.90	3.24		0.14	19.79
India	0.45		0.00	3.76	0.81
Iran, Islamic Republic of	42.15	29.18	589.22	381.90	123.82
Japan	1.00	2.82		37.40	4.50
Russian Federation (Russia)	218.50	693.16			

TABLE C.8 Fresh blackberries, mulberries and loganberries imported in 100 kg into the EU from regions where *Eulecanium giganteum* is known to occur (Source: Eurostat accessed on 7 November 2023).

Country	2018	2019	2020	2021	2022
China	53.91	20.20		0.00	88.39
Iran, Islamic Republic of		0.80		0.06	
Russian Federation (Russia)	4.43				

APPENDIX D

World distribution of Köppen–Geiger climate types that occur in the EU and which occur in countries where *E. giganteum* has been reported (including Cfb climate type)

