

Risk of hepatitis B virus reactivation in oncological patients treated with tyrosine kinase inhibitors: A case report and literature analysis

Francesca Colapietro, Nicola Pugliese, Antonio Voza, Alessio Aghemo, Stella De Nicola

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Lv J, China

Received: December 15, 2023

Peer-review started: December 15, 2023

First decision: January 4, 2024

Revised: January 11, 2024

Accepted: February 18, 2024

Article in press: February 18, 2024

Published online: March 7, 2024



Francesca Colapietro, Nicola Pugliese, Antonio Voza, Alessio Aghemo, Department of Biomedical Sciences, Humanitas University, Milan 20072, Italy

Francesca Colapietro, Nicola Pugliese, Alessio Aghemo, Stella De Nicola, Division of Internal Medicine and Hepatology, Department of Gastroenterology, Humanitas Research Hospital, Milan 20089, Italy

Antonio Voza, Department of Emergency, Humanitas Research Hospital, Milan 20089, Italy

Corresponding author: Francesca Colapietro, MD, Doctor, Department of Biomedical Sciences, Humanitas University, Via Rita Levi Montalcini 4, Milan 20072, Italy.
francesca.colapietro@humanitas.it

Abstract

Hepatitis B virus (HBV) reactivation (HBVr) represents a severe and potentially life-threatening condition, and preventive measures are available through blood test screening or prophylactic therapy administration. The assessment of HBVr traditionally considers factors such as HBV profile, including hepatitis B surface antigen (HBsAg) and antibody to hepatitis B core antigen, along with type of medication (chemotherapy; immunomodulants). Nevertheless, consideration of possible patient's underlying tumor and the specific malignancy type (solid or hematologic) plays a crucial role and needs to be assessed for decision-making process.

Key Words: Chronic hepatitis B; Reactivation; Nucleoside analogue; Tyrosine kinase inhibitors; Onco-hematology

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Hepatitis B virus reactivation (HBVr) is a clinical challenge among patients receiving chemotherapy for solid tumors or hematologic malignancies. The emergence of novel immunosuppressive and immunomodulatory agents requires expertise in delineating the risk of HBVr associated with each drug class. Classifying the risk of HBVr into low (< 1%), intermediate (1%-10%) and high (> 10%) allows physicians to understand in whom nucleos(t)ide analogues (NAs) are required to avoid potential progression to liver failure and death. To note, according to guidelines, patients without immediate indication for NAs should undergo serial monitoring of blood test for transaminases and HBV profile, including hepatitis B surface antigen status and HBV-DNA titer.

Citation: Colapietro F, Pugliese N, Voza A, Aghemo A, De Nicola S. Risk of hepatitis B virus reactivation in oncological patients treated with tyrosine kinase inhibitors: A case report and literature analysis. *World J Gastroenterol* 2024; 30(9): 1253-1256

URL: <https://www.wjgnet.com/1007-9327/full/v30/i9/1253.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i9.1253>

TO THE EDITOR

We were intrigued by the study conducted by Mak *et al*[1], which aimed to provide an updated guidance for monitoring or initiating antiviral prophylaxis in patients at risk of developing hepatitis B virus reactivation (HBVr). When focusing on Bruton tyrosine kinase (BTK) inhibitors, successful in treating various lymphoid malignancies, the authors emphasized the current absence of guidelines for prophylaxis and management of HBVr. Existing data primarily stems from retrospective studies and show HBVr rates ranging up to 8.3% in previously resolved infection. We agree with authors' perspective that administering anti-HBV prophylactic treatment with nucleos(t)ide analogues (NAs) is advisable for patients receiving tyrosine kinase inhibitors (TKIs) for hematologic malignancies who exhibit either positive hepatitis B surface antigen (HBsAg) or resolved HBV infection with detectable HBV DNA.

Papatheodoridis *et al*[2] recently raised awareness on the topic by conducting an analysis of available literature, identifying 4 studies including 268 chronic hepatitis B patients treated with TKI. Overall, 196 HBsAg+ patients and 72 HBsAg-/antibody to hepatitis B core antigen (antiHBc)+ patients were included; to note, HBV DNA status was not reported. The pooled rate of HBVr in HBsAg+ patients was 21/196 (11%), with no HBVr observed in the cohort of HBsAg-/antiHBc+ patients who did not receive NA prophylaxis. Conversely, among HBsAg+ patients who did not receive NA treatment, 16 cases of HBVr-associated hepatitis were observed. No cases of HBVr hepatic decompensation or death were reported. Based on these figures, the authors recommend NA prophylaxis for HBsAg+ patients receiving TKI (high risk of HBVr), while close monitoring and on-demand NA therapy is warranted in HBsAg- (low risk of HBVr)[2].

Next to these papers, several cases of TKI-associated HBVr in HBsAg- patients with hematologic malignancies have been reported (Table 1). Innocenti *et al*[3], in a cohort of 108 chronic lymphocytic leukemia (CLL) patients, reported two cases of HBVr (1.9%) among HBsAg-/anti-HBc+ patients within the initial 6 months of second-line treatment with ibrutinib, a covalent BTK[3]. None of these cases experienced HBVr associated hepatitis, and both individuals responded effectively to entecavir treatment. Additionally, Chiu *et al*[4] analyzed a series of 29 patients treated with TKIs, reporting 3 cases (10%) of HBVr, with 2 occurring in HBsAg-/anti-HBc+ patients receiving BTK inhibitors. Notably, all 3 experienced HBVr hepatitis, with 2 developing liver failure; all cases recovered with anti-HBV therapy[4]. To note, several cases of HBVr have been reported in patients with solid tumors. Lee *et al*[5] described the largest retrospective cohort, comprising 1960 anti-HBc+ patients with lung cancer treated with TKI. Among them, 1594 were HBsAg- and 521 received TKI as first-line treatment. One patient developed HBVr characterized by the reappearance of HBsAg, elevation of alanine aminotransferase (ALT) more than 10 folds upper limit of normal (ULN) and HBV DNA reaching up to 245000 IU/mL (undetectable at baseline).

We present here the case of a 67-year-old Caucasian male patient with CLL treated with acalabrutinib, a Food and Drug Administration-approved next-generation covalent BTK inhibitor[6]. His medical history included prior treatment for colorectal cancer through surgical intervention for bowel obstruction (TNMv8: pT4b N0 R0).

As the patient was HBsAg-/anti-HBc+, close monitoring of liver function blood tests and HBV profile was started. To note, HBV DNA was undetectable before starting treatment. After 8 months of therapy, a mild elevation of transaminases up to three times ULN was observed, with no coagulation impairment or elevation of total bilirubin. Subsequently, one week later the patient developed hepatic liver failure characterized by jaundice (total bilirubin 23 mg/dL), prothrombin time > 1.7 and ascites. At this point HBV DNA was 8.9 log IU/mL, ALT 497 U/L (ULN 50 U/L), aspartate aminotransferase 493 U/L (ULN 50 U/L); nucleos(t)ide analog treatment was started immediately. Two weeks following the onset of HBVr, HBV DNA reduced to 4.9 log IU/mL. Ascites was well controlled, no signs of hepatic encephalopathy developed, but bilirubin persisted at 43 mg/dL and the patient remained hospitalized. Acalabrutinib treatment was stopped.

Based on these reports, HBsAg-/antiHBc+ patients receiving TKI in the onco-hematological setting should be reclassified as intermediate risk for HBVr warranting consideration for antiviral prophylaxis.

We advocate for regular updates on the risk of HBVr associated to specific drug classes. Moreover, we think that type of tumor (solid or hematologic) should be analyzed as a possible risk factor for deciding when to start antiviral prophylaxis.

Table 1 Studies reporting reactivation of hepatitis B virus in hepatitis B surface antigen -/antibody to hepatitis B core antigen + patients treated with tyrosine kinase inhibitors for hematologic malignancies

Ref.	Type of study	TKI	Overall cohort (n)	HBV-r (n)	HBVr-associated hepatitis (n)	Liver failure
Papatheodoridis et al[2], 2022	Meta-analysis	NS	72	0	0	0
Innocenti et al[3], 2022	Retrospective	Ibrutinib	108	2	0	0
Chiu et al[4], 2023	Retrospective	BTKs	29 ¹	2	2	2
Hammond et al[7], 2018	Retrospective	Ibrutinib	21	2	0	0
Tsuruya K et al[8], 2021	Case report	Ibrutinib	/	1	1	0
Lam et al[9], 2023	Case report	Ibrutinib	/	1	1	0

¹Including both past and chronic hepatitis B virus infection.

BTK: Bruton tyrosine kinase; NS: Not specified; TKI: Tyrosine kinase inhibitors; HBV: Hepatitis B virus; HBVr: Hepatitis B virus reactivation.

FOOTNOTES

Author contributions: Colapietro F, Aghemo A, and De Nicola S contributed to the study conception and design; Colapietro F and De Nicola S were involved in the data collection; Colapietro F, Pugliese N, Voza A, Aghemo A, and De Nicola S participated in the analysis and interpretation of results, and draft manuscript preparation; and all authors reviewed the results and approved the final version of the manuscript.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Italy

ORCID number: Francesca Colapietro 0000-0002-7520-744X; Nicola Pugliese 0000-0001-6466-1412; Alessio Aghemo 0000-0003-0941-3226.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Yu HG

REFERENCES

- Mak JWY, Law AWH, Law KWT, Ho R, Cheung CKM, Law MF. Prevention and management of hepatitis B virus reactivation in patients with hematological malignancies in the targeted therapy era. *World J Gastroenterol* 2023; **29**: 4942-4961 [PMID: 37731995 DOI: 10.3748/wjg.v29.i33.4942]
- Papatheodoridis GV, Lekakis V, Voulgaris T, Lampertico P, Berg T, Chan HLY, Kao JH, Terrault N, Lok AS, Reddy KR. Hepatitis B virus reactivation associated with new classes of immunosuppressants and immunomodulators: A systematic review, meta-analysis, and expert opinion. *J Hepatol* 2022; **77**: 1670-1689 [PMID: 35850281 DOI: 10.1016/j.jhep.2022.07.003]
- Innocenti I, Reda G, Visentin A, Coscia M, Motta M, Murru R, Moia R, Gentile M, Pennese E, Quaglia FM, Albano F, Cassin R, Deodato M, Ielo C, Frustaci AM, Picciocchi A, Rughini A, Arena V, Di Sevo D, Tomasso A, Autore F, Del Poeta G, Scarfò L, Mauro FR, Tedeschi A, Trentin L, Pompili M, Foà R, Ghia P, Cuneo A, Laurenti L. Risk of hepatitis B virus reactivation in chronic lymphocytic leukemia patients receiving ibrutinib with or without antiviral prophylaxis. A retrospective multicentric GIMEMA study. *Haematologica* 2022; **107**: 1470-1473 [PMID: 35199505 DOI: 10.3324/haematol.2021.280325]
- Chiu CY, Ahmed S, Thomas SK, Wang LS, Mustafayev K, Fayad LE, Wierda WG, Khawaja F, Torres HA. Hepatitis B Virus Reactivation in Patients Receiving Bruton Tyrosine Kinase Inhibitors. *Clin Lymphoma Myeloma Leuk* 2023; **23**: 610-615 [PMID: 37150651 DOI: 10.1016/j.clml.2023.04.006]
- Lee PH, Huang YH, Hsu YW, Chen KC, Hsu KH, Lin H, Lee TY, Tseng JS, Chang GC, Yang TY. Reactivation of Hepatitis B Virus in Lung Cancer Patients Receiving Tyrosine Kinase Inhibitor Treatment. *J Clin Med* 2022; **12** [PMID: 36615034 DOI: 10.3390/jcm12010231]
- AstraZeneca. Calquence tablet formulation approved in the US across current indications. 2022. [cited 25 September 2023]. Available from: <https://www.astrazeneca.com/media-centre/press-releases/2022/calquence-tablet-formulation-approved-in-the-us-across-current-indications.html>
- Hammond SP, Chen K, Pandit A, Davids MS, Issa NC, Marty FM. Risk of hepatitis B virus reactivation in patients treated with ibrutinib. *Blood* 2018; **131**: 1987-1989 [PMID: 29490923 DOI: 10.1182/blood-2018-01-826495]

- 8 **Tsuruya K**, Anzai K, Shioyama S, Ito A, Arase Y, Hirose S, Tanaka Y, Suzuki H, Kagawa T. Case of hepatitis B virus reactivation after ibrutinib therapy in which the patient remained negative for hepatitis B surface antigens throughout the clinical course. *Hepatol Res* 2021; **51**: 239-244 [PMID: 32978866 DOI: 10.1111/hepr.13575]
- 9 **Lam LK**, Chan TSY, Hwang YY, Mak LY, Seto WK, Kwong YL, Yuen MF. Hepatitis B virus reactivation in seronegative occult hepatitis B patient receiving ibrutinib therapy. *Virol J* 2023; **20**: 168 [PMID: 37528444 DOI: 10.1186/s12985-023-02140-w]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

