



Clinical Article

Transcranial Doppler emboli monitoring for stroke prevention after flow diverting stents

Matias Costa¹, Paul Schmitt², Jaleel N^{1,3}, Matias Baldoncini⁴, Juan Vivanco-Suarez¹, Bipin Chaurasia⁵, Colleen Douville¹, Loh Yince¹, Akshal Patel¹, Stephen Monteith¹

¹Department of Neurosurgery, Swedish Neuroscience Institute, Seattle, Washington, USA

²Department of Neurosurgery, University of Texas Medical Branch, Texas, USA

³Division of Neurosurgery, Dartmouth-Hitchcock Medical Center, New Hampshire, USA

⁴Department of Neurological Surgery, Hospital San Fernando, Buenos Aires, Argentina

⁵Department of Neurosurgery, Neurosurgery Clinic, Birgunj, Nepal

J Cerebrovasc Endovasc Neurosurg.
2024 March;26(1):23-29

Received: 11 May 2023

Revised: 7 July 2023

Accepted: 7 July 2023

Correspondence to
Matias Costa

Department of Neurosurgery, Swedish
Neuroscience Institute, 550 16th Ave, Seattle,
Washington 98122, USA

Tel +1-206-8253810

E-mail matiascostamd@gmail.com

ORCID <https://orcid.org/0000-0003-1833-4589>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective: Flow diverting stents (FDS) are increasingly used for the treatment of intracranial aneurysms. While FDS can provide flow diversion of parent vessels, their high metal surface coverage can cause thromboembolism. Transcranial Doppler (TCD) emboli monitoring can be used to identify subclinical embolic phenomena after neurovascular procedures. Limited data exists regarding the use of TCDs for emboli monitoring in the perioperative period after FDS placement. We evaluated the rate of positive TCDs microembolic signals and stroke after FDS deployment at our institution.

Methods: We retrospectively evaluated 105 patients who underwent FDS treatment between 2012 and 2016 using the Pipeline stent (Medtronic, Minneapolis, MN, USA). Patients were pretreated with aspirin and clopidogrel. All patients were therapeutic on clopidogrel pre-operatively. TCD emboli monitoring was performed immediately after the procedure. Microembolic signals (mES) were classified as “positive” (<15 mES/hour) and “strongly positive” (>15 mES/hour). Clinical stroke rates were determined at 2-week and 6-month post-operatively.

Results: A total of 132 intracranial aneurysms were treated in 105 patients. TCD emboli monitoring was “positive” in 11.4% (n=12) post-operatively and “strongly positive” in 4.8% (n=5). These positive cases were treated with heparin drips or modification of the antiplatelet regimen, and TCDs were repeated. Following medical management modifications, normalization of mES was achieved in 92% of cases. The overall stroke rates at 2-week and 6-months were 3.8% and 4.8%, respectively.

Conclusions: TCD emboli monitoring may help early in the identification of thromboembolic events after flow diversion stenting. This allows for modification of medical therapy and, potentially, prevention of escalation into post-operative strokes.

Keywords Transcranial Doppler Ultrasonography, Endovascular stent, Flow diverter, Intracranial aneurysm

INTRODUCTION

Flow diversion is becoming an increasingly used strategy for the treatment of complex intracranial aneurysms.⁴⁽⁶⁾¹²⁾¹⁵⁾¹⁷⁾ The pipeline embolization device (PED, Medtronic, Minneapolis, MN, USA) is a flow diverting stent that provides up to 35% metal coverage over the parent vessel. It works by disrupting intra-aneurysmal blood flow, therefore promoting stasis and aneurysmal thrombosis. Over time, the stent becomes endothelialized resulting in aneurysm neck obliteration and a durable aneurysm cure.⁶⁾⁷⁾

The high metal surface of a PED is associated with thrombo-embolic complications.³⁾ Dual antiplatelet agents (aspirin in combination with one of the P2Y12 inhibitors - clopidogrel, prasugrel, or ticagrelor) have been used to inhibit platelet activation.¹⁰⁾ Aspirin and clopidogrel response assays can be used before and after interventions to confirm platelet inhibition. However, the association between extent of platelet inhibition and thromboembolic events is not clear. As such, there is significant variability in the practice patterns of neuro-interventionalists in the use of platelet inhibition assays prior to deployment of PEDs.¹⁰⁾¹¹⁾

In this study, we analyzed the use of transcranial Doppler (TCD) monitoring to evaluate microembolic events after flow diversion.¹⁾ Intracranial microembolic signals (mES) are predictive of strokes both de novo⁹⁾²²⁾ and following neurointerventional procedures.¹³⁾¹⁹⁾ For instance, mES are higher in patients with carotid disease and cardioembolic sources when compared to healthy individuals.⁸⁾ Additionally, microemboli monitoring has been used to minimize stroke during and after carotid endarterectomy and cardiac valve surgeries.¹⁶⁾ No data exists regarding the use of TCDs for emboli monitoring in the periprocedural period after flow diverting stent placement. In this study, we present our institutional experience with microemboli monitoring and stroke rates after pipeline embolization of intracranial aneurysms.

MATERIALS AND METHODS

Study design and population

Medical records of our institute were retrospectively reviewed from 110 consecutive patients with intracranial aneurysms treated with the PED between 2012 and 2016. Demographic data and clinical data were analyzed.

Inclusion criteria included all patients with unruptured, or previously coiled aneurysms treated with Pipeline stent (Medtronic, Minneapolis, MN, USA). To minimize confounding variables in the study, we excluded patients who underwent PED treatment for aneurysmal subarachnoid hemorrhage, pseudoaneurysms, or residual aneurysms that were previously stented. Based on this criterion, we excluded five patients and analyzed a total of 105 patients.

Statistical analysis was performed with Prism 9.0.1 software (GraphPad Software, Inc., San Diego, CA, USA). For continuous variables the Mann-Whitney test was used. For categorical variables, chi-square test was employed. A p value <0.05 was considered significant.

Treatment protocol

Our PED deployment protocol has been described previously.¹⁵⁾ Patients were pre-treated with aspirin (81 mg) and clopidogrel (75 mg) for at least 10 days prior to treatment. Therapeutic levels were determined using the Accumetrics VerifyNow P2Y12 assay. P2Y12 reaction units (PRU) <208 were considered therapeutic on clopidogrel. Patients with PRU >208 were considered clopidogrel-non-responders and started on ticagrelor. Baseline and post-clopidogrel/ticagrelor treatment PRU values were used to calculate % platelet inhibition: $1 - (\text{treatment PRU} / \text{baseline PRU})$. The platelet inhibition goal was 30-80% activity.

All patients were adequately heparinized during the procedure with activated clotting times greater than 280 seconds.

Patients were kept on a regimen of aspirin for life, and clopidogrel or ticagrelor for at least 6 months. PEDs were deployed using a standard triaxial system. Control digital subtraction angiography was performed imme-

diately following PED placement. A follow-up angiography was performed at 6 months post-treatment.

Transcranial Doppler studies

TCD emboli monitoring was performed 6 hours and prior to 24-hours post-operatively. The Transcranial Power M-mode transcranial Doppler system (PMD 100 or ST3; Spencer Technologies, Seattle, WA, USA) with a 2-MHz digital Doppler was used.²¹⁾ All exams were performed by registered vascular sonographers specializing in neurovascular sonography. Criteria for mES identification were: 1) characteristic acoustic properties, 2) short duration (<0.3 ms), and 3) random appearance during cardiac cycle. Emboli monitoring was performed continuously for at least 20 minutes in the arterial segment distal to the aneurysm.

Management of patients with positive TCD emboli

All cases with positive mES underwent repeated monitoring withing 6-24 hours. If the initial two scans showed <15 mES/hr, no further studies were performed. If “strongly positive” rates (>15 mES/hr) were noted, TCDs were repeated within 6-hours after medical treatment modification consisting of either heparinization, additional clopidogrel load, or pharmacological switch from clopidogrel to ticagrelor. TCDs were repeated until two stable scans showed <15 mES/hr.

RESULTS

Between October 2012 and July 2020, 110 patients were treated with PEDs for unruptured or previously coiled intracranial aneurysms. Aneurysmal subarachnoid hemorrhage (SAH) and pseudoaneurysms were excluded (n=5). Based on these criteria, we studied 105 patients with 132 aneurysms. The mean patient age was 59.7 years, with the majority (78.1%) being female (Table 1). The mean aneurysm size was 7.8±6.4 mm. The paraclinoid region was the most frequent aneurysm location (Table 2). Two aneurysms in the petrous segment were treated due to intradural aneurysmal extension.

Table 1. Patient demographics

Characteristics	Value
Patients analyzed	105
Mean age (years)	59.7
Gender	
Female	82 (78.1%)
Male	23 (21.9%)

Table 2. Aneurysm characteristics

Variable	Value
Number of aneurysm treated	132
Mean size (mm)	7.8±6.4
Location (n)	
Petrous	2
Cavernous	15
OphA	37
Paraclinoid/sup hypo	40
PCOM	23
Vertebrobasilar	15
Type	
Saccular	124
Fusiform	8

OphA, ophthalmic artery; PCOM, posterior communicating artery

Most cases involved deployment of a single PED (N=93, 88.6%). In twelve cases (11.4%), two PEDs were deployed. Adjunctive aneurysm coiling was performed in fifteen cases (14.3%) (Table 3), and was associated with a higher incidence of positive TCD results (not statistically significant).

Table 4 summarizes the TCD emboli monitoring results from all cases along with associated aneurysm size, number of PEDs deployed, treatment PRU as well as

Table 3. Aneurysm treatment

#PEDs deployed	No. of patients (%)	% with positive TCD
1	93 (88.6)	11.8
2	12 (11.4)	8.3
3	0 (0)	0
Coiling+PED	16 (15.2)	25
PED only	89 (84.8)	9.0

PED, pipeline embolization device; TCD, transcranial Doppler

Table 4. TCD emboli monitoring and perioperative stroke after PED aneurysm treatment

TCD result	N	%	Aneurysm size	#PEDs	PRU	% inhibition	Stroke (N)	
							Peri-op	Follow-up
Positive	12	11.4	11.0±6.7	1.1±0.3	144.1±46.6	38.2±20.1	0	0
<15	7	6.7	11.0±6.4	1.1±0.4	142.1±57.7	36.6±22.1	1	0
15-50	2	1.9	16.5±12.0	1±0	164.5±48.8	28.6±2.1	0	0
>50	3	2.9	7.2±0.8	1±0	138.7±33.5	50.1±19.1	0	0
Negative	86	81.9	8.9±6.8	1.1±0.3	113.2±59.0	58.3±20.6	2	1
Inadequate windows	7	6.7	7.5±3.4	1±0	114.6±45.9	65.8±3.5	1	1

TCD, transcranial Doppler; PED, pipeline embolization device; PRU, P2Y12 reaction units

% platelet inhibition. Inadequate transcranial windows prevented TCD emboli monitoring in seven cases (6.5%). Patients with positive TCD emboli (N=12, 11.4%) had an average pre-procedure PRU of 144.1±46.6 and % platelet inhibition of 38.2±20.1%. Patients with negative TCD emboli (N=86, 81.9%) had an average pre-procedure PRU of 113.2±59.0 with a % platelet inhibition of 58.3±20.6%. Significantly less % platelet inhibition was evident in patients with positive mES when compared to patients with negative mES (38.2±20.1% vs 58.3±20.6%, P<0.02). Positive TCD results did not correlate with aneurysm size or number of PEDs deployed (Tables 3 and 4). Table 4 also shows that, after medical interventions were initiated to prevent on-going mES in patients with positive

TCDs, no strokes were evident upon follow-up.

Table 5 presents the 12 cases of positive mES after PED deployment along with the patient demographics, aneurysm size and treatment details. The magnitude of embolic signals ranged between 3 to 303 mES/hr. Patients with embolic signals less than 15 mES/hr were closely followed with repeat TCDs only and no additional monitoring was done in cases when repeat TCDs showed no further emboli.

In cases of strongly positive TCD signals (>15 mES/hr), patients were either started on heparin and/or reloaded with clopidogrel or switched to ticagrelor. TCDs were again checked within 6-hours to confirm cessation of embolic signals. In one case (patient 10), embolic signals

Table 5. Summary of cases that showed positive TCD embolic signals after PED deployment

Pt	Age	Gender	Aneurysm size	#PED	Coiling	PRU	% inhibition	Initial		Followup		Final		Outcome
								mES/h	Mgmt	mES/h	Mgmt	mES/h	Mgmt	
1	71	1	20	1	N	74	18.7	3	1	0	1	0	0	intact
2	72	2	17	2	Y	199	34.1	3	1	0	1	0	1	intact
3	47	1	3.5	1	N	145	39.1	3	1	0	1	0	1	intact
4	51	1	11	1	Y	119	60.2	3	1	0	1	0	1	intact
5	42	1	8	1	N	172	33.3	6	1	297	4	0	1	intact
6	63	2	8	1	Y	199	27.1	6	1	18	4	0	1	intact
7	42	2	8	1	N	152	8.4	6	1	0	1	0	1	intact
8	44	1	3.5	1	N	78	70.1	9	1	0	1	0	1	intact
9	70	2	14	1	N	217	20.8	9	4	0	1	0	1	intact
10	30	2	25	1	Y	130	30.1	18	2	12	5	18	5	intact
11	49	1	7.2	1	N	139	46.1	42	2	51	2	0	1	intact
12	48	1	6.5	1	N	105	70.8	303	2,4	0	1	0	1	intact

Mgmt: 1=Repeat TCD only, 2=Heparin, 3=Ticagrelor, 4=Reload ASA/Piavix, 5=Multiple Meds
TCD, transcranial Doppler; PED, pipeline embolization device; PRU, P2Y12 reaction units

persisted despite changes in antiplatelet regimen, and an in-stent thrombosis evolved necessitating mechanical thrombectomy. Of note, this patient was a responder to clopidogrel prior to the stent placement. Surprisingly, he transitioned to be a non-responder on immediate post-operative testing. The patient was ultimately placed on ticagrelor and coumadin which aborted further mES. This patient's outcome was favorable with a full recovery and no neurologic deficits.

Table 6 presents the five strokes (4.8%) that occurred in the 105 patients in this study. Patient 1 as described above was a clopidogrel non-responder and had persistent TCD emboli until stabilized on ticagrelor and coumadin. The next two patients had neurologic deficits in the immediate post-operative period despite negative TCDs. Patient 2 had a posterior putaminal stroke relating to a thrombotic event linked to balloon angioplasty of in-stent stenosis, but made a full recovery at 6-months follow-up. Patient 3 died within a 2-week period from a severe gastrointestinal bleeding. Of note, this patient had a very tortuous arch anatomy and underwent flow diverting stents (FDS) of a giant cavernous segment aneurysm via a cervical carotid cut down. Patient 4 had inadequate temporal windows for TCD monitoring and recovered from endovascular treatment with left hemiparesis relating to an embolic event from the use of a Simmons

catheter to navigate a difficult Type III aortic arch. Finally, patient 5 had a thrombotic complication at 5-months after she stopped clopidogrel and aspirin with resultant stent thrombosis. This patient had no TCD windows.

DISCUSSION

PEDs are widely used for complex and wide-necked intracranial aneurysms.⁴⁻⁶⁾¹²⁾¹⁵⁾¹⁷⁾ They promote flow stagnation, aneurysmal thrombosis, and parent vessel remodeling through endothelialization of the stent. Both thrombotic and embolic complications can occur after PED deployment.³⁾¹¹⁾ Flow stagnation at branch vessels covered by PEDs can cause thrombosis and occlusion of the perforating arteries.¹⁸⁾ Embolic complications can occur with dislodgement of platelet aggregates from the stent surface causing distal vessel occlusion.

TCD emboli monitoring has been shown to prevent ischemic complications during cardiac surgery¹⁶⁾ and carotid endarterectomy procedures.²⁰⁾²¹⁾²³⁾ Similar results have also been shown for emboli monitoring after intracranial aneurysm coiling.¹³⁾¹⁹⁾ These studies demonstrate that reduced microembolic events is associated with reduced ischemic events. Schubert et al. showed that

Table 6. Summary of all strokes instances after PED deployment in our series

Pt	Age	Gender	Aneurysm size	#PEDs	Coiling	PRU	TCD results	MRS				Notes
								Preop	D/C	2 wks	6 months	
1	30	2	25	1	Y	130	1	0	2	2	0	Plavix non-responder, required mechanical thrombectomy
2	72	1	10.4	1	N	163	2	0	3	2	0	Required balloon angioplasty of in stent stenosis during procedure
3	76	1	28	2	N	49	2	1	3	8	6	Underwent carotid cutdown for PED placement due to tortuous arch
4	76	1	10	1	N	81	3	0	1	1	0	Had difficult arch requiring Simmons catheter and long exchange wire
5	39	1	3	1	N	124	3	1	0	1	4	Stopped plavix at 5 months with subsequent in stent thrombosis

PED, pipeline embolization device; PRU, P2Y12 reaction units; TCD, transcranial Doppler

early heparinization was associated with reduced TCD emboli as well as reduced neurologic deficits post-procedure.¹⁹⁾

In this study, we found that 11.4% of cases (12 of 105 patients) shower positive mES after PED deployment. In 5 of 12 patients, the emboli resolved with repeat testing and no intervention. In 6 of 12 patient's additional medical interventions (including further heparinization, additional clopidogrel administration, or change in antiplatelet agent) were needed to reduce the microembolic events. In one of the twelve patients, who was clopidogrel-resistant and ticagrelor super-responder, mechanical thrombectomy was needed due to rapid in-stent thrombosis. In this patient, TCD emboli monitoring was pivotal in establishing the appropriate antiplatelet regimen. All twelve patients with microemboli after PED deployment had cessation of TCD emboli prior to hospital discharge and were neurologically intact at 30-days. Comparison of this group of patients to the group with no TCD emboli demonstrated that % platelet inhibition was the only significantly different variable predicting ischemic events.

Our study suggests that close monitoring of % platelet inhibition peri-procedurally results in reduced ischemic events.¹⁴⁾ The routine use of platelet function assays is debated in the neurovascular community.¹⁰⁾ Approximately 11.5% of neuro-interventionalists do not routinely perform platelet function testing after PED deployment. This likely reflects the absence of an established association between stroke and degree of platelet inhibition. For instance, Heller et al. reported no association between strength of antiplatelet therapy and procedural embolism after PED deployment.¹¹⁾ Whereas Brasiliense et al. and Delgado-Almandoz et al. showed that the preprocedure PRU values were most predictive of thromboembolic events.

Finally, reduction in mES is associated with reduced risk of ischemic events and stroke.⁹⁾ The overall stroke rate in our study was 3.8% at two-weeks and 4.8% at six-months, somehow comparable to the 5.6% stroke and mortality rate in the PUFs trial.²⁾ We believe that our stroke rates could have been higher if no TCD

emboli monitoring had been available, since abortion of ongoing embolic events was achieved in most cases after positive mES were detected.

CONCLUSIONS

TCD emboli monitoring represents a powerful non-invasive tool for early identification of thromboembolic events after flow diversion stenting. Early detection of microembolic signals allows for modification of post-operative medical therapy and, potentially, prevention of post-operative stroke.

Disclosure

The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

REFERENCES

1. Babikian VL, Wijman CA. Brain embolism monitoring with transcranial doppler ultrasound. *Curr Treat Options Cardio-vasc Med.* 2003 Jul;5(3):221-32.
2. Beckske T, Kallmes DF, Saatci I, McDougall CG, Szikora I, Lanzino G, et al. Pipeline for uncoilable or failed aneurysms: Results from a multicenter clinical trial. *Radiology.* 2013;267:858-68.
3. Brasiliense LB, Stanley MA, Grewal SS, Cloft HJ, Sauvageau E, Lanzino G, et al. Silent ischemic events after pipeline embolization device: A prospective evaluation with mr diffusion-weighted imaging. *J Neurointerv Surg.* 2016 Nov;8(11):1136-9.
4. Brinjikji W, Murad MH, Lanzino G, Cloft HJ, Kallmes DF. Endovascular treatment of intracranial aneurysms with flow diverters: A meta-analysis. *Stroke.* 2013 Feb;44(2):442-7.
5. Delgado Almandoz JE, Crandall BM, Scholz JM, Fease JL, Anderson RE, Kadkhodayan Y, et al. Pre-procedure P2Y12 reaction units value predicts perioperative thromboembolic and hemorrhagic complications in patients with cerebral aneurysms treated with the pipeline embolization device. *J Neurointerv Surg.* 2013 Nov;5 Suppl 3:iii3-10.
6. Fiorella D, Lylyk P, Szikora I, Kelly ME, Albuquerque FC,

- McDougall CG, et al. Curative cerebrovascular reconstruction with the pipeline embolization device: The emergence of definitive endovascular therapy for intracranial aneurysms. *J Neurointerv Surg*. 2018 Jul;10(Suppl 1):i9-18.
7. Fiorella D, Woo HH, Albuquerque FC, Nelson PK. Definitive reconstruction of circumferential, fusiform intracranial aneurysms with the pipeline embolization device. *Neurosurgery*. 2008 May;62(5):1115-20; discussion 1120-1.
 8. Georgiadis D, Lindner A, Manz M, Sonntag M, Zunker P, Zerkowski HR, et al. Intracranial microembolic signals in 500 patients with potential cardiac or carotid embolic source and in normal controls. *Stroke*. 1997 Jun;28(6):1203-7.
 9. Goertler M, Blaser T, Krueger S, Hofmann K, Baeumer M, Wallesch CW. Cessation of embolic signals after antithrombotic prevention is related to reduced risk of recurrent arterioembolic transient ischaemic attack and stroke. *J Neurol Neurosurg Psychiatry*. 2002 Mar;72(3):338-42.
 10. Gupta R, Moore JM, Griessenauer CJ, Adeeb N, Patel AS, Youn R, et al. Assessment of dual-antiplatelet regimen for pipeline embolization device placement: A survey of major academic neurovascular centers in the United States. *World Neurosurg*. 2016 Dec;96:285-92.
 11. Heller RS, Dandamudi V, Lanfranchi M, Malek AM. Effect of antiplatelet therapy on thromboembolism after flow diversion with the pipeline embolization device. *J Neurosurg*. 2013 Dec;119(6):1603-10.
 12. Kan P, Siddiqui AH, Veznedaroglu E, Liebman KM, Binning MJ, Dumont TM, et al. Early postmarket results after treatment of intracranial aneurysms with the pipeline embolization device: A U.S. multicenter experience. *Neurosurgery*. 2012 Dec;71(6):1080-7; discussion 1087-8.
 13. Klötzsch C, Nahser HC, Henkes H, Kühne D, Berlit P. Detection of microemboli distal to cerebral aneurysms before and after therapeutic embolization. *AJNR Am J Neuroradiol*. 1998 Aug;19(7):1315-8.
 14. Ku JC, Chavda V, Palmisciano P, Pasarikovski CR, Yang VX, Kiwan R, et al. Endovascular treatment for anterior inferior cerebellar artery-posterior inferior cerebellar artery (AICA-PICA) common trunk variant aneurysms: Technical note and literature review. *J Cerebrovasc Endovasc Neurosurg*. 2023 Apr.
 15. Nelson PK, Lylyk P, Szikora I, Wetzel SG, Wanke I, Fiorella D. The pipeline embolization device for the intracranial treatment of aneurysms trial. *AJNR Am J Neuroradiol*. 2011 Jan;32(1):34-40.
 16. Padayachee TS, Parsons S, Theobald R, Linley J, Gosling RG, Deverall PB. The detection of microemboli in the middle cerebral artery during cardiopulmonary bypass: A transcranial Doppler ultrasound investigation using membrane and bubble oxygenators. *Ann Thorac Surg*. 1987 Sep;44(3):298-302.
 17. Puffer RC, Kallmes DE, Cloft HJ, Lanzino G. Patency of the ophthalmic artery after flow diversion treatment of paraclinoid aneurysms. *J Neurosurg*. 2012 Apr;116(4):892-6.
 18. Rangel-Castilla L, Munich SA, Jaleel N, Cress MC, Krishna C, Sonig A, et al. Patency of anterior circulation branch vessels after pipeline embolization: Longer-term results from 82 aneurysm cases. *J Neurosurg*. 2017 Apr;126(4):1064-9.
 19. Schubert GA, Thome C, Seiz M, Douville C, Eskridge J. Microembolic signal monitoring after coiling of unruptured cerebral aneurysms: An observational analysis of 123 cases. *AJNR Am J Neuroradiol*. 2011 Sep;32(8):1386-91.
 20. Siebler M, Sitzer M, Rose G, Bendfeldt D, Steinmetz H. Silent cerebral embolism caused by neurologically symptomatic high-grade carotid stenosis. Event rates before and after carotid endarterectomy. *Brain*. 1993 Oct;116(Pt 5):1005-15.
 21. Spencer MP, Thomas GI, Nicholls SC, Sauvage LR. Detection of middle cerebral artery emboli during carotid endarterectomy using transcranial doppler ultrasonography. *Stroke*. 1990 Mar;21(3):415-23.
 22. Valton L, Larrue V, le Traon AP, Massabau P, Geraud G. Microembolic signals and risk of early recurrence in patients with stroke or transient ischemic attack. *Stroke*. 1998 Oct;29(10):2125-8.
 23. van Zuilen EV, Moll FL, Vermeulen FE, Mauser HW, van Gijn J, Ackerstaff RG. Detection of cerebral microemboli by means of transcranial doppler monitoring before and after carotid endarterectomy. *Stroke*. 1995 Feb;26(2):210-3.