

# Impact of different genetic mutations on granulocyte development and G-CSF responsiveness in congenital neutropenia

Xin Meng,<sup>1,\*</sup> Hai Zhang,<sup>2,\*</sup> Lulu Dong,<sup>1</sup> Qing Min,<sup>2</sup> Meiping Yu,<sup>2</sup> Yaxuan Li,<sup>1</sup> Lipin Liu,<sup>2</sup> Wenjie Wang,<sup>2</sup> Wenjing Ying,<sup>2</sup> Jinqiao Sun,<sup>2</sup> Ji-Yang Wang,<sup>1-3</sup> Jia Hou,<sup>2</sup> and Xiaochuan Wang<sup>2,4</sup>

<sup>1</sup>Department of Immunology, School of Basic Medical Sciences, Fudan University, Shanghai, China; <sup>2</sup>Department of Clinical Immunology, Children's Hospital of Fudan University, National Children's Medical Center, Shanghai, China; <sup>3</sup>Shanghai Sci-Tech Inno Center for Infection & Immunity, Shanghai, China; and <sup>4</sup>Shanghai Institute of Infectious Disease and Biosecurity, Shanghai, China

## Key Points

- Different forms of congenital neutropenia exhibit disruptions in granulocyte development at distinct stages.
- There is a strong correlation between the stage and severity of granulocyte development disruption and the efficacy of G-CSF therapy.

Congenital neutropenia (CN) is a genetic disorder characterized by persistent or intermittent low peripheral neutrophil counts, thus increasing susceptibility to bacterial and fungal infections. Various forms of CN, caused by distinct genetic mutations, exhibit differential responses to granulocyte colony-stimulating factor (G-CSF) therapy, with the underlying mechanisms not fully understood. This study presents an in-depth comparative analysis of clinical and immunological features in 5 CN patient groups (severe congenital neutropenia [SCN]1, SCN3, cyclic neutropenia [CyN], warts, hypogammaglobulinaemia, infections and myelokathexis [WHIM], and Shwachman-Bodian-Diamond Syndrome [SBDS]) associated with mutations in *ELANE*, *HAX1*, *CXCR4*, and *SBDS* genes. Our analysis led to the identification of 11 novel mutations in *ELANE* and 1 each in *HAX1*, *CXCR4*, and *G6PC3* genes. Investigating bone marrow (BM) granulopoiesis and blood absolute neutrophil count after G-CSF treatment, we found that SCN1 and SCN3 presented with severe early-stage disruption between the promyelocyte and myelocyte, leading to a poor response to G-CSF. In contrast, CyN, affected at the late polymorphonuclear stage of neutrophil development, showed a strong G-CSF response. WHIM, displaying normal neutrophil development, responded robustly to G-CSF, whereas SBDS, with moderate disruption from the early myeloblast stage, exhibited a moderate response. Notably, SCN1 uniquely impeded neutrophil development, whereas SCN3, CyN, WHIM, and SBDS also affected eosinophils and basophils. In addition, SCN1, SCN3, and CyN presented with elevated serum immunoglobulins, increased BM plasma cells, and higher A Proliferation-Inducing Ligand levels. Our study reveals a strong correlation between the stage and severity of granulocyte development disruption and the efficacy of G-CSF therapy.

Submitted 13 November 2023; accepted 18 January 2024; prepublished online on *Blood Advances* First Edition 29 January 2024; final version published online 28 March 2024. <https://doi.org/10.1182/bloodadvances.2023012171>.

\*X.M. and H.Z. contributed equally to this work.

Additional data related to this study, as well as specific materials and protocols, can be obtained from the corresponding author, Ji-Yang Wang ([wang@fudan.edu.cn](mailto:wang@fudan.edu.cn)) upon reasonable request.

The full-text version of this article contains a data supplement.

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

## Introduction

Congenital neutropenia (CN) comprises a range of disorders that are either isolated or part of a complex genetic disease, characterized by a persistent or intermittent, low peripheral neutrophil count (severe  $<0.5 \times 10^9/L$  and mild  $0.5 - 1.5 \times 10^9/L$ ).<sup>1</sup> The inherited forms of CN often presents with more severe clinical manifestations than those caused by chemotherapy, viral infections, or drug reactions.<sup>2</sup> Life-threatening bacterial infections mostly occur in the first year of life for patients with severe congenital neutropenia (SCN) who have not received granulocyte colony-stimulating factor (G-CSF) treatment.<sup>3</sup>

Granulocyte development proceeds in a stepwise fashion in the bone marrow (BM) and is regulated by various growth factors and cytokines.<sup>4</sup> Granulocyte-monocyte progenitor cells, the direct precursors to granulocytes and monocytes, first differentiate into myeloblasts (immature granulocytes) and then into promyelocytes.<sup>5</sup> The promyelocytes further differentiate into myelocytes that begin to exhibit distinct characteristics of specific granulocyte lineages, including neutrophils, eosinophils, or basophils.<sup>6</sup> The myelocytes mature into metamyelocytes and subsequently into fully formed granulocytes, including band cells and polymorphonuclear leukocytes (PMNs). These cells then leave the BM and migrate to the periphery.<sup>7</sup> Patients with SCN exhibit a maturation arrest at the stages of promyelocytes and myelocytes during granulocyte development in the BM.<sup>8</sup> However, it is incompletely understood why granulopoiesis is disrupted at specific stage in various types of CN caused by different genetic mutations.

For CN cases with known genetic defects, mutations in the *ELANE* gene are most prevalent, accounting for 55.6% cases.<sup>9</sup> The *ELANE* (or *ELA2*) gene encodes the neutrophil elastase protein. A diagnosis of *ELANE*-related neutropenia is made when a heterozygous pathogenic variant is found in a proband with suggestive clinical findings. Clinical manifestations of *ELANE*-related neutropenia, which include SCN or cyclic neutropenia (CyN), consist of recurrent fever, skin and oropharyngeal inflammation, and cervical lymphadenitis.<sup>10</sup> CN can also arise because of mutations in other genes, including *HAX1*, *CXCR4*, and Shwachman-Bodian-Diamond Syndrome (*SBDS*). These patients require regular G-CSF treatment to prevent frequent infections, but with a risk of ~15% to 25% of developing myelodysplasia (MDS) or acute myelogenous leukemia.<sup>10</sup> Furthermore, patients with CN exhibit heterogeneous responses to G-CSF therapy, but the underlying mechanisms are not well-understood.<sup>1</sup>

Because *ELANE* is the most prevalent pathogenic gene in CN, most studies have focused primarily on *ELANE*-related neutropenia, with few distinguishing it from CN caused by mutations in other genes. To understand the pathogenesis of CN resulting from different genetic defects, we conducted a retrospective analysis of patients with CN at our center and compared the clinical, immunological, and BM characteristics of following 5 CN groups: SCN1 and CyN (both caused by mutations in *ELANE*), SCN3 (associated with *HAX1* mutations), warts, hypogammaglobulinaemia, infections and myelokathexis (WHIM, linked to *CXCR4* mutations), and *SBDS*, related to mutations in the *SBDS* gene. During this analysis, we discovered 11 novel mutations in *ELANE* and 1 mutation each in the *HAX1*, *CXCR4*, and *G6PC3* genes. Notably, we observed distinctive differences in the granulocyte development, G-CSF responsiveness, and antibody

production among these 5 groups. Specifically, we identified a strong correlation between the stage and degree of granulocyte development disruption and the efficacy of G-CSF therapy.

## Methods

### Patients and G-CSF therapy

In this retrospective study conducted at the Department of Clinical Immunology, Children's Hospital of Fudan University, a total of 349 patients diagnosed with chronic neutropenia were recruited over a period of 8 years (2014-2021). Chronic neutropenia is defined as a decrease in the absolute number of neutrophils in peripheral blood for  $>3$  months. Of these, 56 cases had a diagnosed gene defect. These patients were categorized into 5 groups based on gene screening results and clinical manifestations: SCN1, SCN3, CyN, WHIM, and *SBDS*. The cases with the causative genes *WAS*, *G6PC3*, and *SPR54* were not included in the comparative analysis because there was only 1 case each. Typically, patients with SCN were found to have an absolute neutrophil count (ANC)  $<0.5 \times 10^9/L$  for  $>6$  months within the first year of life, accompanied by abnormal granulopoiesis in the BM and frequent monthly infections.

For the diagnosis of CN, routine laboratory examinations were conducted to rule out other potential causes, such as active infections, autoimmune conditions, metabolic disorders, and marrow failure. Complete blood counts were obtained 3 times per week for 6 weeks for CN diagnosis, and the mean neutrophil count was calculated based on ~10 blood tests, encompassing a comprehensive range of the clinical condition, including any periods of fever or suspected infections. To exclude the influence of G-CSF, the blood routine results of patients in rest state were determined as the mean values of the indicators obtained during their regular follow-up blood routine either before or at least 10 days after the administration of G-CSF. BM aspirations were conducted upon the recommendation of their physicians after diagnosis and before G-CSF treatment. The timing of these aspirations was determined based on the availability and consent of the patients' parents, thus resulting in random timing relative to the neutrophil cycle. At the time of BM aspiration, the peripheral ANCs in these patients in CyN group were all  $<1.5 \times 10^9$  cells per L threshold for neutropenia diagnosis. The ANCs at the closest time point of the BM sample collection, along with the fluctuation ranges for the 4 patients in CyN group, were as follow: P5 had an ANC of  $1.16 \times 10^9$  cells per L with a fluctuation range of  $0.23 \times 10^9$  to  $1.84 \times 10^9$  cells per L; P11 had an ANC of  $0.57 \times 10^9$  cells per L with a range of  $0.06 \times 10^9$  to  $2.12 \times 10^9$  cells per L; P16 had an ANC of  $0.67 \times 10^9$  cells per L with a range of  $0.10 \times 10^9$  to  $2.00 \times 10^9$  cells per L; P21 had an ANC of  $0.46 \times 10^9$  cells per L with a range of  $0.26 \times 10^9$  to  $3.73 \times 10^9$  cells per L. For each patient, BM aspiration was performed once, and typically 3 BM smears was prepared. A pathologist then counted 200 cells on 1 well-made smear to ascertain the percentages of various cell types. To ensure accuracy, the results were independently verified by another experienced pathologist.

Recombinant human G-CSF (rhG-CSF) was administered at 5 to 20  $\mu\text{g}/\text{kg}$  body weight according to patient willingness and risk of infection. Initially, we administered G-CSF at a dose of 5  $\mu\text{g}/\text{kg}$  body weight per day for ~1 week. Subsequently, we increased the doses, typically to 7.5 to 10  $\mu\text{g}$ , based on the initial response observed in patients. In a few cases in which there was no significant increase in neutrophil counts after these doses, we further

increased the G-CSF dose, with a maximum dose of 20 µg. The ANC was analyzed before and after G-CSF administration. To address G-CSF responsiveness, we focused on the peak ANC (pANC) achieved within 10 days after G-CSF treatment as our primary criterion.

This study was carried out in accordance with the recommendations of the Ethics Committee of the Children's Hospital of Fudan University (Shanghai, China) (approval number, 2019-048). Written informed consent was obtained from the parent or guardian, and the child's assent was secured before any study-related procedures.

## Data collection

Clinical and laboratory information was collected via electronic chart review, which included (1) general data, such as gender, age, place of origin, and family history; (2) personal history that focused on the patient's infection status, including the frequency, type, location, severity, and infection cycle patterns; (3) physical examinations with specific attention to characteristics associated with CN, including skeletal abnormalities, malformation characteristics, albinism, psychomotor development, warts, cardiac function, hepatosplenic lymph node size, and neurological symptoms; (4) laboratory tests encompassed common laboratory screening items, such as blood smear analysis, C-reactive protein levels, serum electrolytes, liver and kidney function, serological and/or DNA or RNA analysis of viral pathogens (cytomegalovirus, Epstein-bar virus, etc), fasting blood glucose, autoantibodies, and BM cytomorphological examination; (5) immunological examinations included lymphocyte subgroup analysis, serum Ig levels, neutrophil respiratory burst function tests, immunoglobulin G (IgG) subclasses, and assessment of specific antibodies; and (6) additional examinations were conducted as needed, including (i) metabolism-related tests, such as urine organic acids, and urine and serum amino acids, to rule out metabolic diseases associated with neutropenia, including glycogen storage disease Ib, organic acidosis, tyrosinemia, Barth syndrome, and Gaucher disease and (ii) microbiological tests based on the clinical symptoms of infection, such as urine culture, throat wipe culture, blood culture, and/or rapid molecular PCR for bacterial and fungal DNA detection.

## Genomic DNA sequencing

Genomic DNA was extracted and sequenced as described.<sup>11,12</sup> The concentration and quantity of the DNA samples were measured using a NanoDrop ultraviolet spectrophotometer (Thermo Fisher Scientific, Waltham, MA). Next-generation sequencing was carried out using an immunodeficiency gene panel, which included genes associated with CN, such as *AK2*, *AP3B1*, *CD40LG*, *CLPB*, *CSF3R*, *CXCR2*, *CXCR4*, *DNAJC21*, *DNM2*, *DOCK2*, *EFL1*, *EIF2AK3*, *ELANE*, *G6PC3*, *GATA1*, *GATA2*, *GFI1*, *GINS1*, *HAX1*, *IRAK4*, *JAGN1*, *KAT6A*, *KRAS*, *LAMTOR2*, *LYST*, *MYD88*, *PGM3*, *PSTPIP1*, *RAB27A*, *RAC2*, *SBDS*, *SEC61A1*, *SLC37A4*, *SMARCD2*, *SRP54*, *STK4*, *TAZ*, *TCIRG1*, *TCN2*, *TLR8*, *USB1*, *VPS13B*, *VPS45*, *WAS*, *WDR1*, and *WIPF1*. Alternatively, whole-exome sequencing was performed. The genomic DNA fragments were enriched by adapter ligations and sequenced on an Illumina HiSeq 2000 platform (Illumina, San Diego, CA). The sequence data were annotated using ANNOVAR and variant effect predictor software, and variant pathogenicity was predicted with SIFT, PolyPhen-2, and

MutationTaster tools. For each family, we used either panel sequencing or whole-exome sequencing to identify genetic mutations, and the same method was used for both the patient and their parents. Finally, Sanger sequencing was used to confirm pathogenic mutations.


## Statistical analysis

Data were presented using standard parameters, such as the mean, median, interval range, absolute number, standard deviation, and percentile. Categorical variables were shown as n (%), normally distributed continuous variables as mean ± standard deviation, and nonnormally distributed continuous variables as median (interquartile range). For multigroup comparisons, the 1-way analysis of variance test was applied for normally distributed continuous variables, and the Kruskal-Wallis test was used for nonnormally distributed continuous variables in CyN, SCN1, SCN3, WHIM, and SBDS. The 2-way analysis of variance test was used when comparing 2 categorical variables. Data analysis was performed using SPSS 16.0 software or GraphPad Prism 8 software. Statistical significance was evaluated based on the *P* value (\**P* < .05; \*\**P* < .01; \*\*\**P* < .001; and \*\*\*\**P* < .0001). Gray areas represent reference ranges. For age-related indicators, these ranges are shown as median age reference ranges. The reference ranges used for complete blood counts, BM morphologic subgroups, immunoglobulin subgroups, and lymphocytes are based on our institution's established norms. For absolute values of lymphocyte subpopulations, refer to the work by Ding et al.<sup>13</sup> In addition, the overall proportions of each BM lineage are based on the data provided by Sovani et al.<sup>14</sup>

## Results

### Genetic and clinical features of CN

Through a retrospective analysis of patients hospitalized in the Department of Clinical Immunology of the Children's Hospital of Fudan University from 2014 to 2021, a total of 349 patients with the diagnosis of "chronic neutropenia" were identified. The clinical assessments of our patients were primarily based on their infection histories and routine blood tests. These assessments were conducted both during hospitalization and in outpatient settings. Of the 349 patients, only 56 cases (16%) had a clearly identified causative gene (Figure 1A; Table 1). Patients were categorized into 5 groups based on gene screening results, clinical manifestations, and the number of cases: SCN1 and CyN, both caused by mutations in *ELANE*; SCN3, associated with *HAX1* mutations; WHIM, linked to *CXCR4* mutations; and SBDS, related to mutations in the *SBDS* gene. Cases with the causative genes *WAS*, *G6PC3*, and *SPR54*, which were found in only 1 case each, were excluded from the comparison. The onset age and diagnosis age for each patient group are depicted in supplemental Figure 1A. The majority of these cases were due to *ELANE* mutations (*n* = 36, 64.29%), followed by *SBDS* (*n* = 814.29%), *CXCR4* (*n* = 5, 8.93%), and *HAX1* (*n* = 4, 7.14%), with only a single case involving mutations in other genes (*WAS*, *G6PC3*, and *SPR54*) (1.79%). We identified 11 novel mutations in *ELANE* and 1 new mutation each in the *HAX1*, *CXCR4*, and *G6PC3* genes (Figure 1B, red; Table 2).<sup>15-18</sup> De novo mutations emerged as the primary source of *ELANE* variants, representing 87% of patients with *ELANE* mutations. This proportion of de novo mutations is consistent with percentages



**Figure 1. Genetic and clinical features of CN.** (A) A total of 56 cases had a clearly identified causative gene. Among these, 36 had mutations in the *ELANE* gene, which included SCN1 (n = 32, 57.14%) and CyN (n = 4, 7.14%). The remaining 20 patients had mutations in other genes, which included *SBDS* (n = 8, 14.29%), *CXCR4* (n = 5, 8.93%), *HAX1* (n = 4, 7.14%), and *WAS*, *G6PC3*, *SPR54* with 1 patient each (each accounting for 1.79%). (B) Location of the mutations on the *ELANE*, *HAX1*, *CXCR4*, and *SBDS* proteins. Each dot represents a case with that particular mutation. Highlighted in red are novel mutations. On the *ELANE* protein, mutations higher on the diagram are associated with SCN and those lower with CyN. (C) This panel shows the tissues/organs infected or types of infectious agents involved in the medical history of patients in these 5 groups. Red squares indicate a history of infection at the specific site or with the pathogen; White squares denote no reported infection at the site or with the pathogen. BH1 and BH2, Bcl-2 protein homology domains; FYSH, Fungal, Yhr087w, and Shwachman domains; HTH, helix-turn-helix domain; PEST, region rich in proline (P), glutamic acid (E), serine (S), and threonine (T); RRM, RNA recognition motif; TM, transmembrane-like domain.

reported in a previous study.<sup>19</sup> In addition, *ELANE* mutations occurred either in the first amino acid (M1) or the mature protease region (Figure 1B). Although variants of other genes were mainly inherited within families, only 4 patients (P7, P19-20, and P26) inherited the *ELANE* variants from their parents (Table 2;

supplemental Figure 1B). P7 inherited the p.C55Y variant of *ELANE* from her father. This variant, which is also present in P6 as a de novo mutation, had been previously reported.<sup>15</sup> Siblings P19 and P20 inherited the p.V197Sfs\*15 variant from their father. This mutation was previously reported by Makaryan et al in 2015.<sup>16</sup> P26

**Table 1. Clinical features of patients with CN with identified genetic defects diagnosed at Children’s Hospital of Fudan University**

No.	Gene	Gender	Age of onset (mo)	Age of diagnosis (mo)	Infected tissues/organs
1	ELANE	F	120	152	1, 7, 11
2	ELANE	M	2	3	1, 2, 3, 4, 5, 7, 8
3	ELANE	M	3	21	1, 3, 4, 7, 11
4	ELANE	M	0.4	13	1, 2, 7
5	ELANE(CyN)	F	9	58	1, 2, 4, 6
6	ELANE	F	16	17	1, 11
7	ELANE	F	2	33	1, 3, 5, 7, 8, 13
8	ELANE	F	0.2	17	2, 7, 9, 11
9	ELANE	M	5	26	1, 7
10	ELANE	F	2	3	1, 2, 3, 6, 7, 10, 11
11	ELANE(CyN)	F	24	72	2, 8
12	ELANE	F	0.2	15	1, 2, 3, 5, 7, 8, 10
13	ELANE	M	0.1	18	1, 2, 10
14	ELANE	F	1	162	1, 2, 4, 6, 13
15	ELANE	M	2	18	1, 2, 7
16	ELANE(CyN)	F	24	120	2, 6, 7
17	ELANE	M	1	36	1, 3, 7, 11
18	ELANE	M	0.8	6	1, 4, 5, 7
19	ELANE	F	120	152	1, 2, 6, 7
20	ELANE	M	2	3	2, 10
21	ELANE(CyN)	F	3	21	3, 6
22	ELANE	M	0.4	13	1, 7, 9, 10, 12
23	ELANE	F	9	58	1, 2
24	ELANE	F	16	17	1, 2, 4, 7
25	ELANE	F	2	33	1, 2, 4, 7, 8, 9, 11, 12
26	ELANE	M	0.2	17	1
27	ELANE	F	5	26	1, 2, 3, 5, 8, 11
28	ELANE	M	2	3	1, 2, 7
29	ELANE	M	24	72	1, 2, 5, 7
30	ELANE	M	0.2	15	1, 5, 7, 8, 12
31	ELANE	M	0.1	18	1, 3, 7
32	ELANE	M	1	162	1, 2, 3, 5, 7
33	ELANE	F	2	18	1, 2
34	ELANE	F	24	120	1
35	ELANE	M	1	36	1, 2, 7
36	ELANE	M	0.8	6	1, 8, 9, 13
37	HAX1	M	3	110	1, 7, 10
38	HAX1	F	24	69	2, 7
39	HAX1	M	25	34	1, 2, 3, 7
40	HAX1	M	1	8	1, 2, 3, 5, 6, 11
41	CXCR4	M	24	24	12
42	CXCR4	M	2	17	1, 5, 8
43	CXCR4	M	0.7	13	1, 5
44	CXCR4	F	58	62	1, 5
45	CXCR4	F	1	24	1

P19 and P20, as well as P28 and P29, are from 2 separate twin families.

Infected tissues/organs: 1, pulmonary; 2, oral; 3, gums; 4, lymph nodes; 5, middle ear; 6, tonsils; 7, skin; 8, gastrointestinal tract; 9, blood; 10, urinary tract; 11, soft tissues; 12, abdominal cavity; 13, brain/nerves.

F, female. M, male.



**Table 1 (continued)**

No.	Gene	Gender	Age of onset (mo)	Age of diagnosis (mo)	Infected tissues/organs
46	SBDS	F	120	152	1, 7, 11
47	SBDS	M	2	3	1, 8
48	SBDS	F	3	21	1, 8
49	SBDS	F	0.4	13	1, 8
50	SBDS	F	9	58	1, 4
51	SBDS	F	16	17	1, 8
52	SBDS	F	2	33	1, 8, 9
53	SBDS	M	0.2	17	1, 8, 9, 13
54	WAS	M	5	26	1
55	G6PC3	F	2	3	1, 2, 7, 8
56	SPR54	M	24	72	1, 2, 4, 11

P19 and P20, as well as P28 and P29, are from 2 separate twin families.

Infected tissues/organs: 1, pulmonary; 2, oral; 3, gums; 4, lymph nodes; 5, middle ear; 6, tonsils; 7, skin; 8, gastrointestinal tract; 9, blood; 10, urinary tract; 11, soft tissues; 12, abdominal cavity; 13, brain/nerves.

F, female. M, male.

inherited the p.C208R variant from his father, who had a history suggestive of neutropenia, although exact values were not recorded. The patient's elder brother also had recurrent respiratory infections in early childhood, but there is no available information about an *ELANE* mutation.

Recurrent multisite infections by multiple pathogens are the most common clinical manifestations in patients with CN.<sup>8,20</sup> In terms of infected organs or tissues, pulmonary infections were present in >80% of cases across all groups except for CyN (Figure 1C, upper panel). Skin infections (mostly perianal abscesses) and oral ulcers occurred more frequently in SCN1, SCN3, and CyN, whereas SBDS cases manifested more gastrointestinal infections (including diarrhea) and WHIM cases exhibited more otitis media. Patients who exhibited skin infections or oral ulcers were prescribed antibiotics without hospitalization. With respect to the pathogens involved (Figure 1C, lower panel), bacterial infections were the most dominant pathogens, and fungal infections occurring more frequently in SCN1 and SCN3. The most commonly identified bacteria species were *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Escherichia coli* (supplemental Figure 1C). *Candida albicans* was the most commonly identified fungus, and *Mycoplasma pneumoniae* was the predominant mycoplasma identified. There was no specific trend observed for viral infections. Among the 56 cases studied, 40 cases received G-CSF treatment. Partly because of the relatively short follow-up period after admission, we were unaware of any cases of leukemia development. However, there was 1 patient who died of severe infections.

### Comparative analysis of immune cells and granulocyte subsets in peripheral blood and BM

Firstly, we analyzed the absolute counts of white blood cells (WBCs) along with the 3 major lineages (granulocytes, monocytes, and lymphocytes) in peripheral blood. Given the functional nature of CXCR4 as a chemokine,<sup>21</sup> CXCR4 mutation severely affects the release of BM lymphocytes into the periphery, as observed in WHIM syndrome.<sup>22,23</sup> Indeed, as depicted in Figure 2A, WHIM displayed a markedly low WBC count because of a significant

reduction in monocytes and lymphocytes in addition to low levels of granulocytes. In addition, SCN1 and SBDS showed slightly higher peripheral blood WBC counts than SCN3 and CyN. This could be attributable to an increased count of lymphocytes in SCN1 and SBDS (Figure 2A). Monocytes were increased in SCN1 and SCN3 (Figure 2A), which exceeded the normal reference range. Within the granulocytes, neutrophil counts were higher in CyN and SBDS, whereas eosinophils and basophils were increased in SCN1 (Figure 2B). These observations suggest that SCN1 primarily results in a significant decrease in neutrophils but not in eosinophils and basophils, whereas other genetic variants affect all types of granulocytes. One patient in CyN group (P21, mean ANC =  $1.7 \times 10^9$  cells per L) and 3 patients in SBDS group (P47, mean ANC =  $1.6 \times 10^9$  cells per L; P50, mean ANC =  $2.05 \times 10^9$  cells per L; P52, mean ANC =  $1.57 \times 10^9$  cells per L) had neutrophil counts slightly  $>1.5 \times 10^9$  cells per L. The *ELANE* mutation in P21 (ANC range,  $0.26\text{-}3.73 \times 10^9$  cells per L) is novel. However, this mutation, c.597+4G>A, is in close proximity to other mutations (c.597+1G>A, c.597+5G>A) that have been associated with CyN or SCN in the ClinVar database, suggesting a potential pathogenic nature for the P21 mutation. All 3 SBDS mutations have been previously reported as pathogenic. The relevant details can be found in Table 2 (PMID 33607811, 37885353, 12496757, 22935661, 21695142, 14749921, 15769891, and 32412173).

To explore the mechanisms underlying the differential distribution of peripheral immune cell subsets, we next performed a detailed analysis of the development of distinct immune cell lineages in BM. We analyzed the proportions of the 3 lineages (granulocytes, monocytes, and lymphocytes) in the BM (Figure 2C). Both SCN1 and SCN3 showed significantly lower proportions of the granulocyte lineage compared with CyN and WHIM (Figure 2C), predominantly because of a drastic decrease in the neutrophil lineage (Figure 2D). In contrast, CyN and WHIM showed higher proportions of the granulocyte lineages than other subgroups (Figure 2C), largely because of the normal development of neutrophilic granulocytes in the BM (Figure 2D). Within the granulocyte lineage, the proportion of neutrophil lineage was reduced, whereas those of eosinophil and basophil lineages increased in

**Table 2. Detailed gene variants information in genetically defined CN**

No.	Gene	Location	Nucleotide change	Protein level	Inheritance	Novel mutation*	Clinical significance†	Reference (PMID‡)
1	ELANE	Exon1	c.1A>G	p.M1V	De novo	No	Pathogenic	24184683
2	ELANE	Exon1	c.1A>G	p.M1V	De novo	No	Pathogenic	24184683
3	ELANE	Exon1	c.3G>A	p.M1I	De novo	No	Pathogenic	24184683
4	ELANE	Exon2	c.140T>G	p.L47R	De novo	Yes	Pathogenic	NA§
5	ELANE	Exon2	c.158A>G	p.H53R	De novo	No	Pathogenic	NA§
6	ELANE	Exon2	c.164G>A	p.C55Y	NA§	No	Pathogenic	16986121
7	ELANE	Exon2	c.164G>A	p.C55Y	Paternal	No	Pathogenic	16986121
8	ELANE	Exon2	c.212G>T	p.C71F	De novo	No	Pathogenic	31839986
9	ELANE	Exon3	c.242G>C	p.R81P	De novo	No	Likely Pathogenic	16986121
10¶	ELANE	Exon3	c.269_277del	p.R91Afs*175	De novo	Yes	NA§	NA§
	ELANE	Exon3	c.278A>G	p.E93G	De novo	Yes	Pathogenic	NA§
11	ELANE	Exon3	c.308G>T	p.R103L	De novo	No	Likely Pathogenic	17761833
12	ELANE	Exon3	c.362T>C	p.L121P	De novo	No	Likely Pathogenic	NA§
13	ELANE	Exon4	c.368T>A	p.L123H	NA	Yes	Pathogenic	NA§
14	ELANE	Exon4	c.377C>T	p.S126L	De novo	No	Pathogenic	28492532, 11001877, 14962902, 16079102, 16737875, 18611981, 20582973, 22758217, 23463630, 16551967, 26567890
15	ELANE	Exon4	c.401_415del	p.Q134_L138del	De novo	Yes	NA§	NA§
16	ELANE	Exon4	c.416C>T	p.P139L	NA	No	Pathogenic	11001877, 23463630, 21425445, 14962902, 30040071, 16079102, 31321910, 31248972
17	ELANE	Exon4	c.452G>A	p.C151Y	NA§	No	Pathogenic	24523240, 25427142, 11675333, 23463630
18	ELANE	Exon4	c.452G>C	p.C151S	De novo	No	Pathogenic	23463630
19	ELANE	Exon4	c.588delC	p.V197Sfs*15	Paternal	No	NA§	25427142
20	ELANE	Exon4	c.588delC	p.V197Sfs*15	Paternal	No	NA§	25427142
21	ELANE	Exon4	c.597+4C>G	splicing	NA§	Yes	NA§	NA§
22	ELANE	Exon5	c.601del	p.D201Tfs*11	De novo	Yes	NA§	NA§
23	ELANE	Exon5	c.607G>C	p.G203R	NA§	No	Pathogenic	23463630
24	ELANE	Exon5	c.608G>A	p.G203D	De novo	Yes	Pathogenic	NA§
25	ELANE	Exon5	c.608G>A	p.G203D	De novo	Yes	Pathogenic	NA§
26	ELANE	Exon5	c.622T>C	p.C208R	paternal	Yes	Pathogenic	NA§
27	ELANE	Exon5	c.640G>A	p.G214R	De novo	No	Pathogenic	11001877, 15657182, 16079102, 28073911, 30386760, 3229910
28	ELANE	Exon5	c.640G>A	p.G214R	De novo	No	Pathogenic	11001877, 15657182, 16079102, 28073911, 30386760, 3229910
29	ELANE	Exon5	c.640G>A	p.G214R	De novo	No	Pathogenic	11001877, 15657182, 16079102, 28073911, 30386760, 3229910
30	ELANE	Exon5	c.640G>A	p.G214R	De novo	No	Pathogenic	11001877, 15657182, 16079102, 28073911, 30386760, 3229910
31	ELANE	Exon5	c.641G>A	p.G214E	De novo	No	Likely Pathogenic	NA§
32	ELANE	Exon5	c.651delC	p.F218fs	De novo	Yes	NA	NA§
33	ELANE	Exon5	c.658delC	p.R220fs*20	De novo	No	Pathogenic	26174650
34	ELANE	Exon5	c.661G>T	p.G221X	De novo	No	Uncertain Significance	11001877
35	ELANE	Exon5	c.684C>G	p.Y228X	De novo	No	NA§	32054657

\*Novel variant denotes that the variant was not reported in the NCBI website ClinVar database and GnomAD database. We also referred to *ELANE* mutations published previously.<sup>15-18</sup>

†Clinical significance represents the assessment of the variant in the ClinVar database or the predicted pathogenicity using polyphen-2.

‡Reference PMID represents the number of the reported or referenced literature in Pubmed for the mutation.

§NA, not available.

||Cyclic neutropenia (CyN).

¶Compound heterozygous mutations. P19 and P20, as well as P28 and P29, are from 2 separate twin families. The father of P19 and P20 has the same compound heterozygous mutations and presented with neutropenia, recurrent respiratory infections, and oral ulcers. P26 inherited the p.C208R variant from his father, whose blood tests were suggestive of neutropenia, although the exact neutrophil count was not recorded.

**Table 2 (continued)**

No.	Gene	Location	Nucleotide change	Protein level	Inheritance	Novel mutation*	Clinical significance†	Reference (PMID‡)
36	<i>ELANE</i>	Exon5	c.704T>G	p.V235G	De novo	Yes	Pathogenic	NA
37	<i>HAX1</i>	Exon3	c.430dupG	p.V144Gfs*5	Maternal	No	Pathogenic	28492532, 17187068, 18337561, 20065084, 20220065, 22102707, 24482108
	<i>HAX1</i>	Exon5	c.557-1G>C	splicing	Paternal	No	Likely Pathogenic	16199547, 17187068
38	<i>HAX1</i>	Exon2	c.216_217insC	p.I73Hfs*6	Maternal	No	Pathogenic	17187068
	<i>HAX1</i>	Exon3	c.430dupG	p.V144Gfs*5	Paternal	No	Pathogenic	28492532, 17187068, 18337561, 20065084, 20220065, 22102707, 24482108
39	<i>HAX1</i>	Exon3	c.430dupG	p.V144Gfs*5	Maternal	No	Pathogenic	28492532, 17187068, 18337561, 20065084, 20220065, 22102707, 24482108
	<i>HAX1</i>	Exon3	c.430dupG	p.V144Gfs*5	Paternal	No	Pathogenic	28492532, 17187068, 18337561, 20065084, 20220065, 22102707, 24482108
40	<i>HAX1</i>	Exon3	c.486_487dup	p.Q163Pfs*52	Maternal	Yes	NA§	NA§
	<i>HAX1</i>	Exon3	c.430dupG	p.V144Gfs*5	Paternal	No	Pathogenic	28492532, 17187068, 18337561, 20065084, 20220065, 22102707, 24482108
41	<i>CXCR4</i>	Exon2	c.1000C>T	p.R334X	NA	No	Pathogenic	31313072, 12692554, 25662009, 31493092
42	<i>CXCR4</i>	Exon2	c.1000C>T	p.R334X	NA	No	Pathogenic	31313072, 12692554, 25662009, 31493092
43	<i>CXCR4</i>	Exon2	c.1000C>T	p.R334X	NA	No	Pathogenic	31313072, 12692554, 25662009, 31493092
44	<i>CXCR4</i>	Exon2	c.1032dupT	p.E345X	De novo	Yes	NA§	NA§
45	<i>CXCR4</i>	Exon2	c.1000C>T	p.R334X	De novo	No	Pathogenic	31313072, 12692554, 25662009, 31493092
46	<i>SBDS</i>	Exon2	c.184A>T	p.K62X	Maternal	No	Pathogenic/likely pathogenic	33607811 37885353
	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
47	<i>SBDS</i>	Exon2	c.184A>T	p.K62X	Maternal	No	Pathogenic/likely pathogenic	33607811 37885353
	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
48	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Maternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
49	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Maternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
50	<i>SBDS</i>	Exon2	c.183_184TA>CT	p.K62X	Maternal	No	Pathogenic	12496757, 15769891, 32412173
	<i>SBDS</i>	Exon2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
51	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Maternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
	<i>SBDS</i>	Exon2	c.183_184TA>CT	p.K62X	Paternal	No	Pathogenic	12496757, 15769891, 32412173
52	<i>SBDS</i>	Exon2	c.184A>T	p.K62X	Maternal	No	Pathogenic/likely pathogenic	33607811 37885353
	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
53	<i>SBDS</i>	Exon2	c.183_184TA>CT	p.K62X	Maternal	No	Pathogenic	12496757, 15769891, 32412173
	<i>SBDS</i>	Intron2	c.258+2T>C	splicing	Paternal	No	Pathogenic	12496757, 22935661, 21695142, 14749921
54	<i>WAS</i>	Exon9	c.881T>C	p.I294T	Maternal	No	Pathogenic/likely pathogenic	11242115, 19006568
55	<i>G6PC3</i>	Exon3	c.394C>T	p.Q132X	Maternal	Yes	NA	NA
56	<i>SPR54</i>	Exon5	c.349_351del	p.T117del	De novo	No	Pathogenic	29914977, 36159802

\*Novel variant denotes that the variant was not reported in the NCBI website ClinVar database and GnomAD database. We also referred to *ELANE* mutations published previously.<sup>15-18</sup>

†Clinical significance represents the assessment of the variant in the ClinVar database or the predicted pathogenicity using polyphen-2.


‡Reference PMID represents the number of the reported or referenced literature in Pubmed for the mutation.

§NA, not available.

||Cyclic neutropenia (CyN).

¶Compound heterozygous mutations. P19 and P20, as well as P28 and P29, are from 2 separate twin families. The father of P19 and P20 has the same compound heterozygous mutations and presented with neutropenia, recurrent respiratory infections, and oral ulcers. P26 inherited the p.C208R variant from his father, whose blood tests were suggestive of neutropenia, although the exact neutrophil count was not recorded.





**Figure 2. Comparative analysis of BM or peripheral blood immune cell subsets across 5 groups using routine blood tests and cell morphology assays of BM smears.** (A) Total number of WBC, granulocytes, monocytes, and lymphocytes in the peripheral blood. (B) Absolute numbers of granulocytes (neutrophils, eosinophils, and basophils) in the peripheral blood. The brown line in neutrophil indicates the cut-off value of  $1.5 \times 10^9$  cells per L of neutropenia. (C) Proportion of granulocytes, monocytes,

SCN1 (Figure 2D). These observations are in agreement with the findings in the peripheral blood (Figure 2B).

We further examined the specific stages of neutrophil development in the BM (Figure 2E-G). SCN1 and SCN3 exhibited severe dysplasia beginning at the commitment to the neutrophilic lineage (myelocytes) and persisting through to the PMNs (Figure 2F). In contrast, CyN displayed relatively normal development of neutrophilic granulocytes up to the band cell stage, but with a moderate reduction in PMNs (Figure 2F). Neutrophil development in SBDS was intermediate between these 2 patterns. WHIM showed normal neutrophil development from the myelocyte all the way through to the PMNs. These results demonstrate that neutrophilic granulocyte development is variably affected among these groups. Unlike the neutrophilic granulocytes, eosinophilic granulocyte development in SCN1 was almost unaffected, whereas it was severely impaired in WHIM and SBDS, from eosinophilic myelocytes to the eosinophilic PMNs (Figure 2G). Meanwhile, eosinophil development in SCN3 and CyN was normal until the band cell stage but became affected at the PMN stage. Thus, there are entirely opposing effects on the development of neutrophilic vs eosinophilic granulocytes in SCN1, resulting in reduced peripheral neutrophils but normal eosinophils (Figure 2B).

### Distinct responsiveness to G-CSF among different types of CN

Considering the different impacts of various genetic mutations on the counts of BM and peripheral neutrophils, we further analyzed the effects of G-CSF administration on patients in different groups (Table 3). G-CSF was administered as described in the "Methods." To address G-CSF responsiveness, we chose to focus on the pANC achieved within 10 days after G-CSF treatment as our primary criterion. It was found that all patients with CN exhibited significantly higher pANC after G-CSF treatment than baseline ANC (bANC) before G-CSF initiation. However, although ANCs were uniformly very low in all patient groups before G-CSF administration, pANC reached different levels in different groups. Notably, pANC was higher in the CyN, WHIM, and SBDS groups, suggesting better responsiveness to G-CSF, whereas only half of the SCN1 and SCN3 were able to elevate their pANC  $>1.5 \times 10^9$  cells per L (Figure 3A; Table 4). This indicates that although G-CSF is effective for all patients with CN, the degree of responsiveness varies among different types of CN.

To investigate the relationship between G-CSF responsiveness and BM neutrophil development in different patient groups, we statistically analyzed the correlation between the development of the neutrophilic lineage in BM and pANC. We observed a significant correlation that patients with poor BM neutrophil development exhibited reduced responsiveness to G-CSF (Figure 3B). Specifically, SCN1 and SCN3, which had lower percentages of BM neutrophilic lineage cells, also showed lower pANC after G-CSF administration (Figure 3C). In contrast, CyN, WHIM, and SBDS, which had higher percentages of BM neutrophilic cells, showed higher pANC. This suggests a reverse relationship between the

degree of disruption in granulocyte development and the effectiveness of G-CSF treatment.

We further categorized BM neutrophil lineage cells into the stem cell pool (myeloblast), the mitotic cell pool (promyelocyte and myelocyte) and the postmitotic cell pool (metamyelocyte, band cell, and PMN) as described.<sup>24</sup> Our analysis revealed a significant correlation between pANC and the populations of both mitotic and postmitotic pools (Figure 3D). However, no such correlation was observed between pANC and the stem cell pool (Figure 3D). This finding highlights the role of G-CSF in promoting early granulocyte mitosis and mobilizing cells from the postmitotic pool, and its relation to the increase in pANC.

We also conducted a Spearman analysis to assess the relationship between bANC before G-CSF treatment and the pANC after treatment. Our analysis did not reveal a significant correlation between these 2 measures (supplemental Figure 2). It appears that bANC does not predict responsiveness to G-CSF treatment.

### Elevated serum Ig levels in SCN1, SCN3, and CyN correlate with increased BM plasma cells and serum APRIL concentrations

Adaptive immunity in patients with CN, who are characterized primarily by a neutrophil-mediated immune deficiency, has received little attention. We found that serum levels of IgG, IgM, and IgA were significantly elevated in the SCN1, SCN3, and CyN compared with those in the WHIM and SBDS, with SCN1 exceeding the reference range (Figure 4A). We also found a strong correlation between Ig levels and the percentages of BM PC cells in individual patients (Figure 4B). Analysis of lymphocyte subpopulations revealed diminished CD3<sup>+</sup> T, CD4<sup>+</sup> T, CD8<sup>+</sup> T, and CD19<sup>+</sup> B cells in patients in the WHIM group because of their impaired migration (supplemental Figure 3). Serum Igs are primarily secreted by long-lived plasma cells in the BM, which adhere to niches formed by BM stromal cells via CXCL12 and survive for decades by interacting with APRIL and interleukin-6.<sup>25</sup> Consistently, we found a higher proportion of BM plasma cells in SCN1, SCN3, and CyN than in WHIM and SBDS (Figure 4C). In addition, serum APRIL concentrations were higher in patients in SCN1 group than those in age-matched healthy controls (Figure 4D). However, recent infections could influence Ig levels. Therefore, further studies are needed to elucidate the reasons behind the elevated Ig levels in SCN1, SCN3, and CyN.

## Discussion

In this study, we conducted a comprehensive, retrospective systematic analysis of 56 patients with CN with 7 identified genes. Notably, we observed regional variations in the population distribution of CN pathogenic genes. Specifically, *CXCR4* mutations were more prevalent in mainland China, whereas *CLPB* and *G6PT* mutations, commonly found in other regions,<sup>8,26</sup> were not frequent in China. The overall infection

**Figure 2 (continued)** and lymphocytes in BM. (D) Proportion of specific granulocyte types (neutrophils, eosinophils, and basophils) in BM. (E-G) Proportions of BM neutrophilic (E-F) and eosinophilic (E,G) lineage cells at different developmental stages for each patient group. \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ ; \*\*\*\* $P < .0001$ . Kruskal-Wallis test was used for comparison among 5 groups. Bars represent means  $\pm$  standard deviation [SD]. The gray area indicates reference range of each index, as described in the Methods section.

**Table 3. G-CSF administration and follow-up of patients with CN**

No.	Mutated gene	Age at first visit (y)	Follow-up period (mo)	Body weight (kg)	G-CSF				HSCT*
					Injection period (d)	Dosage (µg/kg)	bANC (×10 <sup>9</sup> cells per L)†	pANC (×10 <sup>9</sup> cells per L)‡	
1	ELANE	13.5	1.6	33.5	9	7	0.53	2.37	No
2	ELANE	1.1	2.0	10.0	59	10-20	0.03	0.45	No
3	ELANE	1.8	20.8	12.5	14	10	0.65	2.01	Yes
4	ELANE	1.1	3.0	10	–	–	–	–	No
5	ELANE(CyN)	4.8	1.7	15	–	–	–	–	No
6	ELANE	1.4	2.3	11.5	–	–	–	–	No
7	ELANE	2.7	6.5	12	11	10	0.32	1.45	Yes
8	ELANE	1.4	2.4	9.5	9	5-10	0.34	2.41	No
9	ELANE	1.0	80.3	11.5	5	5-7.5	0.08	0.30	No
10	ELANE	0.2	8.1	6.6	15	7.5	0.47	5.01	Yes
11	ELANE(CyN)	6.7	0.5	23	–	–	–	–	No
12	ELANE	0.2	113.4	10	7	5-20	0.24	1.37	No
13	ELANE	1.5	7.8	NA	–	–	–	–	No
14	ELANE	13.4	12.0	NA	–	–	–	–	Yes
15	ELANE	1.5	29.9	12.5	53	7.5-15	0.91	2.11	Yes
16	ELANE(CyN)	10.1	2.1	26	10	5	0.45	10.33	No
17	ELANE	3.1	33.1	18	3	5	0.40	2.57	Yes
18	ELANE	0.5	0.5	6	9	10-15	0.04	0.90	No
19	ELANE	9.1	0.6	32	3	5	0.68	4.17	No
20	ELANE	5.7	0.6	20	11	5-10	0.26	1.32	No
21	ELANE(CyN)	0.4	33.5	15	1	5	0.46	8.38	No
22	ELANE	0.4	11.4	8	9	5	0.33	1.94	No
23	ELANE	2.5	0.9	11.5	–	–	–	–	No
24	ELANE	1.4	0.3	10	–	–	–	–	Yes
25	ELANE	1.0	60.0	9	93	5-20	0.05	0.98	Yes
26	ELANE	5.0	51.4	17	3	5	0.40	2.24	No
27	ELANE	2.5	19.6	12	50	5-7.5	0.18	1.71	Yes
28	ELANE	0.5	12.3	6	3	5	0.96	1.30	No
29	ELANE	0.8	1.5	8	4	5	0.90	1.07	No
30	ELANE	1.8	38.5	10	46	5-7.5	0.55	0.92	Yes
31	ELANE	6.5	15.4	16	9	5	0.42	0.55	Yes
32	ELANE	3.0	1.6	13	1	5	0.20	–	No
33	ELANE	1.1	1.4	10	10	5-10	0.11	3.81	Yes
34	ELANE	13.2	0.6	12	6	5-10	0.43	1.34	No
35	ELANE	6.6	3.3	22	7	5-10	0.18	2.37	Yes
36	ELANE	0.1	15.3	3	7	10-15	0.08	1.50	No
37	HAX1	1.0	13.7	10	9	5-15	0.06	1.19	No
38	HAX1	3.7	1.2	11	2	5	0.03	0.51	No
39	HAX1	10.9	56.5	31	5	5	0.22	1.68	No
40	HAX1	0.3	94.6	19.5	6	5	0.47	2.17	No
41	CXCR4	8.9	59.4	35	–	–	–	–	No
42	CXCR4	0.9	68.3	8.5	10	5	0.15	5.25	No
43	CXCR4	5.1	1.9	19	8	5	0.10	7.39	No
44	CXCR4	0.1	45.2	5	1	5	0.32	6.70	No

–, not receiving G-CSF treatment.

\*HSCT, hematopoietic stem cell transplantation.

†bANC, last absolute baseline neutrophil count (×10<sup>9</sup> cells per L) before G-CSF initiation.

‡pANC, peak ANC (×10<sup>9</sup> cells per L) within 10 days following G-CSF initiation.

**Table 3 (continued)**

No.	Mutated gene	Age at first visit (y)	Follow-up period (mo)	Body weight (kg)	G-CSF				HSCT*
					Injection period (d)	Dosage (µg/kg)	bANC (×10 <sup>9</sup> cells per L)†	pANC (×10 <sup>9</sup> cells per L)‡	
45	<i>CXCR4</i>	7.3	3.7	22	1	5	0.92	1.36	No
46	<i>SBDS</i>	0.6	33.2	6.4	–	–	–	–	No
47	<i>SBDS</i>	5.1	26.2	15	10	7.5	0.42	22.16	No
48	<i>SBDS</i>	0.3	7.2	6	–	–	–	–	No
49	<i>SBDS</i>	1.7	4.1	9	–	–	–	–	No
50	<i>SBDS</i>	0.9	0.3	8.5	1	5	0.52	3.14	No
51	<i>SBDS</i>	0.3	1.4	5	4	5	0.36	1.32	No
52	<i>SBDS</i>	0.4	29.4	6	–	–	–	–	No
53	<i>SBDS</i>	0.6	0.8	7.8	1	5	0.37	2.39	No
54	<i>WAS</i>	6.4	6.5	NA	–	–	–	–	No
55	<i>G6PC3</i>	0.8	13.1	8	–	–	–	–	No
56	<i>SPR54</i>	3.1	2.9	14.5	–	–	–	–	No

–, not receiving G-CSF treatment.

\*HSCT, hematopoietic stem cell transplantation.

†bANC, last absolute baseline neutrophil count (×10<sup>9</sup> cells per L) before G-CSF initiation.

‡pANC, peak ANC (×10<sup>9</sup> cells per L) within 10 days following G-CSF initiation.

characteristics of patients were similar across different CN groups, indicating that the reduction in neutrophil numbers resulting from different genetic variants was the dominant cause of infections in patients with CN. The higher occurrence of oral ulcers or skin infections in the SCN1 may result from a higher incidence of bacterial or fungal infections. On the other hand, the higher incidence of gastrointestinal symptoms (including diarrhea) in the SBDS group may be a consequence of exocrine pancreatic dysfunction.<sup>27</sup>

Our findings revealed that the development of neutrophils in the BM was affected much earlier and more severely in both SCN1 and SCN3 than in the CyN, WHIM, and SBDS groups, which was generally consistent with the number of neutrophils in the peripheral blood circulation. This suggests that patients in SCN1 and SCN3 group have poor BM neutrophil development, directly leading to severely reduced peripheral neutrophils. In contrast, patients in CyN and SBDS group have less severely affected BM neutrophil development, and consequently have higher neutrophil counts in the peripheral blood. Although BM neutrophil development is normal in WHIM, these patients have low circulating neutrophils because of the inability of BM neutrophils to be released into the periphery. Moreover, SCN1 mainly affects neutrophil development with little impact on eosinophils, presumably because of the specific expression of *ELANE* in the neutrophilic granulocyte lineage in BM.<sup>28</sup>

G-CSF is widely used in the treatment of CN and has achieved remarkable therapeutic outcomes.<sup>29</sup> However, there is a lack of studies comparing the efficacy of different CN types to G-CSF administration, because patient responsiveness to G-CSF treatment has not been well-quantified. Dale et al compared the G-CSF responsiveness among CN, CyN, and autoimmune or idiopathic groups.<sup>20</sup> They observed that the CN group had the lowest

pre-G-CSF neutrophil counts. Upon G-CSF treatment, both the mean and median ANC increased in all groups. Our data are consistent with their findings and further elucidate the relationship between G-CSF responsiveness and granulocyte development. We found that the efficacy of G-CSF treatment inversely correlated with the extent of BM granulocyte development disruption in the patients. Our study reveals that different types of patients with CN have distinct responsiveness to G-CSF; SCN1 and SCN3, which display a maturation arrest at an early stage during BM neutrophil development, show poor response to G-CSF, whereas CyN, WHIM, and SBDS, in which BM neutrophil development is either disrupted at a late stage, unaffected, or moderately disrupted, responded more effectively to G-CSF therapy. This corresponds to the degree of their neutrophil development impairment (as detailed in Figure 5). Our findings establish a clear link between the severity of granulocyte development disruption and responsiveness to G-CSF therapy.

Although hypergammaglobulinemia has been described in patients with SCN1, the underlying mechanisms are not clear.<sup>3</sup> A few studies have suggested an association between neutrophil counts, their elastase activity, and humoral immunity. For example, neutrophil numbers have been shown to negatively correlate with IgA levels.<sup>30</sup> More recently, *ELANE* protein inhibitors were shown to stimulate mouse B cells to differentiate into plasma cells that produce IgG or IgA, and can also upregulate the transcription of *AID*, *interleukin-10*, *BAFF*, and *APRIL* coding genes.<sup>31</sup> We observed that Ig levels and the proportion of lymphocyte subsets varied among patients with CN. Specifically, we noted elevated serum levels of IgG, IgA, and IgM and increased BM plasma cells in SCN1 caused by *ELANE* mutations. In addition, a higher serum *APRIL* concentration was found in SCN1 compared with healthy controls. Our results suggest that the increase in serum Ig levels in SCN1 is a result of an increased number of BM plasma cells,






**Table 4. ANC after G-CSF administration in different CN groups**


CN group	G-CSF responsiveness				Total cases (n)
	pANC ≥3.0	1.5 ≤ pANC<3.0	1.0 ≤ pANC<1.5	pANC<1.0	
SCN1 n (%)	3 (12%)	10 (40%)	6 (24%)	6 (24%)	25
SCN3 n (%)	0 (0%)	2 (50%)	1 (50%)	1 (50%)	4
CyN n (%)	2 (100%)	0 (0%)	0 (0%)	0 (0%)	2
WHIM n (%)	3 (75%)	0 (0%)	1 (50%)	0 (0%)	4
SBDS n (%)	2 (50%)	1 (25%)	1 (25%)	0 (0%)	4

pANC, peak ANC ( $\times 10^9$  cells per L) within 10 days after G-CSF initiation.



**Figure 4. SCN hyperimmunoglobulinemia is associated with elevated plasma cell counts and APRIL levels.**

(A) Serum Igs levels were measured by immunoturbidimetry in the patients. (B) Correlation between serum levels of each Ig isotype and BM PC in each patient, using Spearman correlation test. (C) Proportion of plasma cells in the BM, determined by cell morphology assays. Kruskal-Wallis test was applied to evaluate differences among 5 groups. (D) Serum APRIL levels in patients with SCN1 and age-matched healthy control (HC), measured by enzyme-linked immunosorbent assay. The gray area indicates normal reference range for each parameter. The Mann-Whitney test was used for SCN1 and HC comparison;  $*P < .05$ . Bars represent means  $\pm$  SD.



**Figure 5. Correlation between the stage and severity of neutrophil development impairment and responsiveness to G-CSF therapy.** SCN1 and SCN3, which display a severe maturation arrest at an early stage (between promyelocyte and myelocyte) during BM neutrophil development, show a poor response to G-CSF. CyN, WHIM, and SBDS, where BM neutrophil development is disrupted at a late stage, unaffected, or partially interrupted, respond well to G-CSF therapy. An 'x' indicates a developmental block or migration defect; The line thickness reflects the degree of normal neutrophil development.

elevated serum Ig levels, and a poor response to G-CSF therapy. CyN presented with moderately low peripheral neutrophil counts, well-developed BM granulocytes, normal serum Ig levels, and good responsiveness to G-CSF administration. Patients with WHIM, while displaying lower counts of peripheral WBCs, granulocytes, lymphocytes, and monocytes, along with lower Ig levels, presented with normal BM neutrophil development and responded well to G-CSF treatment. Patients with SBDS were more prone to diarrhea and exhibited moderately low peripheral neutrophil counts, partial impairment in BM neutrophil development, reduced Ig levels, and positive response to G-CSF therapy.

## Acknowledgments

The authors thank the clinicians in the Department of Clinical Immunology, Children's Hospital of Fudan University and the members in Wang laboratory at the Department of Immunology of School of Basic Medical Sciences for their helpful discussions.

This work was supported by the Major Research Plan of the National Natural Science Foundation of China (grant 32330033; J.-Y.W.), the National Natural Science Foundation of China (grant 32270932; J.-Y.W.), Projects of International Cooperation and Exchanges NSFC (grant 82011540008; J.-Y.W.), Shanghai Municipal Science and Technology Major Project (grant ZD2021CY001; X.W.), National Natural Science Foundation of China for Young Scholar (grant 82202013; Q.M.), and China Postdoctoral Science Foundation Grant (grant 2022M720782; Q.M.).

## References

1. Badolato R, Fontana S, Notarangelo LD, Savoldi G. Congenital neutropenia: advances in diagnosis and treatment. *Curr Opin Allergy Clin Immunol*. 2004;4(6):513-521.
2. Lazzareschi I, Rossi E, Curatola A, et al. Assessment of congenital neutropenia in children: common clinical sceneries and clues for management. *Mediterr J Hematol Infect Dis*. 2022;14(1):e2022008.
3. Donadieu J, Fenneteau O, Beaupain B, Mahlaoui N, Chantelot CB. Congenital neutropenia: diagnosis, molecular bases and patient management. *Orphanet J Rare Dis*. 2011;6:26.

## Authorship

Contribution: X.M. and J.H. designed the study, and collected and analyzed the clinical and immunological data; X.M. provided a draft of the manuscript; H.Z., L.L., and L.D. participated in data collection and analysis; Q.M. provided suggestions for the criteria for grouping; Y.L. provided support in healthy control samples and flow cytometry; J.H., L.L., W.W., W.Y., J.S., and X.W. diagnosed and treated patients and helped collect the clinical data of the patients; Q.M., H.Z., L.D., Y.L., and M.Y. corrected the manuscript; X.W. and J.-Y.W. supervised the study; and J.-Y.W. reviewed and revised the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: M.Y., [0000-0001-8698-9611](https://orcid.org/0000-0001-8698-9611); J.-Y.W., [0000-0003-3038-3826](https://orcid.org/0000-0003-3038-3826).

Correspondence: Ji-Yang Wang, Department of Immunology, Fudan University, Room 921, West #13 Building, Dong'an Rd, Shanghai 200032, China; email: [wang@fudan.edu.cn](mailto:wang@fudan.edu.cn); and Jia Hou, Department of Clinical Immunology, Children's Hospital of Fudan University, National Children's Medical Center, Fudan University, 399 Wanyuan Rd, Shanghai 201102, China; email: [doctorhoujia@hotmail.com](mailto:doctorhoujia@hotmail.com); and Xiaochuan Wang, Department of Clinical Immunology, Children's Hospital of Fudan University, National Children's Medical Center, Fudan University, 399 Wanyuan Rd, Shanghai 201102, China; email: [xchwang@shmu.edu.cn](mailto:xchwang@shmu.edu.cn).

4. Ward AC, Loeb DM, Soede-Bobok AA, Touw IP, Friedman AD. Regulation of granulopoiesis by transcription factors and cytokine signals. *Leukemia*. 2000;14(6):973-990.
5. Kwok I, Becht E, Xia Y, et al. Combinatorial single-cell analyses of granulocyte-monocyte progenitor heterogeneity reveals an early uni-potent neutrophil progenitor. *Immunity*. 2020;53(2):303-318.e5.
6. Berdnikovs S. The twilight zone: plasticity and mixed ontogeny of neutrophil and eosinophil granulocyte subsets. *Semin Immunopathol*. 2021;43(3):337-346.
7. Evrard M, Kwok IWH, Chong SZ, et al. Developmental analysis of bone marrow neutrophils reveals populations specialized in expansion, trafficking, and effector functions. *Immunity*. 2018;48(2):364-379.e8.
8. Skokowa J, Dale DC, Touw IP, Zeidler C, Welte K. Severe congenital neutropenias. *Nat Rev Dis Primers*. 2017;3(1):17032.
9. Xia J, Bolyard AA, Rodger E, et al. Prevalence of mutations in ELANE, GFI1, HAX1, SBDS, WAS and G6PC3 in patients with severe congenital neutropenia. *Br J Haematol*. 2009;147(4):535-542.
10. Dale DC, Makaryan V. ELANE-related neutropenia. In: Adam MP, Everman DB, Mirzaa GM, et al, eds. *GeneReviews*. University of Washington; 1993:1-2.
11. Wang J, Yu H, Zhang VW, et al. Capture-based high-coverage NGS: a powerful tool to uncover a wide spectrum of mutation types. *Genet Med*. 2016;18(5):513-521.
12. Sun J, Yang L, Lu Y, et al. Screening for primary immunodeficiency diseases by next-generation sequencing in early life. *Clin Transl Immunology*. 2020;9(5):e1138.
13. Ding Y, Zhou L, Xia Y, et al. Reference values for peripheral blood lymphocyte subsets of healthy children in China. *J Allergy Clin Immunol*. 2018;142(3):970-973.e8.
14. Sovani V. Normal bone marrow, its structure and function. *Diagn Histopathol*. 2021;27(9):349-356.
15. Thusberg J, Vihinen M. Bioinformatic analysis of protein structure-function relationships: case study of leukocyte elastase (ELA2) missense mutations. *Hum Mutat*. 2006;27(12):1230-1243.
16. Makaryan V, Zeidler C, Bolyard AA, et al. The diversity of mutations and clinical outcomes for ELANE-associated neutropenia. *Curr Opin Hematol*. 2015;22(1):3-11.
17. Bellanné-Chantelot C, Clauin S, Leblanc T, et al. Mutations in the ELA2 gene correlate with more severe expression of neutropenia: a study of 81 patients from the French Neutropenia Register. *Blood*. 2004;103(11):4119-4125.
18. Horwitz MS, Corey SJ, Grimes HL, Tidwell T. ELANE mutations in cyclic and severe congenital neutropenia: genetics and pathophysiology. *Hematol Oncol Clin North Am*. 2013;27(1):19-41. vii.
19. Ancliff PJ, Gale RE, Liesner R, Hann IM, Linch DC. Mutations in the ELA2 gene encoding neutrophil elastase are present in most patients with sporadic severe congenital neutropenia but only in some patients with the familial form of the disease. *Blood*. 2001;98(9):2645-2650.
20. Dale DC, Bolyard AA, Shannon JA, et al. Outcomes for patients with severe chronic neutropenia treated with granulocyte colony-stimulating factor. *Blood Adv*. 2022;6(13):3861-3869.
21. Bianchi ME, Mezzapelle R. The chemokine receptor CXCR4 in cell proliferation and tissue regeneration. *Front Immunol*. 2020;11:2109.
22. Lagane B, Chow KY, Balabanian K, et al. CXCR4 dimerization and beta-arrestin-mediated signaling account for the enhanced chemotaxis to CXCL12 in WHIM syndrome. *Blood*. 2008;112(1):34-44.
23. Hernandez PA, Gorlin RJ, Lukens JN, et al. Mutations in the chemokine receptor gene CXCR4 are associated with WHIM syndrome, a combined immunodeficiency disease. *Nat Genet*. 2003;34(1):70-74.
24. Hong CW. Current understanding in neutrophil differentiation and heterogeneity. *Immune Netw*. 2017;17(5):298-306.
25. Nutt SL, Hodgkin PD, Tarlinton DM, Corcoran LM. The generation of antibody-secreting plasma cells. *Nat Rev Immunol*. 2015;15(3):160-171.
26. Warren JT, Bolyard AA, Kelley ML, Makaryan V, Dale DC, Link DC. Spectrum of pathogenic genetic variants in a large cohort of North American congenital and cyclic neutropenia patients: a report from the Severe Chronic Neutropenia International Registry. *Blood*. 2021;138(suppl 1):2059.
27. An WB, Liu C, Wan Y, Chang LX, Chen XY, Zhu XF. [Clinical features and gene mutations of children with Shwachman-Diamond syndrome and malignant myeloid transformation]. *Zhongguo Dang Dai Er Ke Za Zhi*. 2020;22(5):460-465.
28. Takahashi H, Nukiwa T, Basset P, Crystal RG. Myelomonocytic cell lineage expression of the neutrophil elastase gene. *J Biol Chem*. 1988;263(5):2543-2547.
29. Dale DC, Bonilla MA, Davis MW, et al. A randomized controlled phase III trial of recombinant human granulocyte colony-stimulating factor (filgrastim) for treatment of severe chronic neutropenia. *Blood*. 1993;81(10):2496-2502.
30. Jee J, Bonnegarde-Bernard A, Duverger A, et al. Neutrophils negatively regulate induction of mucosal IgA responses after sublingual immunization. *Mucosal Immunol*. 2015;8(4):735-745.
31. Attia Z, Rowe JC, Kim E, et al. Inhibitors of elastase stimulate murine B lymphocyte differentiation into IgG- and IgA-producing cells. *Eur J Immunol*. 2018;48(8):1295-1301.