

Selected markers of ovarian cancer and their relation to targeted therapy (Review)

ANNA MARKOWSKA¹, ZBIGNIEW KOJS², DAMIAN TWARDAWA³,
JOANNA PIETRAS¹ and JANINA MARKOWSKA⁴

¹Department of Perinatology and Women's Diseases, Poznan University of Medical Sciences, 60-535 Poznan; ²Department of Gynecology and Obstetrics with Gynecologic Oncology, Ludwik Rydygier Memorial Specialized Hospital, 31-826 Kraków; ³Medical Department, Bausch Health Poland, 02-674 Warsaw; ⁴Gynecological Oncology Center Poznań, 60-850 Poznan, Poland

Received October 31, 2023; Accepted March 15, 2024

DOI: 10.3892/etm.2024.12523

Abstract. Despite advances in surgical treatment techniques and chemotherapy-including anti-angiogenic and immune poly (ADP-ribose) polymerase inhibitors, the 5-year survival rate in ovarian cancer (OC) remains low. The reasons for this are the diagnosis of cancer in advanced clinical stages, chemoresistance and cancer recurrence. New therapeutic approaches are being developed, including the search for new biomarkers that are also targets for targeted therapy. The present review describes new molecular markers with relevance to targeted therapy, which to date have been studied only in experimental research. These include the angiogenic protein angiopoietin-2, the transmembrane glycoprotein ectonucleotide pyrophosphatase/phosphodiesterase 1, the adhesion protein E-cadherin, the TIMP metalloproteinase inhibitor 1 and Kruppel-like factor 7. Drugs affecting cancer stem cells (CSCs) in OC, such as metformin and salinomycin, as well as inhibitors of CSCs markers aldehyde dehydrogenase 1 (with the drug ATRA) and the transcription factor Nanog homeobox (microRNA) are also discussed. A new approach to prevention and possible therapies under investigation such as development of vaccines containing a subpopulation of CD117(+) and CD44(+) stem cells with a promising option for use in women with OC was described.

Contents

1. Introduction
2. Angiopoietin 2 (Ang-2)
3. Sodium-dependent phosphate transport protein 2B (NaPi2b)
4. Ectonucleotide pyrophosphatase/phosphodiesterase 1 (ENPP1)
5. E-cadherin
6. TIMP metalloproteinase inhibitor 1 (TIMP-1)
7. Endothelial cell-specific molecule 1 (ESM1)
8. Cancer stem cells (CSCs)
9. Kruppel like factor 7 (KLF7)

1. Introduction

Ovarian cancer (OC) is the gynecological malignancy with the worst prognosis. According to GLOBOCAN, in 2020, 314,000 women were diagnosed with OC, while a total of 207,000 deaths due to OC were recorded. In 2040, a total of 428,000 new OC cases and 307,000 deaths from OC are predicted. In early stage of OC, overall survival (OS) is ~92% (1). However, >70% of the cases are diagnosed at an advanced clinical stage, which is associated with an unsuccessful disease course. After successful primary therapy, ~80% of patients are found to have recurrence while OS is only 29%. OC does not only show clinical diversity, but is also histologically and molecularly heterogeneous which influences chemoresistance in therapy. The most aggressive type of OC is type II high-grade serous ovarian carcinoma (HGSOC).

Although anti-angiogenic treatment with bevacizumab and poly (ADP-ribose) polymerase inhibitors have been introduced to standard surgery and platinum- and taxane-based chemotherapy in recent years, the recurrence rate, which is the main cause of death, is still high. There are no effective markers for early detection of HGSOC. Transvaginal ultrasound and determination of serum cancer antigen 125 (CA125) levels are used in practice. CA125 and risk of ovarian malignancy algorithm assay incorporating the CA125 serum level and human epididymis protein 4 (according to menopausal status) are not

Correspondence to: Dr Joanna Pietras, Department of Perinatology and Women's Diseases, Poznan University of Medical Sciences, Polna 33, 60-535 Poznan, Poland
E-mail: fechner.joanna@gmail.com

Key words: ovarian cancer, markers, angiopoietin-2, sodium-dependent phosphate transport protein 2B, ectonucleotide pyrophosphatase/phosphodiesterase 1, E-cadherin, TIMP metalloproteinase inhibitor 1, endothelial cell-specific molecule 1, cancer stem cells, aldehyde dehydrogenase 1 family member A1, Nanog homeobox, vaccines containing CSCs, Kruppel like factor 7

sufficiently reliable in diagnosis to screen for early-stage OC. It is therefore necessary to introduce new biomarkers that are also effective for targeted therapies (2-5).

Potential molecular markers and their relationship to therapeutic benefit in patients with OC, most of which are undergoing *in vitro* and *in vivo* testing, are described below. Estimation of new molecular markers is essential in the development and monitoring of novel treatment options. For the present review, the following electronic databases were used: Medline (<https://www.ebsco.com/products/research-databases/medline>), Scopus (<https://www.elsevier.com/products/scopus>) and Web of Science (<https://webofscience.help.clarivate.com/en-us/Content/home.htm/>). Selected markers of OC are described in Table I.

2. Angiopoietin 2 (Ang-2)

Ang-2 is an angiogenic protein expressed in endothelial cells at the site of vascular remodeling. It is a ligand of the TEK receptor tyrosine kinase. Regulation of its activity is mediated by hypoxia-inducible factor-1 α and VEGF. It shares 60% identical amino acid sequence with Ang-1, but the functions of the two angiopoietins are opposite; Ang-2 is a Tie-2 receptor antagonist while Ang-1 is a Tie-2 receptor agonist. In addition, Ang-2 is not bound to the extracellular matrix, therefore it can be active at different sites in the body (6).

A previous study examining the cancer tissue and serum of 138 patients with OC detected the higher expression of Ang-2 in patients with retroperitoneal spread compared with patients with intraperitoneal disease ($P=0.039$). High Ang-2 expression levels were significantly correlated with longer OS ($P=0.017$) and OS in patients receiving bevacizumab ($P=0.013$). Ang-2 may serve as a molecular marker for patients with OC with early spread to lymph nodes and for patients who receive maintenance targeted therapy with bevacizumab (7).

3. Sodium-dependent phosphate transport protein 2B (NaPi2b)

NaPi2b is located at the cell surface and regulates phosphate homeostasis under physiological conditions. It is encoded by the SLC34A2 gene and expressed in ~80-90% of OC cases. Additionally, it is detected in OC tissues via biopsy. Expression of the SLC34A2 gene was determined at the transcriptional and translational level in 41 OC samples considering different clinicopathological features (clinical grade, neoadjuvant chemotherapy and presence of ascites). The expression of this gene was found to be downregulated in patients receiving neoadjuvant chemotherapy. Thus, expression of this gene may be a marker for predicting response to such therapy (8).

A different study described for the first time the results of treatment in patients with platinum-resistant OC with a humanized anti-NaPi2b antibody conjugated to the antimetabolic drug monomethylauristatin E-lifastuzumab vedotin (LIFA). A total of 47 patients received LIFA and 48 patients (representing the second arm of platinum-resistant patients) received pegylated doxorubicin (PLD). Progression-free survival (PFS) was found to be prolonged in patients receiving LIFA compared with the PLD group (5.3 vs. 3.1 months, respectively); however, these values were not statistically significant (9).

A subsequent multicenter study involving 41 patients with recurrent platinum-sensitive OC assessed the safety and tolerability of LIFA. All patients were administered LIFA with carboplatin, and 12 of them were additionally treated with bevacizumab. All patients experienced ≥ 1 side effects (adverse events); the most common was neutropenia. Side effects, according to the investigators, were acceptable. A complete or partial response to the treatment was observed in 59% of the patients. Thus, this study was encouraging regarding the use of LIFA (10). A careful analysis of the literature review established that NaPi2b fulfils the conditions of an OC biomarker and may serve as a target for therapy in this cancer (3).

4. Ectonucleotide pyrophosphatase/phosphodiesterase 1 (ENPP1)

ENPP1 is a type II trans-membrane glycoprotein with pyrophosphatase and phosphodiesterase capabilities. It is a regulator of extracellular ATP and GTP signalling. It plays a key role in phosphate balance and bone mineralization but is also frequently overexpressed in various cancers (breast cancer, glioma), including OC (11).

A study by Wang *et al* (12) of ENPP1 in 241 patients with OC revealed that its expression was significantly increased ($P<0.05$) in HGSOC (85.4%) and was low in serous adenocarcinoma (1.03%); however, it was not detected in normal ovarian epithelium. Similar expression of ENPP1 was found in established OC cell lines. After downregulation of ENPP1 expression (RNA interference), disease progression exponents such as proliferation, migration and invasion decreased significantly, and expression of proapoptotic caspase 3 increased significantly; epithelial-mesenchymal transition (EMT) was also inhibited, as expressed by decreased expression of the EMT-E-cadherin marker (12).

Subsequent findings (13) confirmed the value of ENPP1 as a molecular marker and target for targeted therapy. In addition to expression in cancer, ENPP1 was found to induce strong immune remodeling (it is expressed in neutrophils and macrophages) promoting cancer progression using the STING pathway associated with antitumor defense mechanisms.

In their review of preclinical studies with ENPP1, Ruiz-Fernández de Córdoba *et al* (13) mentioned several of its inhibitors such as: i) STF-1623, which acts through Zn²⁺ chelation (CM-3163; Angarus Therapeutics), ii) AVA-NP-695 (Avammune Therapeutics), iii) ZX-8177, which can be administered with anti-programmed death-ligand 1 and iv) RBS2418 (Riboscience LLC). Thus, increased expression of ENPP1 may, according to the researchers, be associated with the occurrence of HGSOC and indicate a poor prognosis, and at the same time, may serve as a therapeutic target (13).

5. E-cadherin

E-cadherin belongs to a family of transmembrane proteins that maintain cell adhesion. It participates in signal transduction affecting cell proliferation, survival and differentiation. As aforementioned, E-cadherin is downregulated during EMT, which is related to the mesenchymal phenotype. The EMT process in tumors can be initiated and promoted by multiple signaling pathways, including those originating from hypoxia

Table I. Selected OC markers.

Markers	Characteristics	(Refs.)
Ang-2	Ligand of tyrosine kinase 2 receptor	(6,7)
NaPi2b	1. Regulates phosphate homeostasis under physiological conditions 2. Downregulated expression in patients receiving neoadjuvant chemotherapy	(3,8-10)
ENPP1	Increased expression associated with the occurrence of HGSOc	(11-13)
E-cadherin	Low concentration associated with poor prognosis	(14-16)
TIMP-1	1. High levels in patients with advanced OC associated with reduced OS 2. Associated with platinum resistance	(17-19)
ESM1	1. Participates in the progression of OC 2. Expression correlated with progression-free survival and OS	(20,21)
CSCs	Play a role in progression, multidrug resistance and the formation of metastases and recurrences	(22-28)
ALDH1	High expression associated with platinum resistance in serous carcinomas	(35-38)
NANOG-CSCs transcription factor	Marker of CSCs pluripotency (it promotes tumor growth, metastasis, invasion and chemoresistance)	(39-43)
KLF7	Regulator of the CSCs pluripotency (regulation of the transcription factors OCT4 and NANOG)	(47,48)

Ang-2, angiopoietin-2; NaPi2b, sodium-dependent phosphate transport protein 2B; ENPP1, ectonucleotide pyrophosphatase/phosphodiesterase 1; HGSOc, high-grade serous ovarian carcinoma; TIMP-1, TIMP metalloproteinase inhibitor 1; OC, ovarian cancer; OS, overall survival; ESM1, endothelial cell-specific molecule 1; CSCs, cancer stem cells; ALDH1, aldehyde dehydrogenase 1 family member A1; NANOG, Nanog homeobox; KLF7, Kruppel-like factor 7; OCT4, octamer-binding transcription factor 4.

and the microenvironment of the developing cancer. A number of messengers have been shown to be disrupted, including the E-cadherin binding partner, β -catenin and phosphatidylinositol 3 kinase (PI3K) (14).

Based on the results of 18 cell lines and the evaluation of E-cadherin expression using microarrays and immunofluorescence in 101 patients with OC, it was revealed that low levels of E-cadherin and high levels of keratin 7 predict poor response to treatment in HGOC. E-cadherin together with keratin 7 (a component of the cytoskeleton fibres) assay can be an independent prognostic marker of response for HGOC treatment and also for OS ($P=0.031$ and $P=0.041$ for E-cadherin and keratin 7, respectively) (15).

Investigation on cell lines and E-cadherin mRNA expression in 20 cases of existing ascites in patients with OC patients revealed that loss of E-cadherin expression is associated with OC progression and aggressiveness, and that E-cadherin levels are associated with CA125 levels.

Options were provided for targeted therapies that stimulate E-cadherin expression. These include: i) α -Solanine (a glycoalkaloid extract of *Solanum nigrum*), which lowers the expression of matrix metalloproteinases (MMPs), ii) simvastatin and metformin which increase E-cadherin values and iii) numerous other synthetic peptides used in cancer therapy, such as ADH-1 (antidiuretic hormone) (16).

6. TIMP metalloproteinase inhibitor 1 (TIMP-1)

TIMP-1 is a tissue inhibitor of MMP-9, a member of the gelatinases. MMPs are a group of >20 zinc-dependent proteolytic enzymes. They are involved in matrix remodeling and degradation processes. TIMP-1 has been found to model matrix

metalloproteinase activity and play a role in OC progression by mediating metastasis through colonization, migration and invasion of cancer cells. It is present in both ascites and plasma. It was found from comparative studies (70 ascites and 20 plasma samples) that its concentration is higher in ascites and correlates with its volume. Serum levels in patients with OC (stage I-IV) are the highest after cytoreductive surgery and the lowest after treatment (17). Sonogo *et al* (18) observed that high TIMP-1 levels in patients with OC at stage III and IV were associated with reduced OS especially if patients were treated with platinum or bevacizumab. This indicated that TIMP-1 is associated with platinum resistance regulation and progression and there is potential for its use as a novel biomarker of platinum resistance. According to the researchers, this demonstrates the possibility of using TIMP-1 as a target for targeted therapy. This was confirmed by a study of TIMP-1 in circulating tumor cells (CTCs) in 38 patients with HGSOc (depending on the method used, CTCs are detected in 18-88% of patients with OC). It was identified that, in addition to the stem cell markers CD24 and CD44, TIMP-1 promotes cancer promotion, suggesting its therapeutic target (19).

7. Endothelial cell-specific molecule 1 (ESM1)

ESM1 gene is located on chromosome 5q11.2 and is involved in migration, proliferation, invasion, angiogenesis and escape from apoptosis. It regulates signaling of the conservative, carcinogenesis-related Wnt/ β catenin pathway and is associated with the AKT/mTOR pathway promoting cancer cell proliferation and inhibiting apoptosis (20,21).

A molecular study of 379 OC samples demonstrated that ESM1 expression was higher in cancer tissues compared with

peritumor tissues and healthy ovarian tissues. ESM1 expression was closely correlated with clinical parameters such as lymph node metastasis and cancer recurrence. The authors concluded that ESM1 plays an important role in OC development and progression; it was positively correlated with PFS, but negatively with OS. It may be considered as a novel OC biomarker and therapeutic target (21).

8. Cancer stem cells (CSCs)

CSCs are a subpopulation of cancer cells usually representing <2-5% of the tumour mass, involved in cancer growth, multidrug chemoresistance, metastasis formation and recurrence formation (22-24).

CSCs have been shown to possess properties that allow them to survive under adverse conditions. These include the ability of self-renewal, asymmetric division, staying in an inactive state (as 'dormant cells' in the G0/G1 phase), the ability to repair DNA, overexpression of ATP-binding cassette family genes associated with chemoresistance, lack of apoptosis and the ability to utilize different signaling pathways such as Wnt/ β catenin, Notch and Hedgehog which are associated with survival and self-renewal (22,23).

Molecular markers of CSCs have been identified, both superficial, such as CD44, CD133 (Promin), CD111 (c-Kit), CD24, CD117, and intracellular, such as aldehyde dehydrogenase 1 family member A1 (ALDH1), octamer-binding transcription factor 4 (OCT4), SRY-Box transcription factor 2 (SOX2) and Nanog homeobox (NANOG). Determining their expression is necessary to understand the efficacy and monitoring of targeted therapy (22,24-27). In 2005, Bapat *et al* (28) detected CSCs markers in OC.

Most applied trials of therapies against CSCs *in vitro* and *in vivo* and also in women with OC affect the expression of not only one CSCs marker, but also the signaling pathway. Such known drugs affecting CSCs include metformin and salinomycin.

Metformin, used in the treatment of type 2 diabetes, exerts beneficial effects in OC. An electronic database study on >700 patients with OC revealed an association of metformin use with longer survival (P=0.03) and synergistic activity with carboplatin. Metformin has an inhibitory effect on the AKT/mTOR pathway (29). Furthermore, in a phase I clinical trial, no adverse effects were reported using metformin in combination with carboplatin and paclitaxel (30).

A previous *in vitro* and *in vivo* study detected that metformin selectively inhibits the CD44(+) and CD117(+) populations, also glycoproteins of CSCs affecting cancer progression, including: Cell adhesion, cell growth, migration and resistance to chemotherapy (31).

Salinomycin, an ionophore antibiotic isolated from *Streptomyces albus*, selectively eradicates CSCs (32-34). Previous studies on OC cell lines (OVCAR3) have revealed an inhibitory effect of salinomycin in combination with paclitaxel on CD44(+) and CD117(+) expression (32). This was confirmed by a study by Lee *et al* (33) on isolated CD44(+) and CD117(+) cells from ascites of women with OC. Salinomycin in combination with paclitaxel decreased their viability and promoted apoptosis.

Constructing a carrier mimicking high-density lipoprotein (HDL) in combination with salinomycin (S-HDL) has been

demonstrated to have potent anticancer effects in OC as it inhibits translation of the CSCs proteins c-Myc, NANOG, OCT4 and SOX2 which are associated with chemoresistance and OC recurrence (34). The authors of the investigation on the effects of metformin and salinomycin consider that these two drugs will be an effective future therapy in improving survival and preventing recurrence of OC.

ALDH1. High expression of ALDH1 has been demonstrated to be associated with platinum resistance in serous carcinomas. This was based on a study of 124 patients with stage III and IV serous OC with high malignancy (35). A different study on OC cell lines found that high expression of ALDH1A1 was associated with resistance to paclitaxel and topotecan. At the same time, it was associated with overexpression of P-glycoprotein (P-gp), known as the multidrug resistance protein responsible for removing the 'pumping out' of the cytostatic from the cell. The use of all-trans retinoic acid (ATRA) decreased ALDH1A1 expression and reduced P-gp expression (36).

Kim *et al* (37), who investigated A2780 cells isolated from chemotherapy-resistant OC of female patients, confirmed that high ALDH1 expression is associated with drug resistance and facilitated growth of OC. Furthermore, high expression of ALDH1 was associated with increased expression of nuclear factor erythroid 2-related factor 2 (NRF2). Furthermore, ATRA treatment decreased ALDH1 expression and inhibited NRF2 activation. The effect was to attenuate the malignancy of CSCs in OC.

A different study on cells isolated from ascites of untreated or receiving neoadjuvant therapy patients with OC also showed their resistance to platinum and taxanes, accompanied by high ALDH1 expression. It was revealed that several compounds including EGFR/mTOR-PI3K inhibitors are candidates for targeting the ALDH1 cell population (38).

NANOG (CSCs transcription factor). NANOG is a marker of CSCs pluripotency. It promotes tumour growth, metastasis, invasion and chemoresistance. Using different signaling pathways such as JAK/STAT, Notch, Hedgehog and canonical Wnt/ β -catenin, it participates in the reciprocal regulation of other transcription factors such as SOX2 and OCT4, and its expression is related to several microRNAs such as miR-214, whose downregulation reduces NANOG expression and induces cancer cell apoptosis. High NANOG expression is present in 69.7% of OC cases, particularly HGSO. It is recognized as a key marker of progression. Although gene therapy with microRNA is in the experimental phase, NANOG is a diagnostic marker and a future target for personalized therapy (39-43).

Vaccines containing CSCs. In 2015, the efficacy of a vaccine containing CSCs was described for the first time in a mouse model. Cell lines containing the CD117(+) and CD44(+) subpopulations were isolated from the SKOV3 human OC line. When administered in mice [SKOV3 containing CSCs, CD117(+) and CD44(+) subpopulations], it induced the desired effect: Inhibition of xenograft tumor growth and reduction of CD117(+) and CD44(+) cell subpopulations (44). The same group of researchers produced a vaccine from a patient-derived cell line (HO8910) that also contained CD117,

CD44 and receptor tyrosine kinase-like orphan receptor 1 (ROR1). It was identified that high expression of ROR1 was strongly correlated with vaccine efficacy, while downregulation of ROR1 expression by small interfering RNA (siRNA) reduced vaccine efficacy. It was concluded that in OC immuno-prophylaxis, ROR1 expression should be high (45). A subsequent study confirmed the efficacy of both H08910 and D8 (a cell line isolated in mice) regarding the inhibition of tumor growth and prolongation of survival. According to the authors, it is possible to convert a CSC-containing vaccine into an immunotherapeutic approach (46).

9. Kruppel-like factor 7 (KLF7)

KLF7 is one of 17 KLFs identified in humans and animals and is encoded on chromosome 2q33.3. It is a transcription factor with a broad spectrum of regulatory functions. It is involved in physiological and pathological processes such as: Cardiovascular disease, hematological disease and metabolic disease (type 2 diabetes), and is also a regulator of the maintenance of CSCs pluripotency that involves transcription factors OCT4 and NANOG which in turn regulate KLF7 expression (47). Mao *et al* (47) studied the expression of KLF7 on cell lines and animal models. They discovered chemical substances that downregulate KLF7 expression, including catechin (polyphenol, antioxidant), pitavastatin (lipid-lowering drug) and trametinib (MEK inhibitor).

KLF7 is involved in the development and progression of numerous cancers in various localizations, including OC. A novel bioinformatics meta-analysis of transcriptome data in 2 cohorts of patients with advanced stages of HGSOE (III/IV) including 185 patients from the Gene Expression Omnibus (GEO) database and 266 patients from the National Cancer Institute Genomic Data Commons (NCI-GDC) Data Portal detected that KLF7 is significantly associated with this cancer. A high expression of KLF7 is significantly related to shorter OS (for GEO multivariate analysis, $P < 0.0001$; for NCI-GDC, $P = 0.03$). *In vitro* studies (OV-90 and PEO1 cells) have determined that KLF7 may play a role as an oncogene to stimulate tumor growth and invasion, and is involved in maintaining the pluripotency and the self-renewal characteristics of stem cells.

Previous results have suggested that KLF7 is a promising prognostic marker and therapeutic target in HGSOE (48). The study by De Donato *et al* (48) revealed the first evidence that high expression of KLF7 in HGSOE is connected to shorter OS. The authors attempted to silence KLF7 expression in OV-90 and PEO1 cells by using TransFectin Lipid Reagent and specific siRNAs, but it may be possible to apply the repositioning drugs mentioned by Mao *et al* (47) in *in vivo* studies and clinical trials in an attempt to lower KLF7 expression in HGSOE.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

Not applicable.

Authors' contributions

AM: Originator of the study and writing of the abstract, introduction and chapter on Ang-2. ZK: Writing of chapters on NaPi2b and ENPPI. DT: E-cadherin, TIMP-1. JP: Writing of chapters on ESM1 and KLF7. JM: Writing of chapter on CSCs. All authors read and approved the final version of the manuscript. Data authentication is not applicable.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Cabasag CJ, Fagan PJ, Ferlay J, Vignat J, Laversanne M, Liu L, van der Aa MA, Bray F and Soerjomataram I: Ovarian cancer today and tomorrow: A global assessment by world region and Human Development Index using GLOBOCAN 2020. *Int J Cancer* 151: 1535-1541, 2022.
- Richardson DL, Eskander RN and O'Malley DM: Advances in ovarian cancer care and unmet treatment needs for patients with platinum resistance: A narrative review. *JAMA Oncol* 9: 851-859, 2023.
- Banerjee S, Drapkin R, Richardson DL and Birrer M: Targeting NaPi2b in ovarian cancer. *Cancer Treat Rev* 112: 102489, 2023.
- Guo L, Wang J, Li N, Cui J and Su Y: Peptides for diagnosis and treatment of ovarian cancer. *Front Oncol* 13: 1135523, 2023.
- Zhang R, Siu MKY, Ngam HYS and Chan KKL: Molecular biomarkers for the early detection of ovarian cancer. *Int J Mol Sci* 23: 120941, 2022.
- Yu X and Ye F: Role of angiopoietins in development of cancer and neoplasia associated with viral infection. *Cells* 9: 457, 2020.
- Volk A, Legler K, Hamester F, Kuerti S, Eylmann K, Rossberg M, Schmalfeldt B and Oliveira-Ferrer L: Ang-2 is a potential molecular marker for lymphatic metastasis and better response to bevacizumab therapy in ovarian cancer. *J Cancer Res Clin Oncol* 149: 15957-15967, 2023.
- Nurgaliewa AK, Popov VE, Skripova VS, Bulatova LF, Savenkova DV, Vlasenkova RA, Safina SZ, Shakirova EZ, Filonenko VV, Bogdanov MV and Kiyamova RG: Sodium-dependent phosphate transporter NaPi2b as a potential predictive marker for targeted therapy of ovarian cancer. *Biochem Biophys Rep* 28: 101104, 2021.
- Banerjee S, Oza AM, Birrer MJ, Hamilton EP, Hasan J, Leary A, Moore KN, Mackowiak-Matejczyk B, Pikiel J, Ray-Coquard I, *et al*: Anti-NaPi2b antibody-drug conjugate lifastuzumab vedotin (DNIB0600A) compared with pegylated liposomal doxorubicin in patients with platinum-resistant ovarian cancer in a randomized, open-label, phase II study. *Ann Oncol* 29: 917-923, 2018.
- Moore KN, Birrer MJ, Marsters J, Wang Y, Choi Y, Royer-Joo S, Lemahieu V, Armstrong K, Cordova J, Samineni D, *et al*: Phase 1b study of anti-NaPi2b antibody-drug conjugate lifastuzumab vedotin (DNIB0600A) in patients with platinum-sensitive recurrent ovarian cancer. *Gynecol Oncol* 158: 631-639, 2020.

11. Cogan D and Bakhoun SF: Re-awakening innate immune signaling in cancer: The development of highly potent ENPPI inhibitors. *Cell Chem Biol* 27: 1327-1328, 2020.
12. Wang H, Ye F, Zhou C, Cheng Q and Chen H: High expression of ENPPI in high-grade serous ovarian carcinoma predicts poor prognosis and as a molecular therapy target. *PLoS One* 16: e0245733, 2021.
13. Ruiz-Fernández de Córdoba B, Martínez-Monge R and Lecanda F: ENPPI immunobiology as a therapeutic target. *Clin Cancer Res* 29: 2184-2193, 2023.
14. Kaszak I, Witkowska-Piłaszewicz O, Niewiadomska Z, Dworecka-Kaszak B, Ngosa Toka F and Jurka P: Role of cadherins in cancer-a review. *Int J Mol Sci* 21: 7624, 2020.
15. Communal L, Roy N, Cahuzac M, Rahimi K, Köbel M, Provencher DM and Mes-Masson AM: A Keratin 7 and E-Cadherin signature is highly predictive of tubo-ovarian high-grade serous carcinoma prognosis. *Int J Mol Sci* 22: 5325, 2021.
16. Rosso M, Majem B, Devis L, Lapyckyj L, Besso MJ, Llauradó M, Abascal MF, Matos ML, Lanau L, Castellví J, *et al*: E-cadherin: A determinant molecule associated with ovarian cancer progression, dissemination and aggressiveness. *PLoS One* 12: e0184439, 2017.
17. Wróblewski M, Szewczyk-Golec K, Hołyńska-Iwan I, Wróblewska J and Woźniak A: Characteristics of selected adipokines in ascites and blood of ovarian cancer patients. *Cancers (Basel)* 13: 4702, 2021.
18. Sonogo M, Poletto E, Pivetta E, Nicoloso MS, Pellicani R, Vinciguerra GLR, Citron F, Sorio R, Mongiat M and Baldassarre G: IMP-1 is overexpressed and secreted by platinum resistant epithelial ovarian cancer cells. *Cells* 9: 6, 2019.
19. Abreu M, Cabezas-Sainz P, Alonso-Alconada L, Ferreirós A, Mondelo-Macia P, Lago-Lestón RM, Abalo A, Díaz E, Palacios-Zambrano S, Rojo-Sebastian A, *et al*: Circulating tumor cells characterization revealed TIMP1 as a potential therapeutic target in ovarian cancer. *Cells* 9: 1218, 2020.
20. Vilà-González M, Kelaini S, Magee C, Caines R, Campbell D, Eleftheriadou M, Cochrane A, Drehmer D, Tsifaki M, O'Neill K, *et al*: Enhanced function of induced pluripotent stem cell-derived endothelial cells through ESM1 signaling. *Stem Cells* 37: 226-239, 2019.
21. Li YK, Zeng T, Guan Y, Liu J, Liao NC, Wang MJ, Chen KX, Luo XY, Chen CY, Quan FF, *et al*: Validation of ESM1 related to ovarian cancer and the biological function and prognostic significance. *Int J Biol Sci* 19: 258-280, 2023.
22. Ma H, Tian T and Cui Z: Targeting ovarian cancer stem cells: A new way out. *Stem Cell Res Ther* 14: 28, 2023.
23. García-Rubiño MF, Lozano-López C and Campos JM: Inhibitors of cancer stem cells. *Anticancer Agents Med Chem* 16: 12301-1239, 2016.
24. Wilczyński JR, Wilczyński M and Paradowska E: Cancer stem cells in ovarian cancer-a source of tumor success and a challenging target for novel therapies. *Int J Mol Sci* 23: 2496, 2022.
25. Keyvani V, Farshchian M, Esmaeili SA, Yari H, Moghbeli M, Nezhad SK and Abbaszadegan MR: Ovarian cancer stem cells and targeted therapy. *J Ovarian Res* 12: 120, 2019.
26. Muñoz-Galván S and Carnero A: Targeting cancer stem cells to overcome therapy resistance in ovarian cancer. *Cells* 9: 1402, 2020.
27. Markowska A, Sajdak S, Huczyński A, Rehlis S and Markowska J: Ovarian cancer stem cells: A target for oncological therapy. *Adv Clin Exp Med* 27: 1017-1020, 2018.
28. Bapat SA, Mali AM, Koppikar CB and Kurrey NK: Stem and progenitor-like cells contribute to the aggressive behavior of human epithelial ovarian cancer. *Cancer Res* 65: 3025-3029, 2005.
29. Wen KC, Sung PL, Wu ATH, Chou PC, Lin JH, Huang CF, Yeung SJ and Lee MH: Neoadjuvant metformin added to conventional chemotherapy synergizes anti-proliferative effects in ovarian cancer. *J Ovarian Res* 13: 95, 2020.
30. Broekman KE, Hof MAJ, Touw DJ, Gietema JA, Nijman HW, Lefrandt JD, Reyners AKL and Jalving M: Phase I study of metformin in combination with carboplatin/paclitaxel chemotherapy in patients with advanced epithelial ovarian cancer. *Invest New Drugs* 38: 1454-1462, 2020.
31. Zhang R, Zhang P, Wang H, Hou D, Li W, Xiao G and Li C: Inhibitory effects of metformin at low concentration on epithelial-mesenchymal transition of CD44(+) CD117(+) ovarian cancer stem cells. *Stem Cell Res Ther* 6: 262, 2015.
32. Chung H, Kim YH, Kwon M, Shin SJ, Kwon SH, Cha SD and Cho CH: The effect of salinomycin on ovarian cancer stem-like cells. *Obstet Gynecol Sci* 59: 261-268, 2016.
33. Lee HG, Shin SJ, Chung HW, Kwon SH, Cha SD, Lee JE and Cho CH: Salinomycin reduces stemness and induces apoptosis on human ovarian cancer stem cell. *J Gynecol Oncol* 28: e14, 2017.
34. Zou M, Yin X, Zhou X, Niu X, Wang Y and Su M: Salinomycin-Loaded high-density lipoprotein exerts promising anti-ovarian cancer effects by inhibiting epithelial-mesenchymal transition. *Int J Nanomedicine* 17: 4059-4071, 2022.
35. Roy M, Connor J, Al-Niaimi A, Rose SL and Mahajan A: Aldehyde dehydrogenase 1A1 (ALDH1A1) expression by immunohistochemistry is associated with chemo-refractoriness in patients with high-grade ovarian serous carcinoma. *Hum Pathol* 73: 1-6, 2018.
36. Januchowski R, Wojtowicz K, Sterzyńska K, Sosifka P, Andrzejewska M, Zawierucha P, Nowicki M and Zabel M: Inhibition of ALDH1A1 activity decreases expression of drug transporters and reduces chemotherapy resistance in ovarian cancer cell lines. *Int J Biochem Cell Biol* 78: 248-259, 2016.
37. Kim D, Choi BH, Ryoo IG and Kwak MK: High NRF2 level mediates cancer stem cell-like properties of aldehyde dehydrogenase (ALDH)-high ovarian cancer cells: Inhibitory role of all-trans retinoic acid in ALDH/NRF2 signaling. *Cell Death Dis* 9: 896, 2018.
38. Kaipio K, Chen P, Roering P, Huhtinen K, Mikkonen P, Östling P, Lehtinen L, Mansuri N, Korpela T, Potdar S, *et al*: ALDH1A1-related stemness in high-grade serous ovarian cancer is a negative prognostic indicator but potentially targetable by EGFR/mTOR-PI3K/aurora kinase inhibitors. *J Pathol* 250: 159-169, 2020.
39. Kenda Suster N, Virant-Klun I, Frkovic Grazio S and Smrkolj S: The significance of the pluripotency and cancer stem cell-related marker NANOG in diagnosis and treatment of ovarian carcinoma. *Eur J Gynaecol Oncol* 37: 604-612, 2016.
40. Mahalaxmi J, Devi SM, Kaavya J, Arul N, Balachandrar V and Santhya KS: New insight into NANOG: A novel therapeutic target for ovarian cancer (OC). *Eur J Pharmacol* 852: 51-57, 2019.
41. Vasefifar P, Motafakkerzad R, Maleki LA, Najafi S, Ghobaninezhad F, Najafzadeh B, Alemohammad H, Amini M, Baghbazadeh A and Baradaran B: Nanog, as a key cancer stem cell marker in tumor progression. *Gene* 827: 146448, 2022.
42. Alemohammad H, Asadzadeh Z, Motafakker Azad R, Hemmat N, Najafzadeh B, Vasefifar P, Najafi S and Baradaran B: Signaling pathways and microRNAs, the orchestrators of NANOG activity during cancer induction. *Life Sci* 260: 118337, 2020.
43. Xu CX, Xu M, Tran L, Yang H, Permeth-Wey J, Kruk PA, Wenham RM, Nicosia SV, Lancaster JM, Sellers TA and Cheng JQ: MicroRNA MiR-214 regulates ovarian cancer cell stemness by targeting p53/Nanog. *J Biol Chem* 291: 22851, 2016.
44. Wu D, Wang J, Cai Y, Ren M, Zhang Y, Shi F, Zhao F, He X, Pan M, Yan C and Dou J: Effect of targeted ovarian cancer immunotherapy using ovarian cancer stem cell vaccine. *J Ovarian Res* 8: 68, 2015.
45. Wu D, Yu X, Wang J, Hui X, Zhang Y, Cai Y, Ren M, Guo M, Zhao F and Dou J: Ovarian cancer stem cells with high ROR1 expression serve as a new prophylactic vaccine for ovarian cancer. *J Immunol Res* 2019: 9394615, 2019.
46. Hu H, Zhao F, Wu D, Zhang Y, Bao X, Shi F, Cai Y and Dou J: Eliciting effective tumor immunity against ovarian cancer by cancer stem cell vaccination. *Biomed Pharmacother* 161: 114547, 2023.
47. Mao Y, Chen Y and Zhang Z: Molecular function of Krüppel-like factor 7 in biology. *Acta Biochim Biophys Sin (Shanghai)* 55: 713-725, 2023.
48. De Donato M, Babini G, Mozzetti S, Buttarelli M, Ciucci A, Arduini G, De Rosa MC, Scambia G and Gallo D: KLF7: A new candidate biomarker and therapeutic target for high-grade serous ovarian cancer. *J Exp Clin Cancer Res* 39: 265, 2020.

