

# Molecular Cloning of *GA 2-Oxidase3* from Spinach and Its Ectopic Expression in *Nicotiana sylvestris*<sup>1</sup>

Dong Ju Lee and Jan A.D. Zeevaart\*

Department of Energy Plant Research Laboratory (D.J.L., J.A.D.Z.) and Department of Plant Biology (J.A.D.Z.), Michigan State University, East Lansing, Michigan 48824-1312

Previous work has shown that 13-hydroxylated gibberellins (GAs) are predominant in the long-day (LD) plant spinach (*Spinacia oleracea*; GA<sub>53</sub>, GA<sub>44</sub>, GA<sub>19</sub>, GA<sub>20</sub>, GA<sub>1</sub>, GA<sub>8</sub>, and GA<sub>29</sub>). Also present in spinach are 2 $\beta$ -hydroxylated C<sub>20</sub>-GAs: GA<sub>97</sub>, GA<sub>98</sub>, GA<sub>99</sub>, and GA<sub>110</sub>. Levels of the most abundant GA, GA<sub>97</sub>, decreased when plants were transferred from short photoperiods (SD) to LD. When [<sup>14</sup>C]GA<sub>53</sub> was fed to spinach plants, more GA<sub>53</sub> was converted to GA<sub>97</sub> in SD than in LD, and more radioactive GA<sub>20</sub> was formed in LD than in SD. *SoGA2ox3*, encoding a GA 2-oxidase, was isolated from spinach. The recombinant protein converted only two C<sub>20</sub>-GA precursors, GA<sub>12</sub> and GA<sub>53</sub>, to their respective products, GA<sub>110</sub> and GA<sub>97</sub>. *GA2ox3* competes with *GA2ox1* for their common substrate, GA<sub>53</sub>. In SD, deactivation to GA<sub>97</sub> prevails, whereas in LD conversion to GA<sub>20</sub> is favored. Transcript levels of *SoGA2ox3* were higher in shoot tips than in blades, petioles, and young leaves. Ectopic expression of *SoGA2ox3* in the long-day plant *Nicotiana sylvestris* showed a range of dwarf phenotypes, such as reduced germination, short hypocotyl and stem, dark-green leaves, and late flowering, but normal flowers and seed production. The levels of GA<sub>53</sub> and GA<sub>1</sub> were 3- to 5-fold lower in transgenic plants than in wild type, whereas the levels of GA<sub>97</sub> and GA<sub>110</sub> increased 3- to 6-fold in transgenic plants. It is concluded that genetic manipulation of plant stature by increasing deactivation of precursors of active GA is more advantageous than increased deactivation of bioactive GA<sub>1</sub> itself.

Long-day (LD) rosette plants, such as spinach (*Spinacia oleracea*), grow vegetatively and do not produce a stem when grown under short photoperiods (SD). Upon transfer to LD, stems elongate and flowering is initiated. In spinach, LD-induced stem elongation is dependent on GA-regulated processes. Although 136 different GAs (<http://www.plant-hormones.info/gibberellins.htm>) have been identified from natural sources, most of them are precursors or deactivated catabolites of a few biologically active GAs (Hedden and Phillips, 2000; Yamaguchi and Kamiya, 2000; Olszewski et al., 2002). The major endogenous GAs of spinach belong to the early-13-hydroxylation pathway (GA<sub>53</sub>, GA<sub>44</sub>, GA<sub>19</sub>, GA<sub>20</sub>, GA<sub>1</sub>, GA<sub>8</sub>, and GA<sub>29</sub>; Fig. 1A; Talon et al., 1991). Of these GAs, only GA<sub>1</sub> is active per se (Zeevaart et al., 1993). A small gene family of 2-oxidases catalyzes deactivation of GA<sub>1</sub> by 2 $\beta$ -hydroxylation to GA<sub>8</sub>. Genes encoding 2-oxidases have been isolated from several species: *Phaseolus coccineus* and *Arabidopsis* (*Arabidopsis thaliana*; Thomas et al., 1999), pea (*Pisum sativum*; Lester et al., 1999; Martin et al., 1999), rice (*Oryza sativa*; Sakamoto et al., 2001), spinach (Lee and Zeevaart, 2002), and poplar (*Populus* spp.; Busov et al., 2003). These GA

2-oxidases introduce a 2 $\beta$ -hydroxyl group to the bioactive GAs GA<sub>1</sub> and GA<sub>4</sub> and their respective precursors, GA<sub>9</sub> and GA<sub>20</sub>. The two recombinant GA 2-oxidases (*SoGA2ox1* and *SoGA2ox2*) isolated from spinach 2 $\beta$ -hydroxylated the C<sub>19</sub>-GAs GA<sub>1</sub> and GA<sub>20</sub> to GA<sub>8</sub> and GA<sub>29</sub>, respectively. In addition, *SoGA2ox1* also converted the C<sub>20</sub>-GA GA<sub>53</sub> to GA<sub>97</sub> (Lee and Zeevaart, 2002). In *Arabidopsis*, two GA 2-oxidases that 2 $\beta$ -hydroxylated GA<sub>12</sub> and GA<sub>53</sub> were identified by activation tagging. Increased expression of these genes resulted in decreased levels of active GAs and corresponding dwarf phenotypes in *Arabidopsis* and tobacco (*Nicotiana tabacum*; Schomburg et al., 2003).

In spinach, GA<sub>53</sub> is at a branch point and can be converted to bioactive GA<sub>1</sub> via the early-13-hydroxylation pathway, or it can be 2 $\beta$ -hydroxylated to GA<sub>97</sub> (Fig. 1A). In addition to GA<sub>97</sub>, several other 2 $\beta$ -hydroxylated GAs are found in spinach: GA<sub>98</sub> (2 $\beta$ -hydroxy-GA<sub>44</sub>), GA<sub>99</sub> (2 $\beta$ -hydroxy-GA<sub>19</sub>), and GA<sub>110</sub> (2 $\beta$ -hydroxy-GA<sub>12</sub>; Mander et al., 1996; Owen et al., 1998). Gilmour et al. (1986) found that cell-free extracts from spinach leaves are capable of converting [<sup>14</sup>C]GA<sub>12</sub> to [<sup>14</sup>C]2 $\beta$ -hydroxy-GA<sub>12</sub>. These results led us to postulate that an early-2 $\beta$ -hydroxylation pathway operates in spinach and that a 2-oxidase in spinach specifically 2 $\beta$ -hydroxylates C<sub>20</sub>-GAs (Fig. 1A).

Here, we report the molecular cloning and characterization of *SoGA2ox3* from spinach that 2 $\beta$ -hydroxylates two C<sub>20</sub>-GAs, GA<sub>12</sub> and GA<sub>53</sub>. Ectopic expression of *SoGA2ox3* in *Nicotiana sylvestris* produced a high level of GA<sub>97</sub> with a concomitant decrease in the levels of active GA<sub>1</sub>. Transgenic plants overexpressing this 2-oxidase exhibited GA-deficient

<sup>1</sup> This work was supported by the U.S. Department of Agriculture (grant no. 2002-35304-1269) and by the U.S. Department of Energy (grant no. DE-FG02-91ER20021).

\* Corresponding author; e-mail zeevaart@msu.edu; fax 517-353-9168.

Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.104.056499](http://www.plantphysiol.org/cgi/doi/10.1104/pp.104.056499).



this GA 2-oxidase and its control at the molecular-genetic level.

### Cloning of *SoGA2ox3* cDNA

First-strand cDNA was synthesized with random primers and poly(A<sup>+</sup>) RNA isolated from spinach grown under SD conditions as a template. All combinations of degenerate primers were used in the PCR reactions with first-strand cDNA. The combination of degenerate primers JZ598 and JZ600 (Table I) yielded a 611-bp fragment of a putative GA 2-oxidase (data not shown). The predicted partial amino acid sequence of this reverse transcription (RT)-PCR product shares more identity with the amino acid sequences of AtGA2ox7 and AtGA2ox8 (Schomburg et al., 2003) than with those of the *SoGA2ox1*, *SoGA2ox2*, *AtGA2ox1*, *AtGA2ox2*, and *AtGA2ox3* proteins (Thomas et al., 1999; Lee and Zeevaart, 2002). 5'- and 3'-RACE were performed with sequence-specific primers (JZ616, JZ617, JZ618, and JZ627).

The coding region of *SoGA2ox3* was obtained with primers JZ651 and JZ656 (Table I). The full-length cDNA clone of *SoGA2ox3* (1,242 bp, GenBank accession no. AY935713) has an open reading frame of 1,128 bp, encoding a putative protein of 375 amino acids with a 79-bp 5'-untranslated sequence and a 35-bp 3'-untranslated sequence. The predicted molecular mass of the protein is 43 kD, with a pI of 8.42. Southern-blot analysis using the *SoGA2ox3* cDNA as a probe indicated that there is at least one more related gene in the spinach genome (data not shown). Figure 2A shows an alignment of amino acid sequences of GA 2-oxidases from spinach, Arabidopsis, and rice. When compared with other 2-oxoglutarate-dependent dioxygenases, the deduced amino acid sequence of *SoGA2ox3* fits best in the family of GA 2-oxidases (Fig. 2B). The predicted amino acid sequence shares 59.3%, 42.4%, 46%, and 45.3% identity with AtGA2ox8, AtGA2ox7, OsGA2ox5, and OsGA2ox6, respectively. However, it shares only 23.1% and 22.1% identity with the *SoGA2ox1* and *SoGA2ox2* proteins (Lee and Zeevaart, 2002). The amino acid sequence of the *SoGA2ox3* protein contains gene-specific motifs (located at positions 143–149, 158–164, and 358–371)

that are different from other GA dioxygenases, and a Thr homopolymer (located at positions 49–58). Three underlined regions show unique amino acid sequences that are highly conserved in *SoGA2ox3*, *AtGA2ox7*, *AtGA2ox8*, *OsGA2ox5*, and *OsGA2ox6* but not in other GA dioxygenases, including *SoGA2ox1* and *SoGA2ox2* (Fig. 2, A and B). The amino acid sequence of *SoGA2ox3* shares several conserved regions with other GA dioxygenases, such as putative 2-oxoglutarate-binding sites (Arg-319 and Ser-321) and iron-binding sites (His-253, Asp-255, and His-309; Thomas et al., 1999). However, the amino acid sequence (L-S-W-S-E-A, positions 157–162) shares only 50% identity with the L-P-W-K-E-T sequence, which is conserved in all GA 2-oxidases and which has been proposed to be involved in binding the substrates GA<sub>12</sub> and/or GA<sub>53</sub> (Wu et al., 1996; Kang et al., 1999; Sakamoto et al., 2004). *SoGA2ox3* contains one intron of 781 bp (AY935714), which is located at the same position as the second intron of the *AtGA2ox7* and *AtGA2ox8* genes (Schomburg et al., 2003).

The amino acid sequences of all GA 2-oxidases were aligned using the ClustalW 1.8 Multiple Sequence Alignment program, and a phylogenetic tree was generated (Fig. 2B). This analysis divides the GA 2-oxidase family into three different clades. Members of classes I and II catabolize C<sub>19</sub>-GAs. However, as far as tested, members of class III can only 2β-hydroxylate C<sub>20</sub>-GAs (*SoGA2ox3*, *AtGA2ox7*, and *AtGA2ox8*). The two GA 2-oxidases of rice in class III, *OsGA2ox5* and *OsGA2ox6*, have not been characterized, but their amino acid sequences predict that they are able to 2β-hydroxylate C<sub>20</sub>-GAs.

### Heterologous Expression of *SoGA2ox3* in *Escherichia coli*



To demonstrate that the *SoGA2ox3* cDNA clone encodes a GA 2-oxidase, we heterologously expressed the coding region as a fusion protein of the glutathione S-transferase fusion vector in *E. coli* strain BL21pLysS. The size of the fusion protein is about 69 kD. Soluble protein extracts were used for assays of GA 2-oxidase activity with several radioactive GAs as substrates. The reaction products were separated by reverse-phase HPLC with an on-line radioactivity detector

**Table I.** Primers used for amplification of the GA 2-oxidase3 gene from spinach

Single-letter codes for nucleotides were assigned as follows: Y (C/T), N (A/T/G/C), R (A/G), D (A/G/T), and W (A/T).

Purpose	Primer	Primer Sequence (5' to 3')	Orientation
RT-PCR	JZ598	GARTGGGGNTTYTTCARRT	S
RT-PCR	JZ600	GCYTGRAATAWRTCNCDDATRTT	AS
3'-RACE	JZ617	GCATGATGAGCTGACCTTTATTG	S
3'-RACE	JZ618	AGATACCCACCATGCCCTAAAT	S
5'-RACE	JZ616	TTGAATGGCTCCCTAAACACTT	AS
5'-RACE	JZ627	ACCGTTGAGCATAGATTGCTGA	AS
Coding region	JZ651	TCCGGATCCAAATGGCTTCTACCAAGGTAG	S
Coding region	JZ656	CTGAGAATTCCTAGTTCGAAATGAGGAAG	AS

**Figure 2.** Comparison of the deduced amino acid sequences of SoGA2ox3 with other GA 2-oxidases. A, Sequence alignment of the amino acid sequence of SoGA2ox3 cDNA with related GA 2-oxidases. The amino acid sequences were aligned using BCM Search Launcher. The three underlined regions show the unique amino acid sequences that are highly conserved in the SoGA2ox3, AtGA2ox7, AtGA2ox8, OsGA2ox5, and OsGA2ox6 proteins. Asterisks indicate putative amino acid residues that are supposed to bind iron. The symbol ‡ indicates the putative 2-oxoglutarate-binding sites. B, Phylogenetic tree based on the comparison of GA 2-oxidase genes. The amino acid sequences form three classes (I, II, and III). This unrooted phylogenetic tree was created using ClustalW (<http://www.ebi.ac.uk/clustalw/>). GenBank accession numbers of proteins are (in parentheses): AtGA2ox1 (CAB41007), AtGA2ox2 (CAB41008), AtGA2ox3 (CAB41009), AtGA2ox4 (AAG51528), AtGA2ox6 (AAG00891), AtGA2ox7 (AAG50945), AtGA2ox8 (CAB79120), LsGA2ox1 (BAB12442), OsGA2ox1 (BAB40934), OsGA2ox2 (BAC16751), OsGA2ox3 (BAC16752), OsGA2ox4 (AAU03107), OsGA2ox5 (BAC10398), OsGA2ox6 (CAE03751), PcGA2ox1 (CAB41036), PsGA2ox1 (AAF08609), PsGA2ox2 (AAD45424), SoGA2ox1 (AAN87571), SoGA2ox2 (AAN87572), and SoGA2ox3 (AAX14674). The scale value of 0.1 indicates 0.1 amino acid substitutions per site.



(Lee and Zeevaart, 2002). Retention times (Rts) of the products were compared with those of standard <sup>14</sup>C-labeled GAs for tentative identification. The recombinant SoGA2ox3 protein converted the radioactive C<sub>20</sub>-GAs GA<sub>12</sub> (Rt = 37.6 min) and GA<sub>53</sub> (Rt =

31.8 min) to products with Rts of 28.8 and 14.9 min, corresponding to those of GA<sub>110</sub> and GA<sub>97</sub>, respectively. In some experiments, in addition to GA<sub>97</sub>, a minor unidentified product of GA<sub>53</sub> was found at Rt = 19.6 min. We used 17,17-<sup>2</sup>H<sub>2</sub>GAs as substrates

**Table II.** Identification of products formed after incubation of recombinant GA 2-oxidase (SoGA2ox3) from spinach with GA<sub>12</sub> or GA<sub>53</sub>

Substrate	Product	Mass Spectra of Products <sup>a</sup>	
		<i>m/z</i> (% relative abundance)	
17,17-[ <sup>2</sup> H <sub>2</sub> ]GA <sub>12</sub>	17,17-[ <sup>2</sup> H <sub>2</sub> ]GA <sub>110</sub>	M <sup>+</sup> 450 (5), 435 (6), 418 (25), 390 (44), 375 (4), 360 (2), 328 (6), 318 (7), 300 (100), 285 (66), 274 (40), 260 (26), 259 (22), 241 (98), 225 (24), 201 (17), 197 (10), 145 (34)	
17,17-[ <sup>2</sup> H <sub>2</sub> ]GA <sub>53</sub>	17,17-[ <sup>2</sup> H <sub>2</sub> ]GA <sub>97</sub>	M <sup>+</sup> 538 (35), 523 (9), 506 (6), 479 (9), 448 (3), 389 (10), 373 (5), 329 (11), 299 (2), 239 (44), 210 (65), 209 (100), 195 (11), 179 (16), 147 (3), 119 (14)	

<sup>a</sup>As the methyl ester trimethylsilyl ethers.

to identify the products of GA<sub>12</sub> and GA<sub>53</sub> catabolism by gas chromatography-mass spectrometry (GC-MS; Table II). Recombinant SoGA2ox3 did not convert [<sup>14</sup>C]GA<sub>44</sub> (closed lactone ring) and [<sup>14</sup>C]GA<sub>19</sub> to 2β-hydroxy-[<sup>14</sup>C]GA<sub>44</sub> (=GA<sub>98</sub>) and 2β-hydroxy-[<sup>14</sup>C]GA<sub>19</sub> (=GA<sub>99</sub>), respectively. Also, SoGA2ox3 was not able to convert C<sub>19</sub>-GAs, such as GA<sub>20</sub> and GA<sub>1</sub>. Thus, SoGA2ox3 belongs to the class of 2-oxidases that specifically deactivates the C<sub>20</sub>-GA precursors GA<sub>12</sub> and GA<sub>53</sub> (Schomburg et al., 2003), which are also the substrates for GA 20-oxidases.

#### Effects of Daylength on Gibberellin Levels in Spinach

The GA contents of spinach plants growing in SD or LD were determined by GC-MS-selected ion monitoring (Table III). As reported previously (Wu et al., 1996), transfer of spinach from SD to LD caused an increase in all GAs of the early-13-hydroxylation pathway, except for GA<sub>53</sub> in mature leaves and petioles. By contrast, the levels of the 2β-hydroxylated C<sub>20</sub>-GAs GA<sub>97</sub> and GA<sub>110</sub> showed a decrease after transfer from SD to LD, which corroborates the results obtained with feeding [<sup>14</sup>C]GA<sub>53</sub> (Fig. 1B). GA<sub>97</sub> is by far the most abundant GA in spinach, whereas GA<sub>110</sub> is present in levels comparable to those of the other GAs. This striking difference in contents of GA<sub>97</sub> and GA<sub>110</sub> raises the question whether SoGA2ox3 has a higher affinity for GA<sub>53</sub> than for GA<sub>12</sub>. To answer this question, we

incubated mixtures of the two substrates at equimolar concentrations with recombinant SoGA2ox3 for 1 h. GA<sub>12</sub> and GA<sub>53</sub> were equally well catabolized over a range of substrates from 30 to 300 pmol (data not shown), indicating that the much higher content of GA<sub>97</sub> than of GA<sub>110</sub> is probably not due to different affinities of the enzyme for the two substrates.

#### Expression of SoGA2ox3 in Spinach in SD and LD

Two sizes of transcripts of *SoGA2ox3* were observed around 1.3 kb and 0.7 kb in northern-blot analysis (data not shown). The larger size was assumed to be the mature message because it is the same size as the full-length cDNA (see above).

To investigate how expression of the *SoGA2ox3* gene is regulated by the photoperiod, we isolated total RNA from spinach plants grown in SD and after 8 LD. Figure 3A shows that the levels of *SoGA2ox3* transcripts were relatively high in petioles and shoot tips but low in blades and young leaves in both SD and LD. These expression patterns of *SoGA2ox3* are correlated with the levels of GA<sub>97</sub> in various organs of spinach (Fig. 3A; Table III).


To determine the time course of changes in *SoGA2ox3* expression, spinach plants were harvested at different times after transfer from SD to LD. The levels of *SoGA2ox3* transcripts were slightly increased in

**Table III.** Comparison of GA levels in various organs of spinach in SD and after 8 LD

Spinach plants were grown in SD for 6 weeks, then transferred to LD for 8 d. GA content was determined by GC-MS-selected ion monitoring, using deuterated GAs as internal standards.

Plant Part	GA <sub>53</sub>	GA <sub>19</sub>	GA <sub>20</sub>	GA <sub>1</sub>	GA <sub>29+81</sub>	GA <sub>8</sub>	GA <sub>110</sub>	GA <sub>97</sub>
Mature blades								
SD	4.6 <sup>a</sup>	31.8	2.7	1.7	13.5	1.3	8.5	191.1
8 LD	2.1	23.5	23.6	2.7	36.4	3.2	1.3	73.0
Petioles								
SD	24.6	51.3	1.7	1.6	2.7	2.9	18.9	477.5
8 LD	21.7	145.3	36.7	4.1	42.3	32.3	4.7	446.0
Young leaves								
SD	7.2	54.4	3.0	4.5	10.1	1.3	16.2	286.3
8 LD	31.1	138.4	43.4	19.4	84.7	9.9	2.4	178.7
Shoot tips								
SD	24.7	61.3	2.2	3.8	16.0	3.8	18.3	880.0
8 LD	66.6	173.5	31.4	16.4	66.7	37.1	13.4	696.2

<sup>a</sup>Values are in ng g<sup>-1</sup> dry weight.



**Figure 3.** Northern analysis of *SoGA2ox3* in spinach. **A**, Levels of *SoGA2ox3* transcripts in spinach grown in SD and after 8 LD. **B**, Levels of *SoGA2ox3* transcripts in petioles and tips with increasing numbers of LD. **C**, Levels of *SoGA2ox3* transcripts in male (♂) and female (♀) flowers, stems (St), and shoot tips (ShT). Spinach plants were harvested 4 weeks after transfer to LD. The shoot tips included the upper 1 cm of shoots; stems were the next 1 cm of shoots. Total RNA (30  $\mu$ g) was loaded for each sample. The blots were probed with *SoGA2ox3* or *SoActin*.

blades and young leaves (data not shown) and petioles in a time-dependent manner, but the levels of *SoGA2ox3* transcripts decreased slightly in shoot tips with increasing duration of LD (Fig. 3B).

Expression patterns of *SoGA2ox3* were also investigated in inflorescences. Male and female flowers, stems, and shoot tips were harvested from spinach plants that had been exposed to LD for 4 weeks (Fig. 3C). The *SoGA2ox3* transcripts were more abundant in the upper 2 cm of stems than in shoot tips, and expression was relatively low in male and female flowers.


#### Ectopic Expression of the *SoGA2ox3* Gene in *N. sylvestris*

To determine whether ectopic expression of the *SoGA2ox3* gene would produce  $GA_{97}$  and cause a GA-deficient phenotype in another species, we introduced the 35S::*SoGA2ox3* cDNA construct into *N. sylvestris* by *Agrobacterium tumefaciens*-mediated transformation. Ten homozygous lines expressing *SoGA2ox3* were selected (Fig. 4), which exhibited a profound range of dwarfed phenotypes (data not shown). To ascertain the presence of the *SoGA2ox3* gene in the transgenic plants, we performed genomic PCR amplification with a pair of primers, JZ651 and JZ656 (Table I). The PCR products (1.1 kb) were detected without any other PCR products in all transgenic lines (data not shown). Phenotypes of transgenic lines 6 to 10 were very mild, but transgenic lines 1 to 5 showed severe dwarfed phenotypes (data not shown). The severity of the phenotype was correlated with the accumulation of *SoGA2ox3* transcripts (Fig. 4). The expression patterns of *NsGA20ox1* of *N. sylvestris* were similar to those of *SoGA2ox3* transcripts (Fig. 4), indicating that feedback regulation to maintain GA homeostasis operated in the transgenic plants.


Transgenic line 2 was chosen for further characterization (Fig. 5). In SD, the leaves were slightly shorter and darker green than those of the wild type. The leaves of wild-type plants were smooth, whereas those of transgenic plants were somewhat wrinkled (Fig. 5A). The inset in Figure 5B shows that *SoGA2ox3* was highly expressed in the transgenic plant. The time course of stem elongation (Fig. 5C) shows that the transgenic plants did not start to elongate until the wild-type plants had already reached their final height and were in full bloom. The final height of the transgenic plants was less than half of that of the wild-type plants, and the number of leaves was almost doubled (Table IV). Flower buds in the transgenic plants appeared 30 d later than those of the wild type (Fig. 5C; Table IV). However, the interval between appearance of flower buds and anthesis was the same in wild-type and transgenic plants, indicating that the rate of flower bud development was the same in both types of plants. Fruit and seed production were normal in the transgenic plants. This may indicate that these aspects of plant development require a lower  $GA_1$  level than stem growth. As in leaves (Fig. 4), *SoGA2ox3* was highly expressed in flower buds and mature flowers (Fig. 6). Expression of native *NsGA20ox1* and *NsGA20ox2* in young flower buds was up-regulated as compared to their expressions in the wild type. However, expression of native 20-oxidases in mature flowers was below the limit of detection in northern blots (Fig. 6). Thus, at an early stage, GA 20-oxidase could channel sufficient  $GA_{53}$  through the early-13-hydroxylation pathway for normal flower development. It should be kept in mind, however, that flowers are heterotrophic, so that GAs, instead of being synthesized in situ, may be supplied by green tissue.

#### Effect of Ectopic Expression of *SoGA2ox3* on Gibberellin Content

As expected, GA levels were affected by ectopic expression of *SoGA2ox3* in transgenic plants (Table V). The level of  $GA_{53}$ , which is the substrate for both GA 20-oxidase and GA 2-oxidase3, showed a marked decrease in transgenic plants in both SD and LD. The levels of  $GA_{19}$  and  $GA_{20}$ , the products of GA 20-oxidase, also decreased in transgenic plants in LD with a concomitant 5-fold decrease in bioactive  $GA_1$ . Overexpression of *SoGA2ox3* did not affect the  $GA_{97}$  content



**Figure 4.** Northern analysis of *SoGA2ox3* and *NsGA20ox1* expression in transgenic *N. sylvestris*. Expression levels of the *SoGA2ox3* transgene parallel those of the *NsGA20ox1* gene. For each sample, 30  $\mu$ g of total RNA was loaded. The blots were probed with *SoGA2ox3*, *NsGA20ox1*, or *NsActin*.



**Figure 5.** Ectopic expression of *SoGA2ox3* suppresses stem growth in *N. sylvestris*. **A**, Transgenic plant and wild-type plant (WT) show a similar phenotype in SD. **B**, WT and transgenic plants grown in LD following germination. Inset in **B** shows expression of *SoGA2ox3* in the transgenic plant. **C**, Stem growth of WT and transgenic plants grown in LD following germination. Black arrows indicate the average number of days until appearance of flower buds. White arrows indicate the average number of days until anthesis.  $n = 12$ .

in SD, but it caused a 3- to 6-fold increase in transgenic plants in LD. As in spinach,  $GA_{110}$  was present in much lower amounts than  $GA_{97}$ ; its levels also increased in LD. These data indicate that ectopic expression of *SoGA2ox3* increased the levels of both  $GA_{97}$  and  $GA_{110}$  in transgenic plants and kept the GAs of the early-13-hydroxylation pathway at a low level. Thus, GA 2-oxidase3 in transgenic plants competes effectively with GA 20-oxidase for the common substrates

$GA_{12}$  and  $GA_{53}$ , resulting in a reduced  $GA_1$  content and a dwarfed phenotype.

#### Applied Gibberellin Restores a Normal Phenotype in 35S::SoGA2ox3 Plants

Treatment with  $GA_1$  reversed the phenotype of dwarf transgenic plants to that of wild type. In SD, leaves were elongated and the plants had started to bolt after 25 d. Although grown in SD, the  $GA_1$ -treated plants had the appearance of wild-type plants exposed to LD (Fig. 7A). In LD, the transgenic plants treated with  $GA_1$  showed essentially the same phenotype as wild-type plants (Fig. 7B). These results show that  $GA_1$  treatment effectively increased leaf and stem growth of transgenic plants both in SD and LD, indicating that application of  $GA_1$  could fully compensate for the decreased endogenous  $GA_1$  content of transgenic plants.

#### Effect of Ectopic Expression of *SoGA2ox3* on Germination

Because severely GA-deficient mutants do not germinate in the absence of exogenous GAs (Koornneef et al., 1983), we tested whether overexpression of *SoGA2ox3* affects germination. As shown in Figure 8, a high percentage of wild-type seeds had germinated after 4 d, whereas germination of transgenic seeds did not begin until day 5. Treatment with  $GA_3$  accelerated germination and also caused a higher percentage of seeds to germinate than was observed for seeds without  $GA_3$  treatment (Fig. 8). Thus, although overexpression of *SoGA2ox3* decreases the content of bioactive GA, transgenic seeds are still able to germinate, albeit with some delay.


#### Effect of Ectopic Expression of *SoGA2ox3* on Hypocotyl and Root Growth

A role of GA in the regulation of root growth has been established by the use of GA-deficient mutants (Yaxley et al., 2001; Fu and Harberd, 2003). It was of interest, therefore, to investigate whether root growth is affected in seedlings with a low  $GA_1$  content due to overexpression of *SoGA2ox3*. As shown in Figure 9, ectopic expression of the *SoGA2ox3* gene markedly affected both hypocotyl and root growth. The hypocotyls of wild-type seedlings were elongated in darkness, but those of transgenic seedlings remained very

**Table IV.** Characterization of transgenic plants overexpressing *SoGA2ox3* in *N. sylvestris* (Ns)

Observed Trait	Wild-Type Ns	35S::SoGA2ox3
Days to flower buds	67.3 ± 0.92 <sup>a</sup>	97.8 ± 0.86
Days to anthesis	83.5 ± 0.97	114.5 ± 0.96
Number of leaves to inflorescence	19.0 ± 0.35	35.8 ± 0.35
Final plant height	122.6 ± 2.8	56.1 ± 1.01

<sup>a</sup>SE;  $n = 12$ .



**Figure 6.** Northern analysis of *SoGA2ox3*, *NsGA20ox1* (GenBank accession no. AF494087), and *NsGA20ox2* (accession no. AF494088) in young flower buds and mature flowers of transgenic *N. sylvestris*. For each sample, 30  $\mu\text{g}$  of total RNA was loaded. The blots were probed with *SoGA2ox3*, *NsGA20ox1*, *NsGA20ox2*, or *NsActin*.

short (Fig. 9A). Treatment with 1 mg L<sup>-1</sup> GA<sub>3</sub> completely restored hypocotyl length of transgenic seedlings to that of wild type (Fig. 9B). Differences in root growth between wild-type and transgenic seedlings were most pronounced under continual light (Fig. 9C). Treatment with GA<sub>3</sub> could only partially restore root growth in transgenic seedlings, but hypocotyl growth was promoted to the same extent in both types of seedlings (Fig. 9D).

## DISCUSSION

In this study, we isolated a full-length cDNA clone encoding a GA 2-oxidase from spinach. Heterologous expression in *E. coli* showed that the product of *SoGA2ox3* catalyzes 2 $\beta$ -hydroxylation of the C<sub>20</sub>-GAs GA<sub>12</sub> and GA<sub>53</sub> to GA<sub>110</sub> and GA<sub>97</sub>, respectively. The same activities have previously been found for AtGA2ox7 and AtGA2ox8 (Schomburg et al., 2003). The recombinant proteins from other GA 2-oxidases characterized so far 2 $\beta$ -hydroxylate C<sub>19</sub>-GAs. The only exceptions reported are for PcGA2ox1 and AtGA2ox2, which 2 $\beta$ -hydroxylated GA<sub>15</sub>, a C<sub>20</sub>-GA (Thomas et al., 1999); and OsGA2ox1, which converted GA<sub>44</sub> to GA<sub>98</sub> (Sakamoto et al., 2001). Spinach produces GA<sub>98</sub> and GA<sub>99</sub>, but recombinant *SoGA2ox3* did not metabolize their possible precursors, GA<sub>44</sub> and GA<sub>19</sub>. In transgenic *N. sylvestris* overexpressing *SoGA2ox3*, there was an

increase in GA<sub>97</sub> (Table V), but no GA<sub>98</sub> and GA<sub>99</sub> could be detected in these plants (J.A.D. Zeevaart, unpublished data). Considering these combined results, it is unlikely that *SoGA2ox3* is involved in the synthesis of GA<sub>98</sub> and GA<sub>99</sub>. Another possibility is that GA 20-oxidase converts GA<sub>97</sub> to GA<sub>98</sub> and GA<sub>99</sub>, as indicated in the hypothetical pathway in Figure 1A. But GA<sub>97</sub> was not converted by recombinant *SoGA20ox* (J.A.D. Zeevaart, unpublished data). Perhaps recombinant *SoGA2ox1* can 2 $\beta$ -hydroxylate GA<sub>44</sub> and GA<sub>19</sub>, as it was able to convert GA<sub>53</sub> to GA<sub>97</sub> (Lee and Zeevaart, 2002). Finally, the possibility remains that there are additional GA 2-oxidases that catalyze the formation of GA<sub>98</sub> and GA<sub>99</sub>.

GA<sub>97</sub> is much more abundant than GA<sub>110</sub> in spinach (Table III), and the same is true for *N. sylvestris* in both wild-type and transgenic plants (Table V). Given the results of the experiment in which the two precursors GA<sub>12</sub> and GA<sub>53</sub> were mixed, it is unlikely that GA 2-oxidase3 has a higher affinity for GA<sub>53</sub> than for GA<sub>12</sub>. The cellular location of the GA 13-oxidase that converts GA<sub>12</sub> to GA<sub>53</sub> is not known, but there may be compartmentation of the substrates in such a manner that GA<sub>53</sub> is more readily available to GA 2-oxidase3 than is GA<sub>12</sub>. Further metabolism of GA<sub>97</sub> is not known, so that its very high level may not reflect a high rate of synthesis but rather accumulation of a stable inactive end product. A relatively low rate of synthesis of GA<sub>97</sub> is also indicated by the low levels of *SoGA2ox3* transcripts in spinach (Fig. 3). In addition, antiserum raised against recombinant *SoGA2ox3* protein did not detect any *SoGA2ox3* protein from spinach or transgenic *N. sylvestris* in western blots, indicating that the level of *SoGA2ox3* protein in vivo is very low (D.J. Lee, unpublished data).


GA<sub>53</sub> is at a branch point in GA metabolism in spinach (Fig. 1A). It can be converted to bioactive GA<sub>1</sub> via the early-13-hydroxylation pathway, or it can be deactivated by 2 $\beta$ -hydroxylation to GA<sub>97</sub>. The latter step appears to prevail in SD, whereas in LD most GA<sub>53</sub> will be channeled through the early-13-hydroxylation pathway. As shown in Figure 3, the levels of *SoGA2ox3* transcripts undergo little change when spinach plants are shifted from SD to LD. Neverthe-

**Table V.** GA content of *N. sylvestris* (Ns) and transgenic plants overexpressing *SoGA2ox3* grown under various photoperiodic conditions

Plants were grown in SD for 80 d and then transferred to LD for the number of days indicated prior to harvest. GA content was determined by GC-MS-selected ion monitoring, using deuterated GAs as internal standards. nd, Not detectable.


Sample	Photoperiodic Treatment	GA <sub>53</sub>	GA <sub>19</sub>	GA <sub>20</sub>	GA <sub>1</sub>	GA <sub>97</sub>	GA <sub>110</sub>
Ns	SD	0.7 <sup>a</sup>	3.8	1.2	0.9	11.7	nd
<i>Ns/35S::SoGA2ox3</i>	SD	0.1	2.1	0.3	0.8	11.6	0.5
Ns	14 LD	13.2	32.4	16.7	4.9	16.9	nd
<i>Ns/35S::SoGA2ox3</i>	14 LD	4.4	4.7	0.8	0.9	40.3	7.2
Ns	20 LD	32.3	54.3	27.0	5.2	14.1	1.5
<i>Ns/35S::SoGA2ox3</i>	20 LD	6.3	7.5	1.8	1.0	85.3	9.1
<i>Ns/35S::SoGA2ox3</i>	30 LD	9.7	9.4	2.3	1.1	94.9	10.2

<sup>a</sup>Values are in ng g<sup>-1</sup> dry weight.




**Figure 7.** Reversal of dwarf phenotype of transgenic plants by GA treatment. Plants were sprayed twice weekly with a solution of  $10 \text{ mg L}^{-1}$   $\text{GA}_1$ . GA treatment of transgenic plants in SD (A) and in LD (B). Photographs were taken 25 d after first GA application.

less, the levels of  $\text{GA}_{97}$  decreased, presumably because expression of GA 20-oxidase was strongly up-regulated in LD (Lee and Zeevaart, 2002). Thus, the fate of  $\text{GA}_{53}$  will be determined by the relative activities of the two enzymes GA 2-oxidase3 and GA 20-oxidase. In LD, 20-oxidase activity is high, so that  $\text{GA}_{53}$  is preferentially converted to  $\text{GA}_{20}$ . By contrast, in SD the activity of 20-oxidase is low, and more  $\text{GA}_{53}$  will be deactivated via  $2\beta$ -hydroxylation to  $\text{GA}_{97}$ . Whereas SoGA2ox1 and SoGA2ox2 can directly regulate the level of active  $\text{GA}_1$ , SoGA2ox3 can only indirectly regulate the level of active  $\text{GA}_1$  by reducing the amount of precursor that is channeled through the early-13-hydroxylation path-



**Figure 8.** Effect of overexpression of SoGA2ox3 on germination in *N. sylvestris*. Seeds of wild-type (WT) or transgenic plants were spread on agar plates containing half-strength Hoagland solution. Treatment with GA partially restored germination of transgenic seeds.  $n = 60$ .



**Figure 9.** Effect of overexpression of SoGA2ox3 on hypocotyl and root growth in *N. sylvestris* seedlings. A, Hypocotyl length of seedlings grown in darkness. B, Treatment with GA restored to normal the hypocotyl growth of transgenic seedlings in darkness. C, Root growth of transgenic seedlings grown under continuous light. D, Treatment with GA partially restored the root growth of transgenic seedlings under continuous light. Arrows indicate transition from root to hypocotyl. Photographs taken 15 d after sowing.  $n = 15$ .

way. Thus,  $2\beta$ -hydroxylation of GA precursors of bioactive  $\text{GA}_1$  adds an additional step at which the levels of active GAs can be regulated.

Overexpression of SoGA2ox3 in *N. sylvestris* had a profound effect on growth and development from germination to flowering (Figs. 5, 7, 8, and 9). Analysis of the endogenous GAs showed a reduced  $\text{GA}_1$  level in transgenic plants and an increase in  $\text{GA}_{97}$  and  $\text{GA}_{110}$ .

Wild-type *N. sylvestris* also produced GA<sub>97</sub>, although not at the high levels found in spinach. The reduction in active GA<sub>1</sub> resulted in reduced germination, shorter radicles and hypocotyls of seedlings, reduced stature, and later flowering. The GA-deficient phenotype of older plants could be readily overcome by applied GA<sub>1</sub> (Fig. 7).

Application of GA to the LD rosette plant *N. sylvestris* in SD induces stem elongation and, ultimately, flower formation (Lang, 1989). As shown in Table V, the level of GA<sub>1</sub> increased 5- to 6-fold when plants were transferred from SD to LD. This increase indicates that a high level of bioactive GA is a prerequisite for stem elongation and subsequent flowering. Conversely, a reduced level of GA<sub>1</sub> in transgenic plants delayed stem elongation and flower formation (Fig. 5; Table IV). This retardation in flowering is most vividly expressed by the number of leaves formed prior to the inflorescence in transgenic plants compared to wild-type plants (Table IV).

Semidwarfism is a desirable trait in agriculture (Busov et al., 2003; Hedden, 2003). Overexpression of GA 2-oxidase is an easy way to reduce GA levels in transgenic plants, but ectopic expression of *SoGA2ox1* and *SoGA2ox2* in *N. sylvestris* caused severely dwarfed plants that produced few seeds (D.J. Lee, unpublished results) because active GA was probably deactivated as soon as it was produced. Singh et al. (2002) also reported that ectopic expression of class II GA 2-oxidases (*PsGA2ox2*) caused defective pollen tube growth and seed abortion. Overexpression of *AtGA2ox1* in tobacco reduced the number of capsules 40% to 70% (Biemelt et al., 2004). In rice, ectopic expression of *OsGA2ox1* inhibited not only stem elongation but also development of reproductive organs. In a more direct approach, *OsGA2ox1* was expressed under the control of the promoter of *OsGA3ox2*, which is specifically expressed in the shoot apex. This caused a semidwarf phenotype with normal flowering and grain development (Sakamoto et al., 2003). When *SoGA2ox3* was overexpressed, GA precursors were deactivated, but GA 2-oxidase could still convert a small amount of the precursor to GA<sub>1</sub> (Table V), which could act without being quickly deactivated. This finding demonstrates that we can manipulate plant stature by overexpressing a gene that deactivates precursors of GA<sub>1</sub> rather than active GA<sub>1</sub> itself. The result is plants with reduced GA<sub>1</sub> levels, short stature, and delayed flowering but that otherwise retain desired traits, such as normal flowering and seed production.

## MATERIALS AND METHODS

### Plant Material and Growing Conditions

Spinach (*Spinacia oleracea*, Savoy Hybrid 612) plants were grown in a SD growth chamber for approximately 6 or 7 weeks as described earlier and then transferred to a LD growth chamber for experimental use (Lee and Zeevaart, 2002). *Nicotiana sylvestris* Spegaz. et Comes was grown under the same conditions. Plants used for GA extraction were grown in SD for approximately 3 months before being exposed to LD. All plants were watered with half-

strength Hoagland solution daily. SD consisted of 8 h of light from fluorescent tubes and incandescent bulbs of approximately 300  $\mu\text{mol m}^{-2} \text{s}^{-1}$  at 23°C, followed by 16 h darkness at 20°C. In the case of LD, the 8-h main light period was followed by 16 h of weak light from incandescent bulbs of 10  $\mu\text{mol m}^{-2} \text{s}^{-1}$ .

### Feeding of [<sup>14</sup>C]GA<sub>53</sub> to Spinach in SD and LD

[<sup>14</sup>C]GA<sub>53</sub> (specific activity 127.5 mCi/mmol) was dissolved in a small amount of acetone, to which water containing 0.05% (v/v) Tween 20 was added to give a 75% (v/v) aqueous solution. This solution was injected into petioles and midribs of spinach leaves by means of a 100- $\mu\text{L}$  Hamilton syringe. Each plant received  $0.3 \times 10^6$  dpm of [<sup>14</sup>C]GA<sub>53</sub> (=0.36  $\mu\text{g}$  GA<sub>53</sub>). Spinach plants were either kept in SD or had been exposed to 8 LD. The plants, 4 per treatment, were harvested after 2 d, frozen in liquid N<sub>2</sub>, and lyophilized. The dried material was extracted with methanol, and an acidic fraction was prepared from each sample (Zeevaart et al., 1993; Schomburg et al., 2003). The acidic fractions were analyzed by reverse-phase HPLC with on-line radioactivity detection. Radioactive fractions were collected, the derivatized materials were identified, and their specific activities determined by GC-MS (Zeevaart et al., 1993).

### RT-PCR

RT-PCR was used for the cloning of GA 2-oxidase. First-strand cDNA from 1  $\mu\text{g}$  poly(A<sup>+</sup>) RNA was synthesized by SuperScript II RNase H<sup>-</sup> reverse transcriptase (Invitrogen, Carlsbad, CA), using oligo(dT) primers at 42°C for 1 h. The cDNA product was then used in PCR reactions containing *Taq* polymerase and several pairs of degenerate primers (Table I). For GA 2-oxidase, the degenerate primers were synthesized based on conserved regions of *AtGA2ox7* and *AtGA2ox8* (Schomburg et al., 2003). These primers were used in all combinations in PCR reactions with first-strand cDNA as a template. The PCR amplifications were performed with a DNA thermal cycler (RTC-200; MJ Research, Waltham, MA) in a 50- $\mu\text{L}$  reaction mixture containing 1 $\times$  PCR buffer, 1.5 mM MgCl<sub>2</sub>, 200  $\mu\text{M}$  dNTP, 2.5 pmol of each primer, and 1 unit of *Taq* DNA polymerase (Invitrogen). The reaction mixtures were heated to 94°C for 2 min and then subjected to 30 cycles of 94°C for 1 min, 48°C for 1 min, and 72°C for 45 s. A final extension was performed at 72°C for 10 min. A 5- $\mu\text{L}$  volume of the PCR reaction products was analyzed by 1.2% agarose gel electrophoresis, purified with a Wizard PCR purification system (Promega, Madison, WI), and then cloned into the pGEM-T Easy vector (Promega).

### RACE and Cloning

5'- and 3'-RACE of spinach GA 2-oxidase cDNA were performed using the SMART RACE cDNA amplification kit from CLONTECH (Palo Alto, CA). Poly(A<sup>+</sup>) RNA was isolated from spinach using the PolyATtract mRNA Isolation System IV (Promega) according to the manufacturer's instructions. Following RT with primers supplied by CLONTECH, the first-strand cDNA was used directly in 5'- and 3'-RACE PCR reactions. Primary PCR amplification reactions were achieved using a high-fidelity enzyme (*Pfu* Turbo polymerase; Stratagene, La Jolla, CA) and gene-specific primers JZ616 and JZ627 or JZ617 and JZ618 for GA2ox3 to generate the 5'- or 3'-cDNA RACE fragments, respectively. The PCR reaction consisted of the first denaturation for 3 min at 94°C, a series of 30 cycles (1 min at 94°C, 1 min at 54°C or 55°C, 1 min at 72°C), and a final extension for 5 min at 72°C using a thermal cycler (MJ Research). A 5- $\mu\text{L}$  aliquot of the RT-PCR and RACE reaction solution was analyzed by 1.2% agarose gel electrophoresis. PCR products were purified and cloned into the pCR-Script Cam SK(+) cloning vector from Stratagene. These constructs were sequenced.

The National Center for Biotechnology Information Blast program was used to search genes homologous with the *SoGA2ox3* gene. Analysis of the DNA sequences was carried out using the DNASTAR program (DNASTAR, Madison, WI). Multiple sequence alignment of the amino acid sequences was performed using BCM Search Launcher (<http://searchlauncher.bcm.tmc.edu/multi-align/multi-align.html>) and printed with BOXSHADE 3.21 ([http://www.ch.embnet.org/software/BOX\\_form.html](http://www.ch.embnet.org/software/BOX_form.html)). The phylogenetic tree was created by means of the ClustalW program (<http://www.ebi.ac.uk/clustalw/>).

### Northern-Blot Analysis

Leaf blades, petioles, young leaves, shoot tips, and other organs were harvested, frozen immediately in liquid N<sub>2</sub>, and stored at -80°C (Lee and

Zeevaert, 2002). Northern blots were prepared by electrophoresis of 30  $\mu\text{g}$  of total RNA in the presence of formaldehyde (Sambrook et al., 1989), and RNA was transferred to nitrocellulose. Full-length cDNAs were labeled with [ $^{32}\text{P}$ ]dCTP by the Random Primers DNA labeling system (Invitrogen) and used to probe northern and Southern blots. Hybridization was carried out at 42°C using a 50% formamide system (Sambrook et al., 1989). Membranes were washed twice in  $2 \times \text{SSC}$  at room temperature for 10 min and then twice in  $0.2 \times \text{SSC}$  with 0.1% SDS at 55°C for 10 min. At low stringency, membranes were washed once in  $2 \times \text{SSC}$  at room temperature for 10 min and then in  $2 \times \text{SSC}$  at 55°C for 10 min. Each blot was first probed with *SoGA2ox3* cDNA, stripped, and reprobed with *SoActin*. Each experiment was repeated at least three times with similar results.

### Expression of Recombinant GA 2-Oxidase3 Protein

The coding region of GA 2-oxidase3 was produced by PCR with the following primer set designed from the RACE product: JZ651 and JZ656 for the mature *GA2ox3* cDNA (*Bam*HI and *Eco*RI sites inserted to facilitate cloning). The resulting PCR fragments were cloned into the pGEM-T Easy cloning vector, digested with *Bam*HI/*Eco*RI, and subcloned into the corresponding restriction sites of the pGEX-5X-2 vector (Amersham Biosciences, Piscataway, NJ). This full-length cDNA clone (pGEXGA2ox3) was transformed into *Escherichia coli* strain BL21pLysS. Fifty mL of freshly cultured cells were added to 1 L of Luria-Bertani medium with 100  $\text{mg L}^{-1}$  ampicillin and incubated at 37°C with vigorous shaking. When the optical density at 600 nm reached 0.6, isopropyl- $\beta$ -D-thiogalactopyranoside was added to give a final concentration of 3 mM, and the culture was incubated for another 2 h. The cells were harvested, suspended, and incubated in lysis buffer (100 mM Tris-HCl, pH 7.5, and 10  $\text{mg L}^{-1}$  lysozyme) at room temperature for 10 min. Crude extracts were briefly sonicated by using a Sonifier Cell Disruptor 200 (Branson Ultrasonics, Danbury, CT), carried out for 10 cycles (10  $\times$  10 s) on ice. The lysates were submerged in liquid  $\text{N}_2$  for 2 min and then thawed in an ice bath for 15 min (Johnson and Hecht, 1994). The lysates were centrifuged at 13,000 rpm for 30 min, and the supernatant was stored at  $-80^\circ\text{C}$  until needed for enzyme assays.

### Enzyme Assays and Product Identification

Enzyme assays with recombinant GA 2-oxidase3 were performed with approximately 30,000 dpm of [ $^{14}\text{C}$ ]-labeled GAs. The assays and methods for product identification have been described (Lee and Zeevaert, 2002; Schomburg et al., 2003).

### Gibberellin Extraction and Quantification

The procedures for extraction, purification, and quantification of endogenous GAs with deuterated GAs as internal standards were the same as described (Talon and Zeevaert, 1990; Zeevaert et al., 1993). All GA measurements were repeated at least twice with similar results. The four prominent ions monitored for  $\text{GA}_{97}$  were mass-to-charge ratio ( $m/z$ ) 477/479 and 536/538, and for  $\text{GA}_{110}$   $m/z$  298/300 and 388/390. Deuterated  $\text{GA}_{97}$  and  $\text{GA}_{110}$  were prepared by means of incubating 50  $\mu\text{g}$  of 17,17- $^{2}\text{H}_2$ GA $_{53}$  and 17,17- $^{2}\text{H}_2$ GA $_{12}$ , respectively, with 2.5 mL of recombinant *SoGA2ox3* protein. The GC-MS conditions were as described by Schomburg et al. (2003).

### Construction of 35S::SoGA2ox3 Plasmid and Transformation

The full-length *SoGA2ox3* cDNA was introduced into the pGEM-T Easy vector, and then reintroduced into the *Bam*HI and *Sac*I sites of the pBI121 vector after removing the *GUS* gene. This vector, pBI*SoGA2ox3*, was introduced into *Agrobacterium tumefaciens* strain LBA4404, for transformation of *N. sylvestris* according to Horsch et al. (1988). Leaf discs of *N. sylvestris* were incubated with *A. tumefaciens* for 10 min and then transferred to Murashige and Skoog medium containing 1  $\text{mg L}^{-1}$  6-benzylaminopurine and 0.1  $\text{mg L}^{-1}$  1-naphthaleneacetic acid (NAA) in darkness. After coculture for 3 d, leaf discs were transferred to selection medium (Murashige and Skoog, 1  $\text{mg L}^{-1}$  6-benzylaminopurine, 0.1  $\text{mg L}^{-1}$  NAA, 100  $\text{mg L}^{-1}$  kanamycin, and 500  $\text{mg L}^{-1}$  carbenicillin). When shoots appeared, individual shoots were cut and transferred to root inducing medium (Murashige and Skoog, 0.1  $\text{mg L}^{-1}$  NAA, 100  $\text{mg L}^{-1}$  kanamycin, and 500  $\text{mg L}^{-1}$  carbenicillin). After rooting, plantlets

( $T_0$ ) were planted in soil and grown in a greenhouse. Independent transgenic lines were selected on half-strength Hoagland medium containing 50  $\text{mg L}^{-1}$  kanamycin. The  $T_1$  generations were self-pollinated to get  $T_2$  generations. Each line showing a 3:1 segregation (resistant:sensitive) was selected to obtain homozygous  $T_3$  plants; these homozygous lines were used in further experiments.

### Analysis of Transgenic Plants by PCR

Isolation of genomic DNA from transgenic plants was performed as described (Lee and Kim, 1998). The PCR amplification reaction was for 25 cycles of 30 s at 94°C, 60 s at 50°C, and 60 s at 72°C with two primers (JZ651 and JZ656; Table I) after an initial denaturation step for 2 min at 94°C. The 5  $\mu\text{L}$  of reaction products was analyzed by 1.2% agarose gel electrophoresis.

### Germination Assays

Sixty seeds were sown in petri dishes containing half-strength Hoagland medium in the presence or absence of 1  $\text{mg L}^{-1}$  GA $_3$  and incubated in a growth room at 23°C under continual light from fluorescent lamps (50–80  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). Germination was defined as radicle emergence and was scored daily.

### Hypocotyl and Root Growth

Growth of hypocotyls and roots was determined by sowing seeds on half-strength Hoagland medium containing 0.8% phytoagar with or without 1  $\text{mg L}^{-1}$  GA $_3$ . The petri dishes were vertically oriented so that the seedlings could grow vertically along the agar surface. Wild-type and transgenic lines were sown in the same petri dish at 23°C. Seedlings were kept in darkness to determine hypocotyl length, or kept under continuous light (same source as used for germination assays) to determine root growth. Both hypocotyl length and root length were measured until 12 d after sowing.

### Distribution of Materials

Upon request, all novel materials described in this publication will be made available in a timely manner for noncommercial research purposes, subject to the requisite permission from any third-party owners of all or parts of the material. Obtaining any permissions will be the responsibility of the requestor.

Sequence data from this article have been deposited with the EMBL/GenBank data libraries under accession numbers AY935713 (*SoGA2ox3*, cDNA) and AY935714 (intron of *SoGA2ox3*).

### ACKNOWLEDGMENTS

We thank Dr. Lew Mander of the Australian National University for providing [ $^2\text{H}_2$ ]GAs, and Bev Chamberlin (Michigan State University Mass Spectrometry Facility) for assistance with GC-MS.

Received November 24, 2004; returned for revision January 11, 2005; accepted January 16, 2005.

### LITERATURE CITED

- Biemelt S, Tschiersch H, Sonnewald U (2004) Impact of altered gibberellin metabolism on biomass accumulation, lignin biosynthesis, and photosynthesis in transgenic tobacco plants. *Plant Physiol* **135**: 254–265
- Busov VB, Meilan R, Pearce DW, Ma C, Rood SB, Strauss SH (2003) Activation tagging of a dominant gibberellin catabolism gene (*GA 2-oxidase*) from poplar regulates tree stature. *Plant Physiol* **132**: 1283–1291
- Fu X, Harberd NP (2003) Auxin promotes *Arabidopsis* root growth by modulating gibberellin response. *Nature* **421**: 740–743
- Gilmour SJ, Zeevaert JAD, Schwenen L, Graebe JE (1986) Gibberellin metabolism in cell-free extracts from spinach leaves in relation to photoperiod. *Plant Physiol* **82**: 190–195
- Hedden P (2003) The genes of the green revolution. *Trends Genet* **19**: 5–9
- Hedden P, Phillips AL (2000) Gibberellin metabolism: new insights revealed by the genes. *Trends Plant Sci* **5**: 523–530

- Horsch RB, Fry J, Hoffmann N, Neidermeyer J, Rogers SG, Fraley RT (1988) Leaf disc transformation. In SB Gelvin, RA Schilperoort, eds, *Plant Molecular Biology Manual*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp A5: 1–9
- Johnson BH, Hecht MH (1994) Recombinant proteins can be isolated from *Escherichia coli* cells by repeated cycles of freezing and thawing. *Bio-technology (N Y)* **12**: 1357–1360
- Kang HG, Jun SH, Kim J, Kawaide H, Kamiya Y, An G (1999) Cloning and molecular analyses of a gibberellin 20-oxidase gene expressed specifically in developing seeds of watermelon. *Plant Physiol* **121**: 373–382
- Koorneef M, van Eden J, Hanhart CJ, de Jongh AMM (1983) Genetic fine-structure of the *GA-1* locus in the higher plant *Arabidopsis thaliana* (L.) Heynh. *Genet Res Cambridge* **41**: 57–68
- Lang A (1989) *Nicotiana*. In AH Halevy, ed, *Handbook of Flowering*, Vol VI. CRC Press, Boca Raton, FL, pp 427–483
- Lee DJ, Kim SS (1998) The regulation of 5' upstream regions of a Korean radish cationic peroxidase gene by gibberellic acid and abscisic acid. *Plant Sci* **139**: 105–115
- Lee DJ, Zeevaart JAD (2002) Differential regulation of RNA levels of gibberellin dioxygenases by photoperiod in spinach. *Plant Physiol* **130**: 2085–2094
- Lester DR, Ross JJ, Smith JJ, Elliott RC, Reid JB (1999) Gibberellin 2-oxidation and the *SLN* gene of *Pisum sativum*. *Plant J* **19**: 65–73
- Mander LN, Owen DJ, Croker SJ, Gaskin P, Hedden P, Lewis MJ, Talon M, Gage DA, Zeevaart JAD, Brenner ML, et al (1996) Identification of three C<sub>20</sub>-gibberellins: GA<sub>97</sub> (2 $\beta$ -hydroxy-GA<sub>53</sub>), GA<sub>98</sub> (2 $\beta$ -hydroxy-GA<sub>44</sub>) and GA<sub>99</sub> (2 $\beta$ -hydroxy-GA<sub>10</sub>). *Phytochemistry* **43**: 23–28
- Martin DN, Proebsting WM, Hedden P (1999) The *SLENDER* gene of pea encodes a gibberellin 20-oxidase. *Plant Physiol* **121**: 775–781
- Olszewski N, Sun T-p, Gubler F (2002) Gibberellin signaling: biosynthesis, catabolism, and response pathways. *Plant Cell (Suppl)* **14**: S61–S80
- Owen DJ, Mander LN, Storey JMD, Huntley RP, Gaskin P, Lenton JR, Gage DA, Zeevaart JAD (1998) Synthesis and confirmation of structure for a new gibberellin, 2 $\beta$ -hydroxy-GA<sub>12</sub> (GA<sub>110</sub>), from spinach and oil palm. *Phytochemistry* **47**: 331–337
- Sakamoto T, Kobayashi M, Itoh H, Tagiri A, Kayano T, Tanaka H, Iwahori S, Matsuoka M (2001) Expression of a gibberellin 2-oxidase gene around the shoot apex is related to phase transition in rice. *Plant Physiol* **125**: 1508–1516
- Sakamoto T, Morinaka Y, Ishiyama K, Kobayashi M, Itoh H, Kayano T, Iwahori S, Matsuoka M, Tanaka H (2003) Genetic manipulation of gibberellin metabolism in transgenic rice. *Nat Biotechnol* **21**: 909–913
- Sakamoto T, Miura K, Itoh H, Tatsumi T, Ueguchi-Tanaka M, Ishiyama K, Kobayashi M, Agrawal GK, Takeda S, Abe K, et al (2004) An overview of gibberellin metabolism enzyme genes and their related mutants in rice. *Plant Physiol* **134**: 1642–1653
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Plainview, NY
- Schomburg FM, Bizzell CM, Lee DJ, Zeevaart JAD, Amasino RM (2003) Overexpression of a novel class of gibberellin 2-oxidases decreases gibberellin levels and creates dwarf plants. *Plant Cell* **15**: 151–163
- Singh DP, Jermakow AM, Swain SM (2002) Gibberellins are required for seed development and pollen tube growth in *Arabidopsis*. *Plant Cell* **14**: 3133–3147
- Talon M, Zeevaart JAD (1990) Gibberellins and stem growth as related to photoperiod in *Silene armeria* L. *Plant Physiol* **92**: 1094–1100
- Talon M, Zeevaart JAD, Gage DA (1991) Identification of gibberellins in spinach and effects of light and darkness on their levels. *Plant Physiol* **97**: 1521–1526
- Thomas SG, Philips AL, Hedden P (1999) Molecular cloning and functional expression of gibberellin 2-oxidases, multifunctional enzymes involved in gibberellin deactivation. *Proc Natl Acad Sci USA* **96**: 4698–4703
- Wu K, Li L, Gage DA, Zeevaart JAD (1996) Molecular cloning and photoperiod-regulated expression of gibberellin 20-oxidase from the long-day plant spinach. *Plant Physiol* **110**: 547–554
- Yamaguchi S, Kamiya Y (2000) Gibberellin biosynthesis: its regulation by endogenous and environmental signals. *Plant Cell Physiol* **41**: 251–257
- Yaxley JR, Ross JJ, Sherriff LJ, Reid JB (2001) Gibberellin biosynthesis mutations and root development in pea. *Plant Physiol* **125**: 627–633
- Zeevaart JAD, Gage DA, Talon M (1993) Gibberellin A<sub>1</sub> is required for stem elongation in spinach. *Proc Natl Acad Sci USA* **90**: 7401–7405