



PA28 γ –20S proteasome is a proteolytic complex committed to degrade unfolded proteins

Jean-Yves Alejandro Frayssinhes¹ · Fulvia Cerruti¹ · Justine Laulin² · Angela Cattaneo³ · Angela Bachi⁴ · Sebastien Apcher² · Olivier Coux⁵ · Paolo Cascio¹

Received: 6 November 2020 / Revised: 15 November 2021 / Accepted: 17 November 2021 / Published online: 16 December 2021
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2021

Abstract

PA28 γ is a nuclear activator of the 20S proteasome that, unlike the 19S regulatory particle, stimulates hydrolysis of several substrates in an ATP- and ubiquitin-independent manner and whose exact biological functions and molecular mechanism of action still remain elusive. In an effort to shed light on these important issues, we investigated the stimulatory effect of PA28 γ on the hydrolysis of different fluorogenic peptides and folded or denatured full-length proteins by the 20S proteasome. Importantly, PA28 γ was found to dramatically enhance breakdown rates by 20S proteasomes of several naturally or artificially unstructured proteins, but not of their native, folded counterparts. Furthermore, these data were corroborated by experiments in cell lines with a nucleus-tagged myelin basic protein. Finally, mass spectrometry analysis of the products generated during proteasomal degradation of two proteins demonstrated that PA28 γ does not increase, but rather decreases, the variability of peptides that are potentially suitable for MHC class I antigen presentation. These unexpected findings indicate that global stimulation of the degradation of unfolded proteins may represent a more general feature of PA28 γ and suggests that this proteasomal activator might play a broader role in the pathway of protein degradation than previously believed.

Keywords Proteasome activator · Proteasome gate · Protein degradation · ATP-independent proteolysis · Intrinsically disordered proteins (IDP) · PA28 $\alpha\beta$ · PSME 3

Introduction

The vast majority of intracellular proteins in eukaryotic cells are degraded by the 26S proteasome, a large (2.4 MDa) multimeric protease abundantly expressed in the nucleus and cytosol [1]. The 26S proteasome is composed of the 20S core particle, which in its internal cavity harbors the

proteolytic sites and the 19S regulatory particle, which is used for recognition, unfolding, and translocation of protein substrates within the protease [2]. The 20S proteasome is the central core of this proteolytic macromolecular machine, has a cylindrical structure with a molecular weight of ~700 kDa, and is formed by four overlapping rings of seven subunits each. The two outer rings are constituted of α subunits, and the two inner ones of β subunits [3]. The catalytic active sites of the constitutive 20S proteasomes are located on the β 1, β 2, and β 5 subunits. However, under the stimulus of γ -interferon or other pro-inflammatory cytokines, new catalytic subunits are synthesized (β 1i, β 2i, and β 5i) which replace those constitutive in so-called newly assembled 20S immunoproteasomes, which are optimized to generate the antigenic peptides (or epitopes) that are presented on the cell surface in association with MHC class I complexes [4]. In general, proteasomes can cleave any peptide bond except those at the C-terminus of proline; peptidase activities defined by fluorogenic substrates recognize three cleavage specificities: trypsin-like (i.e., hydrolysis after

✉ Paolo Cascio
paolo.cascio@unito.it

¹ Department of Veterinary Sciences, University of Turin, Largo P. Braccini 2, 10095 Grugliasco, Turin, Italy

² Université Paris-Saclay, Institut Gustave Roussy, Inserm, Immunologie Des Tumeurs et Immunothérapie, Villejuif, France

³ Cogentech SRL Benefit Corporation, 20139 Milan, Italy

⁴ The FIRC Institute of Molecular Oncology (IFOM), 20139 Milan, Italy

⁵ Centre de Recherche de Biologie Cellulaire de Montpellier (CRBM), CNRS UMR 5237, Université de Montpellier, 1919 Route de Mende, 34293 Montpellier, France

basic residues), chymotrypsin-like (i.e., hydrolysis after hydrophobic residues), and caspase-like (i.e., after acidic residues) [5].

Another important family of 20S proteasome regulators is that of the ATP- and ubiquitin-independent PA28 activators (also called 11S, REG or PMSE), which in vertebrates is formed by three highly homologous subunits α , β , and γ [6]. The crystal structure of PA26 (a homolog of PA28 in *Trypanosoma brucei*) clearly reveals that binding with the activator determines the opening of the gate, formed by the N-terminal tails of the α subunits, which normally occludes the access to (and exit from) the internal proteolytic cavity [7]. It is generally assumed that this represents the main molecular mechanism of action of all the other members of this class of proteasome activators [8]. Accordingly, opening of the gate, although with a related but different molecular mechanism, has also been described for a PA28 homolog present in *Plasmodium falciparum*, whose structure in association with the homospecific 20S particle has recently been solved. Interestingly, in this case PA28 was found to bind 20S asymmetrically, strongly engaging subunits on only one side of the core particle [9]. However, strong evidence indicating that PA28 may also act through long distance allosteric modifications of proteasomal proteolytic sites has been recently reported [10–12].

PA28 α and β form a heteroheptameric ring with a subunit stoichiometry of $4\alpha 3\beta$ [13] present in both the nucleus and cytosol of mammalian cells and whose formation is highly induced by γ -interferon [14]. PA28 $\alpha\beta$ in vitro strongly stimulates hydrolysis of short peptides by all three peptidase activities of 20S proteasomes, but not of full-length proteins, regardless of whether they are ubiquitinated, properly folded, or completely denatured [15–18]. However, a specific function of PA28 $\alpha\beta$, predominantly in association with the immunoproteasome in promoting the degradation of oxidized proteins, has been also described, which may indicate that it plays a role in adaptive responses to stressful conditions [19–24]. In vivo PA28 $\alpha\beta$ exerts a clear action of stimulation of MHC class I antigen presentation, although this activity is restricted to only some epitopes [25]. On the contrary, PA28 γ forms homoheptameric rings that are located exclusively in the nucleus and which are not induced (and even reduced) by γ -interferon [26]. PA28 γ was initially reported to exclusively enhance the trypsin-like activity of proteasomes [27, 28], but additional data indicates that it can also stimulate the other two (i.e., chymotrypsin- and caspase-like) peptidase activities [29–31]. Importantly, PA28 γ was shown to increase degradation of some full-length proteins through both direct and indirect mechanisms [32]. Specifically, the list of proteins that are directly targeted by PA28 γ toward proteasomal degradation in an ATP- and ubiquitin-independent process includes the oncogenic proteins steroid receptor coactivator SRC-3 [33], HCV core protein [34],

pituitary tumor-transforming 1 PTTG1 (securin) [35], cyclin-dependent kinase inhibitors p21 [36], p16, and p19 [37]. PA28 γ has also been shown to strongly reduce the stability of the tumor suppressor p53 through an indirect mechanism of stimulus involving ubiquitination by the ubiquitin-ligase MDM2 and subsequent ubiquitin- and ATP-dependent 26S proteasome degradation [38]. The exact biological functions of PA28 γ have not yet been fully elucidated, although this proteasome activator is clearly involved in the regulation of several essential cellular pathways, including cell growth and proliferation [39], transition from G to S phase in the cell cycle [40], inhibition of apoptosis [41], chromosomal stability [42], nuclear dynamics through modulation of the number and size of various nuclear bodies, including Cajal bodies (CBs) [30, 43], nuclear speckles [44], promyelocytic leukemia protein bodies [45], cellular response to DNA double-strand breaks [46], autophagy inhibition [47], spermatogenesis [31], neoplastic transformation [48–50], and MHC class I antigen presentation [51, 52]. Since in previous studies we demonstrated that heteroheptameric PA28 $\alpha\beta$ profoundly modifies the size and composition of amino acid sequences of products generated during proteasomal degradation of unfolded proteins without, however, affecting the overall rates of substrate hydrolysis [18], the present study was undertaken to determine whether similar biochemical behavior is also shared by the homolog PA28 γ .

Materials and methods

Proteins' purification

Recombinant PA28 γ and PA28 $\alpha\beta$ were expressed in *E. coli* and purified as described previously [30, 53]. Human 20S proteasome was purified from extracts of HeLa cells (Ipracell, Mons, BE) using classic chromatographic procedures as reported [30]. All preparations were free of contaminant endo- and exo-proteolytic activities that may interfere with proteasomal degradation experiments. PA28 γ - and PA28 $\alpha\beta$ -20S proteasomes were reconstituted by preincubating 20S particles with a sixfold molar excess of PA28 γ at 37 °C for 30 min in 20 mM HEPES, pH 7.5, and were used immediately for degradation experiments.

Peptidase assays and kinetic analysis

Peptidase activities of 20S proteasomes were measured using specific fluorogenic substrates (Bachem, Bubendorf, CH) at concentrations between 100 and 200 μ M in 20 mM Tris-HCl, pH 7.5, 0.2% BSA as previously described [54]. Briefly, the fluorescence of released amc (excitation, 380 nm; emission, 460 nm) was monitored continuously at 37 °C with a Cary Eclipse spectrofluorometer (VARIAN,

Palo Alto, CA, USA). Assays were calibrated using standard solutions of the free fluorophore, and reaction velocities were calculated from the slopes of the initial linear portions of the curves. Substrate consumption at the end of incubation never exceeded 1%. Kinetic parameters of the PA28 γ -20S complexes were calculated by assessing reaction velocities at different activator concentrations. The best fit was performed using GraphPad prism (version 5) software through nonlinear regression for enzyme kinetics.

Protein degradation and analysis of peptide products

Degradation of β -casein, IGF-1, MBP and α -lactalbumin was performed as previously described [18, 55]. Reduction and carboxymethylation of IGF-1 and α -lactalbumin to ensure denaturation and prevent sulfhydryl bond formation were performed according to published methods [56]. To study the kinetics of substrate degradation using fluorescamine, casein (710 μ M), IGF-1 (830 μ M), MBP (85 μ M), and α -lactalbumin (307 μ M) were incubated with 20S and PA28 γ -20S proteasomes (20 to 80 nM depending on the substrate) in 20 mM HEPES, pH 7.5. Epoxomicin (Enzo Life Sciences, Farmingdale, New York) was used at a final concentration of 20 μ M. To assay peptides generated during protein degradation, at the indicated time points peptide products were separated from undegraded protein by ultrafiltration through a membrane with a 3 kDa cutoff (Pall Corporation, NY, USA), and the appearance of new amino groups was measured using fluorescamine [56]. Consumption of the substrate at the end of the 8-h incubation never exceeded 10%. Degradation of IGF-1 (400 μ M), MBP (90 μ M), and α -lactalbumin (260 μ M) by 20S and PA28-20S proteasomes (120 nM for IGF-1 and α -lactalbumin, 5.6 nM for MBP) for electrophoretic analysis was performed at 37 °C in the same buffer described previously. At 0, 4, 8 and 12 h, aliquots were taken, the undegraded protein was separated on a 12% (for MBP) or 18% (for IGF-1) SDS-PAGE gel, and densitometric analysis of Coomassie-stained bands was performed with a VersaDoc 1000 Imaging System (Bio-Rad Laboratories, Hercules, CA, USA) using Quantity One software (Bio-Rad Laboratories, Hercules, CA, USA). For HP-size exclusion chromatography of β -casein products, peptides generated during an 8h hydrolysis reaction were isolated by ultrafiltration using both 3 and 10 kDa cutoff membranes (Pall Corporation, NY, USA), which gave very similar results in terms of size distribution of proteasome products. After that, equal amounts of peptides were diluted in 0.1 M HEPES, pH 6.8, and separated on a polyhydroxyethyl aspartamide column (0.46 \times 20 cm, Poly LC, Columbia, MD, USA) using an HP1100 HPLC (Hewlett-Packard, Palo Alto, CA, USA) equipped with a fluorometer. The mobile phase was 0.2 M Na₂SO₄, 25% acetonitrile (pH 3.0;

adjusted with phosphoric acid) at a flow rate of 0.125 ml/min. For each analysis, 20 μ l of peptide solution was added to 16 μ l of fluorescamine (dissolved 0.3 mg/ml in acetonitrile). The reaction was terminated after 30 s with 22 μ l of H₂O, and the sample was immediately injected on the HPLC column. The fluorescence of eluted material was monitored continuously and a blank run (corresponding to time 0 h of the degradation reaction) was always subtracted. To determine the apparent molecular mass of peptides eluted, the column was calibrated with 19 standard amino acids and peptides in the 75–3500 Da range that had been derivatized with fluorescamine in the same manner as proteasomal degradation products. Prior control studies showed that retention times of these fluorescamine-derivatized products are highly reproducible and linearly dependent on the logarithm of their molecular weights, and that recovery of amino acids and peptides of different lengths is quantitative [18]. Note that amino acids and peptides eluting from the column are bound to fluorescamine, whose molecular weight must be subtracted to calculate the actual mass of proteasomal products. Mean and median sizes of peptides generated by 20S and PA28 γ -20S proteasomes were calculated from the distributions of products obtained by SEC, assuming an average molecular weight of 110 Da for each residue.

Liquid chromatography–tandem MS (LC–MS/MS) analysis

Samples from the 20S -/+ PA28 γ 6 h degradation experiments containing approximately 5 (for IGF-1) and 30 (for MBP) pmol NH₂/ μ l were loaded onto a StageTips μ C18 column [57]; peptides were eluted in 40 μ l 80% acetonitrile in 0.1% formic acid. The acetonitrile was allowed to evaporate in a Speed-Vac, and samples were then resuspended in 10 μ l of eluent A (see composition below) for nLC–MS/MS analysis. Two microliters of each sample was injected as technical replicates into a nLC–ESI–MS/MS quadrupole Orbitrap QExactive-HF mass spectrometer (Thermo Fisher Scientific). Peptide separation was achieved using a linear gradient from 95% solvent A (2% ACN, 0.1% formic acid) to 50% solvent B (80% acetonitrile, 0.1% formic acid) for 23 min and from 60 to 100% solvent B for 2 min at a constant flow rate of 0.25 μ l/min on a UHPLC Easy-nLC 1200 (Thermo Scientific) connected to a 25-cm fused-silica emitter with an inner diameter of 75 μ m (New Objective, Inc. Woburn, MA, USA), packed in-house with ReproSil-Pur C18-AQ 1.9- μ m beads (Dr. Maisch GmbH, Ammerbuch, Germany) using a high-pressure bomb loader (Proxeon, Odense, Denmark). MS data were acquired using a data-dependent top-15 method for HCD fragmentation. Survey full-scan MS spectra (300–1750 Th) were acquired in the Orbitrap at a resolution of 60,000, an AGC target of 1×10^{66} , and an IT of 120 ms. For HCD spectra, the resolution was

set to 15,000 at m/z 200, with an AGC target of $1e^5$, an IT of 120 ms, an NCE of 28% and an isolation width of 3.0 m/z .

Data processing and analysis

For quantitative proteomics analysis, raw data were processed with MaxQuant (ver. 1.6.0.16) and searched against a database containing the sequence of the IGF-1 and MBP + contaminant fasta included in MaxQuant. No enzyme specificity was selected, and there were no differences between I and L. The mass deviation for MS/MS peaks was set at 20 ppm, and the peptide false discovery rate (FDR) was set at 0.01. The list of identified peptides was filtered to eliminate contaminants. Statistical analyses were performed with Perseus (ver. 1.6.2.3) considering the peptide intensity; normalization based on the Z-score and imputation was applied. Significant peptides were determined with a t test, Benjamini–Hochberg correction, and $FDR < 0.05$. Only significant peptides were used for supervised hierarchical clustering analysis. MS data as raw files, peptides identified with relative intensities and search parameters have been deposited to the ProteomeXchange Consortium via the PRIDE [58] partner repository with the dataset identifier PXD029248.

MBP expression in cell lines

The gene coding for MBP isoform (transcript variant 1) containing in its sequence a nontraditional PY-Nuclear Localization Signal (NLS) was purchased from Origene and insert in pCMV6-Entry eukaryotic expression plasmid. The A375 (ATCC, nCRL-1619) human melanoma cells and the derivative clone A375 Crispr knockout for PA28 γ have been described in detail [52] and were cultured according to ATCC's protocol. Once a month, mycoplasma contamination in cell cultures was assessed using the Venor[®]GeM OneStep mycoplasma detection kit (Minerva biolabs). Cells were used within 4 weeks after thawing (~10 passages) and were transfected with 1 μ g of plasmid DNA along with 2 μ l of JetPrime according to the manufacturer's protocol (Ozyme). Importantly, no difference in growth rates was observed between WT A375 and A375 Crispr cells. For Western blots, cells were lysed in RIPA buffer supplemented with Complete[™] Protease Inhibitor (Roche, #11697498001) and centrifuged at 16,000 $\times g$ for 20 min at 4 °C. Protein concentration of the supernatant was determined by BCA assay using the Pierce[™] BCA Protein Assay Kit (Thermo Fisher Scientific). 15 μ g of proteins was denatured for 10 min at 95 °C in 1X Laemmli sample buffer containing 2.5% β -mercaptoethanol. Proteins were loaded on 4–15% acrylamide gels and run at 100 V in 1X TGS buffer. Proteins were then transferred onto PVDF membranes using a Trans-Blot Semi-Dry Transfer Cell (Bio-Rad). Membranes

were blocked for 1 h with 5% milk solution and incubated with primary antibody overnight at 4 °C. The day after, the blot was incubated with a secondary antibody coupled with HRP for 1 h at RT. An anti- β -actin antibody (Santa Cruz, #sc47778) coupled with HRP was used as a loading control. Depending on the experiment, the following primary and secondary antibodies were used: anti-Myc-tag (9B11) mouse mAb (Cell Signaling, #2276), anti-PA28 γ recombinant rabbit mAb (Thermo Fisher Scientific, #700180), anti-p21 Waf1/Cip1 (12D1) rabbit mAb (Cell Signaling, #2947), anti- β -actin antibody coupled with HRP (Santa Cruz, #sc47778), polyclonal swine anti-rabbit immunoglobulins/HRP (Dako, #P0217) and polyclonal rabbit anti-mouse immunoglobulins/HRP (Dako, #P0260). Membrane images were acquired with ChemiDoc[™] MP and ImageLab software; no photographic editing was performed after acquisition.

Statistical analyses

To compare average measurements of generation of amino group and protein degradation and expression levels, we adopted a non-parametric Mann–Whitney test.

Results

PA28 γ differently affects 20S proteasome peptidase activities

In a preliminary effort aimed at better characterizing the exact biochemical properties of PA28 γ activator, we assessed its effects on the hydrolysis of a panel of short fluorogenic peptides specifically designed to probe endopeptidase activities. To perform these (and the following) experiments, we utilized a recombinant protein expressed in *E. coli* and purified to homogeneity as described in “Materials and Methods”. Importantly, size exclusion chromatography of the final preparation demonstrated complete conversion of the PA28 γ monomer into a stable high molecular weight complex of apparent MW of ~200 kDa, consistent with that of a heptameric ring (Supplementary Fig. 1A). Initially, we showed that of the 12 fluorogenic peptides tested (4 designed to probe chymotrypsin-like, CT-L, 2 caspase-like, C-L, and 6 trypsin-like T-L, endopeptidase activities), all were cleaved by human 20S particles, although at greatly divergent rates, as expected for substrates that differ in both length and amino acid composition (Supplementary Table 1). Next, we investigated the effect of increasing concentrations of PA28 γ on the degradation rates of these proteasomal substrates. As shown in Supplementary Fig. 2, a strong dose-dependent acceleration of the rate of hydrolysis reactions could be observed for ten peptides, and this stimulation

involved substrates cleaved by all three proteasome active sites. Consequently, the maximal specific activity of 20S particles completely activated by PA28 γ (i.e., 20S bound at both side with a PA28 γ ring) varied from ~2 to ~450 nmol of peptide cleaved per min/mg of enzyme depending on the substrate (Table 1). Interestingly, however, the extent of this PA28 γ -dependent stimulation (compared to the basal activity of 20S in absence of the activator) was not constant, but ranged between 4- and 18-fold, and unexpectedly differed strongly even between peptides cleaved by the same proteasomal active site (Table 1). Furthermore, careful analysis of the mechanism of 20S activation with increasing amounts of PA28 γ showed a kinetic trend consistent with a rectangular hyperbola, with a concentration of the activator in the nanomolar range required to fully activate picomolar concentrations of proteasome (Supplementary Fig. 2). Of note, the non-sigmoid shape of the activation kinetics of hydrolysis of the fluorogenic peptide indicates the absence of cooperative binding between the two PA28 γ molecules that associate with the two opposite ends of the 20S particle. In contrast with these results, hydrolysis by proteasomal trypsin-like activity of two dipeptides (LR and FR), unlike that of a third one which is very similar (RR), was completely unaffected by the presence of high concentrations of PA28 γ (Supplementary Fig. 3A). Of note this took place when all three peptides were tested at a concentration (i.e., 100 μ M) at which the catalytic sites of the 20S are not saturated by the substrate, and therefore below V_{max} (Supplementary Fig. 3B).

Table 1 PA28 γ kinetic parameters. Maximum PA28 γ stimulation of 20S proteasome peptidase activities was calculated as described in the Materials and Methods from the curves shown in Supplementary Fig. 2

Proteasomal activity	Substrate	Maximum stimulation of 20S activities by PA28 γ	
		Fold	20S maximum specific activity (nmol peptide cleaved /mg * min)
Chymotrypsin-like	Suc-LLVY-amc	18.2	223.7
	Suc-LY-amc	4.3	62.6
	Suc-GGL-amc	3.8	368.6
	Suc-AAF-amc	6.7	264.1
Caspase-like	Suc-YVAD-amc	10.2	29.0
	Suc-DEVD-amc	6.7	7.0
Trypsin-like	Bz-VGR-amc	7.3	444.6
	Z-ARR-amc	4.3	134.8
	Boc-LRR-amc	16.5	411.0
	Z-RR-amc	6.1	1.9

PA28 γ enhances the rates of peptide bond cleavage in the full-length protein β -casein

If on the one hand, short fluorogenic peptides represent useful tools to investigate the biochemical properties of proteasomal peptidase sites, on the other hand their hydrolysis only roughly mirrors the complex process of degradation of real, physiological substrates of proteasomes, which in vivo mainly degrade proteins rather than short polypeptides [59, 60]. For this reason, we decided to investigate the effect of PA28 γ on degradation of full-length proteins. To this end, we initially chose β -casein, widely used for biochemical studies as a proteasome substrate, which has little tertiary structure and thus does not require artificial denaturation to be degraded at detectable rates by 20S particles [56, 61–64]. As shown in Fig. 1A, the rates of appearance of new amino groups, generated as a consequence of the hydrolysis of peptide bonds in the protein and assessed by means of fluorescamine, were linear for up to 8 h for both proteasome species analyzed (i.e., 20S and PA28 γ -20S). Moreover, as demonstration of the absolute dependence on the proteasome of the hydrolysis reaction, NH_2 generation was completely absent in the presence of the more specific proteasome inhibitor epoxomicin (Fig. 1A) and when the substrate was incubated alone or with only PA28 γ (Supplementary Fig. 4A and B). More importantly, however, the PA28 γ -20S particles reproducibly released threefold more NH_2 at each time point than unligated 20S (Fig. 1A). This finding was surprising because PA28 γ has generally been reported to be able to stimulate proteasomal degradation of small peptides, but only very few full-length proteins [32, 65] and, to the best of our knowledge, not for β -casein.

Association of PA28 γ with 20S does not affect size distribution of products generated during hydrolysis of casein

Theoretically, the higher rate of generation of primary amino groups by PA28 γ -20S proteasomes might be caused by either enhanced substrate hydrolysis induced by the activator or, alternatively, by a significant reduction of size distribution of peptide products generated (i.e., even in the absence of modifications in the rates of β -casein consumption), as in the case of the homolog PA28 $\alpha\beta$ [18]. To discriminate between these two alternatives, we analyzed the size distribution of products generated during hydrolysis of casein by 20S and PA28 γ -20S particles using an HP-size exclusion chromatographic method that allows linear separation and accurate quantification of peptides in the range of 1–30 residues (Supplementary Fig. 5).

In particular, the protein was degraded at linear rates and under conditions ensuring that peptides released by proteasomes do not re-enter the degradative particle, and therefore




Fig. 1 Increased rates of peptide bond cleavage and size distribution of peptides generated during hydrolysis of casein by PA28 γ -20S proteasomes. **A** β -casein was incubated with 20S and PA28 γ -20S proteasomes and the amino groups released, as a consequence of the degradation of the substrate, were measured with the fluorescamine at the indicated time points. Data are the average of three independent experiments (four for time 8 h) \pm SEM. Appearance of amino groups was totally abrogated in presence of 20 μ M of the highly specific proteasome inhibitor epoxomicin. *** P < 0.001. **B** Equal amounts of peptides generated during degradation of casein by 20S and PA28 γ -20S proteasomes were reacted with fluorescamine and immediately fractionated by HP-SEC. Data are representative of four independent experiments. **C** Mean and median peptide sizes generated by 20S and PA28 γ -20S proteasomes from casein were calculated using the product distributions obtained by HP-SEC, assuming an average molecular weight of 110 Da for each residue. Values are the average of four experiments (\pm SEM)


not subjected to a second round of hydrolysis (i.e., the substrate was present in large excess and not more than 10% was degraded at the end of the incubation) [55]. When analyzed by this approach, peptides generated from casein by the 20S and PA28 γ -20S proteasomes were found to fall into

a continuum of size distribution ranging from 1 to 30 residues (Fig. 1B) that appeared to fit a lognormal distribution, which is in agreement with previous analyses of different substrates and proteasome species [62, 64, 66]. Specifically, the chromatographic profile of 20S proteasome products was characterized by a large peak that culminates at the size of two to three residues with a small shoulder corresponding to less than 3% of the total fluorescence signal eluting from the column, in correspondence of sizes between 30 and 16 residues (Fig. 1B). Of interest, a nearly superimposable profile was also detected for peptides generated by PA28 γ -20S particles (Fig. 1B). In this case, the only minor differences in size distribution involved a reduction of about two-thirds in the minor peak of larger (> 16) residues accompanied by a small, but reproducible, shift toward a high molecular weight of the peak of the main products when PA28 γ associated with 20S particles (Fig. 1B). Moreover, if samples fractionated by SEC-HPLC were not normalized by the amount of peptides generated, equal volumes of degradation reaction carried out with equimolar amounts of enzymes (i.e., 20S and PA28 γ -20S proteasomes) were analyzed, the profile of the peak of PA28 γ -20S products did not change. This is in good agreement with the kinetic data shown in Fig. 1A, and its height and area increased about threefold (data not shown). As a result, the means of product lengths calculated from HPLC size distributions were the same (4 residues) for 20S and PA28 γ -20S proteasomes and the medians (~3 residues) were also very similar (Fig. 1C).

PA28 γ , but not PA28 $\alpha\beta$, consistently stimulates hydrolysis of different unfolded proteins by the 20S proteasome

To assess whether the unexpected finding that PA28 γ is able to promote proteasomal degradation of the naturally disordered substrate β -casein might imply a more general, and so far unrecognized, property of this activator, we challenged its role in degradation of several unrelated, unfolded proteins by the 20S particle. To this end we utilized another natively unfolded substrate, myelin basic protein (MBP), and two chemically denatured proteins, insulin-like growth factor 1 (IGF-1) and α -lactalbumin. As shown in Fig. 2, the rates of peptide bond cleavage by the 20S proteasome were dramatically enhanced for all three substrates in the presence of PA28 γ . Furthermore, as for β -casein, the dependence on the proteasome of the hydrolysis reaction was demonstrated by the absence of NH₂ generation in the presence of epoxomicin (Fig. 2) or when the substrates were incubated alone or only with PA28 γ (Supplementary Fig. 6). Moreover, the absolute specificity for unfolded proteins of the PA28 γ -induced proteolytic stimulation was unambiguously shown by directly comparing the rates of proteasomal hydrolysis of chemically unfolded

Fig. 2 Increased rates of peptide bond cleavage during hydrolysis of different unfolded proteins by PA28 γ -20S proteasomes. MBP, IGF-1, and α -lactalbumin were incubated with 20S and PA28 γ -20S proteasomes and the amino groups released were measured with the fluorescamine as in Fig. 1. Data are the average of three to six independent experiments \pm SEM. *** $P < 0.001$



IGF-1 and α -lactalbumin with those of their native, folded counterparts. In fact, the rates of proteasomal hydrolysis of folded IGF-1 and α -lactalbumin were nearly undetectable (data not shown) and only minimally affected by the presence of PA28 γ , in striking contrast with what observed for their corresponding denatured forms (Fig. 3).

Further direct and unequivocal evidence of the enhancement of proteins turnover rates induced by PA28 γ was subsequently obtained by following the disappearance of undigested substrates incubated for several hours in the presence of 20S proteasome alone or conjugated with PA28 γ . These experiments clearly demonstrated that the presence of PA28 γ strongly accelerated the kinetics of full-length unfolded proteins hydrolysis by the 20S proteasome (Fig. 4). This result appears even more remarkable in light of the fact that, in parallel experiments, PA28 $\alpha\beta$ was unable to activate denatured protein degradation (Fig. 4), in accordance with what has already been published for other full length substrates [15, 16, 18]. Moreover, as demonstration of the absolute dependence on the proteasomes of the hydrolysis reaction, disappearance of substrate was completely prevented when PA28 γ -20S particles were inhibited by epoxomicin or when the substrates were incubated alone (Supplementary Fig. 7).

PA28 γ modifies the patterns of peptides produced by 20S proteasome during hydrolysis of full-length proteins

To understand the modifications in the 20S proteasome proteolytic properties induced by the binding of PA28 γ , the peptides generated from IGF-1 and MBP were analyzed by tandem mass spectrometry (MS/MS). By this approach, 171 and 279 individual peptides were identified from IGF-1 and MBP, respectively, ranging in length from 8 to 25 residues (shorter products were eliminated to reduce false-positive identification) and are listed in the Supplementary Table 2. Importantly, this analysis allowed accurate identification of products with the correct size to potentially bind (directly or after trimming by aminopeptidases) to MHC class I heterodimers, and therefore is of particular interest to evaluate the possible role of PA28 γ in the context of cell-mediated immune responses.

Through this experimental approach, we also demonstrated that some peptides are exclusively released by one of the two proteasomal forms (i.e., 20S or PA28 γ -20S particle). Remarkably, and somewhat unexpectedly (see Discussion), in the range of lengths taken into consideration, almost all




Fig. 3 PA28 γ promotes proteasomal degradation of unfolded proteins but not of their native counterparts. Native and denatured IGF-1 (upper panel) and α -lactalbumin (lower panel) were incubated with PA28 γ -20S proteasomes and the amino groups released, as a consequence of the degradation of the substrate, were measured with fluorescamine at the indicated time points. Data are the average of three to five independent experiments \pm SEM. *** P < 0.001


the peptides generated in a specific way by only one form of proteasome were released by the 20S particle, while very few were produced exclusively in the presence of PA28 γ (Supplementary Table 2). Thereafter, our analysis focused on more detailed characterization of the peptides generated in common by both 20S and PA28 γ -20S particles. To this end, the relative amount of each product was assessed by comparing the corresponding ion intensities measured in sequential MS/MS analyses, according to a method we have validated previously [18, 52]. Briefly, ion intensities were used to quantify the relative amounts of single fragments generated from IGF-1 and MBP by 20S proteasome

alone or when associated with PA28 γ . The results of this analysis are shown in Fig. 5 and show that several peptide products are released in vastly different amounts by the 20S and PA28 γ -20S particles. In this case, the association with PA28 γ was also found to cause a consistent reduction in the number of individual peptides released in greater quantities during degradation of IGF-1 and MBP by the 20S proteasome. Although this effect of PA28 γ concerns the peptides of the entire range of lengths analyzed, a marked reduction in the generation of longer fragments can be noted, especially for IGF-1 (Fig. 5 and Supplementary Table 2), which is in agreement with the decrease of high molecular weight peptides released from casein in the presence of PA28 γ seen in Fig. 1B. Furthermore, to investigate the modifications in the cleavage specificities caused by the association of PA28 γ with the 20S particle, we analyzed the relative frequency of the amino acids present on the two sides (positions from P4 to P4') of the peptide bonds, whose hydrolysis generated the peptides quantified by MS/MS. To analyze this, we compared the amino acid sequences involved in the generation (both at the N and at the C-ter) of peptides produced in greater amounts by 20S alone or by PA28 γ -20S (Fig. 5). Although limited to peptides longer than seven residues, this analysis did not highlight any enrichment of basic amino acids in the P1 position in the presence of PA28 γ (Supplementary Fig. 8), in good agreement with the data obtained with fluorogenic peptides, which show that PA28 γ does not stimulate only tryptic 20S activity.

PA28 γ regulates intracellular level of ectopic MBP

To verify if what was observed in vitro was also reflected in cell lines, we took advantage of a particular 21.5 kDa isoform of MBP (i.e., isoform 3 or MBP1) which has been shown to be an intrinsically disordered protein, such as the most common 18.5 kDa isoform (isoform 5 or MBP3). The 21.5 kDa isoform also contains a nontraditional PY-nuclear localization signal (NLS) in its sequence that allows its transport and accumulation at the nucleus [67, 68]. The expression of the protein (carrying a Myc-tag to facilitate its detection by western blot) was ectopically induced from a plasmid in eukaryotic cells characterized by different levels of PA28 γ expression. Specifically, we utilized the A375 human melanoma cell line, which expresses high basal amounts of PA28 γ , and its derivative Cas9-A375 PA28 γ -knockout line (A375 Crispr), obtained by the CRISPR/Cas9 system and characterized in detail previously [52]. As expected, the A375 Crispr clone shows no detectable amounts of PA28 γ , regardless of whether MBP is present (Fig. 6A). Crucially, when expressed in parallel in the two different clones (i.e., A375 WT and A375 Crispr), MBP accumulates over time to much higher levels in cells that do not express PA28 γ compared to those expressing it. This effect is evident within 24 hs after transfection, but becomes

Fig. 4 Enhanced rates of MBP, IGF-1, and α -lactalbumin proteasomal hydrolysis induced by PA28 γ , but not by PA28 $\alpha\beta$. MBP **A**, IGF-1 **B**, and α -lactalbumin **C** were incubated as indicated in the figure, and the undegraded protein was separated by SDS-PAGE. Note that in the MBP gel the bands corresponding to enzymes cannot be appreciated since they were used at lower concentrations (see [Materials and Methods](#) for further details). **D** Densitometric quantification of the residual protein. Data are the average of three to four independent experiments (\pm SD). *** P < 0.001



very strong after 48 h (Fig. 6B and Supplementary Fig. 9). As independent demonstration that PA28 γ -dependent proteasomal activity is impaired in A375 Crispr cells, the intracellular amounts of p21, a well characterized PA28 γ -20S proteasome substrate, were similarly increased in these cells (Fig. 6C). Although the biological relevance of this observation needs still to be investigated in greater detail, this in cell lines data confirm the findings of our in vitro experiments.

Discussion

To shed light on the molecular mechanism of PA28 γ activity, we initially assessed the effect of PA28 γ on the hydrolysis of a panel of different fluorogenic short peptides (2–4 amino acids long) thus probing all three 20S proteasome active sites. Although initial reports indicated that PA28 γ





Fig. 5 MS/MS differential analysis of peptides generated during hydrolysis of IGF-1 and MBP by 20S and PA28 γ -20S particles. Heatmap comparison of the abundance of significant peptides generated after 6 h of proteasomal degradation of IGF-1 and MBP in the 20S (biological and technical replicates of samples designated **A**, **B**, and **C** and PA28 γ -20S (biological and technical replicates of samples designated 1, 2, and 3) samples. Samples were analyzed by nLC-

MS/MS and processed by MaxQuant against the database IGF-1 or MBP+contaminant sequences. Differences and similarities in peptide intensities (normalized to the Z-score) are shown; green indicates decreased levels, and red indicates increased levels. Data were obtained from supervised hierarchical clustering analysis by applying a *t* test, Benjamini–Hochberg correction, and a *p* value of 0.05

exerts its stimulatory effect only towards 20S trypsin-like activity [27, 28], subsequent studies also described strong activation of the other two proteasomal peptidase activities both in vitro [29, 30] and in vivo [31]. In accordance with these latter findings, our study showed that hydrolysis rates of ten fluorogenic substrates were highly enhanced when PA28 γ associated with the 20S particle. Importantly,

this stimulatory effect was not restricted to tryptic activity, but clearly affected chymotryptic and caspasic proteasomal cleavage specificities. Moreover, the extent of 20S activation varied between 4- and 18-fold, with no clear preference for a proteasome catalytic site or a specific substrate size.

Fig. 6 Knockout of PA28 γ results in accumulation of a nucleus-targeted MBP isoform in cell. **A** The expression of PA28 γ was assessed by western blotting in A375 and A375 Crispr melanoma cells. The intensity of each band was normalized to that of the corresponding β -actin band, and the value related to that of the control (A375 expressing MBP) set as one. Experiments were performed in triplicate and data are expressed as mean \pm SE. **B** Expression levels of MBP was assessed by WB with an anti-Myc-tag antibody, and data are presented as in (A). **C** Expression of p21 was verified by WB and indicated as in (A). * $P < 0.05$



The most surprising finding of our study, however, arises from experiments characterizing the effects of PA28 γ on the hydrolysis of several naturally or chemically unstructured proteins. It is known that, by virtue of the absence of tightly folded higher-order structures, denatured proteins freely diffuse inside the proteolytic cavity of latent 20S particles, whose axial pores, through which substrates access the internal catalytic lumen and products exit from the particle, are obstructed by the N-terminal tails of the α subunits [69]. However, the latency of the 20S core particle not associated with activators (i.e., 19S, PA28, PA200) is not complete, because, even without artificial conditions that result in its activation [3], the 20S proteasome can degrade full-length proteins [18, 61], likely because of transitory and/or only limited gate opening [62]. Accordingly, when unfolded substrate proteins were incubated with 20S proteasomes the

rates of appearance of new amino groups, generated as a consequence of hydrolysis of the peptide bonds in the substrate, were linear for up to 8 h. The same linearity was also evident when substrates were degraded by PA28 γ -20S particles, but in this case an impressive higher generation of new amino groups (varying between three and tenfold, depending on the substrate) systematically took place at each time point. Crucially, this enhanced release of primary amino groups by PA28 γ -20S proteasomes might result from either accelerated substrate hydrolysis caused by the activator or by a significant reduction in the size distribution of the peptide products generated, as in the case of the homolog PA28 $\alpha\beta$ [18]. Analysis of the size distributions of proteasomal products released during degradation of β -casein in the presence and absence of PA28 γ , however, clearly disproves the second hypothesis. In fact, the chromatographic profiles

of 20S and PA28 γ -20S proteasome products appear nearly superimposable and, accordingly, the means and medians of product lengths are almost the same (4 and \sim 3 residues).

Since binding of PA28 γ does not significantly modify the size distribution of peptide products released from casein by 20S proteasomes, it is therefore clear that the higher generation rates of NH₂ groups seen in Fig. 1A must be truly indicative of an accelerated casein breakdown induced by the activator. Direct evidence that PA28 γ strongly stimulates the rates of unfolded protein hydrolysis by 20S proteasome was obtained by monitoring the disappearance of undigested substrates in time course in vitro degradation kinetics reactions. Such experiments unambiguously showed that binding of PA28 γ transforms relatively latent proteases such as free 20S into an efficient proteolytic enzyme that hydrolyzes unfolded proteins at greatly accelerated rates. Importantly, the stimulatory activity of PA28 γ appears restricted only towards unfolded substrates, as shown by its inability to sustain degradation of native IGF-1 and α -lactalbumin. This last observation is not totally unexpected, since, unlike the 19S activator, PA28 γ lacks the ATPase activity necessary to denature tightly folded higher-order structures in protein substrates to allow access to the internal proteolytic cavity of the 20S degradative particle. In line with these in vitro findings, a nucleus-targeted isoform of MBP has been shown to accumulate over time at much higher levels in a melanoma cell line knocked out for PA28 γ compared to the parental one that markedly expresses the activator. Although the biological relevance of this observation needs further investigation, this data is in agreement with the results of our in vitro experiments, and indicates that PA28 γ may promote proteasomal degradation of MBP in vivo, as also suggested by a recent investigation on the stability of MBP in human embryonic kidney 293 (HEK293) cells [70].

Of note, several studies have demonstrated that PA28 γ is able to increase in vitro and in vivo hydrolysis by 20S particles of some specific proteins such as SRC-3, p21, p16, p19, PTTG1, and HCV core protein [33–37]. Although all these proteins share the same characteristic of being completely or in large part intrinsically unstructured [65, 71, 72], it is generally believed that some very specific, yet to be identified, features in their amino acid sequence or in their residual folded structure may be responsible for their recognition and interaction with PA28 γ [32]. Therefore, the enhanced degradation of these proteins induced by PA28 γ is generally seen as an exception rather than as the normal consequence of an intrinsic feature of this proteasomal activator. On the contrary, the finding that proteasomal degradation of several totally unrelated proteins, which differ in terms of molecular weight, amino acid composition, and chemical properties (e.g., isoelectric point and hydrophobicity), but share the same absence of folded structures, is potently improved by PA28 γ , strongly suggests that stimulation of hydrolysis of

unstructured proteins might represent a more general property of this activator. This molecular model appears even more reinforced by the observation that PA28 γ does not enable 20S degradation of the folded counterparts of the denatured proteins. In view of the fact that several analyses indicate that a substantial fraction of eukaryotic proteome is composed of proteins which are completely or in large part intrinsically unstructured [73], and whose degradation seems to involve ubiquitin-independent mechanisms [74], it is therefore reasonable to speculate that PA28 γ might play a much more significant and broad role in the pathway of protein breakdown than previously believed. Further in vivo studies, possibly with selective inhibitors of PA28 γ that are currently underway will be required to clarify this important point.

The present data are even more surprising in the light of results we obtained previously in a similar investigation characterizing the biochemical functions of PA28 $\alpha\beta$ [18]. In that study, PA28 $\alpha\beta$ was found to strongly reduce the mean and median sizes of proteasome products and to profoundly modify the sequences of peptides released without, however, increasing the overall rates of protein substrate breakdown. Therefore, the biochemical effects on protein degradation by 20S particles of PA28 γ and PA28 $\alpha\beta$ seems to be rather opposed. Since PA28 γ and PA28 α and β share a high degree of homology, the molecular reasons underlying these differences in biochemical properties are likely to rely on the few divergent regions, namely the so-called homolog-specific inserts that connect helix 1 with helix 2 and which are not resolved in the X-ray structure of PA28 α , presumably because they are flexible [6]. Furthermore, recent phylogenetic analyses indicate that PA28 γ is most similar to the common ancestor of the PA28 activator family, and most likely retains its original functions, while PA28 α and PA28 β appeared later and evolved very rapidly to perform new tasks related to the γ -interferon-inducible MHC class I system [75], although their role in adaptive responses to stressful conditions (e.g., oxidative stress) also seems very likely [19–24]. Moreover, biochemical characterization of a PA28 γ homolog in *Dictyostelium discoideum* led to the conclusion that PA28 γ -20S proteasomes could represent early unique nuclear proteases of eukaryotic cells [76]. On the basis of these considerations, one would be tempted to speculate that PA28 γ could have maintained the ancient property of promoting degradation of unstructured long polypeptides and proteins, while the phylogenetically more recent PA28 α and β lost the ability to stimulate protein hydrolysis to gain a more pronounced capacity to modify the spectrum of peptides released by proteasomes in a functional way to favor MHC class I antigen presentation. This hypothesis is corroborated by the results of mass spectrometry analysis of products in our study. In fact, the association of PA28 γ with the 20S proteasome determines an evident reduction in the

variability of peptide products that are potentially available for class I presentation, while PA28 $\alpha\beta$ activity is always correlated with enhancing the diversity of proteasomal products, in a way that makes it more likely that an appropriate CTL response is elicited [18, 77, 78]. In this regard, it is worth noting that in the presence of PA28 γ there is a general reduction in the production of different peptides throughout the entire range of lengths analyzed by mass spectrometry. However, especially in the case of IGF-1 (and to a somewhat lesser extent in the case of MBP), PA28 γ leads to lower production of longer fragments, which has been shown to play an important role in MHC class I antigen presentation following trimming by aminopeptidases [79, 80].

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00018-021-04045-9>.

Acknowledgements We thank Francesco Turci and Francesco Ferrini for help in preparing figures, Patrick Moore for assistance in preparation of the manuscript and Massimo Coletta for insightful discussions.

Author contributions J-YAF acquisition, analysis and interpretation of data; FC acquisition, analysis and interpretation of data; JL acquisition, analysis and interpretation of data; AC acquisition, analysis and interpretation of data; AB analysis and interpretation of data; AS, analysis and interpretation of data; OC analysis and interpretation of data; PC conception and design, acquisition of data, analysis and interpretation of data, drafting the article.

Funding Ricerca Locale (ex 60%) to P.C.

Availability of data and material MS data as raw files, peptides identified with relative intensities and search parameters have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD029248.

Code availability Not applicable.

Declarations

Conflicts of interest The authors declare no conflict of interest.

References

- Voges D, Zwickl P, Baumeister W (1999) The 26S proteasome: a molecular machine designed for controlled proteolysis. *Annu Rev Biochem* 68:1015–1068. <https://doi.org/10.1146/annurev.biochem.68.1.1015>
- Baumeister W, Walz J, Zuhl F, Seemuller E (1998) The proteasome: paradigm of a self-compartmentalizing protease. *Cell* 92(3):367–380. [https://doi.org/10.1016/s0092-8674\(00\)80929-0](https://doi.org/10.1016/s0092-8674(00)80929-0)
- Coux O, Tanaka K, Goldberg AL (1996) Structure and functions of the 20S and 26S proteasomes. *Annu Rev Biochem* 65:801–847. <https://doi.org/10.1146/annurev.bi.65.070196.004101>
- Goldberg AL, Cascio P, Saric T, Rock KL (2002) The importance of the proteasome and subsequent proteolytic steps in the generation of antigenic peptides. *Mol Immunol* 39(3–4):147–164. [https://doi.org/10.1016/s0161-5890\(02\)00098-6](https://doi.org/10.1016/s0161-5890(02)00098-6)
- Harris JL, Alper PB, Li J, Rechsteiner M, Backes BJ (2001) Substrate specificity of the human proteasome. *Chem Biol* 8(12):1131–1141. [https://doi.org/10.1016/s1074-5521\(01\)00080-1](https://doi.org/10.1016/s1074-5521(01)00080-1)
- Rechsteiner M, Realini C, Ustrell V (2000) The proteasome activator 11 S REG (PA28) and class I antigen presentation. *Biochem J* 345:1–15. <https://doi.org/10.1042/0264-6021:345001>
- Whitby FG, Masters EI, Kramer L, Knowlton JR, Yao Y, Wang CC et al (2000) Structural basis for the activation of 20S proteasomes by 11S regulators. *Nature* 408(6808):115–120. <https://doi.org/10.1038/35040607>
- Stadtmueller BM, Hill CP (2011) Proteasome activators. *Mol Cell* 41(1):8–19. <https://doi.org/10.1016/j.molcel.2010.12.020>
- Xie SC, Metcalfe RD, Hanssen E, Yang T, Gillett DL, Leis AP et al (2019) The structure of the PA28-20S proteasome complex from *Plasmodium falciparum* and implications for proteostasis. *Nat Microbiol* 4(11):1990–2000. <https://doi.org/10.1038/s41564-019-0524-4>
- Chen JH, Wang YF, Xu C, Chen KJ, Zhao QY, Wang ST et al (2021) Cryo-EM of mammalian PA28 alpha beta-iCP immunoproteasome reveals a distinct mechanism of proteasome activation by PA28 alpha beta. *Nat Commun*. <https://doi.org/10.1038/s41467-021-21028-3>
- Lesne J, Locard-Paulet M, Parra J, Zivkovic D, Menneteau T, Bousquet MP et al (2020) Conformational maps of human 20S proteasomes reveal PA28-and immunodependent inter-ring cross-talks. *Nat Commun*. <https://doi.org/10.1038/s41467-020-19934-z>
- Yu ZL, Yu YD, Wang F, Myasnikov AG, Coffino P, Cheng YF (2020) Allosteric coupling between alpha-rings of the 20S proteasome. *Nat Commun*. <https://doi.org/10.1038/s41467-020-18415-7>
- Huber EM, Groll M (2017) The mammalian proteasome activator PA28 forms an asymmetric alpha(4)beta(3) complex. *Structure*. 25(10):14730–+. <https://doi.org/10.1016/j.str.2017.07.013>
- Cascio P (2014) PA28 $\alpha\beta$: the enigmatic magic ring of the proteasome? *Biomolecules* 4(2):566–584. <https://doi.org/10.3390/biom4020566>
- Dubiel W, Pratt G, Ferrell K, Rechsteiner M (1992) Purification of an 11-s regulator of the multicatalytic protease. *J Biol Chem* 267(31):22369–22377
- Ma CP, Slaughter CA, Demartino GN (1992) Identification, purification, and characterization of a protein activator (pa28) of the 20-s proteasome (macropain). *J Biol Chem* 267(15):10515–10523
- Kuehn L, Dahlmann B (1996) Proteasome activator PA28 and its interaction with 20 S proteasomes. *Arch Biochem Biophys* 329(1):87–96. <https://doi.org/10.1006/abbi.1996.0195>
- Raule M, Cerruti F, Benaroudj N, Migotti R, Kikuchi J, Bachi A et al (2014) PA28 alpha beta reduces size and increases hydrophilicity of 20S immunoproteasome peptide products. *Chem Biol* 21(4):470–480. <https://doi.org/10.1016/j.chembiol.2014.02.006>
- Li J, Powell SR, Wang XJ (2011) Enhancement of proteasome function by PA28 alpha overexpression protects against oxidative stress. *FASEB J* 25(3):883–893. <https://doi.org/10.1096/fj.10-160895>
- Pickering AM, Linder RA, Zhang HQ, Forman HJ, Davies KJA (2012) Nrf2-dependent induction of proteasome and Pa28 alpha beta regulator are required for adaptation to oxidative stress. *J Biol Chem* 287(13):10021–10031. <https://doi.org/10.1074/jbc.M111.277145>
- Pickering AM, Davies KJA (2012) Differential roles of proteasome and immunoproteasome regulators Pa28 alpha beta, Pa28 gamma and Pa200 in the degradation of oxidized proteins. *Arch Biochem Biophys* 523(2):181–190. <https://doi.org/10.1016/j.abb.2012.04.018>
- Hernebring M, Fredriksson A, Liljevald M, Cvijovic M, Norrman K, Wiseman J et al (2013) Removal of damaged proteins during

- ES cell fate specification requires the proteasome activator PA28. *Sci Rep*. <https://doi.org/10.1038/srep01381>
23. Raynes R, Pomatto L, Davies K (2016) Degradation of oxidized proteins by the proteasome: distinguishing between the 20S, 26S, and immunoproteasome proteolytic pathways. *Mol Aspects Med* 50:41–55. <https://doi.org/10.1016/j.mam.2016.05.001>
 24. Lobanova ES, Finkelstein S, Li J, Travis AM, Hao Y, Klingeborn M et al (2018) Increased proteasomal activity supports photoreceptor survival in inherited retinal degeneration. *Nat Commun*. <https://doi.org/10.1038/s41467-018-04117-8>
 25. Sijts A, Sun YC, Janek K, Kral S, Paschen A, Schadendorf D et al (2002) The role of the proteasome activator PA28 in MHC class I antigen processing. *Mol Immunol* 39(3–4):165–169. [https://doi.org/10.1016/s0161-5890\(02\)00099-8](https://doi.org/10.1016/s0161-5890(02)00099-8)
 26. Cascio P (2011) PA28 gamma: new insights on an ancient proteasome activator. *Biomolecules*. <https://doi.org/10.3390/biom11020228>
 27. Realini C, Jensen CC, Zhang ZG, Johnston SC, Knowlton JR, Hill CP et al (1997) Characterization of recombinant REG alpha, REG beta, and REG gamma proteasome activators. *J Biol Chem* 272(41):25483–25492. <https://doi.org/10.1074/jbc.272.41.25483>
 28. Zhang ZG, Clawson A, Rechsteiner M (1998) The proteasome activator 11 S regulator or PA28—contribution by both alpha and beta subunits to proteasome activation. *J Biol Chem* 273(46):30660–30668. <https://doi.org/10.1074/jbc.273.46.30660>
 29. Wilk S, Chen WE, Magnusson RP (2000) Properties of the beta subunit of the proteasome activator PA28 (11S REG). *Arch Biochem Biophys* 384(1):174–180. <https://doi.org/10.1006/abbi.2000.2112>
 30. Jonik-Nowak B, Menneteau T, Fesquet D, Baldin V, Bonne-Andrea C, Mechali F et al (2018) PIP30/FAM192A is a novel regulator of the nuclear proteasome activator PA28 gamma. *Proc Natl Acad Sci USA* 115(28):E6477–E6486. <https://doi.org/10.1073/pnas.1722299115>
 31. Huang L, Haratake K, Miyahara H, Chiba T (2016) Proteasome activators, PA28 gamma and PA200, play indispensable roles in male fertility. *Sci Rep*. <https://doi.org/10.1038/srep23171>
 32. Mao I, Liu J, Li X, Luo H (2008) REGgamma, a proteasome activator and beyond? *Cell Mol Life Sci* 65(24):3971–3980. <https://doi.org/10.1007/s00018-008-8291-z>
 33. Li XT, Lonard DM, Jung SY, Malovannaya A, Feng G, Qin J et al (2006) The SRC-3/AIB1 coactivator is degraded 14 in a ubiquitin- and ATP-independent manner by the REG gamma proteasome. *Cell* 124(2):381–392. <https://doi.org/10.1016/j.cell.2005.11.037>
 34. Moriishi K, Okabayashi T, Nakai K, Moriya K, Koike K, Murata S et al (2003) Proteasome activator PA28 gamma-dependent nuclear retention and degradation of hepatitis C virus core protein. *J Virol* 77(19):10237–10249. <https://doi.org/10.1128/jvi.77.19.10237-10249.2003>
 35. Ying H, Furuya F, Zhao L, Araki O, West BL, Hanover JA et al (2006) Aberrant accumulation of PTTG1 induced by a mutated thyroid hormone beta receptor inhibits mitotic progression. *J Clin Invest* 116(11):2972–2984. <https://doi.org/10.1172/jci28598>
 36. Li XT, Amazit L, Long W, Lonard DM, Monaco JJ, O'Malley BW (2007) Ubiquitin- and ATP-independent proteolytic turnover of p21 by the REG gamma-proteasome pathway. *Mol Cell* 26(6):831–842. <https://doi.org/10.1016/j.molcel.2007.05.028>
 37. Chen XY, Barton LF, Chi Y, Clurman BE, Roberts JM (2007) Ubiquitin-independent degradation of cell-cycle inhibitors by the REG gamma proteasome. *Mol Cell* 26(6):843–852. <https://doi.org/10.1016/j.molcel.2007.05.022>
 38. Zhang Z, Zhang RW (2008) Proteasome activator PA28 gamma regulates p53 by enhancing its MDM2-mediated degradation. *EMBO J* 27(6):852–864. <https://doi.org/10.1038/emboj.2008.25>
 39. Murata S, Kawahara H, Tohma S, Yamamoto K, Kasahara M, Nabeshima Y et al (1999) Growth retardation in mice lacking the proteasome activator PA28 gamma. *J Biol Chem* 274(53):38211–38215. <https://doi.org/10.1074/jbc.274.53.38211>
 40. Masson P, Lundgren J, Young P (2003) Drosophila proteasome regulator REG gamma: transcriptional activation by DNA replication-related factor DREF and evidence for a role in cell cycle progression. *J Mol Biol* 327(5):1001–1012. [https://doi.org/10.1016/s0022-2836\(03\)00188-8](https://doi.org/10.1016/s0022-2836(03)00188-8)
 41. Barton LF, Runnels HA, Schell TD, Cho YJ, Gibbons R, Tevethia SS et al (2004) Immune defects in 28-kDa proteasome activator gamma-deficient mice. *J Immunol* 172(6):3948–3954. <https://doi.org/10.4049/jimmunol.172.6.3948>
 42. Zannini L, Lecis D, Buscemi G, Carlessi L, Gasparini P, Fontanella E et al (2008) REG gamma proteasome activator is involved in the maintenance of chromosomal stability. *Cell Cycle* 7(4):504–512. <https://doi.org/10.4161/cc.7.4.5355>
 43. Cioce M, Boulon S, Matera AG, Lamond AI (2006) UV-induced fragmentation of Cajal bodies. *J Cell Biol* 175(3):401–413. <https://doi.org/10.1083/jcb.200604099>
 44. Baldin V, Militello M, Thomas Y, Doucet C, Fic W, Boireau S et al (2008) A novel role for PA28 gamma-proteasome in nuclear speckle organization and SR protein trafficking. *Mol Biol Cell* 19(4):1706–1716. <https://doi.org/10.1091/mbc.E07-07-0637>
 45. Zannini L, Buscemi G, Fontanella E, Lisanti S, Delia D (2009) REG gamma/PA28 gamma proteasome activator interacts with PML and Chk2 and affects PML nuclear bodies number. *Cell Cycle* 8(15):2399–2407. <https://doi.org/10.4161/cc.8.15.9084>
 46. Levy-Barda A, Lerenthal Y, Davis AJ, Chung YM, Essers J, Shao ZP et al (2011) Involvement of the nuclear proteasome activator PA28 gamma in the cellular response to DNA double-strand breaks. *Cell Cycle* 10(24):4300–4310. <https://doi.org/10.4161/cc.10.24.18642>
 47. Dong SX, Jia CF, Zhang SP, Fan GJ, Li YB, Shan PP et al (2013) The REG gamma proteasome regulates hepatic lipid metabolism through inhibition of autophagy. *Cell Metab* 18(3):380–391. <https://doi.org/10.1016/j.cmet.2013.08.012>
 48. Chen DB, Yang XS, Huang LY, Chi P (2013) The expression and clinical significance of PA28 gamma in colorectal cancer. *J Investig Med* 61(8):1192–1196
 49. Li L, Dang YY, Zhang JS, Yan WJ, Zhai WL, Chen H et al (2015) REG gamma is critical for skin carcinogenesis by modulating the Wnt/beta-catenin pathway. *Nat Commun*. <https://doi.org/10.1038/ncomms7875>
 50. He J, Cui L, Zeng Y, Wang GQ, Zhou P, Yang YY et al (2012) REG gamma is associated with multiple oncogenic pathways in human cancers. *Bmc Cancer*. <https://doi.org/10.1186/1471-2407-12-75>
 51. Yao LF, Zhou L, Xuan Y, Zhang P, Wang XS, Wang TZ et al (2019) The proteasome activator REG gamma counteracts immunoproteasome expression and autoimmunity. *J Autoimmun*. <https://doi.org/10.1016/j.jaut.2019.05.010>
 52. Boulpicante M, Darrigrand R, Pierson A, Salgues V, Rouillon M, Gaudineau B et al (2020) Tumors escape immunosurveillance by overexpressing the proteasome activator PSME3. *Oncimmunology*. <https://doi.org/10.1080/2162402x.2020.1761205>
 53. Le Feuvre A, Dantas-Barbosa C, Baldin V, Coux O (2009) High yield bacterial expression and purification of active recombinant PA28 alpha beta complex. *Protein Expr Purif* 64(2):219–224. <https://doi.org/10.1016/j.pep.2008.10.014>
 54. Cerruti F, Martano M, Petterino C, Bollo E, Morello E, Bruno R et al (2007) Enhanced expression of interferon-gamma-induced antigen-processing machinery components in a spontaneously occurring cancer. *Neoplasia* 9(11):960–969. <https://doi.org/10.1593/neo.07649>
 55. Raule M, Cerruti F, Cascio P (2014) Enhanced rate of degradation of basic proteins by 26S immunoproteasomes. *BBA-Mol Cell*

- Res 1843(9):1942–1947. <https://doi.org/10.1016/j.bbamcr.2014.05.005>
56. Akopian TN, Kisselev AF, Goldberg AL (1997) Processive degradation of proteins and other catalytic properties of the proteasome from *Thermoplasma acidophilum*. *J Biol Chem* 272(3):1791–1798. <https://doi.org/10.1074/jbc.272.3.1791>
 57. Rappsilber J, Mann M, Ishihama Y (2007) Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat Protoc* 2(8):1896–1906. <https://doi.org/10.1038/nprot.2007.261>
 58. Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu D et al (2019) The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* 47(D1):D442–D450. <https://doi.org/10.1093/nar/gky1106>
 59. Dolenc I, Seemuller E, Baumeister W (1998) Decelerated degradation of short peptides by the 20S proteasome. *FEBS Lett* 434(3):357–361. [https://doi.org/10.1016/s0014-5793\(98\)01010-2](https://doi.org/10.1016/s0014-5793(98)01010-2)
 60. Saric T, Graef CI, Goldberg AL (2004) Pathway for degradation of peptides generated by proteasomes—a key role for thimet oligopeptidase and other metallopeptidases. *J Biol Chem* 279(45):46723–46732. <https://doi.org/10.1074/jbc.M406537200>
 61. Kisselev AF, Akopian TN, Goldberg AL (1998) Range of sizes of peptide products generated during degradation of different proteins by archaeal proteasomes. *J Biol Chem* 273(4):1982–1989. <https://doi.org/10.1074/jbc.273.4.1982>
 62. Kohler A, Cascio P, Leggett DS, Woo KM, Goldberg AL, Finley D (2001) The axial channel of the proteasome core particle is gated by the Rpt2 ATPase and controls both substrate entry and product release. *Mol Cell* 7(6):1143–1152. [https://doi.org/10.1016/s1097-2765\(01\)00274-x](https://doi.org/10.1016/s1097-2765(01)00274-x)
 63. Emmerich NPN, Nussbaum AK, Stevanovic S, Priemer M, Toes REM, Rammensee HG et al (2000) The human 26 S and 20 S proteasomes generate overlapping but different sets of peptide fragments from a model protein substrate. *J Biol Chem* 275(28):21140–21148. <https://doi.org/10.1074/jbc.M000740200>
 64. Kisselev AF, Akopian TN, Woo KM, Goldberg AL (1999) The sizes of peptides generated from protein by mammalian 26 and 20 S proteasomes - Implications for understanding the degradative mechanism and antigen presentation. *J Biol Chem* 274(6):3363–3371. <https://doi.org/10.1074/jbc.274.6.3363>
 65. Zhou PB (2006) REG gamma: a shortcut to destruction. *Cell* 124(2):256–257. <https://doi.org/10.1016/j.cell.2006.01.003>
 66. Cascio P, Hilton C, Kisselev AF, Rock KL, Goldberg AL (2001) 26S proteasomes and immunoproteasomes produce mainly N-extended versions of an antigenic peptide. *EMBO J* 20(10):2357–2366. <https://doi.org/10.1093/emboj/20.10.2357>
 67. Smith G, Seymour L, Boggs J, Harauz G (2012) The 21.5-kDa isoform of myelin basic protein has a non-traditional PY-nuclear-localization signal. *Biochem Biophys Res Commun* 422(4):670–5. <https://doi.org/10.1016/j.bbrc.2012.05.051>
 68. Harauz G, Ladizhansky V, Boggs JM (2009) Structural polymorphism and multifunctionality of myelin basic protein. *Biochemistry* 48(34):8094–8104. <https://doi.org/10.1021/bi901005f>
 69. Groll M, Bajorek M, Kohler A, Moroder L, Rubin DM, Huber R et al (2000) A gated channel into the proteasome core particle. *Nat Struct Biol* 7(11):1062–1067. <https://doi.org/10.1038/80992>
 70. Kudriaeva A, Kuzina ES, Zubenko O, Smirnov IV, Belogurov A (2019) Charge-mediated proteasome targeting. *FASEB J* 33(6):6852–6866. <https://doi.org/10.1096/fj.201802237R>
 71. Sanchez-Puig N, Veprintsev DB, Fersht AR (2005) Human full-length Securin is a natively unfolded protein. *Protein Sci* 14(6):1410–1418. <https://doi.org/10.1110/ps.051368005>
 72. Duvignaud JB, Savard C, Fromentin R, Majeau N, Leclerc D, Gagne SM (2009) Structure and dynamics of the N-terminal half of hepatitis C virus core protein: an intrinsically unstructured protein. *Biochem Biophys Res Commun* 378(1):27–31. <https://doi.org/10.1016/j.bbrc.2008.10.141>
 73. Peng ZL, Mizianty MJ, Kurgan L (2014) Genome-scale prediction of proteins with long intrinsically disordered regions. *Proteins Struct Funct Bioinform* 82(1):145–158. <https://doi.org/10.1002/prot.24348>
 74. Tsvetkov P, Reuven N, Shaul Y (2009) The nanny model for IDPs. *Nat Chem Biol* 5(11):778–781. <https://doi.org/10.1038/nchembio.233>
 75. Fort P, Kajava AV, Delsuc F, Coux O (2015) Evolution of proteasome regulators in eukaryotes. *Genome Biol Evol* 7(5):1363–1379. <https://doi.org/10.1093/gbe/evv068>
 76. Masson P, Lundin D, Soderbom F, Young P (2009) Characterization of a REG/PA28 proteasome activator homolog in *Dictyostelium discoideum* indicates that the ubiquitin- and atp-independent reg gamma proteasome is an ancient nuclear protease. *Eukaryot Cell* 8(6):844–851. <https://doi.org/10.1128/ec.00165-08>
 77. Cascio P, Call M, Petre BM, Walz T, Goldberg AL (2002) Properties of the hybrid form of the 26S proteasome containing both 19S and PA28 complexes. *EMBO J* 21(11):2636–2645. <https://doi.org/10.1093/emboj/21.11.2636>
 78. Groettrup M, Ruppert T, Kuehn L, Seeger M, Standera S, Koszinowski U et al (1995) The interferon-gamma-inducible 11 S regulator (PA28) and the LMP2/LMP7 subunits govern the peptide production by the 20 S proteasome in vitro. *J Biol Chem* 270(40):23808–23815. <https://doi.org/10.1074/jbc.270.40.23808>
 79. Serwold T, Gonzalez F, Kim J, Jacob R, Shastri N (2002) ERAAP customizes peptides for MHC class I molecules in the endoplasmic reticulum. *Nature* 419(6906):480–483. <https://doi.org/10.1038/nature01074>
 80. Saric T, Chang SC, Hattori A, York IA, Markant S, Rock KL et al (2002) An IFN-gamma-induced aminopeptidase in the ER, ERAAP1, trims precursors to MHC class I-presented peptides. *Nat Immunol* 3(12):1169–1176. <https://doi.org/10.1038/ni859>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.