




Carex duanensis (*Carex* sect. *Rhomboidales*), a new species of Cyperaceae from limestone areas of Guangxi, China

Zhao-Cen Lu¹, Yi-Fei Lu², Shi-Li Chang^{1,3}, Ming-Lin Mo^{1,3}, Xiao-Feng Jin²

¹ Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guilin, 541006, Guangxi, China

² School of Forestry and Biotechnology, Zhejiang A&F University, Hangzhou, 311300, Zhejiang, China

³ College of Life Sciences, Guangxi Normal University, Guilin, 541006, Guangxi, China

Corresponding author: Xiao-Feng Jin (docxfjin@163.com)

Abstract

Carex duanensis Z.C.Lu, Y.F.Lu & X.F.Jin, a new species in limestone areas of Guangxi, China, was discovered and described. The morphology showed that *C. duanensis* is similar to *C. calcicola*, but differs in having culms central, leaf blades 3–5.5 mm wide, bracts longer than spikes, utricles 4–5 mm long, shorter, and nutlets abruptly contracted into an erect beak at apex. SEM microphotographs of utricles and nutlets are provided for the new and related species, *C. calcicola*.

Key words: *Carex duanensis*, Cyperaceae, morphology and micromorphology, nutlet, taxonomy, utricle

Introduction

Cyperaceae (sedges), containing 5600+ species in 95 genera worldwide, are the third largest monocot family. It is distributed from tropical to Polar regions, from alpine meadows to tropical rainforests, from Gobi deserts to swampy wetlands, and plays a crucial role in various ecosystems (Larridon et al. 2021). The genus *Carex* L., comprising ca. 2000 species, stands as the largest genus within Cyperaceae and is distributed almost worldwide (Reznicek 1990; Starr and Ford 2009; Roalson et al. 2021). Recent phylogenetic studies have provided a framework for identifying morphologically diagnosable lineages within *Carex*, organizing it into six subgenera: *Carex* subg. *Siderosticta*, *C.* subg. *Psyllophorae*, *C.* subg. *Euthyceras*, *C.* subg. *Uncinia*, *C.* subg. *Vignea*, and *C.* subg. *Carex*. Moreover, the genus was subdivided into 62 formally named Linnean sections plus 49 informal groups (Roalson et al. 2021). *Carex* is one of the largest genera of angiosperms in China, boasting representation of more than 700 species (Chen and Zhang 2018).

Carex section *Rhomboidales* was established by Kükenthal in his worldwide monograph of *Carex* (Kükenthal 1909). Jin and Zheng (2013) revised this section and recognized 40 species, along with six subspecies and four varieties. Since then, some new species of *C.* sect. *Rhomboidales* have been described and published (Chen and Jin 2015; Yang et al. 2015a, 2015b, 2016). Recent phy-



Academic editor: Peter de Lange

Received: 17 February 2024

Accepted: 20 April 2024

Published: 2 May 2024

Citation: Lu Z-C, Lu Y-F, Chang S-L, Mo M-L, Jin X-F (2024) *Carex duanensis* (*Carex* sect. *Rhomboidales*), a new species of Cyperaceae from limestone areas of Guangxi, China. *PhytoKeys* 241: 221–228. <https://doi.org/10.3897/phytokeys.241.121098>

Copyright: © Zhao-Cen Lu et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

logenetic studies revealed that *C.* sect. *Rhomboidales* is monophyletic (Roalson et al. 2021), comprising 50+ species mainly distributed in East and Southeast Asia, with a few species occurring from Europe to western Asia (Kükenthal 1909; Dai et al. 2000, 2010; Jin and Zheng 2013; Roalson et al. 2021).

During the investigation from May 2021 to March 2023, we collected specimens of *Carex* with mature nutlets from limestone evergreen broad-leaved forests and shrubs in Du'an County and Yizhou City, Guangxi, China. After carefully checking the morphological characters of these specimens and comprehensively consulting relevant literature, with comparison of nutlet and utricle micromorphology (Dai et al. 2000, 2010; Jin and Zheng 2013; Jin et al. 2014; Chen and Jin 2015; Yang et al. 2015a, 2015b, 2016; Roalson et al. 2021), we confirmed it as a new species of *C.* sect. *Rhomboidales* and described below.

Materials and methods

Specimens of this new species were collected from Du'an County and Yizhou City, Guangxi, China. After that, we carefully studied relevant literature and the morphological characters of the specimens, which involved measuring and recording the size, shape, and color of rhizomes, culms, leaves, bracts, spikes, glumes, utricles, and nutlets. We examined herbarium specimens at BM, E, HTC, IBK, IBSC, K, KUN, P, PE and ZJFC. The other related species of *Carex* sect. *Rhomboidales* were examined online images from Kew Herbarium Catalogue (<http://apps.kew.org/herbcat/gotoHomePage.do>), JSTOR Global Plants (<https://plants.jstor.org/>) and Chinese Virtual Herbarium (<https://www.cvh.ac.cn/>).

SEM (scanning electron microscope) observations of utricles and nutlets of the new species and the similar species *Carex calcicola* Tang & F.T.Wang in *C.* sect. *Rhomboidales* were carried out. Mature utricles and nutlets were gathered from specimens we collected, the specimens 'W. B. Xu, C. R. Lin & Z. C. Lu 14641' for the new species and 'X. F. Jin & al. 2391' for *C. calcicola* respectively. The utricles were submerged in 50% ethanol to clean for 2 hours, then air dried. The cleaned utricles were mounted on stubs by doubled-sided adhesive tape and coated with gold. The nutlets were initially soaked in a solution of concentrated sulfuric acid and acetic anhydride (volume ratio = 1:9) for 16 hours, then rinsed in acetic acid for 10 min and water for 5 min, and placed in a bath-type ultrasonic cleaner for 30 min with 70% ethanol to remove the cuticle and outer periclinal wall of the epidermis (Jin et al. 2014). After air drying, the nutlets were also mounted on stubs using double-sided adhesive tape, and directly coated with a layer of gold. The coated utricles and nutlets were observed and photographed under a GEMINI-300 scanning electron microscope (SEM).

Taxonomic treatment

***Carex duanensis* Z.C.Lu, Y.F.Lu & X.F.Jin, sp. nov.**

urn:lsid:ipni.org:names:77340986-1

Figs 1A–F, 2A–G

Chinese name: dū ān tái cǎo (都安薹草)

Diagnostic description. This new species is similar to *Carex calcicola* Tang & F.T.Wang, but differs in having culms central (vs. culms lateral), leaf blades

3–5.5 mm wide (vs. 7–15 mm wide), bracts longer than spikes (vs. shorter than spikes or nearly equal in length), utricles 4–5 mm long (vs. 5.5–6.5 mm long), shorter, and nutlets abruptly contracted into an erect beak at apex (vs. curved or coiled beak at apex).

Type. CHINA. Guangxi: Du'an County, National Geopark in Dongmiao Town, 23°59'41"N, 107°59'14"E, limestone slope, alt. 240 m, 5 May 2021, *W. B. Xu, C. R. Lin & Z. C. Lu 14641* (holotype: IBK! barcode IBK00457898; isotypes: IBK! barcode IBK00457899, ZJFC!).

Description. Perennial herbs. Rhizomes short, woody, thick. Culms central, caespitose, 20–50 cm tall, trigonous, slightly scabrous, base with dark brown fibrous sheaths. Leaves longer than culms; blades 3–5.5 mm wide, leathery, margin slightly involuted, scabrous on margins and abaxial leaf surfaces. Bracts leaf-like, longer than spikes, sheathed; sheaths 10–30 mm long. Spikes 3–6; terminal spike staminate, clavate-cylindrical, 3–7 cm long, 2–3 mm wide, base with a 1.5–2.5 cm long peduncle; lateral spikes androgynous, rarely pistillate, cylindrical, 3–5(–8) cm long, 5–7 mm wide, staminate part 1–3 cm long, densely flowered, pistillate part 1.5–4 cm long, sparsely flowered, base with a 1–5 cm long peduncle; peduncles enclosed or slightly exerted from bract sheaths. Staminate glumes oblong-lanceolate, 4–5.5 mm long, yellow-brown, obtuse or acuminate at apex, pale yellow-brown 3-veined dorsal costa. Pistillate glumes (basal part) broadly ovate, ca. 4 mm long, yellow-white, acuminate at apex, yellow 3-veined dorsal costa excurrent into a scabrous awn, middle and upper ones ovate, 3–3.5 mm long, yellow-white, acuminate at apex, yellow 3-veined dorsal costa excurrent into a 0.5–1 mm long scabrous awn. Utricles yellow-green, ovoid (excluding beak), obtusely trigonous, 4–5 mm long (including beak), longer than pistillate glumes, chartaceous, distinctly thinly veined, glabrous, base cuneate, apex gradually contracted into a 2–2.2 mm long beak, orifice 2-lobed with short teeth. Nutlets tightly enveloped, rhombic-ovoid, trigonous, castaneous, 2.5–3 mm long, with 3 angles constricted at middle, sides concave above and below, base with a curved stipe, apex abruptly contracted into a ca. 1 mm long beak, beak erect or slightly curved, annulate at orifice; style thickened at base; stigmas 3.

Etymology. The specific epithet '*duanensis*' refers to the type locality of this new species.

Phenology. Flowering and fruiting mid-March to early May.

Distribution and Habitat. *Carex duanensis* has only been collected from limestone areas of Guangxi, China. It is currently known to grow sporadically in forests or shrubs on limestone slopes, at an elevation of 200–500 m.

Additional specimens examined (paratypes). CHINA. Guangxi: Du'an County, National Geopark in Dongmiao Town, 23°59'41"N, 107°59'14"E, limestone slope, alt. 240 m, 5 May 2021, *W. B. Xu, C. R. Lin & Z. C. Lu 14640* (IBK!, ZJFC!); the same locality, 5 May 2021, *W. B. Xu, C. R. Lin & Z. C. Lu 14642* (IBK!, ZJFC!); Nongshui, Yu'an Village of Dongmiao Town, 23°57'47.37"N, 107°56'39.37"E, in limestone thickets slope, alt. 290 m, 13 March 2023, *W. B. Xu, C. R. Lin, Z. C. Lu & J. Q. Huang 15556* (IBK!); Nongyi, Ditong Village of Dongmiao Town, 23°58'51.52"N, 107°52'06.17"E, in limestone slope, alt. 465 m, 13 March 2023, *W. B. Xu, C. R. Lin, Z. C. Lu & J. Q. Huang 15579* (IBK!); Yizhou City, Latang Village of Anma Town, 24°42'14"N, 108°28'07"E, under limestone forests, alt. 250 m, 1 May 2021, *W. B. Xu, C. R. Lin & Z. C. Lu 14459* (IBK!).




Figure 1. *Carex duanensis* sp. nov. **A** habit **B** lateral spikes **C** utricles **D** nutlets **E** pistillate glumes of middle and upper part **F** staminate glumes.

Conservation status. The new species has been found in three localities in Du'an County and one locality in Yizhou City, Guangxi, China. One locality is in Du'an County, National Geopark in Dongmiao Town, two localities are in the assessment area of Southwest Karst National Park, which is currently being prepared; the populations are in protected areas where they are less threatened. According to the IUCN Red List Categories and Criteria (IUCN 2022), *Carex duanensis* will be considered in the Least Concern (LC) category.

SEM micromorphology. SEM micromorphology uses detailed descriptions which are shown in Figs 1, 2. The utricles of *Carex duanensis* and *C. calcicola* are both obovoid, glabrous, with many longitudinal veins; beak margins prickly-hairy, and short and sharp 2-teethed at the top (Fig. 3), but the utricle length of *C. duanensis* is slightly shorter than those of *C. calcicola*. The nutlet shapes of *C. duanensis* and *C. calcicola* are both rhombic-ovoid, with 3 angles constricted at middle and epidermal cells 5- or 6-gonal, rarely 4-gonal, with straight anticlinal walls and solitary rarely 2 silica bodies. The nutlet beak of *C. duanensis* is erect, whereas that of *C. calcicola* is curved (Fig. 4).




Figure 2. *Carex duanensis* sp. nov. **A** habit **B** staminate glume **C** pistillate glume of basal part **D, E** pistillate glumes of middle and upper part **F** utricle **G** nutlet. (Drawn by Xiao-Feng Jin; based on the holotype: *W.B.Xu et al. 14641* in IBK).




Figure 3. SEM micromorphology of utricles of *Carex duanensis* (A–C) and *C. calcicola* (D–F) A, D overview B, E surface C, F beak . Scale bars: 1 mm (A, D); 200 μ m (B, C, E, F)




Figure 4. SEM micromorphology of nutlets of *Carex duanensis* (A, B) and *C. calcicola* (C, D) A, C overview B, D surface, Scale bar: 1 mm (A, C); 50 μ m (B, D)

Notes. *Carex duanensis* has nutlets rhombic-ovoid, obtusely trigonous, with 3 angles constricted at middle, sides concave above and below, apex abruptly contracted into a ca. 1 mm long beak, beak erect or slightly curved, annulate at orifice. Based on these morphological characters, *C. duanensis* belongs to *C. sect. Rhomboidales* and is similar to *C. calcicola* (Dai et al. 2010), but differs from the latter in having culms central, leaf blades 3–5.5 mm wide, bracts longer than spikes, utricles 4–5 mm long, shorter, and nutlets abruptly contracted into an erect beak at apex. The morphological differences of *C. duanensis* and *C. calcicola* are shown in Table 1.

Table 1. Morphological characters distinguishing *C. duanensis* from *C. calcicola*.

Characters	<i>C. duanensis</i>	<i>C. calcicola</i>
1. Culms	Central	Lateral
2. Leaf blades	3–5.5 mm wide	8–15 mm wide
3. Bracts	Longer than spikes	Shorter than spikes
4. Utricles	4–5 mm long	5.5–6.5 mm long
5. Nutlets	Abruptly contracted into an erect or slightly curved beak at apex	Abruptly contracted into a coiled or curved beak at apex

Acknowledgments

The authors are grateful to Prof. Wei-Bin Xu (IBK), Prof. Chun-Rui Lin (IBK) and Mr. Jin-Quan Huang for assistance during the fieldwork.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by the Guangxi Natural Science Foundation (Grant No. 2021JJB130408 to ZCL), the Fundamental Research Fund of Guangxi Institute of Botany (Grant No. GZY 21007 to ZCL), Guangxi Key Laboratory Construction Project (Grant No. 20-065-7) and the start-up project of Zhejiang A&F University (Grant No. 2022LFR011 to XFJ).

Author contributions

Zhao-Cen Lu, Yi-Fei Lu, and Xiao-Feng Jin conceived the study. Zhao-Cen Lu collected the samples. Zhao-Cen Lu, Yi-Fei Lu, Shi-Li Chang, and Ming-Lin Mo conducted the measurements of morphological characters, Yi-Fei Lu, and Xiao-Feng Jin conducted SEM observation. Zhao-Cen Lu, Yi-Fei Lu, and Xiao-Feng Jin wrote the manuscript. All authors approved the final version of the manuscript.

Author ORCIDs

Zhao-Cen Lu  <https://orcid.org/0000-0002-1725-0247>

Yi-Fei Lu  <https://orcid.org/0000-0001-6095-3821>

Shi-Li Chang  <https://orcid.org/0009-0005-9801-4121>

Ming-Lin Mo  <https://orcid.org/0009-0000-8895-5474>

Xiao-Feng Jin  <https://orcid.org/0000-0002-8195-6738>

Data availability

All of the data that support the findings of this study are available in the main text.

References

Chen WJ, Jin XF (2015) *Carex huangshanica* (sect. *Rhomboidales*, Cyperaceae), a new species from Anhui, eastern China. *Phytotaxa* 233(3): 281–287. <https://doi.org/10.11646/phytotaxa.233.3.5>

- Chen WL, Zhang SR (2018) Species Catalogue of China. Vol. 1 Plants, Spermatophytes (II). Science Press, Beijing, 49–91.
- Dai LK, Liang SY, Tang YC, Li PC (2000) *Carex* In: Dai LK, Liang SY (Eds) Flora Reipublicae Popularis Sinicae. Vol. 12. Science Press, Beijing, 56–528.
- Dai LK, Liang SY, Zhang SR, Tang YC (2010) *Carex* In: Wu ZY, Raven PH, Hong DY (Eds) Flora of China. Vol. 23. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, 285–461.
- IUCN (2022) Guidelines for using the IUCN Red List Categories and Criteria. Version 15.1. Prepared by the Standards and Petitions Committee of the IUCN Species Survival Commission. <https://www.iucnredlist.org/resources/redlistguidelines> [accessed on 1 February 2024]
- Jin XF, Zheng CZ (2013) Taxonomy of *Carex* section *Rhomboidales* (Cyperaceae). Science Press, Beijing, 1–237.
- Jin XF, Zhou YY, Hipp AL, Jin SH, Oda J, Ikeda H, Yano O, Nagamasu H (2014) Achene micromorphology of *Carex* sect. *Rhomboidales* sensu Kükenthal (Cyperaceae) and its systematic implications. Botanical Journal of the Linnean Society 175(1): 123–143. <https://doi.org/10.1111/boj.12149>
- Kükenthal G (1909) Cyperaceae-Caricoideae. In: Engler A (Ed.) Das Pflanzenreich (IV), Vol. 20 (Heft. 38). Verlag von Wilhem Engelmann, Leipzig, 1–767.
- Larridon I, Zuntini AR, Lévillé-Bourret É, Barrett RL, Starr JR, Muasya AM, Villaverde T, Bauters K, Brewer GE, Bruhl JJ, Costa SM, Elliott TL, Epiawalage N, Escudero M, Fairlie I, Goetghebeur P, Hipp AL, Jiménez-Mejías P, Kikuchi IABS, Luceño M, Márquez-Corro JI, Martín-Bravo S, Maurin O, Pokorny L, Roalson EH, Semmouri I, Simpson DA, Spalink D, Thomas WW, Wilson KL, Xanthos M, Forest F, Baker WJ (2021) A new classification of Cyperaceae (Poales) supported by phylogenomic data. Journal of Systematics and Evolution 59(4): 852–895. <https://doi.org/10.1111/jse.12757>
- Reznicek AA (1990) Evolution in sedges (*Carex*, Cyperaceae). Canadian Journal of Botany 68(7): 1409–1432. <https://doi.org/10.1139/b90-180>
- Roalson EH, Jiménez-Mejías P, Hipp AL, Benítez-Benítez C, Bruederle LP, Chung KS, Escudero M, Ford BA, Ford K, Gebauer S, Gehrke B, Hahn M, Hayat MQ, Hoffmann MH, Jin XF, Kim S, Larridon I, Lévillé-Bourret É, Lu YF, Luceño M, Maguilla E, Márquez-Corro JI, Martín-Bravo S, Masaki T, Míguez M, Naczi RFC, Reznicek AA, Spalink D, Starr JR, Uzma, Villaverde T, Waterway MJ, Wilson KL, Zhang S-R (2021) A framework infrageneric classification of *Carex* (Cyperaceae) and its organizing principles. Journal of Systematics and Evolution 59(4): 726–762. <https://doi.org/10.1111/jse.12722>
- Starr JR, Ford BA (2009) Phylogeny and evolution in Cariceae (Cyperaceae): Current knowledge and future directions. Botanical Review 75(1): 110–137. <https://doi.org/10.1007/s12229-008-9020-x>
- Yang HB, Li XX, Bai CJ, Wang WQ, Wang QL, Yu DG, Dong RS, Liu GD (2015a) *Carex procumbens* (*Carex* sect. *Rhomboidales*, Cyperaceae), a new species from Hainan, China. Phytotaxa 201(3): 207–213. <https://doi.org/10.11646/phytotaxa.201.3.4>
- Yang HB, Li XX, Wang WQ, Bai CJ, Liu GD (2015b) *Carex jianfengensis* (*Carex* sect. *Rhomboidales*, Cyperaceae), a New Species from Hainan, China. PLoS ONE 10(9): 1–9. <https://doi.org/10.1371/journal.pone.0136373>
- Yang HB, Li XX, Bai CJ, Wang WQ, Liu GD (2016) *Carex concava* (*Carex* sect. *Rhomboidales*, Cyperaceae), a new species from Hainan, China. Phytotaxa 283(2): 155–162. <https://doi.org/10.11646/phytotaxa.283.2.4>