



Consumption of ultra-processed foods and nutritional profile in a health promotion service of primary care

Suellen F Campos, Luana Caroline dos Santos, Mariana S Lopes, Patrícia P de Freitas and Aline CS Lopes*

Nutrition Department, Nursing School, Universidade Federal de Minas Gerais, Center for Studies in Food and Nutrition in Life Cycles, Research Group on Nutrition Interventions, 190 Alfredo Balena Avenue, Santa Efigênia, Belo Horizonte, MG 30130-100, Brazil

Submitted 16 November 2020: Final revision received 29 April 2021: Accepted 12 May 2021: First published online 26 May 2021

Abstract

Objective: Describing the consumption of ultra-processed foods (UPF) and their association with the nutritional profiles among users of a health promotion service in a Brazilian city.

Design: Cross-sectional study.

Setting: Public health promotion service of Primary Health Care in Belo Horizonte, Minas Gerais, Brazil.

Participants: Totally, 3372 participants.

Results: UPF were found to contribute to 27.7 % of the diet's total energy. The highest consumption was associated with higher values for energy intake (1561.8 *v.* 1331.8 kcal/d; $P < 0.01$), energy density (1.7 *v.* 1.4 kcal/g; $P < 0.01$), total (32.5 *v.* 27.3 %; $P < 0.01$) and trans-fats (2.1 *v.* 1.2 %; $P < 0.01$) and Na (1001.6 *v.* 758.9 mg/1000 kcal; $P < 0.01$) and with lower values for proteins (14.9 *v.* 19.6 %; $P < 0.01$), mono-unsaturated fats (16.1 *v.* 20.1 %; $P = 0.02$), *n*-3 (0.9 *v.* 1.1 %; $P < 0.01$) and some vitamins and minerals when comparing individuals in the last quintile of energy contribution from UPF in relation to the first one. The prevalence rate of nutrient inadequacy aimed at preventing non-communicable diseases increased between 30 % and 100 % when compared with the values of the fifth to the first quintile of UPF consumption ($P < 0.001$). However, the participants had lower energy intake, energy density and Na and higher fibre consumption when compared to Brazilian population.

Conclusions: Participants showed a high consumption of UPF, but also positive diet characteristics when compared with the national data. The results suggest the importance of health promotion services to promote healthy food and the need to include approaches to reduce UPF consumption.

Keywords

Primary Health Care
Ultra-processed foods
Quality of diet
Nutrients

Chronic non-communicable diseases

Ultra-processed foods (UPF) are formulations of ingredients used mainly for industrial purposes and created by industrial techniques. UPF are characterised by their high-energy density; low protein, micronutrients and fibre content and high Na, saturated fat and sugar content^(1,2). This composition is extremely palatable and addictive to humans, which can lead to greater energy consumption and relatively lower protein and micronutrient intake⁽²⁾. Furthermore, given that these foods are widely available and require little to no culinary preparation, they are convenient and increasingly pervasive in the diets of many populations⁽¹⁾.

In developed countries, studies of household samples indicate that UPF account for 57.5 %, 56.8 % and 48 % of

energies in the typical diets of individuals in the USA, UK and Canada, respectively^(3–5). In Latin America, 24 % of the energies in the diet came from UPF in Chile⁽⁶⁾, whereas this value was 21.5 % in Brazil⁽⁷⁾. In all studies, the added energies in UPF were negatively correlated to some vitamin and mineral content and positively correlated to the Na and fat content of the diet in question. Moreover, excessive consumption of UPF is associated with obesity^(8,9), arterial hypertension⁽¹⁰⁾, metabolic syndrome⁽¹¹⁾, cancer⁽¹²⁾, heart and cerebrovascular diseases⁽¹³⁾, depression^(14,15) and increased risk of morbidity⁽¹⁶⁾ and mortality^(17–19). These findings highlight the need to intervene on the excessive consumption of UPF and to encourage the consumption

*Corresponding author: Email alinelopesenf@gmail.com



of minimally processed foods aimed at promoting health and preventing and controlling chronic non-communicable diseases (NCD).

In light of these facts, Brazil published its Dietary Guidelines for the Brazilian Population in 2014, which has as its golden rule the following guideline: 'Always prefer fresh or minimally processed foods and culinary preparations to ultra-processed foods'. These guidelines are incorporated into national nutrition policy^(20,21), and it is used as a reference for building policies on food and nutrition, for directing food and nutrition educational activities and guiding the production of materials to the practice of professionals that work in health promotion services of Brazil's Unified Health System (SUS)⁽²²⁻²⁴⁾, which is coordinated by the Primary Health Care (PHC).

The PHC's health professionals develop recommendations of health promotion, disease prevention and rehabilitation based on the principles of universality and equity⁽²⁵⁾. An important initiative at PHC was the creation of the Health Academy Program (Programa Academia da Saúde (PAS)) in 2011, whose objective is to advance general health promotion and prevent NCD. This programme offers free health promotion activities to vulnerable populations with limited access to health promotion services or facilities⁽²⁶⁾. Among the activities carried out are those related to food and nutrition, which address various topics defined according to local demand and Brazilian Dietary guidelines. Currently, PAS has reached 2900 Brazilian municipalities spanning all units of the Federal District⁽²⁷⁾.

Despite these and other efforts, the rate of NCD continues to grow. Studies investigating UPF consumption remain scarce, since the classification of food by its degree of industrial processing (NOVA) was only recently introduced. It is unknown whether regular users of public health promotion services have a different nutritional profile with respect to UPF consumption relative to the general population, and, further, what the relationship might be between the consumption of these foods and overall nutritional profile. Thus, this study aimed to describe the consumption of UPF and analyse its association with the nutritional profile of health promotion service users of PHC in a Brazilian city.

Methods

Data source and sampling

This cross-sectional study of baseline data from a controlled and randomised community trial aimed to assess the impact of intervention to encourage the consumption of fruits and vegetables in users of PAS in the city of Belo Horizonte in Minas Gerais, Brazil⁽²⁸⁾. This city has an estimated population of 2 512 070 inhabitants distributed across nine administrative regions, making it the sixth largest city in the country⁽²⁹⁾.

Sampling of the PAS units was stratified by the nine administrative regions of the municipality⁽²⁸⁾. At the time

of the sampling process (2012), 50 PAS units were in operation in these nine administrative regions, including those with morning operation, which were located in an area of medium and high to very high vulnerability to health problems, due to greater availability of PAS in this municipality. The areas were classified by the Health Vulnerability Index, which is composed of socio-economic and environmental variables and classifies the risk of vulnerability as low risk, medium risk, high risk or very high risk⁽³⁰⁾, and including the units that had not participated in a nutritional intervention study in the past 2 years. Of the forty-two eligible units, two were drawn by municipal region, resulting in 18 (42.8%) total units, two in each administrative region of the municipality. This sample was representative of services in the municipality with medium and high or very high Health Vulnerability Index, with a 95% CI and an error rate of 1.4%⁽²⁸⁾.

In the sampled units, all individuals aged ≥ 20 years who participated in at least one activity in the most recent month were interviewed. The exclusion criteria were being pregnant and having a cognitive impairment that precluded completion of the questionnaire. Of the 3763 eligible individuals, 112 (3%) failed to meet the inclusion criteria and were excluded and 237 (6.3%) declined to complete the questionnaire. The baseline sample comprised 3414 participants⁽²⁸⁾. Additional details about the study's sampling process are described in a previous publication⁽²⁸⁾.

As recommended by Willet⁽³¹⁾, our study's analyses excluded individuals who reported, according to the 24-h dietary recall (R24 h) interview, those who reported a very low (<500 kcal/d) or a very high (> 7000 kcal/d) total daily energy consumption (n 37) and those who returned incomplete questionnaires (n 5). Therefore, the final sample comprised 3372 individuals, accounting for a participation rate of 98.8% relative to the baseline of the community trial.

Data collection

Data were collected between February 2013 and June 2014 via in-person interviews conducted by trained interviewers who were supervised by nutritionists⁽²⁸⁾. Training for data collection occurred every 6 months, and a field manual was drafted for consultation⁽²⁸⁾.

Dietary intake

The R24 h was used because it is indicated for individuals who have a low level of education and for epidemiological studies with large samples^(32,33). To minimise biases, the interviewees were not informed about their R24 h completion date.

The R24 h was administered on two non-consecutive days (one corresponding to a weekend day), and the data were tabulated and processed in the Brazil Nutri software program. The self-reported measurements of food preparations were transformed into grams or millilitres using



food consumption assessment tables and manuals^(34–36) and industrialised food labels and measurements (standardising/weighing). The standardised tabulation and processing of food consumption data were performed by trained and supervised nutritionists.

For the inclusion of food in the Brazil Nutri program's database, the content of sugar added in beverages was considered in cases where the individual reported adding only sugar (10 % of the volume) or adding both sugar and artificial sweeteners (5 % of the volume) (7:35). The inclusion of salt and oil to the culinary preparations was considered, as well as the amount of table salt added to the food. Data from individuals who completed at least one R24 h form (n 196; 5.6 %) were analysed. For the other participants who answered both R24 h forms, the data were averaged.

Classification of foods according to processing

Foods were assigned to one of three groups according to the NOVA food classification system: (1) unprocessed (fresh) or minimally processed foods, including culinary preparations based on one or more natural or minimally processed foods, and processed culinary ingredients; (2) processed foods and (3) UPF. UPF are industrial formulations composed mostly or entirely of substances extracted from food, derived from food constituents, or synthesised in the laboratory based on organic materials (e.g. dyes, flavourings and flavour enhancers)⁽²⁾.

Consumption of UPF and nutritional dietary profile

The total energetic consumption of UPF was expressed as a percentage of the total energetic value of the diet according to self-reported data from R24 h. To assess the nutritional profile, quintiles of the percentage-based contribution of energy from UPF consumption were assessed, with the first quintile referring to the lowest consumption of UPF and the last to the highest consumption.

The macronutrients investigated were carbohydrates, proteins, total fats and fat subtypes: saturated, monounsaturated, polyunsaturated, trans, linolenic (n -3) and linoleic (n -6) fatty acids. They were evaluated according to their percentage-based energetic contribution to the total energetic value of the diet.

The micronutrients investigated were vitamins A, B₃, B₆, B₁₂, folate and C and the minerals Ca, Fe, P, Mg, K and Na. To present these micronutrients and fibre, the measure of nutrient density was used; that is, the amount of nutrient or fibre consumed per every 1000 kcal (g, mg, or μ g/1000 kcal). The energy density of the diet (kcal/g) was calculated by dividing the sum of the energies from the intake of solid foods by their quantity in grams. The recommendations used for this indicator were based on those proposed by the World Cancer Research Foundation⁽³⁷⁾.

Study variables

The socio-demographic variables investigated were age (adult: 20–59 years; elderly: \geq 60 years), sex (female or male), marital status (married or stable union; separated or divorced; single or widowed), education ($<$ 8 and $>$ 8 years of study) and per capita income (categorised in tertiles). To calculate per capita income, the sum of the income of all family members was divided by the number of residents.

The main independent variable was the percentage-based contribution of energy from UPF to the total energetic value of the diet. The dependent variables were total energies (kcal); energy density (kcal/g); energetic distribution (%) of macronutrients and density of fibre, vitamins and minerals (μ g, mg or g/1000 kcal).

To estimate the prevalence of non-recommended intake levels of nutrients (total fat, saturated fat, trans-fat, fibre, K and Na), the recommended levels for the prevention of NCD were adopted^(38–40). They were considered inadequate if their consumption patterns were characterised by \geq 30 % of total energy intake for total fat; \geq 10 % of total energy intake for saturated fats; \geq 1 % of total energy intake for trans fats; \leq 12.5 g/1000 kcal for dietary fibre; \geq 1000 mg/1000 kcal for Na and \leq 1755 mg/1000 kcal for K. The Estimated Average Requirement was used to estimate the prevalence of vitamin and mineral deficiency, as recommended by the Dietary Reference Intakes^(41–44).

Statistical analysis

The contribution of energy from UPF to the diet and its distribution in quintiles were presented according to socio-demographic variables (gender, age, marital status, education and per capita income in tertiles). The consumption of the macro- and micronutrients were presented according to the quintiles of percentage contribution of energy from UPF in the diet and stratified by sex.

Histograms were generated to assess the normality of the variables. The variables that presented a distorted distribution (vitamin A, B₁₂, C, trans fats and n -3) underwent a logarithmic transformation before statistical analysis; however, the original values are presented in the corresponding Tables. Then, parametric tests were chosen for all variables based on their greater statistical power and result accuracy. To assess the direction of the correlations, ANOVA and Scheffé's post hoc tests were used. Trend tests were also performed for socio-demographic and food consumption variables according to quintiles of energy contribution from UPF.

For the analysis of the diet's nutritional profile (macro- and micronutrients), linear regression models were applied to each dependent variable. Unadjusted and adjusted models for age, marital status, education and sex were performed, and standardised regression coefficients (SD) were calculated.

The prevalence of inadequate nutrient intake was presented according to the quintiles of the percentage-based

contribution of energy from UPF in the diet, which was stratified by sex. Trend tests were performed for food consumption inadequacy according to quintiles of energy contribution from UPF, and the difference in the prevalence of inadequacy between the first and last quintiles was assessed using the χ^2 test with Bonferroni correction ($P < 0.005$). To measure the relationship between the prevalence of inadequate nutrient intake and the quintiles of the contribution of energy from UPF in the diet, the prevalence ratio was calculated using Poisson regression. The models were adjusted for sex, age, marital status and education. All analyses were performed with Data Analysis and Statistical Software (STATA) v. 14.0, with a significance level of 5% ($P < 0.05$).

Results

Among the 3372 individuals who participated in the study. About 88.1% were female and 43.6% were elderly. Participants reported an average daily energy consumption of 1435.4 kcal/d (± 559 kcal), with 27.7% coming from UPF. The percentage of UPF consumption was higher among women (28.2% *v.* 23.9% in men), adults (29.3% *v.* 25.6% in the elderly) and those with a higher level of education (29.7% *v.* 25.6% with a lower level of education) and income (28.9% *v.* 27.2% in second tertile *v.* 27.1% in first tertile) (Table 1).

Individuals of either sex in the quintile of highest consumption of UPF had diets with a higher energy density, greater consumption of energy, total, saturated, mono-unsaturated and trans-fats and lower protein intake relative to individuals in the first quintile. Furthermore, a smaller proportion of *n-3* fatty acids was observed in women in the highest quintile of consumption (Table 2).

For both sexes, linear regression analysis showed a direct relationship between the contribution of energy from UPF and the total diet energy, energy density and the consumption of total, saturated, monounsaturated and trans fats; an inverse relationship was found between the proportion of protein and the proportion of *n-3* fatty acids. For women, a direct relationship was also observed between high UPF consumption and energies coming from polyunsaturated fat; for men, an inverse relationship was found between high UPF consumption and energies coming from carbohydrates (Table 2).

For all participants who reported a higher consumption of UPF, we found a lower intake of fibre, Fe, P, Mg and K and a higher intake of Na relative to participants in the first quintile. After the adjustments, an inverse relationship was observed between the coefficients of the contribution of energy from UPF and the consumption of fibre, Fe, P, Mg and K, and a direct relationship was observed to Na consumption for both sexes. An inverse association

Table 1 Average and distribution in quintiles of the percentage of energy from the consumption of ultra-processed foods (UPF) in the total energy of the diet, according to the socio-demographic characteristics of those attending a Brazilian health promotion service. Belo Horizonte, Minas Gerais, Brazil (2013–2014)

Variables	n	%	% UPF consumption	Percent quintiles of UPF energetic contribution/total dietary energy*					P value†	
				1	2	3	4	5		
Sex‡stratified by age range										
Women										
Adults (20–59 years)	1745	51.8	29.4	15.7	18.6	21.4	21.9	22.4		
Elderly (≥ 60 years)	1224	42.2	26.4	22.4	21.9	18.9	18.6	18.1		
Total	2969	88.1	28.2	18.5	19.9	20.4	20.5	20.6		<0.001
Men										
Adults (20–59 years)	158	39.2	27.5	24.7	16.5	17.7	20.2	20.9		
Elderly (≥ 60 years)	245	60.8	21.5	35.5	22.9	16.7	13.1	11.8		
Total	403	11.9	23.9	31.3	20.3	17.1	15.9	15.4		
Marital status‡										
Married/stable relationship	2078	61.7	27.5	20.6	19.9	19.6	20.5	19.3		
Separated/divorced	280	8.3	28.8	19.3	17.9	18.2	19.3	25.4		0.89
Not married	477	14.15	29.3	19.1	17.2	18.7	21.8	23.3		
Widower	536	15.9	26.3	18.9	23.9	23.7	16.6	16.9		
Age range‡										
Adults (20–59 years)	1903	56.4	29.3	16.4	18.4	21.1	21.8	22.3		<0.001
Elderly (≥ 60 years)	1469	43.6	25.6	24.6	22.1	18.6	17.7	17.0		
Education‡										
< 8 years	1686	50	25.6	24.4	21.3	19.3	18.6	16.4		<0.001
≥ 8 years	1686	50	29.7	15.6	18.7	20.7	21.4	23.6		
Per capita income (tertiles)‡										
1 (US\$0–US\$205.8)	1068	34.5	27.1	21.7	20.3	20.1	20.6	17.3		
2 (US\$208.2–US\$370.4)	1006	32.7	27.2	21.4	19.4	19.5	19.8	19.9		<0.001
3 (US\$372.02–US\$6790.1)	1003	32.6	28.9	15.7	19.8	22.1	20.4	22.0		

*Percentage of total energy consumption originated from UPF: average (amplitude): Q1 = 8.4% (0 to 14.4); Q2 = 18.5% (14.4 to 22.6); Q3 = 26.4% (22.6 to 30.4); Q4 = 34.9% (30.4 to 40.6); Q5 = 50.2% (40.6 to 85.7).

†P value <0.05 for linear trend test.

‡They presented a significant difference with percentage of UPF consumption.

Table 2 Averages of energy consumption and macronutrients according to the quintiles of percentage of energy derived from the consumption of ultra-processed foods (UPF) from regulars in a Brazilian health promotion service. Belo Horizonte, Minas Gerais, Brazil (2013–2014)

Variables	Sex	Average	SD	Percent quintiles of UPF energetic contribution/total dietary energy†,*										Standardised regression coefficient§	
				1	SD	2	SD	3	SD	4	SD	5	SD	Crude	Adjusted‡‡
Total energy (kcal/d)	Women†	1397.5	530.8	1283.1 ^a	554.1	1362.1 ^{a,b}	520.7	1384.6 ^{a,b}	470.1	1427.8 ^{b,c}	512.9	1517.1 ^c	566.6	84.32**	71.31**
	Men†	1714.7	663.2	1544.1 ^a	586.9	1671.4 ^{a,b}	661.4	1754.1 ^{a,b}	566.5	1784.5 ^{a,b}	620.3	2002.9 ^b	836.8	162.34**	136.75**
	Total†	1435.4	557.7	1331.8 ^a	569.2	1399.7 ^{a,b}	550.5	1422.3 ^{a,b}	490.3	1461.7 ^b	533.9	1561.8 ^c	612.2	84.08**	80.09**
Energy density (kcal/g)	Women†	1.5	0.4	1.4 ^a	0.3	1.4 ^{a,b}	0.3	1.4 ^{a,b}	0.3	1.5 ^b	0.4	1.7 ^c	0.4	0.14**	0.13**
	Men†	1.5	0.4	1.4 ^a	0.3	1.5 ^a	0.3	1.5 ^a	0.3	1.6 ^a	0.3	1.9 ^b	0.4	0.17**	0.17**
	Total†	1.5	0.4	1.4 ^a	0.3	1.4 ^a	0.3	1.5 ^{a,b}	0.3	1.5 ^b	0.4	1.7 ^c	0.4	0.14**	0.14**
Proteins (% VCT)	Women†	17.4	4.9	19.5 ^a	5.4	18.3 ^b	5.2	17.6 ^b	4.5	16.7 ^c	4.1	14.9 ^d	3.8	-1.57**	-1.58**
	Men†	18.0	4.7	20.0 ^a	5.0	18.5 ^{a,b,c}	4.6	17.8 ^{b,c}	3.9	16.7 ^c	3.9	14.9 ^{c,d}	3.4	-1.81**	-1.84**
	Total†	17.4	4.8	19.6 ^a	5.4	18.4 ^b	5.1	17.6 ^c	4.5	16.7 ^d	4.1	14.9 ^e	3.8	-1.61**	-1.62**
Carbohydrates (% VCT)	Women†	53.4	9.1	54.0 ^a	11.1	53.9 ^a	9.7	54.1 ^a	8.5	53.6 ^a	8.2	53.2 ^a	8.0	-0.36**	-0.29
	Men†	52.3	10.3	54.2 ^a	10.7	52.4 ^a	10.9	51.5 ^a	9.4	51.6 ^a	9.3	49.9 ^a	10.4	-1.36**	-1.35**
	Total†	53.6	9.3	54.1 ^a	11.0	53.7 ^a	9.9	53.8 ^a	8.6	53.4 ^a	8.3	52.9 ^a	8.3	-0.43**	-0.43**
Total fat (% VCT)	Women†	29.8	7.3	27.5 ^a	7.9	28.8 ^b	7.1	29.6 ^b	6.6	30.6 ^c	6.7	32.4 ^d	7.0	1.74**	1.99**
	Men†	29.3	7.9	26.5 ^a	7.7	28.8 ^{a,b}	7.5	29.8 ^{a,b}	7.3	31.6 ^b	7.2	32.9 ^{b,c}	8.2	2.37**	2.26**
	Total†	29.7	7.3	27.3 ^a	7.9	28.8 ^b	7.1	29.8 ^b	6.6	37.0 ^c	6.7	32.5 ^d	7.2	1.83**	1.77**
Saturated fat (% VCT)	Women†	9.7	3.4	8.8 ^a	3.4	9.4 ^a	3.4	9.4 ^a	2.9	10.1 ^b	3.4	10.6 ^b	3.6	0.64**	0.63**
	Men†	9.7	3.8	8.7 ^a	3.4	9.5 ^{a,b}	3.3	10.1 ^{a,b}	4.9	10.3 ^{a,b}	3.3	11.0 ^b	3.6	0.89**	0.82**
	Total†	9.7	3.5	8.8 ^a	3.4	9.4 ^{a,b}	3.4	9.5 ^b	3.1	10.1 ^c	3.4	10.7 ^c	3.7	0.67**	0.65**
Monounsaturated fat (% VCT)	Women†	10.0	3.5	9.3 ^a	3.7	9.7	3.3	9.9 ^{a,b}	3.1	10.4 ^{b,d}	3.3	10.8 ^{c,d}	3.9	0.54**	0.53**
	Men†	10.1	3.8	8.9 ^a	3.3	9.9 ^{a,b}	3.5	10.3 ^{a,b}	3.2	10.9 ^b	3.7	11.6 ^b	4.6	0.90**	0.88**
	Total†	10.0	3.5	9.2 ^a	3.7	9.8 ^{a,b}	3.4	9.9 ^{b,d}	3.1	10.4 ^{c,d}	3.3	10.9 ^{c,e}	3.9	0.58**	0.58**
Polyunsaturated fat (% VCT)	Women†	6.2	2.2	6.1	2.4	6.1 ^a	2.2	6.4 ^a	2.1	6.2 ^a	2.0	6.4 ^a	2.3	0.14**	0.13**
	Men	6.1	2.0	5.9 ^a	2.0	5.9 ^a	2.1	6.2 ^a	2.1	6.3 ^a	1.9	6.2 ^a	2.1	0.12	0.12
	Total†	6.2	2.2	6.1 ^a	2.3	6.1 ^a	2.2	6.3 ^a	2.1	6.2 ^a	2.0	6.4 ^a	2.3	0.14**	0.12**
Trans fat (% VCT)	Women†	1.7	1.4	1.2 ^a	0.9	1.5 ^b	1.2	1.7 ^{b,c}	1.2	1.9 ^{c,d}	1.4	2.1 ^d	1.8	0.32**	0.32**
	Men†	1.7	1.4	1.2 ^a	0.9	1.6 ^{a,b,c}	1.2	2.1 ^{b,c}	1.6	2.1 ^{b,c}	1.7	2.2 ^c	1.8	0.38**	0.37**
	Total†	1.7	1.4	1.2 ^a	0.9	1.5 ^b	1.2	1.8 ^{b,c}	1.3	1.9 ^{c,d}	1.4	2.1 ^d	1.8	0.33**	0.33**
n 6 (% VCT)	Women	5.3	2.1	5.4 ^a	2.5	5.3 ^a	2.1	5.5 ^a	2.0	5.3 ^a	1.8	5.3 ^a	1.9	-0.01	-0.02
	Men	5.3	1.9	5.3 ^a	2.1	5.2 ^a	2.1	5.4 ^a	1.8	5.5	1.7	5.3 ^a	1.8	0.01	0.02
	Total†	5.3	2.1	5.4 ^a	2.5	5.3 ^a	2.1	5.5 ^a	1.9	5.3 ^a	1.8	5.3 ^a	1.9	-0.01	-0.02
n 3 (% VCT)	Women	0.9	0.9	1.0 ^a	1.2	0.9 ^{a,b}	0.9	0.9 ^{a,b}	0.8	0.9 ^b	0.6	0.9 ^b	0.8	-0.06**	-0.06**
	Men	0.9	0.9	1.0 ^a	0.9	0.9 ^a	0.8	0.7 ^a	0.6	0.9 ^a	0.9	0.7 ^a	0.7	-0.09**	-0.11**
	Total	0.9	0.9	1.1 ^a	1.2	0.9 ^{a,b}	0.9	0.9 ^{a,b}	0.7	0.9 ^b	0.7	0.9 ^b	0.7	-0.06**	-0.07**

VCT, total energetic value of the diet.

*Values with the same letter do not differ using the ANOVA and Scheffe test at 5% of probability.

†P value <0.05 for linear trend test of food consumption variables according to ultra-processed food consumption quintiles.

‡Percentage of total energy consumption of ultra-processed foods: average (amplitude): Q1 = 8.4% (0–14.4); Q2 = 18.5% (14.4–22.6); Q3 = 26.4% (22.6–30.4); Q4 = 34.9% (30.4–40.6); Q5 = 50.2% (40.6–85.7).

§Linear regression of dietary nutrients in quintiles of% energy from consumption of ultra-processed foods, expressed in units of sD.

||Recommendation: difference between% total fats and other fats (polyunsaturated, trans, linoleic and linolenic fatty acids), with a range of 1.21 to 31.93% in the sample.

**P value <0.05.

‡‡Adjusted by sex, age, years of study and marital status.



between a higher intake of UPF and Ca was also identified for women (Table 3).

Women in the quintile of highest consumption of UPF were found to consume a smaller amount of all vitamins, except B₁₂. On the other hand, men had a lower consumption of only vitamin B₆. For both sexes, an inverse relationship was found between the coefficients of the contribution of energy from UPF and the consumption of vitamins B₆, C and folate; and the consumption of vitamins A, B₃ and B₁₂ for women (Table 4).

The prevalence of nutrient inadequacy associated with a higher risk of developing NCD was higher in the quintile of highest consumption of UPF relative to the first quintile, except for Na among men. When the prevalence of inadequate consumption was analysed, and considering the first quintile of percentage contribution of energy from UPF as a reference, a 100% increase was found in inadequate Na intake, an 80% increase in inadequate total fat intake, a 70% increase in inadequate fibre and saturated fat intake, a 60% increase in inadequate trans-fat intake and a 30% increase in inadequate K intake among individuals in the last quintile. Among men, no significant increase was found in the prevalence of inadequate consumption of K, considering the quintiles of contribution of energy from UPF (Table 5).

The prevalence of inadequate micronutrient intake was > 50% for vitamins A, B₃, and folate and for the minerals Ca and Mg. The higher consumption of UPF from the fifth to the first quintile was associated with a higher prevalence of inadequate intake of vitamin B₆ among women and vitamins A and C among both sexes and a lower prevalence of inadequate Ca intake among women and vitamin B₃ among men. The inadequate intake prevalence ratios were significantly higher in the last quintile of UPF consumption for vitamins B₆ and C among females, reaching an increase of 64% and 25%, respectively, when compared with the first quintile (Table 6).

Discussion

The consumption of UPF among the participants in a health-promotion service of Brazilian Primary Care was high, especially among women, adults and those with a higher income or education. The higher consumption of UPF was associated with higher energy density and energy consumption, higher content of total and trans fats and lower content of fibre, Fe, P, Mg, K and vitamin B₆. For women in particular, the higher consumption of UPF was associated with lower consumption of monounsaturated and *n*-3 fats and of vitamins A, B₃, B₁₂, folate and C and higher Na content. The prevalence ratios of inadequate nutrient intake associated with a higher risk of NCD ranged from 30% to 100% for the total sample when the highest and first quintile of UPF consumption are compared.

A high consumption of UPF was found that reached 50.2% (40.6–85.7%) of the total energetic intake in the last quintile, even in users of a health-promotion service⁽⁴⁵⁾. This consumption was even higher than that observed in the Brazilian population (27.7% *v.* 21.5%)⁽⁷⁾, but lower than that found in countries such as the USA (57.5%)⁽³⁾ and the UK (56.8%)⁽⁴⁾ and in a study of young adults from a Brazilian city (51.2%)⁽⁴⁶⁾. These discrepancies between national data may be related to differences in the age groups investigated.

Similar to the results found in our study, in general, past studies have broadly found that the consumption of UPF is higher in women⁽⁴⁶⁾, adults^(5,47) and those with higher educational attainment^(46–48) and income^(46–48). The developed countries, which have greater income equality and less inequality in its distribution, also present with higher consumption of UPF (42% in Australia; 47.7% in Canada; 56.8% in the UK and 57.5% in the USA)^(3–5,49) compared with developing countries (29.8% in Mexico; 28.6% in Chile; 21.5% in Brazil and 15.9% in Colombia)^(7,47,50,51). A study carried out in the same city as the present study showed that the energetic contribution of UPF is higher among individuals of high socio-economic status⁽⁴⁸⁾.

The majority of PHC service users in Brazil present an economic classification from medium (54.3%) to low (38%), with 32% receiving some type of financial government aid⁽⁵²⁾. Although some studies^(46,48) have shown that lower income is associated with lower consumption of UPF, a high consumption of UPF is nevertheless found in users of PHC services residing in vulnerable areas. This result may be related to social inequalities such as limited access to establishments that sell healthy food and a high supply of readily accessible UPF⁽⁵³⁾. In addition, marketing and advertisements for UPF are pervasive in Brazil, which is also home to an ineffective food labelling system that may not be easily understood by a wide range of populations⁽⁵⁴⁾.

Similar to results from other studies^(3–5,7,46,47,49–51,55,56), we found that a high consumption of UPF is associated with a poorer diet quality, which can compromise metabolic processes and which constitutes an important risk factor for NCD^(8–13). Corroborating these results, we identified a high prevalence of inadequate consumption of nutrients (total, saturated and trans fats; fibre; K and Na) according to the references for the prevention of NCD, with an upward progression proportional to the contribution of energy derived from UPF. The prevalence of inadequate intake of vitamins B₆ and C among women also has become worse, with the highest contribution of energy derived from UPF, as analysed according to Dietary Reference Intakes guidelines.

Compared to the results of a UK-based study⁽⁴⁾, the prevalence of nutrient inadequacy in our study was lower for saturated fats (43.6% *v.* 74.45%), fibre density (58.8% *v.* 84.02%) and Na (29.4% *v.* 74.92%) and higher for K (85.4% *v.* 74.76%). In relation to an Australian-based study⁽⁵⁷⁾, the prevalence of inadequate intake in our

Table 3 Average consumption of fibre and minerals according to the quintiles of percentage of energy derived from the consumption of ultra-processed foods (UPF) from regulars in a Brazilian health promotion service. Belo Horizonte, Minas Gerais, Brazil (2013–2014)

Nutrients	Sex	Average	SD	Percent quintiles of UPF energetic contribution/total dietary energy†,*										Standardised regression coefficient§	
				1	SD	2	SD	3	SD	4	SD	5	SD	Crude	Adjusted
Fibre (g/1000 kcal)	Women†	12.2	4.4	13.6 ^a	4.9	12.8 ^b	4.3	12.4 ^{b,c}	4.0	11.8 ^c	4.1	10.5 ^d	3.9	-1.04**	-1.03**
	Men†	11.7	4.3	13.5 ^a	4.5	11.9 ^{a,b}	4.2	11.2 ^{a,b}	3.7	10.9 ^b	3.6	8.8 ^c	3.2	-1.60**	-1.59**
	Total†	12.2	4.4	13.6 ^a	4.8	12.8 ^b	4.3	12.4 ^{b,c}	4.0	11.8 ^c	4.1	10.4 ^d	3.9	-1.08**	-1.10**
Minerals															
Ca (mg/1000 kcal)	Women	372.0	173.5	358.3 ^a	170.2	375.5 ^{a,b}	173.6	392.3 ^{b,c}	176.8	378.9 ^{a,b}	178.1	354.1 ^a	165.7	-13.86	-16.71**
	Men	331.2	156.1	337.9 ^a	165.6	342 ^a	160.3	322.5 ^a	136.9	348.9 ^a	176.6	294.7 ^a	120.5	-4.02	-3.83
	Total	367.2	171.9	354.5 ^a	169.3	371.4 ^{a,b}	172.5	385.2 ^b	174.4	376.0 ^{a,b}	177.9	348.6 ^a	163.0	-4.00	-5.32
Fe (mg/1000 kcal)	Women†	5.1	1.5	5.4 ^a	1.6	5.3 ^a	1.5	5.1 ^a	1.4	5.1 ^a	1.5	4.7 ^b	1.5	-0.44**	-0.42**
	Men†	5.4	1.6	5.9 ^a	1.8	5.3 ^{a,b}	1.6	5.2 ^{a,b}	1.2	4.9 ^b	1.2	4.8 ^b	1.4	-0.22**	-0.22**
	Total†	5.1	1.5	5.5 ^a	1.6	5.3 ^{a,b}	1.5	5.1 ^b	1.4	5.1 ^b	1.5	4.7 ^c	1.5	-0.26**	-0.25**
P (mg/1000 kcal)	Women†	574.4	168.9	605.9 ^a	175.6	592.1 ^{a,b}	172.8	598.2 ^a	166.8	566.2 ^b	160.8	513.5 ^c	153.1	-33.32**	-33.63**
	Men†	563.9	167.3	592.3 ^a	172.1	594.8 ^a	178.7	558.7 ^{a,b}	140.7	560.4 ^a	172.6	474.4 ^b	130.3	-40.79**	-42.11**
	Total†	573.2	168.8	603.6 ^a	174.8	591.6 ^a	173.5	595.7 ^a	164.7	564.9 ^b	161.8	510.2 ^c	151.4	-33.78**	-34.81**
Mg (mg/1000 kcal)	Women†	131.8	39.5	141.4 ^a	39.0	137.7 ^a	39.6	135.9 ^a	36.9	129.0 ^b	38.8	116.5 ^c	38.5	-8.92**	-8.88**
	Men†	127.2	37.1	140.6 ^a	37.4	133.1 ^{a,b}	37.7	121.3 ^{b,c}	30.2	124.4 ^{a,c}	35.0	101.7 ^d	30.9	-13.36**	-12.99**
	Total†	131.5	39.3	141.7 ^a	38.7	137.3 ^a	39.4	134.8 ^a	36.5	128.5 ^b	38.5	115.2 ^c	38.0	-9.29**	-9.45**
K (mg/1000 kcal)	Women†	1399.4	398.4	1552.9 ^a	429.1	1495.6 ^{a,b}	388.7	1448.6 ^b	371.8	1349.4 ^c	353.1	1169.6 ^d	335.2	-138.65**	-138.63**
	Men†	1316.8	359.4	1490.5 ^a	373.6	1383.5 ^{a,b}	368.1	1258.6 ^b	232.7	1252.5 ^b	290.1	1006.9 ^c	252.8	-169.19**	-169.65**
	Total†	1389.5	394.8	1541.3 ^a	419.6	1481.9 ^{a,b}	387.9	1429.2 ^b	364.7	1340.2 ^c	348.3	1154.6 ^d	331.5	-139.32**	-142.94**
Na (mg/1000 kcal)	Women†	883.8	430.0	762.8 ^a	411.0	840.9 ^b	440.7	868.2 ^{b,c}	342.4	931.0 ^d	413.2	1002.5 ^d	491.5	83.82**	84.06**
	Men†	838.9	449.3	722.3 ^a	389.4	825.0 ^{a,b,c}	582.7	860.8 ^{a,b,c}	480.1	965.2 ^{b,c}	356.1	935.9 ^c	352.2	90.23**	88.55**
	Total†	878.4	432.5	755.3 ^a	407.1	838.9 ^b	460.3	867.5 ^{b,c}	357.7	934.3 ^{c,d}	408.0	996.2 ^d	480.2	85.59**	84.89**

*Values with the same letter do not differ using the ANOVA and Scheffe test at 5% of probability.

†P value <0.05 for linear trend test of food consumption variables according to UPFS consumption quintiles.

‡Percentage of total energy consumption from UPF: average (amplitude): Q1 = 8.4% (0 to 14.4); Q2 = 18.5% (14.4 to 22.6); Q3 = 26.4% (22.6 to 30.4); Q4 = 34.9% (30.4 to 40.6); Q5 = 50.2% (40.6 to 85.7).

§Obtained by Linear Regression of the amount of dietary nutrients in the quintiles of energy percentage from consumption of UFP and expressed in units of sD.

||Adjusted by sex, age, years of education and marital status.

**P value statistically significant at P < 0.05.

Table 4 Average consumption of vitamins according to the quintiles of percentage of energy from the consumption of ultra-processed foods (UPF) from regulars in a Brazilian health promotion service. Belo Horizonte, Minas Gerais, Brazil (2013–2014)

Vitamins	Sex	Average	SD	Percent quintiles of UPF energetic contribution/total dietary energy‡,*										Standardised regression coefficient§	
				1	SD	2	SD	3	SD	4	SD	5	SD	Crude	Adjusted
A (µg/1000 kcal)	Women†	526.5	853.7	570.0 ^a	928.7	574.2 ^a	1011.1	604.8 ^a	1034.6	498.6 ^a	728.4	391.8 ^b	392.9	-68.38**	-72.23**
	Men	459.4	820.9	505.8 ^a	1106.9	401.7 ^a	487.1	573.4 ^a	864.1	357.4 ^a	618.8	419.9 ^a	641.8	-41.05	-47.79
	Total†	519.2	849.9	560.9 ^a	963.5	552.4 ^a	962.7	605.9 ^a	1019.2	486.8 ^{a,b}	718.7	390.0 ^b	422.2	-62.60**	-69.70**
B ₃ (mg/1000 kcal)	Women†	8.2	3.6	8.6 ^a	4.2	8.3 ^a	3.4	8.5 ^a	3.9	8.2 ^a	3.4	7.7 ^b	3.1	-0.32**	-0.34**
	Men	7.9	3.2	7.8 ^a	3.3	8.0 ^a	3.7	8.2 ^a	2.8	8.4 ^a	3.3	7.2 ^a	2.9	-0.10	-0.07
	Total†	8.2	3.6	8.4 ^a	4.1	8.2 ^a	3.5	8.4 ^a	3.8	8.2 ^{a,b}	3.4	7.6 ^b	3.1	-0.28**	-0.31**
B ₆ (mg/1000 kcal)	Women†	1.3	1.2	1.6 ^a	2.0	1.2 ^b	1.2	1.1 ^b	0.9	1.0 ^{b,c}	0.8	0.8 ^c	0.5	-0.25**	-0.25**
	Men†	1.2	1.1	1.4 ^a	1.5	1.2 ^{a,b}	1.3	0.9 ^{a,b}	0.4	1.0 ^a	0.9	0.8 ^b	0.4	-0.20**	-0.22**
	Total†	1.3	1.1	1.6 ^a	1.9	1.2 ^b	1.2	1.1 ^{b,c}	0.9	1.0 ^{b,c,d}	0.8	0.8 ^d	0.5	-0.24**	-0.25**
B ₁₂ (µg/1000 kcal)	Women†	2.8	5.9	3.0 ^a	6.2	3.2 ^a	7.1	3.3 ^a	7.4	2.6 ^{a,b}	5.0	1.9 ^{a,b}	2.4	-0.41**	-0.43**
	Men†	2.7	5.9	3.2 ^a	7.9	2.0 ^a	2.6	3.6 ^a	6.6	1.9 ^a	4.5	2.4 ^a	4.5	-0.34	-0.37
	Total†	2.8	5.9	3.1 ^{a,b}	6.5	3.0 ^a	6.8	3.4 ^a	7.3	2.5 ^{a,b}	5.0	1.9 ^b	2.7	-0.39**	-0.42**
Folate (µg/1000 kcal)	Women†	152.2	70.7	158.9 ^a	79.9	157.7 ^a	72.2	155.5 ^a	69.9	150.9 ^{a,b}	67.9	138.8 ^b	61.9	-7.43**	-7.29**
	Men†	146.5	69.2	158.9 ^a	79.8	147.9 ^a	75.3	145.9 ^a	55.2	139.3 ^a	59.2	127.2 ^a	57.4	-12.34**	-11.44**
	Total†	151.6	70.6	159.0 ^a	79.8	156.1 ^a	72.6	155.1 ^a	68.7	149.9 ^{a,b}	67.2	137.9 ^b	61.4	-7.84**	-7.87**
C (mg/1000 kcal)	Women†	84.1	114.8	92.9 ^a	107.8	92.1 ^a	124.7	87.1 ^a	118.7	82.1 ^{a,b}	127.6	67.6 ^b	89.2	-9.00**	-9.18**
	Men†	64.5	95.4	73.8 ^{a,b}	104.5	88.8 ^a	141.7	50.6 ^{a,b}	57.1	56.4 ^{a,b}	56.0	37.1 ^b	43.4	-12.53**	-12.01**
	Total†	82.2	112.8	89.7 ^a	107.4	92.4 ^a	126.9	83.9 ^a	114.6	80.1 ^{a,b}	122.8	64.9 ^b	86.4	-8.81**	-9.61**

*Values with the same letter do not differ using the ANOVA and Scheffe test at 5% of probability.

†P value <0.05 for linear trend test of food consumption variables according to ultra-processed food consumption quintiles.

‡Percentage of total energy consumption from ultra-processed foods: average (amplitude): Q1 = 8.4% (0 to 14.4); Q2 = 18.5% (14.4 to 22.6); Q3 = 26.4% (22.6 to 30.4); Q4 = 34.9% (30.4 to 40.6); Q5 = 50.2% (40.6 to 85.7).

§Obtained by linear regression of the amount of dietary nutrients in the quintiles of energy percentage from consumption of ultra-processed foods and expressed in units of sD.

||Adjusted by sex, age, years of education and marital status.

**P values statistically significant at P < 0.05.

**Table 5** Prevalence and prevalence ratio of inadequate nutrient intake with recommendations for the prevention of non-communicable disease (NCD) according to the quintiles of energy percentage derived from the consumption of ultra-processed foods (UPF) among health promotion service users in Brazil, Belo Horizonte, Minas Gerais, Brazil (2013–2014)

Nutrients	Sex	Inadequacy total sample (%)	Percent quintiles of UPF energetic contribution/total dietary energy‡										P value**
			1		2		3		4		5		
			%	PR	%	PR	%	PR	%	PR	%	PR	
Total fat	Women*	48.9	36.1	1.0	43.1	1.2	45.4	1.2†	54.6	1.5†	63.9	1.7†	<0.001
	Men*	43.1	25.4	1.0	40.2	1.6	49.3	1.9†	56.2	2.2†	62.9	2.4†	<0.001
	Total*	48.2	34.1	1.0	42.7	1.2†	45.7	1.3†	54.7	1.6†	63.8	1.8†	<0.001
Saturated fat	Women*	43.7	32.9	1.0	39.3	1.2	39.9	1.2	49.8	1.5†	55.4	1.7†	<0.001
	Men*	43.2	26.4	1.0	47.6	1.7†	46.9	1.6†	48.4	1.7†	60.9	2.2†	0.001
	Total*	43.6	31.7	1.0	40.3	1.3†	40.6	1.2†	49.7	1.5†	55.9	1.7†	<0.001
Trans-fat	Women*	62.2	44.3	1.0	59.6	1.3†	66.2	1.5†	67.7	1.5†	71.6	1.6†	<0.001
	Men*	60.5	42.1	1.0	60.9	1.4	69.6	1.6†	75.0	1.8†	72.6	1.7†	<0.001
	Total*	62.0	43.8	1.0	59.8	1.4†	66.5	1.5†	68.4	1.5†	71.7	1.6†	<0.001
Fibre	Women*	58.3	46.1	1.0	51.7	1.1	56.3	1.2†	63.1	1.4†	72.7	1.6†	<0.001
	Men*	62.5	38.9	1.0	63.4	1.6†	76.8	1.9†	67.2	1.7†	88.7	2.3†	<0.001
	Total*	58.8	44.7	1.0	53.1	1.2†	58.4	1.3†	63.5	1.4†	74.2	1.7†	<0.001
K	Women*	83.3	72.5	1.0	79.2	1.1	82.0	1.1	86.7	1.2†	94.8	1.3†	<0.001
	Men*	86.1	72.2	1.0	82.9	1.1	94.2	1.3	95.3	1.3	100.0	1.4	<0.001
	Total*	85.4	74.9	1.0	82.0	1.1	85.3	1.1†	88.9	1.2†	95.7	1.3†	<0.001
Na	Women*	29.8	21.3	1.0	22.9	1.1	28.5	1.3†	31.8	1.5†	43.1	2.0†	<0.001
	Men*	25.3	19.2	1.0	19.0	0.9	24.2	1.3	34.4	1.8	37.5	1.9†	0.006
	Total*	29.4	20.7	1.0	22.9	1.1	27.7	1.3†	32.8	1.5†	42.6	2.0†	<0.001

PR, prevalence ratio.

*P value <0.05 for linear trend test.

†Statistically significant values for Wald test at 5% significance.

‡Percentage of total energy consumption from ultra-processed foods: average (amplitude): Q1 = 8.4% (0 to 14.4); Q2 = 18.5% (14.4 to 22.6); Q3 = 26.4% (22.6 to 30.4); Q4 = 34.9% (30.4 to 40.6); Q5 = 50.2% (40.6 to 85.7).

** χ^2 test with Bonferroni correction.

study was lower for total fats (48.2% *v.* 60.0%), saturated fats (43.6% *v.* 73.8%), fibre (58.8% *v.* 70.2%), Na (29.4% *v.* 77.9%) and K (85.4% *v.* 86.1%). Both studies took place in developed countries with an average consumption of UPF greater than 40%, which may explain the higher rate of inadequate nutrient intake. Further studies are needed to investigate inadequate nutrient intake associated with NCD in developing countries and in health promotion services.

Unlike national and international studies, our study identified positive characteristics of the participants' diets, even among those with high consumption of UPF. The values for energy consumption (average: 1435 kcal/d) and energy density (average: 1.5 kcal/g) of the diet were lower than those found in a Brazilian population study (1866 kcal/d and 1.7 kcal/g, respectively)⁽⁷⁾. Similarly, average fibre consumption was higher (12.2 *v.* 11.1 g/d for the Brazilian population study)⁽⁷⁾, and Na consumption was lower (average: 877.8 mg/1000 kcal) relative to the national average of approximately 12 000 mg/person⁽⁵⁸⁾.

These positive characteristics of the diet, such as lower energy consumption, energy density and Na and higher fibre consumption, when compared to the national average may be related to the high prevalence of overweight or obesity (64.3%) and chronic diseases, such as arterial hypertension (53.3%) and diabetes (16.1%), identified among PAS users. The presence of NCD can encourage the search for food choices that promote an energetic deficit and participation in regular physical exercise, especially

among women⁽⁵⁹⁾. Data from the Surveillance System risk and protection factors for NCD carried out in the 27 Brazilian capitals (VIGITEL) showed that only 22.9% of Brazilian respondents and 31.3% of respondents from Belo Horizonte presented recommended consumption of fruits and vegetables⁽⁶⁰⁾, which differed from the results of our study that found a high rate of adequate consumption of fruits and vegetables (5.4 servings/d)⁽⁶¹⁾. This fact may explain the higher fibre content and lower energy density in the diet of the health promotion service users studied when compared with national data⁽⁷⁾. In addition, the PAS offers frequent health education activities and services that aim to promote adequate food intake and healthy choices⁽⁶²⁾.

This paper makes a significant contribution to the field by highlighting the importance of health services for improving diet quality and thereby delaying the progression of NCD and their associated risk factors. On the other hand, the high consumption of UPF found in our study population reinforces the importance of access to dietary guidelines that discourage UPF consumption in favour of healthier alternatives and that provides guidance on strategies to limit UPF intake. The new Dietary Guidelines for the Brazilian Population⁽²¹⁾, which provides guidance on food consumption according to the degree and extent of industrial processing, was published after the data collection period in the present study. Thus, further studies are needed to assess changes in the consumption of UPF in

Table 6 Prevalence and prevalence ratio of inadequate consumption of micronutrients according to quintiles of percentage of energy from consumption of ultra-processed foods (UPF) among health promotion service users in Brazil, Belo Horizonte, Minas Gerais, Brazil (2013–2014)

Nutrients	Sex	Inadequacy total sample (%)	Percent quintiles of UPF energetic contribution/total dietary energy†										P value**
			1		2		3		4		5		
			%	PR	%	PR	%	PR	%	PR	%	PR	
Minerals													
Ca (mg/1000 kcal)	Women*	92.3	95.6	1.0	94.1	0.99	89.7	0.94	92.6	0.98	90.1	0.95	<0.001
	Men*	83.5	90.0	1.0	80.2	0.89	86.6	0.99	73.0	0.84	82.2	0.96	0.136
	Total*	91.2	94.6	1.0	92.4	0.98	89.3	0.95	90.6	0.97	89.3	0.96	0.001
Iron (mg/1000 kcal)	Women	34.3	37.9	1.0	32.0	0.86	33.0	0.87	31.0	0.82	37.6	1.00	0.902
	Men	24.6	25.4	1.0	31.7	1.33	21.7	1.00	21.9	1.03	19.3	0.93	0.439
	Total	33.0	35.5	1.0	32.0	0.92	31.8	0.92	30.1	0.87	35.7	1.05	0.940
Phosphorus (mg/1000 kcal)	Women	29.8	34.2	1.0	30.7	0.93	26.2	0.81	26.5	0.83	31.5	1.01	0.327
	Men	17.4	19.8	1.0	14.6	0.79	20.3	1.19	14.1	0.81	16.1	0.99	0.539
	Total	28.3	31.5	1.0	28.8	0.94	25.6	0.86	25.4	0.86	30.1	1.05	0.570
Mg (mg/1000 kcal)	Women	88.2	87.8	1.0	86.8	0.99	86.9	0.99	90.2	1.04	89.2	1.03	0.448
	Men	92.1	92.1	1.0	90.0	0.98	92.7	1.01	95.3	1.03	90.3	0.98	0.688
	Total	88.7	88.6	1.0	87.2	0.99	87.5	0.99	90.6	1.03	89.3	1.02	0.670
Vitamins													
A (µg/1000 kcal)	Women*	62.5	66.3	1.0	64.8	0.99	60.4	0.94	60.9	0.95	60.4	0.95	0.039
	Men	69.5	79.4	1.0	59.7	0.75	68.1	0.87	67.2	0.87	66.1	0.86	0.049
	Total*	63.3	68.7	1.0	64.2	0.95	61.2	0.91	61.6	0.92	60.9	0.92	0.003
B ₃ (mg/1000 kcal)	Women*	59.1	65.9	1.0	59.1	0.91	55.4	0.86	57.0	0.89	58.5	0.93	0.009
	Men*	52.8	66.7	1.0	53.6	0.82	52.2	0.82	42.2	0.67	35.5	0.56†	<0.001
	Total*	58.3	66.1	1.0	58.4	0.89	55.1	0.85†	55.6	0.87†	56.4	0.89	<0.001
B ₆ (mg/1000 kcal)	Women*	39.8	33.5	1.0	34.9	1.07	33.5	1.03	44.4	1.38†	51.8	1.64†	<0.001
	Men	26.0	24.6	1.0	24.4	1.06	23.2	1.07	23.4	1.09	37.1	1.69	0.075
	Total*	38.2	46.7	1.0	33.7	1.09	32.4	1.06	42.4	1.39†	50.4	1.69†	<0.001
B ₁₂ (µg/1000 kcal)	Women	42.6	45.5	1.0	42.6	0.95	39.3	0.89	42.4	0.97	43.6	1.01	0.513
	Men	31.0	30.9	1.0	32.9	1.08	26.1	0.88	35.9	1.20	29.0	0.97	0.788
	Total	41.2	42.8	1.0	41.4	0.98	37.9	0.91	41.8	1.01	42.3	1.04	0.840
Folate (µg/1000 kcal)	Women	87.6	88.2	1.0	88.0	1.00	85.1	0.97	87.2	0.99	89.5	1.02	0.454
	Men	78.2	79.4	1.0	78.0	0.98	75.4	0.96	78.1	1.01	79.0	1.01	0.958
	Total	86.5	86.5	1.0	86.8	1.00	84.1	0.98	86.3	1.00	88.6	1.03	0.250
C (mg/1000 kcal)	Women*	43.5	41.3	1.0	42.6	1.05	42.2	1.04	40.8	1.00	50.2	1.25†	0.006
	Men	56.6	55.5	1.0	47.6	0.87	60.9	1.13	50.0	0.93	72.6	1.39	0.024
	Total*	45.0	44.0	1.0	43.2	1.01	44.1	1.04	41.7	0.97	52.2	1.23†	0.003

PR, prevalence ratio.

*P value <0.05 for linear trend test.

†Statistically significant values for Wald test at 5% significance.

‡Percentage of total energy consumption from ultra-processed foods: average (amplitude): Q1 = 8.4% (0 to 14.4); Q2 = 18.5% (14.4 to 22.6); Q3 = 26.4% (22.6 to 30.4); Q4 = 34.9% (30.4 to 40.6); Q5 = 50.2% (40.6 to 85.7).

** χ^2 test with B.

these services following the implementation of this dietary guideline.

These results also alert to challenges of nutritional care in SUS. The expansion of health promotion programmes such as PAS must occur in tandem with the production and dissemination of materials based on the Dietary Guidelines for the Brazilian Population^(21–24). Ongoing food and nutrition education and professional development for health professionals is also urgently required⁽⁶³⁾. In addition, it is important to promote the expansion of dietitian's presence in PHC. However, since November 2019, the Expanded Nucleus of Family Health and Primary Care, a strategy that expanded the hiring of professionals, including dietitians, no longer has specific financing, and this choice was conditioned to the political organisation of each municipality^(64,65). Concomitantly, the current government approved a freeze on spending on the

health sector and attempted to modify the Brazilian Federal Constitution to remove the minimum investment in health in the country. All these aspects increase the challenges to implement health food promotion policies in Brazilian PHC.

Another important strategy to reduce UPF consumption is the deployment of intersectoral public policies with guaranteed funding, including the National Food and Nutrition Security Policy⁽⁶⁶⁾, National Food and Nutrition Policy⁽⁶⁷⁾, National System Food and Nutrition Surveillance⁽⁶⁸⁾ and programmes as Food Acquisition Program and Bolsa Família, which is a cash transfer programme⁽⁶⁹⁾. Considering the importance of intersectoral policies, Belo Horizonte (the municipality of this study) has several internationally awarded initiatives aimed at improving access to adequate and healthy food, such as the implementation of public farmer's market and open air



markets. However, these initiatives are scarce in their reach⁽⁷⁰⁾, highlighting the importance of policies that expand access and commercialisation of healthy foods in the context of health promotion services.

Policies that meet the regulatory advertising rules and nutritional labelling requirements of foods and that include proposals such as the taxation of UPF are also important for minimising UPF intake. In a study evaluating the immediate effects of adopting front-of-pack nutrition labelling in Uruguay, 58% of respondents changed their typical purchasing decisions, opting for healthier products, presumably due to the simplified nutritional information that was more visible and more easily understood⁽⁷¹⁾. In Brazil, front-of-pack nutrition labelling was approved for nationwide use as early as late 2022⁽⁷²⁾.

Another effective example in which UPF consumption was reduced was the tax on sugar-sweetened beverages adopted in Mexico, which was associated with fewer purchases of the taxed beverages and greater purchases of untaxed beverages⁽⁷³⁾. In a study conducted with a household sample of the Brazilian population, taxation of UPF was suggested as a method for obesity control. A 1% increase in the retail price of UPF is believed to result in a lower prevalence of overweight and obesity by 0.33% and 0.59%, respectively⁽⁷⁴⁾. However, there is no political perspective for taxing UPF in Brazil. So far, there have been few bill proposals for taxation and what few there were failed to be approved in 2020⁽⁷⁵⁾.

Among the study's limitations are possible inherent biases in food-related questionnaires, such as memory errors, underreporting and conscious or subconscious modification of food consumption patterns on recorded days. To minimise these biases, the data collection team was trained every 6 months, the instruments were pre-tested and validated and a field manual was compiled for consultation. In addition, the use of R24 h is advised for epidemiological studies with a large sample size, and especially for individuals with reduced education, because it is less sensitive to the respondent's educational and literacy levels^(28,29).

Another limitation is that added or free sugars were not analysed in this study. Given that self-reports of sugar additions by participants were often inaccurate, thereby leading to possible measurement biases, this analysis was excluded; rather, sugar content was standardised according to the volume ingested. Further studies are needed to uncover the relationship between the consumption of UPF and the sugar content of the diet in the context of preventing NCD.

Furthermore, the external validity of this study is limited because the participants were recruited from health services rather than from the general population. However, the study was conducted in a city that was a pioneer in implementing this health promotion service and that has a larger number of PAS units than most Brazilian municipalities. Finally, data were collected in 2013–2014, suggesting

that the results of the study may not accurately reflect the dynamic fluctuations characteristic of the UPF market. Nevertheless, our results can be used as a baseline for future studies that wish to assess the impact of the Dietary Guidelines for the Brazilian Population, which was published in 2014⁽²¹⁾.

The advantages of this study are the large sample size and the inclusion of the NOVA food system to analyse food consumption in a health promotion programme, even before this classification was proposed in the Dietary Guidelines for the Brazilian Population. Our study is the first study to assess the consumption of UPF and its association with the nutritional profile of the users of a health promotion service, highlighting the challenges yet to be overcome to improve public policies and health promotion strategies.

The results suggest that the actions taken to promote adequate and healthy food consumption developed in a health promotion service of PHC apparently contribute to an adequate nutritional dietary profile. On the other hand, the results indicate the need to incorporate socio-demographic factors and approaches that encourage the reduction of UPF consumption in order to generate more informative data. In addition, addressing issues inherent to the food environment in the design of public policies can contribute to promoting healthy food environments and thereby favour a broader range of interventions offered by health services.

Conclusions

The high consumption of UPF, which was especially observed among women, adults and individuals with higher education or income levels, was associated with a worse diet profile in participants in a health promotion service of Brazilian PHC. The positive characteristics of the diet of the health service users, such as lower energy consumption, energy density and Na and higher fibre consumption, when compared to the national data, suggest the importance of investing in health services that offer regular actions to promote adequate and healthy food consumption, since these diet characteristics configure protective factors for NCD. However, the results also highlight the need for future studies to include approaches that encourage the reduction of UPF consumption and to support these approaches with public policies that promote healthy food consumption.

Acknowledgements

Acknowledgements: To the team of the Research Group on Nutrition Interventions at the Federal University of Minas Gerais who conducted the community trial and who organised the database; to the users of the Health Gym Program and employees of municipal health secretary of the city of

Belo Horizonte who contributed to this study and to Luiza Ali Dinis and Mariana Oliveira Martiniano for reviewing the English. *Financial support:* This research was funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico –CNPq (476686/2013-0; a research productivity scholarship to ACSL e LCS) and Fundação de Amparo à Pesquisa do Estado de Minas Gerais – FAPEMIG (21618/2013). The funders had no role in the design of the study; in the collection, analyses or interpretation of data; in the writing of the manuscript, or in the decision to publish the results. *Conflict of interest:* There are no conflicts of interest. *Authorship:* S.F.C. – designed the research; took care analyses; interpreted the data; wrote the first draft of the manuscript; contributed to the data interpretation and revised each draft for important intellectual content. L.C.D.S. – took care of data collection and management; interpreted the data; contributed to the data interpretation and revised each draft for important intellectual content. M.S.L. – took care of data collection and management; contributed to the data interpretation and revised each draft for important intellectual content. P.P.D.F. – took care of data collection and management; contributed to the data interpretation and revised each draft for important intellectual content; got funding acquisition. A.C.S.L. – designed the research; took care of data collection and management; interpreted the data; wrote the first draft of the manuscript; contributed to the data interpretation and revised each draft for important intellectual content; got funding acquisition; was responsible for conceptualisation and project administration. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving research study participants were approved by the Research Ethics Committees of Universidade Federal de Minas Gerais (0537.0.0203.000-11) and of the Belo Horizonte municipality (0537.0.0203.410-11 A). In addition, it was registered at the Brazilian government's Registry of Clinical Trials (RBR-9h7ckx) in accordance with the criteria required by the International Committee of Medical Journal Editors (ICMJE) and by the WHO. All users of the PAS units signed an informed consent form.

References

1. Monteiro CA, Cannon G, Moubarac JC *et al.* (2018) The UN decade of nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr* **21**, 5–17.
2. FAO (2019) *Ultra-Processed Foods, Diet Quality, and Health Using the NOVA Classification System*. Rome: FAO/WHO.
3. Steele EM, Popkin BM, Swinburn B *et al.* (2017) The share of ultra-processed foods and the overall nutritional quality of diets in the US: evidence from a nationally representative cross-sectional study. *Popul Health Metr* **15**, 11.
4. Rauber F, Da Costa Louzada ML, Steele EM *et al.* (2018) Ultra-processed food consumption and chronic non-communicable diseases-related dietary nutrient profile in the UK (2008–2014). *Nutrients* **10**, 587.
5. Moubarac JC, Batal M, Louzada ML *et al.* (2017) Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite* **108**, 512–520.
6. Crovetto M, Uauy R, Martins AP *et al.* (2014) Household availability of ready-to-consume food and drink products in Chile: impact on nutritional quality of the diet (2006–2007). *Rev Méd Chile* **142**, 850–858.
7. Louzada MLD, Martins APB, Canella DS *et al.* (2015) Ultra-processed foods and the nutritional dietary profile in Brazil. *Rev Saude Publica* **49**, 11.
8. Mendonça RD, Pimenta AM, Gea A *et al.* (2016) Ultraprocessed food consumption and risk of overweight and obesity: the University of Navarra Follow-Up (SUN) cohort study. *Am J Clin Nutr* **104**, 1433–1440.
9. Monteiro CA, Moubarac JC, Levy RB *et al.* (2018) Household availability of ultra-processed foods and obesity in nineteen European countries. *Public Health Nutr* **21**, 18–26.
10. Mendonça RD, Lopes ACS, Pimenta AM *et al.* (2017) Ultra-processed food consumption and the incidence of hypertension in a Mediterranean cohort: the Seguimiento Universidad de Navarra Project. *Am J Hypertens* **30**, 358–366.
11. Steele EM, Juul F, Neri D *et al.* (2017) Dietary share of ultra-processed foods and metabolic syndrome in the US adult population. *Prev Med* **125**, 40–48.
12. Fiolet T, Srour B, Sellem L *et al.* (2018) Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. *BMJ* **360**, k322.
13. Srour B, Fezeu LK, Kesse-Guyot E *et al.* (2019) Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Sante). *BMJ* **365**, 14.
14. Gómez-Donoso C, Sanchez-Villegas A, Martínez-González MA *et al.* (2019) Ultra-processed food consumption and the incidence of depression in a Mediterranean cohort: the SUN Project. *Eur J Nutr* **59**, 1093–1103.
15. Adjibade A, Julia C, Alles B *et al.* (2019) Prospective association between ultra-processed food consumption and incident depressive symptoms in the French NutriNet-Santé cohort. *BMC Med* **17**, 1–13.
16. Sandoval-Insauti H, Blanco-Rojo R, Graciani A *et al.* (2019) Ultra-processed food consumption and incident frailty: a prospective cohort study of older adults. *J Gerontol A Biol Sci Med Sci* **75**, 1126–1133.
17. Kim H, Hu E & Rebbholz C (2019) Ultra-processed food intake and mortality in the USA: results from the Third National Health and Nutrition Examination Survey (NHANES III, 1988–1994). *Public Health Nutr* **22**, 1777–1785.
18. Schnabel L, Kesse-Guyot E, Allès B *et al.* (2019) Association between ultraprocessed food consumption and risk of mortality among middle-aged adults in France. *JAMA Intern Med* **179**, 490–498.
19. Rico-Campà A, Martínez-González MA, Alvarez-Alvarez I *et al.* (2019) Association between consumption of ultra-processed foods and all cause mortality: SUN prospective cohort study. *BMJ* **365**, 1949.
20. World Health Organization (2017) *Noncommunicable Diseases Progress Monitor*. Geneva: World Health Organization.
21. Brasil (2014) *Dietary Guidelines for the Brazilian Population*. Brasília: Ministério da Saúde.
22. Ministério da Saúde (2019) *Instructional Manual: Implementing the Dietary Guidelines for the Brazilian Population in Teams that Work in Primary Health Care*. Brasília: Ministério da Saúde.
23. Ministério da Saúde (2021) *Issue 1 Protocols for Use of the Dietary Guidelines for the Brazilian Population in Food Orientation: Theoretical and Methodological Bases and Protocol for the Adult Population*. Brasília: Ministério da Saúde.
24. Ministério da Saúde (2021) *Issue 2 Protocols for the Use of the Dietary Guidelines for the Brazilian Population in the Food*



- Orientation of the Elderly Population*. Brasília: Ministério da Saúde.
25. Brasil (2017) Ordinance n° 2.436, september 21th, 2017. Approves the National Primary Care Policy, establishing the revision of guidelines for the organization of Primary Care, within the scope of the Unified Health System (SUS); available at https://bvsms.saude.gov.br/bvs/saudelegis/gm/2017/prt2436_22_09_2017 (accessed May 2020).
 26. Brasil (2017) Consolidation Ordinance n° 5, September 28th, 2017 Consolidation of the rules on health actions and services of the Unified Health System; available at http://bvsms.saude.gov.br/bvs/saudelegis/gm/2017/prc0005_03_10_2017 (accessed May 2020).
 27. Brasil (2020) Health Academy Program [Internet]. Primary health care department. Ministério da Saúde; available at <https://aps.saude.gov.br/ape/academia/sobre> (accessed May 2020).
 28. Menezes MCD, Costa BVDL, Ferreira NL *et al.* (2017) Methodological course of a community controlled trial in health care services: translational epidemiological research on nutrition. *Demetra* **12**, 1203–1222.
 29. IBGE (2020) Brasil/Minas Gerais/Belo Horizonte [Internet]. IBGE Cities. IBGE; available at <https://cidades.ibge.gov.br/brasil/mg/belo-horizonte/panorama> (accessed May 2020).
 30. PBH (2018) Health Vulnerability Index 2003 [Internet]. Belo Horizonte: Prefeitura de Belo Horizonte; available at <http://www.pbh.gov.br/smsa/biblioteca/gabinete/risco2003.pdf> (accessed May 2020).
 31. Willett W (2013) *Nutritional Epidemiology*, 3rd ed. New York: Oxford University Press.
 32. Fisberg RM, Martini LA, Slater B *et al.* (2005) Food survey methods. In *Food Surveys: Methods and Scientific Bases*, pp. 1–29 [RM Fisberg, editor]. Barueri: Manole.
 33. Fisberg R, Marchioni DML & Colucci ACA (2009) Assessment of food consumption and nutrient intake in clinical practice. *Arq Bras Endocrinol Metab* **53**, 617.
 34. Pinheiro ABC, Lacerda EMA, Benzecry EH *et al.* (2004) *Table for Evaluation of Food Consumption in Home Measures*, 5th ed. São Paulo: Atheneu.
 35. Brasil (2011) Household Budget Survey 2008–2009. Tables of nutritional composition of foods consumed in Brazil. Rio de Janeiro: Ministério da Saúde; available <https://biblioteca.ibge.gov.br/visualizacao/livros/liv50002.pdf> (accessed May 2020).
 36. Fisberg RM & Marchioni DML (2012) *Food Consumption Assessment Manual in Population Studies: The Health Survey Experience Em São Paulo*. São Paulo, SP: ISA.
 37. World Cancer Research Foundation (2009) *Energy Density: Finding the Balance for Cancer Prevention*. London: World Cancer Research Foundation.
 38. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases: Report of a Joint WHO/FAO Expert Consultation*. Geneva: World Health Organization.
 39. World Health Organization (2013) *WHO Issues New Guidance on Dietary Salt and Potassium*. Geneva: World Health Organization.
 40. World Health Organization (2018) *Healthy Diet. Fact sheet N°394*. Geneva: World Health Organization.
 41. Institute of Medicine (1998) *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline*. Washington (DC): National Academy Press.
 42. Institute of Medicine (2000) *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*. Washington (DC): National Academy Press.
 43. Institute of Medicine (2002) *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*. Washington (DC): National Academy Press.
 44. Institute of Medicine (2005) *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. Washington (DC): National Academy Press.
 45. Lopes ACS, Ferreira AD, Mendonça RDD *et al.* (2016) Health promotion strategy: Health Academy Program of Belo Horizonte. *Rev Bras Atividade Física Saúde* **21**, 379–384.
 46. Bielemann RM, Motta JVS, Minten GC *et al.* (2015) Consumption of ultra-processed foods and their impact on the diet of young adults. *Rev Saúde Pública* **49**, 28.
 47. Marrón-Ponce JA, Sánchez-Pimienta TG, Louzada MLDC *et al.* (2018) Energy contribution of NOVA food groups and sociodemographic determinants of ultra-processed food consumption in the Mexican population. *Public Health Nutr* **21**, 87–93.
 48. Simoes BS, Barreto SM, Molina MCB *et al.* (2018) Consumption of ultra-processed foods and socioeconomic position: a cross-sectional analysis of the Brazilian Longitudinal Study of Adult Health. *Cad Saúde Pública* **34**, 00019717.
 49. Machado PP, Steele EM, Levy RB *et al.* (2019) Ultra-processed foods and recommended intake levels of nutrients linked to non-communicable diseases in Australia: evidence from a nationally representative cross-sectional study. *BMJ* **9**, e029544.
 50. Cediel G, Reyes M, Louzada MLC *et al.* (2017) Ultra-processed foods and added sugars in the Chilean diet. *Public Health Nutr* **21**, 125–133.
 51. Parra DC, da Costa-Louzada ML, Moubarac JC *et al.* (2019) Association between ultra-processed food consumption and the nutrient profile of the Colombian diet in 2005. *Salud Publica Mex* **61**, 147–154.
 52. Guibu IA, Moraes JCD, Guerra Junior AA *et al.* (2017) Main characteristics of patients of primary health care services in Brazil. *Rev Saúde Pública* **51**, Suppl. 2, 17s.
 53. Costa BVL, Oliveira CDL & Lopes ACS (2015) Food environment of fruits and vegetables in the territory of the Health Academy Program. *Cad Saúde Pública* **31**, Suppl. 1, 159–169.
 54. Maia EG, Costa BVDL, Coelho FDS *et al.* (2017) Analysis of TV food advertising in the context of recommendations by the Food Guide for the Brazilian Population. *Cad Saúde Pública* **33**, e00209115.
 55. Steele EM, Baraldi LG, Louzada MLD *et al.* (2016) Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* **6**, e009892.
 56. Marrón-Ponce JA, Flores M, Cediel G *et al.* (2019) Associations between consumption of ultra-processed foods and intake of nutrients related to chronic non-communicable diseases in Mexico. *J Acad Nutr Diet* **119**, 1852–1865.
 57. Machado PP, Steele EM, Levy RB *et al.* (2019) Ultra-processed foods and recommended intake levels of nutrients linked to non-communicable diseases in Australia: evidence from a nationally representative cross-sectional study. *BMJ Open* **9**, e029544.
 58. IBGE (2011) Household Budget Survey 2008–2009. Table of Referred Measures for Foods Consumed in Brazil. Rio de Janeiro: Ministério de Saúde; available at <https://biblioteca.ibge.gov.br/visualizacao/livros/liv50000.pdf> (accessed May 2021).
 59. Assumpção D, Domene SMA, Fisberg RM *et al.* (2017) Differences between men and women in diet quality: a population-based study in Campinas, São Paulo. *Ciência Saúde Coletiva* **22**, 347–358.
 60. Brasil (2020) Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey [Internet]. Brasília: Ministério da Saúde; available at http://bvsms.saude.gov.br/bvs/publicacoes/vigitel_brasil_2019_vigilancia_fatores_risco.pdf (accessed May 2021).



61. Mendonça RD, Lopes MS & Freitas PP (2019) Monotony in the consumption of fruits and vegetables and food environment characteristics. *Re Saúde Pública* **53**, 63.
62. Sá GB, Dornelles GC, Cruz KG *et al.* (2016) The Health Academy Program as a strategy to promote health and healthy lifestyles: the national implementation scenario. *Ciênc Saúde Coletiva* **21**, 1849–1860.
63. Tramontt CR & Jaime PC (2020) Improving knowledge, self-efficacy and collective efficacy regarding the Brazilian dietary guidelines in primary health care professionals: a community controlled trial. *BMC Fam Pract* **21**, 214.
64. Brasil (2014) Family Health Support Center (Primary Care Notebooks, n. 39) [Internet]. Vol. 39. Brasília: Ministério da Saúde; available at http://189.28.128.100/dab/docs/portaldab/publicacoes/caderno_39.pdf (accessed May 2021).
65. Brasil (2019) Ordinance n° 2.698, October, 14th, 2019. It suspends the transfer of financial incentives referring to Community Health Agents (CHA), Family Health Teams (ESF) and Oral Health Teams (ESB), Extended Family Health and Primary Care Center (NASF-AB), Health Teams Street Clinic (eCR), Family Health Team Fluvial / Basic Family Health Unit Fluvial (ESFF / UBSF), Health Teams in the Prison System (ESP) and Mobile Dental Unit (UOM), with no system supply Information System for Primary Care (SISAB); available at <https://www.in.gov.br/en/web/dou/-/portaria-n-2.698-de-14-de-outubro-de-2019-221814122> (accessed May 2021).
66. Brasil (2010) Decree 7.272, August, 25th, 2010. It creates the National System of Food and Nutritional Security – SISAN with a view to ensuring the human right to adequate food, institutes the National Policy of Food and Nutritional Security – PNSAN, establishes the parameters for the preparation of the National Plan for Food and Nutritional Security, and take other measures; available at http://www.planalto.gov.br/ccivil_03/_ato2007-2010/2010/decreto/d7272.htm (accessed May 2021).
67. Brasil (2013) National Food and Nutrition Policy. Brasília: Ministério da Saúde; available at http://bvsms.saude.gov.br/bvs/publicacoes/politica_nacional_alimentacao_nutricao.pdf (accessed May 2021).
68. Brasil (2006) Law n° 11.346, September 15th, 2006. It creates the National System of Food and Nutritional Security – SISAN with a view to ensuring the human right to adequate food and takes other measures; available at http://www.planalto.gov.br/ccivil_03/_ato2004-2006/2006/lei/111346.htm (accessed May 2021).
69. Brasil (2004) Law n° 10.836, January 9th, 2004. It creates the Bolsa Família Program and makes other arrangements; available at http://www.planalto.gov.br/ccivil_03/_ato2004-2006/2004/lei/110.836.htm (accessed May 2021).
70. Lopes ACS, Menezes MCD & Araújo MLD (2017) Food environment and access to fruits and vegetables: “a metropolis into perspective”. *Saúde E Soc* **26**, 10.
71. Ares G, Antúñez L, Gimenez A *et al.* (2020) *Immediate Effects of the Implementation of Frontal Nutritional Labeling in Uruguay*. Montevideo: UNICEF.
72. Brasil (2020) Collegiate Board Resolution – RDC n° 429, October, 8th, 2020 – ANVISA. Provides for the nutritional labeling of packaged foods; available at <https://www.in.gov.br/en/web/dou/-/resolucao-de-diretoria-colegiada-rdc-n-429-de-8-de-outubro-de-2020-282070599> (accessed May 2021).
73. Colchero MA, Popkin BM, Rivera JA *et al.* (2016) Beverage purchases from stores in Mexico under the excise tax on sugar sweetened beverages: observational study. *BMJ* **352**, 6704.
74. Passos CM, Maia EG, Levy RB *et al.* (2020) Association between the price of ultra-processed foods and obesity in Brazil. *Nutr Metab Cardiovasc Dis* **30**, 589–598.
75. Mariath A & Martins A (2020) Ultra-processed food industry regulation for tackling obesity and diet-related non-communicable diseases in the Brazilian legislature: many proposals, no enactments. *Public Health Nutr* **13**, 1–6.