

RESEARCH ARTICLE

Association between particulate air pollution and hypertensive disorders in pregnancy: A retrospective cohort study

Yi Sun^{1,2}, Rashmi Bhuyan^{2,3,4}, Anqi Jiao², Chantal C. Avila⁵, Vicki Y. Chiu⁵, Jeff M. Slezak⁵, David A. Sacks^{5,6}, John Molitor⁷, Tarik Benmarhnia⁸, Jiu-Chuan Chen⁹, Darios Getahun^{5,10}*, Jun Wu²†*

1 Institute of Medical Information, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China, **2** Department of Environmental and Occupational Health, Program in Public Health, University of California, Irvine, California, United States of America, **3** Occupational and Environmental Medicine Residency Program, University of California, Irvine, California, United States of America, **4** Department of Occupational Medicine, Kaiser Permanente Northern California, Antioch, California, United States of America, **5** Department of Research & Evaluation, Kaiser Permanente Southern California, Pasadena, California, United States of America, **6** Department of Obstetrics and Gynecology, University of Southern California, Keck School of Medicine, Los Angeles, California, United States of America, **7** College of Public Health and Human Sciences, Oregon State University, Corvallis, Oregon, United States of America, **8** Scripps Institution of Oceanography, University of California, San Diego, California, United States of America, **9** Departments of Population & Public Health Sciences and Neurology, University of Southern California, Keck School of Medicine, Los Angeles, California, United States of America, **10** Department of Health Systems Science, Kaiser Permanente Bernard J. Tyson School of Medicine, Pasadena, California, United States of America

☞ These authors contributed equally to this work.

† DG and JW also contributed equally to this work.

* Darios.T.Getahun@kp.org (DG); junwu@hs.uci.edu (JW)



OPEN ACCESS

Citation: Sun Y, Bhuyan R, Jiao A, Avila CC, Chiu VY, Slezak JM, et al. (2024) Association between particulate air pollution and hypertensive disorders in pregnancy: A retrospective cohort study. *PLoS Med* 21(4): e1004395. <https://doi.org/10.1371/journal.pmed.1004395>

Academic Editor: Sarah J. Stock, The University of Edinburgh Usher Institute of Population Health Sciences and Informatics, UNITED KINGDOM

Received: July 28, 2023

Accepted: April 4, 2024

Published: April 26, 2024

Copyright: © 2024 Sun et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be shared publicly because of ethical/privacy reasons. Data are available from the Kaiser Permanente Southern California (KPSC) electronic health records for researchers who meet the criteria for access to confidential data. The individual data from KPSC cannot be provided as the scientific data collected and generated under this project utilizes Kaiser Permanente business confidential or proprietary information, which Kaiser Permanente is prohibited from publicly disclosing, including

Abstract

Background

Epidemiological findings regarding the association of particulate matter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$) exposure with hypertensive disorders in pregnancy (HDP) are inconsistent; evidence for HDP risk related to $\text{PM}_{2.5}$ components, mixture effects, and windows of susceptibility is limited. We aimed to investigate the relationships between HDP and exposure to $\text{PM}_{2.5}$ during pregnancy.

Methods and findings

A large retrospective cohort study was conducted among mothers with singleton pregnancies in Kaiser Permanente Southern California from 2008 to 2017. HDP were defined by International Classification of Diseases-9/10 (ICD-9/10) diagnostic codes and were classified into 2 subcategories based on the severity of HDP: gestational hypertension (GH) and preeclampsia and eclampsia (PE-E). Monthly averages of $\text{PM}_{2.5}$ total mass and its constituents (i.e., sulfate, nitrate, ammonium, organic matter, and black carbon) were estimated using outputs from a fine-resolution geoscience-derived model. Multilevel Cox proportional hazard models were used to fit single-pollutant models; quantile g-computation approach

scientific data containing or related to Kaiser Permanente: HealthConnect, health plan enrollment and benefit information, utilization data including diagnoses, procedures, treatments, and pharmacy dispensing records. Further details can be requested by email from KPSC Sponsored Projects Administration (Research-SPA@kp.org).

Funding: This study was supported by the National Institute of Environmental Health Sciences (NIEHS; R01ES030353 to JW and DG). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: ACOG, American College of Obstetricians and Gynecologists; BMI, body mass index; CI, confidence interval; EHR, electronic health record; GH, gestational hypertension; HDP, hypertensive disorders in pregnancy; HELLP, hemolysis, elevated liver enzymes, and low platelet count; HR, hazard ratio; ICD-9/10, International Classification of Diseases-9/10; IQR, interquartile range; KPSC, Kaiser Permanente Southern California; LMP, last menstrual period; PAF, population attributable fraction; PE-E, preeclampsia and eclampsia; PM, particulate matter; US EPA, US Environmental Protection Agency; WHO, World Health Organization.

was applied to estimate the joint effect of PM_{2.5} constituents. The distributed lag model was applied to estimate the association between monthly PM_{2.5} exposure and HDP risk.

This study included 386,361 participants (30.3 ± 6.1 years) with 4.8% (17,977/373,905) GH and 5.0% (19,381/386,361) PE-E cases, respectively. In single-pollutant models, we observed increased relative risks for PE-E associated with exposures to PM_{2.5} total mass [adjusted hazard ratio (HR) per interquartile range: 1.07, 95% confidence interval (CI) [1.04, 1.10] *p* < 0.001], black carbon [HR = 1.12 (95% CI [1.08, 1.16] *p* < 0.001)] and organic matter [HR = 1.06 (95% CI [1.03, 1.09] *p* < 0.001)], but not for GH. The population attributable fraction for PE-E corresponding to the standards of the US Environmental Protection Agency (9 µg/m³) was 6.37%. In multi-pollutant models, the PM_{2.5} mixture was associated with an increased relative risk of PE-E ([HR = 1.05 (95% CI [1.03, 1.07] *p* < 0.001)], simultaneous increase in PM_{2.5} constituents of interest by a quartile) and PM_{2.5} black carbon gave the greatest contribution of the overall mixture effects (71%) among all individual constituents. The susceptible window is the late first trimester and second trimester. Furthermore, the risks of PE-E associated with PM_{2.5} exposure were significantly higher among Hispanic and African American mothers and mothers who live in low- to middle-income neighborhoods (*p* < 0.05 for Cochran's Q test). Study limitations include potential exposure misclassification solely based on residential outdoor air pollution, misclassification of disease status defined by ICD codes, the date of diagnosis not reflecting the actual time of onset, and lack of information on potential covariates and unmeasured factors for HDP.

Conclusions

Our findings add to the literature on associations between air pollution exposure and HDP. To our knowledge, this is the first study reporting that specific air pollution components, mixture effects, and susceptible windows of PM_{2.5} may affect GH and PE-E differently.

Author summary

Why was this study done?

- Although previous studies have explored the associations of particulate matter ≤2.5 µm (PM_{2.5}) exposure with hypertensive disorders in pregnancy (HDP), earlier findings are inconclusive.
- Several unaddressed issues remain, such as lack of simultaneous consideration of PM_{2.5} constituents, mixture effects, windows of susceptibility, and levels of disease severity.

What did the researchers do and find?

- A retrospective cohort study (*N* = 386,361) was conducted using electronic health records from Kaiser Permanente Southern California (2008 to 2017) to examine the associations between HDP (mild HDP: gestational hypertension (GH) versus severe HDP: preeclampsia and eclampsia (PE-E)) and exposure to PM_{2.5}, and identify windows of susceptibility to PM_{2.5} during pregnancy.

- We found higher risks for PE-E associated with exposures to PM_{2.5} total mass, organic matter, and black carbon, but not for GH. The main effect of increased PE-E risk was driven by PM_{2.5} black carbon (71%) and PM_{2.5} organic matter (26%).
- The most influential time windows of PM_{2.5} exposure associated with HDP are the late first trimester and second trimester.
- Hispanic and African American mothers and mothers who live in low- to middle-income neighborhoods may be more vulnerable to air pollution on HDP risk.

What do these findings mean?

- Our findings suggest that different PM_{2.5} chemical constituents and critical exposure windows may affect GH and PE-E differently.
- Our work highlights the need for further research to investigate different components of air pollutants, levels of disease severity, and various susceptible windows, which is important to better understand the underlying mechanisms between air pollution and HDP and develop corresponding interventions.

Introduction

Particulate matter (PM) is a complex mixture of both solid and liquid particles that spans many sizes and consists of various chemical components, including carbon and organic compounds, and inorganic compounds. The different components come from specific sources and can be grouped into primary PM, which is directly emitted from traffic, industrial processes, and burning waste; and secondary PM, which forms in the atmosphere from chemical reactions. PM with an aerodynamic diameter $\leq 2.5 \mu\text{m}$ (PM_{2.5}) has been shown to have an adverse impact on the human body through direct deposits in the lung and, to some extent, by penetrating the alveolar membranes. These tiny particles activate inflammatory processes by releasing mediators resulting in imbalance in the autonomic nervous system and neuroendocrine pathway [1]. There is an increasing body of literature showing the adverse effects of ambient air pollution, predominantly on respiratory and cardiovascular health. Previous studies have established a strong relationship between ambient PM_{2.5} and adverse cardiovascular effects, including elevated blood pressure among the general population [2]; however, their effects on the cardiovascular system during pregnancy, including the occurrence of hypertensive disorders in pregnancy (HDP) remains unclear [3–7].

HDP, which complicates 5% to 10% of pregnancies, is a major cause of maternal and fetal morbidity and mortality [8–10]. During pregnancy, there is an increase in blood volume by 30% to 50% to adapt to the need for increased metabolic demand of the growing fetus [11]. This physiological hypervolemia during pregnancy results in an increased cardiac output. There is also a compensatory increase in heart rate in an effort to increase oxygenation in the face of dilutional hypervolemia. The concomitant increase in the rate of hyperventilation potentially leads to inhalation of more ambient toxic substances during pregnancy, which may contribute to other maternal morbidities, including HDP.

The existing epidemiological findings regarding the association of PM_{2.5} exposure with HDP are inconsistent [3–6]. Previous studies have shown both harmful impacts [12–20] and

negative or null associations [21,22]. Previous studies examining the relationship between $PM_{2.5}$ exposure and preeclampsia also reported inconsistent associations [23–28]. Inconclusive results may be partially due to differences in study populations, sample sizes, methods of exposure assessments, study regions with different $PM_{2.5}$ levels and compositions, climate conditions, and failure to differentiate different magnitudes of HDP.

HDP are classified into 4 types based on the severity of clinical features by the American College of Obstetricians and Gynecologists (ACOG) [29]: chronic hypertension, gestational hypertension (GH), preeclampsia (PE), and chronic hypertension with superimposed preeclampsia and eclampsia (E) (S1 Appendix). The mechanisms linking air pollution and HDP subcategories may be different. For example, the pathophysiology of GH is thought to be distinct from that of PE's in terms of alterations in placental vasculature and angiogenic factors [30]; PE-E involves chronic uteroplacental ischemia from maternal angiogenic imbalance with a changed interaction between vasoactive cytokines, leading to vasospasm [31,32], as opposed to GH. Therefore, examining HDP subcategories separately would be more elucidative about the true associations of the risk factors with HDP and the underlying mechanistic pathways [33].

Additionally, some $PM_{2.5}$ constituents and sources may be more harmful than others to health [34,35]. Previous studies found that different $PM_{2.5}$ constituents and exposure windows were differently associated with several adverse pregnancy outcomes, such as gestational diabetes mellitus [36] and postpartum depression [37], which is important to develop corresponding interventions targeting the main culprits and critical windows. However, very limited studies have investigated the windows of susceptibility to $PM_{2.5}$ and HDP risk [38], and the effects of $PM_{2.5}$ chemical compositions [12]; no prior study has explored the HDP risk related to overall mixture effects of $PM_{2.5}$.

The objective of this study is to investigate the relationship between maternal residential exposures to $PM_{2.5}$ total mass and its constituents (i.e., sulfate, ammonium, nitrate, organic matter, and black carbon) and HDP in a large pregnancy cohort taken from Kaiser Permanente Southern California (KPSC) electronic health record (EHR) data between 2008 and 2017. Specific aims are to (1) examine the associations between HDP and exposure to $PM_{2.5}$ mass and its constituents by the severity of the outcomes (GH versus PE-E); and (2) identify windows of susceptibility to $PM_{2.5}$ exposure during pregnancy. The primary hypothesis is that maternal exposure to $PM_{2.5}$ is associated with an increased risk of developing HDP, while the associations would differ by the severity of HDP, $PM_{2.5}$ constituents, and different exposure windows during pregnancy.

Methods

Study population

The participants were women with singleton pregnancies from January 1, 2008 to December 31, 2017. KPSC EHR is the source of information on demographic characteristics, residential history, individual lifestyles, medical records, and birth records. A total of 386,361 pregnancies were included in PE-E analysis after excluding participants with the following criteria (S2 Appendix): pregnancies who were not KPSC members at the time of pregnancy (i.e., mostly only having labor at KPSC and having no EHRs and outcome measures during pregnancy) or those with gestational age <20 or >43 weeks ($n = 8,408$), those with multiple fetuses ($n = 6,694$), and those without a residential address ($n = 653$). A total of 373,905 pregnancies were included in GH analysis after further excluding pregnancies complicated by chronic hypertension without superimposed preeclampsia ($n = 12,456$). Date of the last menstrual period (LMP) coupled with early pregnancy ultrasonography was utilized to determine the estimated conception date and corresponding gestational age. If there was a discrepancy

between LMP and early pregnancy ultrasound report, the date found on the latter was given preference based on ACOG guidelines [39].

This study was approved by the Institutional Review Board of KPSC and the University of California, Irvine with exemption of informed consent.

Outcome definition

Diagnoses of GH and PE-E were defined by International Classification of Diseases-9/10 (ICD-9/10) diagnostic codes [33]. Pregnant women were screened for hypertension during each prenatal visit. The HDP cases were divided into 2 groups: (1) GH: systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg, on 2 occasions at least 4 h apart after 20 weeks of gestation in a previously normotensive woman; and (2) PE-E: with any one of the following: preeclampsia, preeclampsia superimposed upon chronic hypertension, or eclampsia (S1 Appendix).

Exposure assessment

Monthly concentrations of PM_{2.5} total mass and its constituents (i.e., sulfate, nitrate, ammonium, organic matter, and black carbon) from 2007 to 2017 were obtained from 1-km resolution publicly available data generated by validated geoscience-derived models over North America [40,41] that included chemical transport modeling (GEOS-Chem), satellite remote sensing of aerosol optical depth, and ground-based observations combining with a geographically weighted regression. The PM_{2.5} mass estimates were consistent with ground PM_{2.5} measurements with R² ranging from 0.6 to 0.85. The PM_{2.5} species in our study region of the Southwestern US has moderate to high cross-validated agreement with R² values for the selected PM_{2.5} species: R² sulfate = 0.59, R² nitrate = 0.78, R² ammonium = 0.75, R² organic matter = 0.52, and R² black carbon = 0.42 [40].

KPSC EHR was used to abstract information on residential histories during the entire pregnancy, including residential addresses and residency start and end dates. Monthly concentrations of PM_{2.5} total mass and components at a 1-km resolution were spatiotemporally linked to each woman based on the geocoded residential address history.

Covariates

Covariates and potential confounders were selected from KPSC EHRs based on existing literature [3–6,42–44], including maternal age, race/ethnicity (African/American, non-Hispanic Asians, Hispanic, non-Hispanic white, and others including Pacific Islanders, Native American/Alaskan and mothers with multiple race/ethnicities specified), educational achievement (less than 8th grade, 9th grade to high school graduates, less than 4 years of college, college and more than 4 years of college); median household income at block group level in 2013; exposure to active and passive second-hand smoke during pregnancy; season of conception (warm; May–October; cool; November–April), parity (primiparous versus multiparous); pre-pregnancy body mass index (BMI, kg/m², underweight: <18.5, normal weight: 18.5 to 24.9, overweight: 25.0 to 29.9, obese: ≥ 30.0); health insurance status (Medical or Medicare versus other kinds of health insurance) and year of infant birth. Zip Code Tabulation Areas defined by the US Census Bureau were used to represent zip codes [45]. Potential confounding factors in the relationship between air pollution and HDP were represented by a directed acyclic graph (S3 Appendix).

Statistical analysis

Descriptive statistics were performed with distribution of selected population characteristics and PM_{2.5} exposures. We used chi-square to test the difference between pregnant women with

and without GH or PE-E. Correlation between air pollution metrics was estimated with Pearson's correlation. Multilevel Cox proportional hazard models with zip code as a random effect were applied to examine the associations between each air pollutant (PM_{2.5} total mass and 5 PM_{2.5} constituents) and each outcome during the entire pregnancy. "Entire pregnancy" was defined as the period from the conception date to the date of diagnosis for cases or the date of delivery for non-cases. Per interquartile range (IQR) increment of each air pollutant was used to calculate hazard ratios (HRs) and 95% confidence intervals (CIs). The main model was adjusted for maternal age, race/ethnicity, education, neighborhood household income, season, active and passive smoking during pregnancy, insurance type, and year of infant birth. In addition to the single-pollutant model, air pollutants with hypothesized detrimental effects on the outcome were identified and selected to be further analyzed in the multi-pollutant model. A quantile-based g-computation method from the "qgcomp" package in R was applied to measure the PM_{2.5} mixture effects. This method can estimate the effects of increasing a specific subset of exposures simultaneously controlling for possible confounding from other components in the mixture [46,47] and would be an appropriate approach to assess the joint effect of air pollution mixtures on HDP. To further identify potential windows of susceptibility during pregnancy, distributed lag models incorporating Cox models were applied to estimate the association between monthly exposure to each air pollutant and HDP risk. The lag range was defined as gestational months from the first month to the corresponding month. We considered current exposure at a given time t (month t), past exposure before time t (month₁~month _{$t-1$}), using an inverse weighting approach with weights being calculated based on time to a month t to give more weight to months right before a month t , and potential interactions between past and current exposures (an interaction term of month $t \times$ weighted month₁~month _{$t-1$}).

Additionally, sensitivity analyses were carried out to further examine the influence of adjusting for pre-pregnancy BMI and parity. We also applied the discrete time approach with logit function as a sensitivity analysis. The discrete-time approach is an alternative method to the Cox proportional hazard model but is more flexible without requiring the proportional hazards assumption and may be useful for handling large datasets with time-dependent variables [48,49]. The population attributable fraction (PAF) was used to estimate the proportion of cases in our population that can be attributed to PM_{2.5} exposures corresponding to the air quality standards set by the US Environmental Protection Agency (US EPA, 9 $\mu\text{g}/\text{m}^3$) [50] and the World Health Organization (WHO, 5 $\mu\text{g}/\text{m}^3$) [51]. Further, maternal characteristics and lifestyle behaviors such as maternal age [52], race/ethnicity [53], socioeconomic status [54], smoking [55], obesity, and primiparity [56] could be potential modifiers between air pollution and HDP. Stratified analysis was conducted to explore the differences between population subgroups. Cochran's Q tests were used to measure the heterogeneity among subgroups. SAS version 9.4 software (SAS Institute, Cary, North Carolina, United States of America) and R 4.1.3 were used to conduct all the analyses. A two-sided $P < 0.05$ was considered statistically significant. This study is reported as per the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guideline (S1 STROBE Checklist).

Results

The descriptive statistics of selected demographic and pregnancy characteristics and air pollution levels are presented in Table 1. In total, 4.8% (17,977/373,905) and 5.0% (19,381/386,361) of eligible pregnancies were GH and PE-E cases, respectively. Among PE-E group, 19,334 cases of PE, 3,054 cases of preeclampsia superimposed upon chronic hypertension, and 283 cases of eclampsia were identified.

Table 1. Description of the study population, 2008–2017.

Characteristics	Non-HDP, N = 354,655	GH, N = 17,977	PE-E, N = 19,381	Total births, N = 386,361
Maternal age, years, mean (SD)	30.1(5.8)	30.7(5.9)	30.5(6.3)	30.3(6.1)
Maternal race/ethnicity (N, %)				
African American	26,554 (7.5)	1,484 (8.3)	2,197 (11.3)	29,659 (7.7)
Non-Hispanic Asian	45,061 (12.7)	1,746 (9.7)	2,097 (10.8)	48,276 (12.5)
Hispanic	181,650 (51.2)	7,872 (43.8)	10,453 (53.9)	197,108 (51.0)
Non-Hispanic white	92,449 (26.1)	6,288 (35.0)	4,099 (21.1)	101,367 (26.2)
Multiple/other	8,969 (2.5)	583 (3.2)	532 (2.7)	9,912 (2.6)
Missing	33 (0.1)	4 (0.1)	3 (0.1)	39 (0.1)
Maternal education (N, %)				
≤ 8th Grade	3,890 (1.1)	113 (0.6)	156 (0.8)	4,133 (1.1)
9 Grade to High School	108,911 (30.7)	5,392 (30.0)	6,070 (31.3)	118,631 (31.3)
College (<4 years)	79,696 (22.5)	4,275 (23.8)	4,923 (25.4)	87,494 (23.1)
College (4 years)	109,364 (30.8)	5,646 (31.4)	5,748 (29.7)	119,047 (31.4)
> College	45,806 (12.9)	2,221 (12.4)	2,051 (10.6)	49,437 (13.1)
Missing	6,988 (2.0)	330 (1.8)	433 (2.2)	7,639 (2.0)
Block group median household income in 2013 (N, %)				
<\$43,667	87,920 (24.8)	4,395 (24.4)	5,721 (29.5)	96,428 (25.0)
\$43,667–\$55,929	88,431 (25.0)	4,457 (24.8)	4,961 (25.6)	96,493 (25.0)
\$55,930–\$71,591	88,303 (24.9)	4,510 (25.1)	4,688 (24.2)	96,100 (24.9)
> \$71,591	88,777 (25.0)	4,562 (25.4)	3,960 (20.4)	96,023 (24.9)
Missing	1,224 (0.3)	53 (0.3)	51 (0.3)	1,317 (0.3)
Smoking (N, %)				
Never smoker	295,991 (83.5)	14,236 (79.2)	15,917 (82.1)	321,594 (83.2)
Ever smoker	40,175 (11.3)	2,438 (13.6)	2,332 (12.0)	44,212 (11.4)
Smoking during pregnancy	18,438 (5.2)	1,302 (7.2)	1,130 (5.8)	20,501 (5.3)
Missing	51 (0.1)	1 (0.1)	2 (0.1)	54 (0.1)
Passive smoker (N, %)				
Yes	7,772 (2.2)	444 (2.5)	498 (2.6)	8,572 (2.2)
No	344,644 (97.2)	17,515 (97.4)	18,827 (97.1)	375,483 (97.2)
Missing	2,239 (0.6)	18 (0.1)	56 (0.3)	2,306 (0.6)
Parity (N, %)				
Primiparous	141,266 (39.8)	9,604 (53.4)	10,890 (56.2)	158,534 (41.0)
Multiparous	211,283 (59.6)	8,296 (46.1)	8,360 (43.1)	225,541 (58.4)
Missing	2,106 (0.6)	77 (0.4)	131 (0.7)	2,286 (0.6)
Pre-pregnancy BMI kg/m ² (N, %)				
Underweight (<18.5)	9,239 (2.6)	156 (0.9)	282 (1.5)	9,619 (2.5)
Normal weight (18.5–24.9)	157,454 (44.4)	4,173 (23.2)	5,363 (27.7)	165,768 (42.9)
Overweight (25.0–29.9)	98,435 (27.8)	5,037 (28.0)	5,362 (27.9)	107,243 (27.8)
Obese class 1 (30.0–34.9)	50,782 (14.3)	3,777 (21.0)	4,033 (20.8)	57,348 (14.8)
Obese class 2 (35.0–39.9)	22,708 (6.4)	2,510 (14.0)	2,326 (12.0)	26,762 (6.9)
Obese class 3 (>40.0)	12,862 (3.6)	2,168 (12.1)	1,843 (9.5)	16,235 (4.2)
Missing	3,175 (0.9)	156 (1.0)	174 (0.9)	3,445 (0.9)
Insurance type (N, %)				
MediCal (or Medicaid)	32,224 (9.1)	1,495 (8.3)	1,917 (9.9)	35,123 (9.1)
Other	316,274 (89.2)	16,284 (90.6)	17,206 (88.8)	344,692 (89.2)
Missing	6,157 (1.7)	198 (1.4)	258 (1.3)	6,546 (1.7)

(Continued)

Table 1. (Continued)

Characteristics	Non-HDP, N = 354,655	GH, N = 17,977	PE-E, N = 19,381	Total births, N = 386,361
Season (N, %)				
Warm season (May–October)	175,113 (49.4)	9,087 (50.6)	9,407 (48.5)	190,758 (49.4)
Cool season (November–April)	179,542 (50.6)	8,890 (49.5)	9,974 (51.5)	195,603 (50.6)
Year of infant birth (N, %)				
2008	33,152 (9.4)	1,418 (7.9)	1,402 (7.2)	35,567 (9.2)
2009	31,614 (8.9)	1,406 (7.8)	1,455 (7.5)	34,074 (8.8)
2010	31,445 (8.9)	1,498 (8.3)	1,398 (7.2)	33,928 (8.8)
2011	33,035 (9.3)	1,751 (9.7)	1,467 (7.6)	35,795 (9.3)
2012	35,173 (9.9)	1,799 (10.0)	1,622 (8.4)	38,100 (9.9)
2013	35,738 (10.1)	1,765 (9.8)	1,820 (9.4)	38,786 (10.0)
2014	37,027 (10.4)	1,802 (10.0)	2,059 (10.6)	40,241 (10.4)
2015	38,315 (10.8)	2,039 (11.3)	2,390 (12.3)	42,045 (10.9)
2016	39,495 (11.1)	2,174 (12.1)	2,746 (14.2)	43,641 (11.3)
2017	39,661 (11.2)	2,325 (12.9)	3,022 (15.6)	44,184 (11.4)

Pregnancies complicated by chronic hypertension (N = 12,456) were excluded from GH analysis (N = 373,905).

BMI, body mass index; HDP, hypertensive disorders in pregnancy; GH, gestational hypertension; PE-E, preeclampsia-eclampsia; SD, standard deviation.

<https://doi.org/10.1371/journal.pmed.1004395.t001>

Patients with GH were more frequent among older mothers, African American or non-Hispanic white mothers, mothers with college (≤ 4 years), mothers who live in high-income neighborhoods, ever smokers, smoking or passive smoking mothers, mothers without Medi-Cal/Medicaid insurance, primiparous mothers, and obese mothers; PE-E cases were more frequent among older mothers, African American or Hispanic mothers, mothers with 9 Grade to college < 4 years, mothers who live in low-income neighborhoods, ever smoker, smoking or passive smoking mothers, mothers with MediCal/Medicaid insurance, primiparous mothers, and obese mothers ($p < 0.05$).

Summary statistics and Pearson correlation coefficients between air pollution metrics during the entire pregnancy are presented in Table 2. $PM_{2.5}$ total mass was highly correlated with most $PM_{2.5}$ chemical constituents, including $PM_{2.5}$ organic matter ($r = 0.91$), nitrate ($r = 0.85$), black carbon ($r = 0.79$), and ammonium ($r = 0.75$), and moderately correlated with $PM_{2.5}$ sulfate ($r = 0.48$). Relatively weaker correlations were noticed between sulfate and other $PM_{2.5}$ constituents ($r \leq 0.45$). Moderate to strong correlations were observed between other $PM_{2.5}$

Table 2. Summary statistics and Pearson correlation coefficients between air pollution exposure metrics during the entire pregnancy.

	Mean (SD)	IQR	$PM_{2.5}$ total mass	$PM_{2.5}$ sulfate	$PM_{2.5}$ nitrate	$PM_{2.5}$ ammonium	$PM_{2.5}$ organic matter	$PM_{2.5}$ black carbon
$PM_{2.5}$ total mass	12.86 (2.62)	3.85	1.00					
$PM_{2.5}$ sulfate	1.27 (0.27)	0.35	0.48	1.00				
$PM_{2.5}$ nitrate	2.41 (0.66)	0.94	0.85	0.34	1.00			
$PM_{2.5}$ ammonium	0.95 (0.32)	0.40	0.75	0.45	0.80	1.00		
$PM_{2.5}$ organic matter	5.39 (1.31)	1.78	0.91	0.33	0.65	0.55	1.00	
$PM_{2.5}$ black carbon	1.48 (0.61)	1.05	0.79	0.23	0.54	0.53	0.72	1.00

The units for $PM_{2.5}$ mass and $PM_{2.5}$ constituents are $\mu\text{g}/\text{m}^3$.

IQR, interquartile range; SD, standard deviation.

<https://doi.org/10.1371/journal.pmed.1004395.t002>




Fig 1. Adjusted HRs and 95% CIs of air pollution during pregnancy associated with GH and PE-E. HRs and 95% CIs were calculated for per IQR increment for each air pollutant. Model adjusted for maternal age, race/ethnicity, education, block group household income, smoking and passive smoking status during pregnancy, insurance type, season, and year of infant birth. Zip code was fitted as a random effect. CI, confidence interval; GH, gestational hypertension; HR, hazard ratio; IQR, interquartile range; PE-E, preeclampsia-eclampsia.

<https://doi.org/10.1371/journal.pmed.1004395.g001>

constituents (nitrate, ammonium, organic matter, and black carbon) ($r \geq 0.53$). All the correlation coefficients in Table 2 are statistically significant ($p < 0.05$).

Results from Cox regression models are illustrated in Fig 1. No significant associations were observed between the risks of GH and PM_{2.5} and its constituents; while the relative risks of PE-E were positively associated with PM_{2.5} total mass [HR = 1.07 (95% CI [1.04, 1.10] $p < 0.001$)] and 2 PM_{2.5} constituents: highest for PM_{2.5} black carbon [HR = 1.12 (95% CI [1.08, 1.16] $p < 0.001$)], followed by PM_{2.5} organic matter [HR = 1.06 (95% CI [1.03, 1.09] $p < 0.001$)]. In sensitivity analysis (S4 Appendix), similar results were observed for associations in base models between air pollution and GH or PE-E after further adjusting for pre-pregnancy BMI and primiparity, or applying the discrete time approach models. The PAFs for PE-E corresponding to the standards of the US EPA (9 $\mu\text{g}/\text{m}^3$) and the WHO (5 $\mu\text{g}/\text{m}^3$) were 6.37% and 12.26%, respectively. For example, 6.37% PE-E cases among our study population could be attributed to PM_{2.5} levels above 9 $\mu\text{g}/\text{m}^3$. In the multi-pollutant model (Table 3), β coefficients > 0 indicate positive weights of individual constituents; β coefficients < 0 indicate negative weights of individual constituents. The overall mixture coefficient ψ (log HR) from quantile g-computation is the sum of all β coefficients of the exposures of interest. Simultaneous increase in all of the 5 PM_{2.5} constituents by a quartile was associated with a higher relative risk of PE-E [HR = 1.05 (95% CI [1.03, 1.07] $p < 0.001$)], and black carbon gave the greatest contribution of overall mixture effects (70.63%) among all individual constituents, followed by organic matter (25.59%) and sulfate (3.79%).

In the time window analyses during gestational months 1 to 8 (Fig 2 and S5 Appendix), the GH group showed negative associations with exposure to PM_{2.5} total mass at month 1 and

Table 3. Adjusted HRs and 95% CIs of preeclampsia-eclampsia associated with 1 quartile increase in PM_{2.5} mixture during pregnancy based on quantile-based g computation.

Air pollutants	Contribution to positive/negative effect % ^c	Positive/negative coefficient β	Overall mixture coefficient ψ (log HR) _d 95% CI	Overall mixture effect HR 95% CI
Positive mixture^a				
PM _{2.5} sulfate	3.79	0.00		
PM _{2.5} organic matter	25.59	0.02		
PM _{2.5} black carbon	70.63	0.05	0.05 (0.03, 0.07)	1.05 (1.03, 1.07)
Negative mixture^b				
PM _{2.5} nitrate	59.80	-0.01	<i>P</i> < 0.001	<i>P</i> < 0.001
PM _{2.5} ammonium	40.20	-0.01		

HR, hazard ratio; CI, confidence interval. Model adjusted for maternal age, race/ethnicity, education, block group household income, smoking and passive smoking status during pregnancy, insurance type, season, and year of infant birth. Zip code was fitted as a random effect.

^aThe positive mixture includes pollutants positively associated with the outcome in the model.

^bThe negative mixture includes pollutants negatively associated with the outcome in the model.

^cThe sum of contribution of all positive/negative pollutants is 100.0%.

^dThe overall mixture coefficient is the sum of coefficients of the positive mixture and negative mixture.

<https://doi.org/10.1371/journal.pmed.1004395.t003>

months 7 to 8, positive associations at month 3, and nonsignificant associations for other gestational months. On the other hand, elevated risks of PE-E were associated with PM_{2.5} exposure in months 2 to 6, with the highest risk identified at month 4 [HR = 1.05 (95% CI [1.03, 1.07] *p* < 0.001)]. In terms of PM_{2.5} constituents, only exposure to PM_{2.5} sulfate (months 1 to 3) and organic matter (months 3 to 6) were associated with higher GH risks. For PE-E, windows of increased risks occurred during late first trimester and second trimester for all PM_{2.5} constituents of interest.

Subgroup analysis (S6 Appendix) was conducted stratified by maternal age, race/ethnicity, household income, smoking status during pregnancy, pre-pregnancy BMI, and primiparity. No substantial increased GH risks associated with PM_{2.5} were observed in stratified analyses. In the PE-E analyses, positive associations were significantly stronger among Hispanic and African American mothers and mothers who lived in low- to middle-income neighborhoods (*p* < 0.05), but no significant evidence of heterogeneity for other potential effect modifiers.




Fig 2. Monthly associations between maternal exposure to PM_{2.5} during pregnancy and hypertensive disorders of pregnancy. *N* = 373,905 for GH cohort; *N* = 386,361 for PE-E cohort. HRs and 95% CIs were calculated for per IQR increment for each air pollutant. Models adjusted for maternal age, race/ethnicity, education, household income, smoking and passive smoking status during pregnancy, parity, insurance type, season, and year of infant birth. CI, confidence interval; GH, gestational hypertension; HR, hazard ratio; IQR, interquartile range; PE-E, preeclampsia-eclampsia.

<https://doi.org/10.1371/journal.pmed.1004395.g002>

Discussion

This is a large retrospective cohort study of 386,361 pregnant women residing in Southern California from 2008 to 2017. Our results show that exposures to PM_{2.5} total mass, organic matter, and black carbon were associated with an increased risk of PE-E, but not with GH. In multi-pollutant models, PM_{2.5} black carbon contributed the most to the overall mixture effects among all PM_{2.5} constituents. The late first trimester and second trimester are likely the most influential time windows for air pollution and PE-E. Furthermore, Hispanic and African American mothers and mothers who live in low- to middle-income neighborhoods were associated with higher risks for PE-E on exposure to PM_{2.5}.

To the best of our knowledge, it is the first study to examine the associations between PM_{2.5} constituents and mixtures and GH and PE-E separately in a large and diverse population in regions with relatively low air pollution levels. Two recent studies in China showed positive associations between PM_{2.5} total mass and GH with null association for PE-E [12,14]. Interestingly, our study observed opposite results of positive associations between PM_{2.5} and PE-E with null associations for GH. Factors that may partially explain the inconsistent results include differences in exposure levels, exposure assessment methods, criteria for HDP diagnosis, and study region and population. For example, the PM_{2.5} concentrations in those studies were approximately 4 to 5 times (around 50 $\mu\text{g}/\text{m}^3$) that of our study. They estimated air pollution exposure based on a hospital's address rather than individual maternal address that may lead to more exposure misclassification. Furthermore, in those studies, the outcome was measured with physician-filled questionnaires instead of using diagnostic codes, which might introduce recall bias. Our findings of positive association of PE-E with exposure to PM_{2.5} are supported by a Swedish study with similar air pollution levels as that of our study (OR = 1.35 (95% CI [1.11, 1.63]) per 5 $\mu\text{g}/\text{m}^3$ increment of PM_{2.5}) [57]. A meta-analysis also reported that exposure to PM_{2.5} enhanced the risk of PE, but not for GH [5]. Our finding of positive associations between PE-E and PM_{2.5} black carbon is also consistent with previous studies [12,57].

Among HDP types (S1 Appendix), HDP with isolated elevated blood pressure with no systemic involvement is GH, and HDP with organ involvement or systemic manifestations (i.e., proteinuria, involvement of kidney, elevated liver enzymes, or decreased platelet count) constitute PE, that also includes superimposed preeclampsia, and hemolysis, elevated liver enzymes, and low platelet count (HELLP) syndrome, whereas, preeclampsia with neurological manifestations is considered as eclampsia. In our analyses, we observed that exposures to PM_{2.5} total mass, black carbon, and organic matter are positively associated with the severe categories in the HDP spectrum (i.e., PE-E), while the milder form of HDP (i.e., GH) may not be affected as such (S4 and S5 Appendix). The differential associations between air pollution and HDP subcategories may be explained by different mechanisms involved with mild and severe varieties of HDP. Increased expression of cytochrome P-450 and induction of stress response enzymes by PM_{2.5} have been documented [58]. The cytochrome P-450 liver enzymes induction pathway could result in rapid clearing of vaso-constrictive cytokines from the system [59]. Therefore, in cases of mild HDP, PM_{2.5} may lead to vasodilatation by inducing breakdown of vasoconstrictive factors and thus not contributing to the risk for GH. The mechanistic pathway for PE-E on the other hand has been suggested to be the result of compromised trophoblast invasion by placental vasculatures, release of placental vasoactive substances leading to marked vasoconstriction and resulting placental hypoxia [60]. PM_{2.5} exposure has been linked to endothelial dysfunction [61]. An imbalance between vasoactive substances, including vasoconstrictive factors (e.g., Thromboxane A2 and endothelin) and vasodilator factors (e.g., prostacyclin and nitric oxide), have been demonstrated as potential mechanisms of PE-E [60]. Therefore, exposure to the ambient pollution could trigger an increased production of placental

vasoconstrictive factors that would overwhelm the cytochrome P-450 enzymes induction system leading to more severe categories of HDP, explaining the positive associations of pollution exposure with PE-E. Previous studies also documented elevated placental biomarker 3-nitrotyrosine and hypomethylated leptin promoters in placental tissue on exposure to PM_{2.5} and black carbon [62,63], which has been linked to high-risk pregnancies such as preeclampsia [64]. These mechanistic pathways are shown in [S7 Appendix](#).

It should be noted that PE was more common among smoking mothers in our study population, which is different from previous studies reporting a protective effect of smoking on preeclampsia risk [65]. It has been proposed that combustion products in tobacco (e.g., carbon monoxide) [66] might be responsible for the protective effect of smoking on PE due to vasodilation. However, the inverse association between smoking and PE might be controverted due to multiple sources of bias, including eligibility criteria, losses to follow-up of women potentially at risk, misclassification, competing events, or incorrect adjustment in previous studies [55]. Further studies are warranted to extricate bias when evaluating the smoking-preeclampsia paradox.

Although several studies examined the associations between air pollution and HDP, studies explored different exposure windows are insufficient; findings are not consistent among studies, including the first, second, or third trimester [4–6]. Overall, the first and the second trimester were more susceptible to PM_{2.5} exposure in our study. Specifically, we found gestational month 3 as the critical exposure period for GH, and months 2 to 6 as the critical exposure period for PE-E. Our results are partially consistent with findings from another study in Southern California, indicating that first trimester exposure to PM_{2.5} was associated with increased odds of HDP [38]. However, this study investigated the PM_{2.5}-HDP relationship among Hispanic population without considering separate HDP outcomes and had a small sample size ($n = 298$). To date, further research to identify the susceptible windows of PM_{2.5} is still needed given the inconclusive results among previous studies.

As both health and economic burdens increase with severity of HDP, more research on the associations between environmental exposure and reproductive health is needed. We used the PAF to assess the public health impact of exposure to PM_{2.5} during pregnancy. Our results showed that a 6.37% absolute reduction of PE-E cases ($N = 1,235$) in our study population would occur if PM_{2.5} exposure levels were reduced to 9 $\mu\text{g}/\text{m}^3$ based on the US EPA air quality standards. Women with PE-E require prolonged hospitalization often including emergent cesarean section, which significantly increase health care costs. In 2003, the average (direct and indirect) PE management cost was estimated at \$11,208 per woman in the United States [67]. PE has been recognized as one of the independent risk factors of future cardiovascular diseases and stroke, it also affects other vital organs, psychological health, and fetal health [68], thus contributing to the long-term social and financial impact of the health care system [69]. Although everyone is affected by air pollution, the risk may disproportionately be higher among women of certain socioeconomic backgrounds, such as women living in poverty and the minority populations in our study [70], subsequently increasing the burden of HDP. Shen and colleagues reported relatively larger effect estimates of PM_{2.5} exposure on HDP among underweight women (BMI <18.5) compared to their counterparts, but the differences between subgroups were not statistically significant [12]. Mobasher and colleagues found that PM_{2.5} exposure was significantly associated with HDP among non-obese women (BMI <30), but not obese women [38]. However, our results did not reveal any heterogeneity between the obese and non-obese subgroups. Further, identifying the main sources of air pollution associated with HDP is also important to initiate actions targeting this highly modifiable risk factor (e.g., reducing fuel emissions) to mitigate those adverse effects on reproductive health and future generations.

The main strengths of this study include the large and diverse study population in Southern California; comprehensive and detailed demographic information and medical records from KPSC EHRs allowed us to control for a number of potential confounders in our analysis; the use of time-to-event models to examine the relationship between HDP and air pollution exposure, and identify critical exposure windows; major PM_{2.5} chemical components from well-validated air pollution models; the use of an innovative statistical method to estimate the joint effects of air pollution mixtures; and consider residential changes during pregnancy, which may enhance the accuracy for the air pollution exposure assessments in this study. Furthermore, while most previous studies used combined GH and PE as the umbrella diagnosis of HDP, our study addressed the gap in the literature to link the effect of air pollution and different expressions of HDP and to understand corresponding pathophysiology.

There are certain limitations in this study. First, air pollution exposure was estimated only for residential outdoors without considering other exposure sources which could also contribute to the observed associations (e.g., indoor exposures, occupational exposures), or activity patterns due to data unavailability, which may lead to exposure misclassification and bias estimated associations in either direction. Future studies utilizing personal air pollution sampling or blood biomarkers for exposure to the pollutants would help alleviate exposure misclassification. Second, identification of disease status solely based on ICD diagnostic codes may lead to potential outcome misclassification [71]. Third, the dates of HDP diagnosis were collected from the EHR based on the routine screening during patients' prenatal visits, which may not actually coincide with the start of the disease process. Nonetheless, the KPSC EHRs include repeated measurements from multiple time points during the entire pregnancy (7 ± 4 times) to minimize the potential bias in the estimates of time-varying exposures and outcomes. Moreover, although several covariates were adjusted, some HDP-related factors were unavailable in our analysis, such as family or personal history of HDP. In addition, further exploration of the severity and different subtypes of HDP (e.g., early-onset PE versus late-onset PE, HELLP syndrome, which is not available in our database) would give additional insights of the effects of air pollution on HDP and mechanisms. Given the complexity of air pollution and related environmental exposures, further research is also warranted to examine the effect of other air pollutants (e.g., nitrogen oxides and sulfur oxides), environmental factors (e.g., traffic noise and meteorological factors) [15,72,73], and their joint effect on HDP. Finally, animal studies examining molecular changes from air pollution exposure may shed light on the causation pathway and pathogenesis of GH and PE.

In conclusion, we found that exposures to PM_{2.5} total mass, organic matter, and black carbon were associated with an increased risk for PE-E, but not for GH. The main effect of increased PE-E risk was driven by PM_{2.5} black carbon (71%) and PM_{2.5} organic matter (26%). The susceptible window of air pollution exposure associated with PE-E is gestational months 2 to 6. Hispanic and African American mothers and mothers who live in low- to middle-income neighborhoods may be more vulnerable to air pollution on HDP risk.

Supporting information

S1 STROBE Checklist. STROBE Statement—Checklist of items that should be included in reports of cohort studies.

(DOCX)

S1 Appendix. American College of Obstetricians and Gynecologists (ACOG) classification of hypertensive disorders in pregnancy (HDP).

(DOCX)

S2 Appendix. Schematic flowchart for the study design with exclusion criteria.
(DOCX)

S3 Appendix. Directed acyclic graph (DAG) conceptualizing the relationship between variables.
(DOCX)

S4 Appendix. Adjusted hazard ratios (HRs) and 95% confidence intervals (CI) of GH and PE-E associated with air pollution during the entire pregnancy in sensitivity analysis.
(DOCX)

S5 Appendix. Monthly associations between hypertensive disorders of pregnancy and maternal exposure to PM_{2.5}.
(DOCX)

S6 Appendix. Adjusted hazard ratios (HRs) and 95% confidence intervals (CI) of GH and PE-E associated with air pollution during the entire pregnancy among population subgroups.
(DOCX)

S7 Appendix. Differential pathway of pollution for GH and PE-E.
(DOCX)

Acknowledgments

The authors thank the patients of Kaiser Permanente for helping to improve care through the use of information collected through our electronic health record systems.

Author Contributions

Conceptualization: Yi Sun, Rashmi Bhuyan, Darios Getahun, Jun Wu.

Data curation: Yi Sun, Vicki Y. Chiu, Darios Getahun, Jun Wu.

Formal analysis: Yi Sun, Rashmi Bhuyan.

Funding acquisition: Darios Getahun, Jun Wu.

Investigation: Yi Sun, Rashmi Bhuyan.

Methodology: Yi Sun, Rashmi Bhuyan, Anqi Jiao, Jeff M. Slezak, David A. Sacks, John Molitor, Tarik Benmarhnia, Jiu-Chiuan Chen, Darios Getahun, Jun Wu.

Project administration: Yi Sun, Chantal C. Avila, Darios Getahun, Jun Wu.

Resources: Darios Getahun, Jun Wu.

Software: Yi Sun, Rashmi Bhuyan, Vicki Y. Chiu, Jun Wu.

Supervision: Yi Sun, Darios Getahun, Jun Wu.

Validation: Yi Sun, Rashmi Bhuyan, Jun Wu.

Visualization: Yi Sun, Rashmi Bhuyan.

Writing – original draft: Yi Sun, Rashmi Bhuyan.

Writing – review & editing: Anqi Jiao, Chantal C. Avila, Vicki Y. Chiu, Jeff M. Slezak, David A. Sacks, John Molitor, Tarik Benmarhnia, Jiu-Chiuan Chen, Darios Getahun, Jun Wu.

References

1. Thangavel P, Park D, Lee Y-C. Recent Insights into Particulate Matter (PM_{2.5})-Mediated Toxicity in Humans: An Overview. *Int J Environ Res Public Health*. 2022; 19(12):7511. <https://doi.org/10.3390/ijerph19127511> PMID: 35742761
2. Fan F, Wang S, Zhang Y, Xu D, Jia J, Li J, et al. Acute Effects of High-Level PM_{2.5} Exposure on Central Blood Pressure. *Hypertension*. 2019; 74(6):1349–1356. <https://doi.org/10.1161/HYPERTENSIONAHA.119.13408> PMID: 31630576.
3. Cao L, Wang L, Wu L, Wang T, Cui X, Yu L, et al. Particulate matter and hypertensive disorders in pregnancy: systematic review and meta-analysis. *Public Health*. 2021; 200:22–32. <https://doi.org/10.1016/j.puhe.2021.08.013> PMID: 34653738
4. Bai W, Li Y, Niu Y, Ding Y, Yu X, Zhu B, et al. Association between ambient air pollution and pregnancy complications: A systematic review and meta-analysis of cohort studies. *Environ Res*. 2020; 185:109471. <https://doi.org/10.1016/j.envres.2020.109471> PMID: 32276169.
5. Sun M, Yan W, Fang K, Chen D, Liu J, Chen Y, et al. The correlation between PM_{2.5} exposure and hypertensive disorders in pregnancy: A Meta-analysis. *Sci Total Environ*. 2020; 703:134985. <https://doi.org/10.1016/j.scitotenv.2019.134985> PMID: 31731170.
6. Alvarado-Jiménez D, Donzelli G, Morales-Suárez-Varela M. A systematic review on the association between exposure to air particulate matter during pregnancy and the development of hypertensive disorders of pregnancy and gestational diabetes mellitus. *Rev Environ Health*. 2023 May 4. <https://doi.org/10.1515/reveh-2022-0258> PMID: 37141623.
7. Decrue F, Townsend R, Miller MR, Newby DE, Reynolds RM. Ambient air pollution and maternal cardiovascular health in pregnancy. *Heart*. 2023:heartjnl-2022-322259. <https://doi.org/10.1136/heartjnl-2022-322259> PMID: 37217298
8. Chappell LC, Cluver CA, Kingdom J, Tong S. Pre-eclampsia. *Lancet*. 2021; 398(10297):341–354. [https://doi.org/10.1016/S0140-6736\(20\)32335-7](https://doi.org/10.1016/S0140-6736(20)32335-7) PMID: 34051884.
9. Ferrazzani S, Luciano R, Garofalo S, D'Andrea V, De Carolis S, De Carolis MP, et al. Neonatal outcome in hypertensive disorders of pregnancy. *Early Hum Dev*. 2011; 87(6):445–449. <https://doi.org/10.1016/j.earlhumdev.2011.03.005> PMID: 21497462.
10. Li F, Wang T, Chen L, Zhang S, Chen L, Qin J. Adverse pregnancy outcomes among mothers with hypertensive disorders in pregnancy: A meta-analysis of cohort studies. *Pregnancy Hypertens*. 2021; 24:107–117. <https://doi.org/10.1016/j.pregphy.2021.03.001> PMID: 33813363.
11. Soma-Pillay P, Nelson-Piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr*. 2016; 27(2):89–94. <https://doi.org/10.5830/CVJA-2016-021> PMID: 27213856
12. Shen Y, Yu G, Liu C, Wang W, Kan H, Zhang J, et al. Prenatal Exposure to PM_{2.5} and Its Specific Components and Risk of Hypertensive Disorders in Pregnancy: A Nationwide Cohort Study in China. *Environ Sci Tech*. 2022; 56(16):11473–11481. <https://doi.org/10.1021/acs.est.2c01103> PMID: 35914180.
13. Su X, Zhao Y, Yang Y, Hua J. Correlation between exposure to fine particulate matter and hypertensive disorders of pregnancy in Shanghai, China. *Environ Health*. 2020; 19(1):101. <https://doi.org/10.1186/s12940-020-00655-1> PMID: 32943053; PMCID: PMC7499904.
14. Yan M, Liu N, Fan Y, Ma L, Guan T. Associations of pregnancy complications with ambient air pollution in China. *Ecotoxicol Environ Saf*. 2022; 241:113727. <https://doi.org/10.1016/j.ecoenv.2022.113727> PMID: 35679731.
15. Pedersen M, Stayner L, Slama R, Sørensen M, Figueras F, Nieuwenhuijsen MJ, et al. Ambient Air Pollution and Pregnancy-Induced Hypertensive Disorders. *Hypertension*. 2014; 64(3):494–500. <https://doi.org/10.1161/hypertensionaha.114.03545> PMID: 24935943.
16. Li C, Xu JJ, Zhou FY, Ge YZ, Qin KZ, Huang HF, et al. Effects of Particulate Matter on the Risk of Gestational Hypertensive Disorders and Their Progression. *Environ Sci Tech*. 2023; 57(12):4930–9. <https://doi.org/10.1021/acs.est.2c06573> PMID: 36913485; PMCID: PMC10061918.
17. Vinikoor-Imler LC, Gray SC, Edwards SE, Miranda ML. The effects of exposure to particulate matter and neighbourhood deprivation on gestational hypertension. *Paediatr Perinat Epidemiol*. 2012; 26(2):91–100. <https://doi.org/10.1111/j.1365-3016.2011.01245.x> PMID: 22324494
18. Yang R, Luo D, Zhang YM, Hu K, Qian ZM, Hu LQ, et al. Adverse Effects of Exposure to Fine Particulate Matters and Ozone on Gestational Hypertension. *Curr Med Sci*. 2019; 39(6):1019–1028. <https://doi.org/10.1007/s11596-019-2137-9> PMID: 31845236.
19. Yuan Z, Wang H-J, Li Q, Su T, Yang J, Chen J, et al. Risk of De Novo Hypertensive Disorders of Pregnancy After Exposure to PM₁ and PM_{2.5} During the Period From Preconception to Delivery: Birth Cohort Study. *JMIR Public Health Surveill*. 2023; 9:e41442. <https://doi.org/10.2196/41442> PMID: 36689262; PMCID: PMC9903185.

20. Xu X, Hu H, Ha S, Roth J. Ambient air pollution and hypertensive disorder of pregnancy. *J Epidemiol Community Health*. 2014; 68(1):13. <https://doi.org/10.1136/jech-2013-202902> PMID: 24022815; PMCID: PMC3878158.
21. Michikawa T, Morokuma S, Fukushima K, Ueda K, Takeuchi A, Kato K, et al. A register-based study of the association between air pollutants and hypertensive disorders in pregnancy among the Japanese population. *Environ Res*. 2015; 142:644–650. <https://doi.org/10.1016/j.envres.2015.08.024> PMID: 26340650.
22. Savitz DA, Elston B, Bobb JF, Clougherty JE, Dominici F, Ito K, et al. Ambient Fine Particulate Matter, Nitrogen Dioxide, and Hypertensive Disorders of Pregnancy in New York City. *Epidemiology*. 2015; 26(5):748–57. <https://doi.org/10.1097/EDE.0000000000000349> PMID: 26237745; PMCID: PMC4719124.
23. Davvand P, Figueras F, Basagana X, Beelen R, Martinez D, Cirach M, et al. Ambient air pollution and preeclampsia: a spatiotemporal analysis. *Environ Health Perspect*. 2013; 121(11–12):1365–71. <https://doi.org/10.1289/ehp.1206430> PMID: 24021707; PMCID: PMC3855505 interests.
24. Lee PC, Roberts JM, Catov JM, Talbott EO, Ritz B. First trimester exposure to ambient air pollution, pregnancy complications and adverse birth outcomes in Allegheny County, PA. *Matern Child Health J*. 2013; 17(3):545–55. <https://doi.org/10.1007/s10995-012-1028-5> PMID: 22544506; PMCID: PMC3636771.
25. Jia L, Liu Q, Hou H, Guo G, Zhang T, Fan S, et al. Association of Ambient air Pollution with risk of pre-eclampsia during pregnancy: a retrospective cohort study. *BMC Public Health*. 2020; 20(1):1663. <https://doi.org/10.1186/s12889-020-09719-w> PMID: 33153479; PMCID: PMC7643463.
26. Wu J, Ren C, Delfino R, Chung J, Wilhelm M, Ritz B. Association between local traffic-generated air pollution and preeclampsia and preterm delivery in the south coast air basin of California. *Environ Health Perspect*. 2009 Nov; 117(11):1773–9. <https://doi.org/10.1289/ehp.0800334> PMID: 20049131; PMCID: PMC2801174.
27. Mendola P, Wallace M, Liu D, Robledo C, Männistö T, Grantz KL. Air pollution exposure and pre-eclampsia among US women with and without asthma. *Environ Res*. 2016; 148:248–255. <https://doi.org/10.1016/j.envres.2016.04.004> PMID: 27085496
28. Weber KA, Yang W, Lyons E, Stevenson DK, Padula AM, Shaw GM. Greenspace, Air Pollution, Neighborhood Factors, and Preeclampsia in a Population-Based Case-Control Study in California. LID—10.3390/ijerph18105127 LID—5127. *Int J Environ Res Public Health*. 2021 May 12; 18(10):5127. <https://doi.org/10.3390/ijerph18105127> PMID: 34066190; PMCID: PMC8151979.
29. ACOG. Gestational Hypertension and Preeclampsia: ACOG Practice Bulletin, Number 222. *Obstet Gynecol*. 2020; 135(6).
30. Melamed N, Ray JG, Hladunewich M, Cox B, Kingdom JC. Gestational Hypertension and Preeclampsia: Are They the Same Disease? *J Obstet Gynaecol Can*. 2014; 36(7):642–647. [https://doi.org/10.1016/S1701-2163\(15\)30545-4](https://doi.org/10.1016/S1701-2163(15)30545-4) PMID: 25184984
31. Ying W, Catov JM, Ouyang P. Hypertensive Disorders of Pregnancy and Future Maternal Cardiovascular Risk. *J Am Heart Assoc*. 2018; 7(17):e009382. <https://doi.org/10.1161/JAHA.118.009382> PMID: 30371154; PMCID: PMC6201430.
32. Phipps EA, Thadhani R, Benzong T, Karumanchi SA. Pre-eclampsia: pathogenesis, novel diagnostics and therapies. *Nat Rev Nephrology*. 2019; 15(5):275–89. <https://doi.org/10.1038/s41581-019-0119-6> PMID: 30792480; PMCID: PMC6472952.
33. Nobles CJ, Williams A, Ouidir M, Sherman S, Mendola P. Differential Effect of Ambient Air Pollution Exposure on Risk of Gestational Hypertension and Preeclampsia. *Hypertension*. 2019; 74(2):384–90. <https://doi.org/10.1161/HYPERTENSIONAHA.119.12731> PMID: 31230552; PMCID: PMC6620155.
34. Bell ML, Ebisu K, Leaderer BP, Gent JF, Lee HJ, Koutrakis P, et al. Associations of PM2.5 constituents and sources with hospital admissions: analysis of four counties in Connecticut and Massachusetts (USA) for persons >= 65 years of age. *Environ Health Perspect*. 2014; 122(2):138–44. <https://doi.org/10.1289/ehp.1306656> PMID: 24213019; PMCID: PMC3915260 interests.
35. Thurston G, Awe Y, Ostro B, Sanchez-Triana E. Are All Air Pollution Particles Equal? How Constituents and Sources of Fine Air Pollution Particles (PM2.5) Affect Health. World Bank Publications—Reports 36269 <https://ideasrepec.org/p/wbk/wboper/36269.html>. 2021.
36. Sun Y, Li X, Benmarhnia T, Chen JC, Avila C, Sacks DA, et al. Exposure to air pollutant mixture and gestational diabetes mellitus in Southern California: Results from electronic health record data of a large pregnancy cohort. *Environ Int*. 2022; 158:106888. <https://doi.org/10.1016/j.envint.2021.106888> PMID: 34563749; PMCID: PMC9022440.
37. Sun Y, Headon KS, Jiao A, Slezak JM, Avila CC, Chiu VY, et al. Association of Antepartum and Postpartum Air Pollution Exposure With Postpartum Depression in Southern California. *JAMA Netw Open*. 2023; 6(10):e2338315. <https://doi.org/10.1001/jamanetworkopen.2023.38315> PMID: 37851440;

PMCID: PMC10585409 institution from Pfizer, Dynavax Technologies, and ALK outside the submitted work. Ms Avila reported receiving grants from Kaiser Permanente Southern California, Department of Research and Evaluation, during the conduct of the study. Ms Chiu reported receiving funding from Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) grants to her institution (Kaiser Permanente Southern California) outside the submitted work. Dr Getahun reported receiving funding from grants to his institution from the NICHD outside the submitted work and grants from Bayer Pharma AG, Johnson & Johnson, and the Garfield Memorial Fund outside the submitted work. No other disclosures were reported.

38. Mobasher Z, Salam MT, Goodwin TM, Lurmann F, Ingles SA, Wilson ML. Associations between ambient air pollution and Hypertensive Disorders of Pregnancy. *Environ Res*. 2013; 123:9–16. <https://doi.org/10.1016/j.envres.2013.01.006> PMID: 23522615
39. ACOG. ACOG/AIUM/SMFM Committee Opinion 700: Methods for Estimating Due Date. 2017.
40. van Donkelaar A, Martin RV, Li C, Burnett RT. Regional Estimates of Chemical Composition of Fine Particulate Matter Using a Combined Geoscience-Statistical Method with Information from Satellites, Models, and Monitors. *Environ Sci Tech*. 2019; 53(5):2595–2611. <https://doi.org/10.1021/acs.est.8b06392> PMID: 30698001.
41. Meng J, Li C, Martin RV, van Donkelaar A, Hystad P, Brauer M. Estimated Long-Term (1981–2016) Concentrations of Ambient Fine Particulate Matter across North America from Chemical Transport Modeling, Satellite Remote Sensing, and Ground-Based Measurements. *Environ Sci Tech*. 2019; 53(9):5071–5079. <https://doi.org/10.1021/acs.est.8b06875> PMID: 30995030.
42. Brown MA, Magee LA, Kenny LC, Karumanchi SA, McCarthy FP, Saito S, et al. Hypertensive Disorders of Pregnancy: ISSHP Classification, Diagnosis, and Management Recommendations for International Practice. *Hypertension*. 2018; 72(1):24–43. <https://doi.org/10.1161/HYPERTENSIONAHA.117.10803> PMID: 29899139.
43. Wilkerson RG, Ogunbodede AC. Hypertensive Disorders of Pregnancy. *Emerg Med Clin N Am*. 2019; 37(2):301–316. <https://doi.org/10.1016/j.emc.2019.01.008> PMID: 30940374.
44. Umesawa M, Kobashi G. Epidemiology of hypertensive disorders in pregnancy: prevalence, risk factors, predictors and prognosis. *Hypertension Res*. 2017; 40(3):213–220. <https://doi.org/10.1038/hr.2016.126> PMID: 27682655.
45. U.S. Census Bureau. ZIP Code Tabulation Areas (ZCTAs). 2010.[accessed on 2022 Jan 31]. Available from: <https://www.census.gov/programs-surveys/geography/guidance/geo-areas/zctas.html>. 2010.
46. Sun Y, Li X, Benmarhnia T, Chen JC, Avila C, Sacks DA, et al. Exposure to air pollutant mixture and gestational diabetes mellitus in Southern California: Results from electronic health record data of a large pregnancy cohort. *Environ Int*. 2021a; 158:106888. <https://doi.org/10.1016/j.envint.2021.106888> PMID: 34563749
47. Keil AP, Buckley JP, O'Brien KM, Ferguson KK, Zhao S, White AJ. A Quantile-Based g-Computation Approach to Addressing the Effects of Exposure Mixtures. *Environ Health Perspect*. 2020; 128(4):47004. <https://doi.org/10.1289/EHP5838> PMID: 32255670; PMCID: PMC7228100.
48. Murray EJ, Caniglia EC, Petito LC. Causal survival analysis: A guide to estimating intention-to-treat and per-protocol effects from randomized clinical trials with non-adherence. *Res Methods Med Health Sci*. 2020; 2(1):39–49. <https://doi.org/10.1177/2632084320961043>
49. Suresh K, Severn C, Ghosh D. Survival prediction models: an introduction to discrete-time modeling. *BMC Med Res Methodol*. 2022; 22(1):207. <https://doi.org/10.1186/s12874-022-01679-6> PMID: 35883032; PMCID: PMC9316420.
50. U.S. Environmental Protection Agency. Ambient Air Quality Standards (NAAQS) for PM. 2024. [accessed on 2024 Feb 28]. Available from: <https://www.federalregister.gov/documents/2024/03/06/2024-02637/reconsideration-of-the-national-ambient-air-quality-standards-for-particulate-matterNational>.
51. WHO. World Health Organization global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Geneva: World Health Organization; 2021.
52. Diel A, Farthmann J. Gestational hypertension and advanced maternal age. *Lancet*. 2015; 386(10004):1627–1628. [https://doi.org/10.1016/S0140-6736\(15\)00532-2](https://doi.org/10.1016/S0140-6736(15)00532-2) PMID: 26595630.
53. Tanaka M, Jaamaa G, Kaiser M, Hills E, Soim A, Zhu M, et al. Racial disparity in hypertensive disorders of pregnancy in New York State: a 10-year longitudinal population-based study. *Am J Public Health*. 2007; 97(1):163–70. <https://doi.org/10.2105/AJPH.2005.068577> PMID: 17138931; PMCID: PMC1716255.
54. Mattsson K, Juárez S, Malmqvist E. Influence of Socio-Economic Factors and Region of Birth on the Risk of Preeclampsia in Sweden. *Int J Environ Res Public Health*. 2022; 19(7):4080. <https://doi.org/10.3390/ijerph19074080> PMID: 35409763
55. Rodríguez-Lopez M, Escobar MF, Merlo J, Kaufman JS. Reevaluating the protective effect of smoking on preeclampsia risk through the lens of bias. *J Hum Hypertens*. 2023; 37(5):338–44. <https://doi.org/10.1038/s41371-023-00827-9> PMID: 37041252; PMCID: PMC10156598.

56. Shen M, Smith GN, Rodger M, White RR, Walker MC, Wen SW. Comparison of risk factors and outcomes of gestational hypertension and pre-eclampsia. *PLoS ONE*. 2017; 12(4):e0175914. <https://doi.org/10.1371/journal.pone.0175914> PMID: 28437461; PMCID: PMC5402970.
57. Mandakh Y, Rittner R, Flanagan E, Oudin A, Isaxon C, Familiari M, et al. Maternal Exposure to Ambient Air Pollution and Risk of Preeclampsia: A Population-Based Cohort Study in Scania, Sweden. *Int J Environ Res Public Health*. 2020;17(5). <https://doi.org/10.3390/ijerph17051744> PMID: 32155988; PMCID: PMC7084298.
58. Ghio AJ, Carraway MS, Madden MC. Composition of air pollution particles and oxidative stress in cells, tissues, and living systems. *J Toxicol Environ Health B Crit Rev*. 2012; 15(1):1–21. <https://doi.org/10.1080/10937404.2012.632359> PMID: 22202227.
59. Zhang JM, An J. Cytokines, inflammation, and pain. *Spring*. 2007; 45(2):27–37. <https://doi.org/10.1097/aia.0b013e318034194e> PMID: 17426506
60. Possomato-Vieira JS, Khalil RA. Mechanisms of Endothelial Dysfunction in Hypertensive Pregnancy and Preeclampsia. *Adv Pharmacol*. 2016; 77:361–431. <https://doi.org/10.1016/bs.apha.2016.04.008> PMID: 27451103; PMCID: PMC4965238.
61. Brook RD, Rajagopalan S. Chronic Air Pollution Exposure and Endothelial Dysfunction: What You Can't See—Can Harm You. *J Am Coll Cardiol*. 2012; 60(21):2167–2169. <https://doi.org/10.1016/j.jacc.2012.08.974> PMID: 23103034
62. Saenen ND, Vrijens K, Janssen BG, Roels HA, Neven KY, Vanden Berghe W, et al. Lower Placental Leptin Promoter Methylation in Association with Fine Particulate Matter Air Pollution during Pregnancy and Placental Nitrosative Stress at Birth in the ENVIRONAGE Cohort. *Environ Health Perspect*. 2017; 125(2):262–268. <https://doi.org/10.1289/EHP38> PMID: 27623604; PMCID: PMC5289914 interests.
63. Saenen ND, Vrijens K, Janssen BG, Madhloum N, Peusens M, Gyselaers W, et al. Placental Nitrosative Stress and Exposure to Ambient Air Pollution During Gestation: A Population Study. *Am J Epidemiol*. 2016; 184(6):442–449. <https://doi.org/10.1093/aje/kww007> PMID: 27601048.
64. Bosco C, Gonzalez J, Gutierrez R, Parra-Cordero M, Barja P, Rodrigo R. Oxidative damage to pre-eclamptic placenta: immunohistochemical expression of VEGF, nitrotyrosine residues and von Willebrand factor. *J Matern-Fetal Neonatal Med*. 2012; 25(11):2339–2345. <https://doi.org/10.3109/14767058.2012.695823> PMID: 22612323.
65. Wang J, Yang W, Xiao W, Cao S. The association between smoking during pregnancy and hypertensive disorders of pregnancy: A systematic review and meta-analysis. *Int J Gynecol Obstet*. 2022; 157(1):31–41. <https://doi.org/10.1002/ijgo.13709> PMID: 33864264
66. Wikstrom AK, Stephansson O, Chattingius S. Tobacco use during pregnancy and preeclampsia risk: effects of cigarette smoking and snuff. *Hypertension*. 2010; 55(5):1254–1259. <https://doi.org/10.1161/HYPERTENSIONAHA.109.147082> PMID: 20231527.
67. Fox A, McHugh S, Browne J, Kenny LC, Fitzgerald A, Khashan AS, et al. Estimating the Cost of Preeclampsia in the Healthcare System: Cross-Sectional Study Using Data From SCOPE Study (Screening for Pregnancy End Points). *Hypertension*. 2017; 70(6):1243–1249. <https://doi.org/10.1161/HYPERTENSIONAHA.117.09499> PMID: 29084880.
68. Lu HQ, Hu R. Lasting Effects of Intrauterine Exposure to Preeclampsia on Offspring and the Underlying Mechanism. *AJP Rep*. 2019; 9(3):e275–e291. <https://doi.org/10.1055/s-0039-1695004> PMID: 31511798; PMCID: PMC6736667.
69. Turbeville HR, Sasser JM. Preeclampsia beyond pregnancy: long-term consequences for mother and child. *Am J Physiol Renal Physiol*. 2020; 318(6):F1315–F1326. <https://doi.org/10.1152/ajprenal.00071.2020> PMID: 32249616
70. ACOG. American College of Obstetricians and Gynecologists. Low-Dose Aspirin Use for the Prevention of Preeclampsia and Related Morbidity and Mortality. Practice Advisory. 2021. [accessed on 2023 Dec 10]. Available from: <https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2021/12/low-dose-aspirin-use-for-the-prevention-of-preeclampsia-and-related-morbidity-and-mortality>.
71. Jacobsen SJ, Demissie K, Strauss JA, Chen W, Fassett MJ, Rhoads GG, et al. Accuracy of reporting maternal and infant perinatal service system coding and clinical utilization coding. *J Med Stat Info*. 2013; 1(1):3. <https://doi.org/10.7243/2053-7662-1-3>
72. NTP. National Toxicology Program. NTP monograph on the systematic review of traffic-related air pollution and hypertensive disorders of pregnancy. NTP. Monograph. 2019;Dec(7). <https://doi.org/10.22427/NTP-MGRAPH-7> PMID: 33560269; PMCID: PMC8054471.
73. Pedersen M, Halldorsson TI, Olsen SF, Hjortebjerg D, Ketzel M, Grandström C, et al. Impact of Road Traffic Pollution on Pre-eclampsia and Pregnancy-induced Hypertensive Disorders. *Epidemiology*. 2017; 28(1):99–106. <https://doi.org/10.1097/EDE.0000000000000555> PMID: 27648591; PMCID: PMC5380106.