

# The Vitamin D Metabolite Diagnostic Ratio Associates With Phenotypic Traits of Idiopathic Hypercalciuria



Nasser A. Dhayat<sup>1,2</sup>, Cédric Mattmann<sup>3</sup>, Harald Seeger<sup>4</sup>, Alexander Ritter<sup>4</sup>, Thomas Ernandez<sup>5</sup>, Catherine Stoermann-Chopard<sup>5</sup>, Florian Buchkremer<sup>6</sup>, Stephan Segerer<sup>6</sup>, Beat Roth<sup>7,13</sup>, Gregoire Wuerzner<sup>8</sup>, Carsten A. Wagner<sup>2,9</sup>, Olivier Bonny<sup>2,8,10,11</sup>, Albrecht W. Popp<sup>12</sup>, Bruno Vogt<sup>3</sup>, Matteo Bargagli<sup>2,3,14</sup> and Daniel G. Fuster<sup>2,3,14</sup>

<sup>1</sup>B. Braun Medical Care AG, Nephrology and Dialysis Care Center, Hochfelden, Zürich, Switzerland; <sup>2</sup>Swiss National Centre of Competence in Research Kidney.CH, Zürich, Switzerland; <sup>3</sup>Department of Nephrology and Hypertension, Inselspital, Bern University Hospital and University of Bern, Bern, Switzerland; <sup>4</sup>Division of Nephrology, University Hospital Zürich, Zürich, Switzerland; <sup>5</sup>Service of Nephrology, Geneva University Hospitals, Geneva, Switzerland; <sup>6</sup>Division of Nephrology, Kantonsspital Aarau, Aarau, Switzerland; <sup>7</sup>Department of Urology, Lausanne University Hospital, CHUV, University of Lausanne, Switzerland; <sup>8</sup>Service of Nephrology and Hypertension, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland; <sup>9</sup>Institute of Physiology, University of Zürich, Zürich, Switzerland; <sup>10</sup>Department of Biomedical Sciences, University of Lausanne, Lausanne, Switzerland; <sup>11</sup>Service of Nephrology, Fribourg State Hospital and University of Fribourg, Fribourg, Switzerland; <sup>12</sup>Department of Osteoporosis, Inselspital, Bern University Hospital and University of Bern, Bern, Switzerland; and <sup>13</sup>Department of Urology, Inselspital, Bern University Hospital and University of Bern, Bern, Switzerland

**Introduction:** Underlying mechanisms for hypercalciuria remain unknown in most cases; thus, the designation “idiopathic.” We hypothesized that the vitamin D-inactivating enzyme, CYP24A1 contributes to the pathogenesis of hypercalciuria in kidney stone formers.

**Methods:** We conducted association analyses between CYP24A1 activity, estimated by the vitamin D metabolite diagnostic ratio (25(OH) vitamin D<sub>3</sub>/total 24,25 (OH)<sub>2</sub> vitamin D ratio; VMDR), and the phenotype of participants in 2 observational cohorts of kidney stone formers, the Swiss Kidney Stone Cohort (SKSC) and the Bern Kidney Stone Registry (BKSR). Circulating 25(OH)- and 24,25 (OH)<sub>2</sub> vitamin D were quantified using a validated liquid chromatography tandem mass spectrometry assay.

**Results:** A total of 974 participants were included in the analysis. We found a positive association of VMDR (and hence negative association of CYP24A1 activity) with total ( $\beta$  0.009 mmol/l; 95% confidence interval [CI]: 0.002, 0.016;  $P = 0.02$ ) and ionized plasma calcium ( $\beta$  0.005 mmol/l; 95% CI: 0.002, 0.008;  $P < 0.01$ ), absolute and fractional excretion of urinary calcium ( $\beta$  0.054 mmol/24h; 95% CI: 0.010, 0.097;  $P = 0.02$  and  $\beta$  0.046%; 95% CI: 0.018, 0.074;  $P < 0.01$ , respectively). Further, VMDR was associated with an increased likelihood of forming calcium oxalate dihydrate stones (Odds ratio [OR] 1.64; 95% CI: 1.22, 2.35;  $P < 0.01$ ) and reduced bone mineral density (BMD) at the femoral neck ( $\beta$   $-0.005$  g/cm<sup>2</sup>; 95% CI:  $-0.010$ ,  $-0.001$ ;  $P = 0.04$ ). The described associations became stronger when the analysis was confined to idiopathic calcium stone formers.

**Conclusion:** Our study reveals that CYP24A1 activity, estimated by VMDR, is associated with clinical traits previously linked to idiopathic hypercalciuria.

*Kidney Int Rep* (2024) 9, 1072–1082; <https://doi.org/10.1016/j.ekir.2024.01.004>

KEYWORDS: calcium; CYP24A1; kidney stones; nephrolithiasis; vitamin D

© 2024 International Society of Nephrology. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

**Correspondence:** Daniel G. Fuster, Department of Nephrology and Hypertension, Inselspital, Bern University Hospital, University of Bern, Freiburgstrasse 15, 3010 Bern, Switzerland. E-mail: [daniel.fuster@insel.ch](mailto:daniel.fuster@insel.ch)

<sup>14</sup>MB and DGF contributed equally.

Received 18 October 2023; revised 5 December 2023; accepted 2 January 2024; published online 10 January 2024

## See Commentary on Page 743

Kidney stone formation is complex and depends on dietary, environmental, and genetic factors.<sup>1</sup> Eighty percent to 90% of stones are composed of calcium oxalate, calcium phosphate, or a mixture of both.<sup>2,3</sup> Hypercalciuria, defined as increased urinary

calcium excretion, is the most frequent proliothogenic abnormality encountered in patients with kidney stones and is present in up to 80% of affected patients.<sup>3,4</sup> Metabolic work-up of hypercalciuric stone formers rarely reveals an underlying systemic cause, and thus in most cases, hypercalciuria is defined as “idiopathic.”<sup>5</sup> The phenotype of idiopathic hypercalciuria can be replicated by exogenous administration of 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> to healthy individuals.<sup>6</sup> In contrast, the P450 inhibitor ketoconazole, which attenuates 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> synthesis, normalized the clinical features of patients with idiopathic hypercalciuria.<sup>7</sup> Circulating 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> is determined by the rate of synthesis from its substrate, 25(OH) vitamin D<sub>3</sub>, through the activity of 25(OH)D-1-hydroxylase (CYP27B1), and its degradation by 1,25(OH)<sub>2</sub>D-24-hydroxylase (CYP24A1).<sup>8</sup> CYP24A1 is a P450 enzyme expressed in target tissues and is strongly up-regulated by its substrate 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub>. In addition to 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub>, CYP24A1 also inactivates its precursor 25(OH) vitamin D. Homozygous pathogenic variants in *CYP24A1* cause idiopathic infantile hypercalcemia, a condition first described in children undergoing vitamin D supplementation characterized by failure to thrive, dehydration, hypercalcemia, and nephrocalcinosis.<sup>9</sup> Subsequently, biallelic pathogenic variants in *CYP24A1* were also discovered in adults with elevated vitamin D, suppressed parathyroid hormone (PTH), hypercalcemia, hypercalciuria, osteopenia, and recurrent calcium nephrolithiasis.<sup>10,11</sup> Individuals with heterozygous pathogenic *CYP24A1* variants exhibit an intermediary phenotype characterized by recurrent calcium nephrolithiasis, normocalcemic hypercalciuria, high-normal 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub>, and low-normal PTH, a constellation typically encountered in idiopathic hypercalciuria, suggesting a gene dosage effect for pathogenic *CYP24A1* variants.<sup>10,12,13</sup>

The ratio between 25(OH)- and 24,25(OH)<sub>2</sub> vitamin D (VMDR) is considered as a proxy of CYP24A1-mediated vitamin D clearance.<sup>13-15</sup> In a cohort of 153 first-time calcium stone formers, increased serum calcium and 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub>, a lower serum 24,25(OH)<sub>2</sub> vitamin D / 25(OH) vitamin D ratio but no difference in urine calcium was observed compared to non-stone formers.<sup>16</sup>

The association of CYP24A1 activity, estimated by the VMDR, with stone composition and comprehensive clinical traits linked to kidney stone disease, has not been studied yet. Therefore, the diagnostic utility of the VMDR in kidney stone formers remains unclear. Given the high phenotypic similarity between carriers of pathogenic *CYP24A1* variants and kidney stone formers with idiopathic hypercalciuria, we hypothesized that reduced CYP24A1 activity may be an

important cause of idiopathic hypercalciuria. To this end, we assessed the association of VMDR with the clinical phenotype of participants in 2 large prospective observational Swiss cohorts of kidney stone formers.

## METHODS

### Study Population

The SKSC is an investigator-initiated prospective, multi-center, observational study of patients with kidney stones recruited at the nephrology outpatient clinics of 6 tertiary care nephrology centers in Switzerland (Aarau, Basel, Bern, Geneva, Lausanne, and Zürich).<sup>17</sup> The BKSr includes kidney stone formers recruited at the nephrology outpatient clinic of the Department of Nephrology and Hypertension at the Bern University Hospital, Bern, Switzerland.<sup>18</sup> SKSC and BKSr adhered to the Declaration of Helsinki and were approved by the responsible cantonal ethical committees (approval #BE 173/13 and #BE 95/06, respectively). Inclusion criteria for both cohorts are written informed consent, age  $\geq 18$  years,  $\geq 2$  past kidney stone events or 1 past stone event combined with additional risk factors for stone recurrence, such as first stone episode  $< 25$  years, positive family history, stones other than calcium oxalate, bilateral or multiple stones or nephrocalcinosis detected by imaging, single kidney or chronic kidney disease (estimated glomerular filtration rate  $< 60$  ml/min), metabolic syndrome, gout, or osteoporosis. A stone event was defined as a symptomatic stone event with visible passage of a stone (with or without accompanying typical symptoms), or urological intervention of a symptomatic or asymptomatic stone. At the baseline visit, demographic and anthropometric data, comorbidities and stone composition analysis results were collected, and a comprehensive blood and 24-hour urine metabolic workup performed. In addition, biobank plasma samples were collected at the baseline visit, immediately frozen and stored at  $-80$  °C. Stone formers with end stage kidney disease were excluded from this study. A total of 974 participants (560 SKSC participants and 414 BKSr participants) met the eligibility criteria and had biobank plasma available for the analysis of vitamin D metabolites.

### Measurements and Definitions

Plasma 25(OH) vitamin D<sub>2</sub> and D<sub>3</sub>, and total plasma 24,25(OH)<sub>2</sub> vitamin D were measured by an established, highly sensitive and specific liquid chromatography tandem mass spectrometry method at the Mayo Clinic Central Laboratories, Rochester, MN.<sup>19,20</sup> 25(OH) vitamin D<sub>2</sub> was not detected in any of the participants, thus the VMDR was calculated as 25(OH) vitamin D<sub>3</sub>/total 24,25(OH)<sub>2</sub> vitamin D. All other blood and urinary parameters of BKSr and SKSC participants

were measured centrally by standard clinical laboratory methods at the Central Laboratory of the Bern University Hospital, Bern, Switzerland. Assay characteristics for the measurements of PTH, C-terminal fibroblast growth factor 23 (cFGF23) and  $1,25(\text{OH})_2$  vitamin  $\text{D}_3$  were previously described.<sup>21</sup> Estimated glomerular filtration rate was determined with the Chronic Kidney Disease-Epidemiology Collaboration 2009 equation.<sup>22</sup> The mean value of two 24-hour urine collections on consecutive days was used to calculate mean 24-hour urinary calcium excretion. Fractional excretion of calcium was determined from blood and urine samples collected on the same day. Osteodensitometry at the lumbar spine and the femoral neck was performed in all BKSr participants at the time point of metabolic work-up at the Department of Osteoporosis of the Bern University Hospital, Bern, Switzerland, by dual-energy X-ray absorptiometry (DEXA; Hologic QDR 4500A, Hologic, Bedford, MA).<sup>23</sup> Kidney stone composition was determined by Fourier-transform infrared spectroscopy. In participants with several stone composition analyses available, the most recent analysis prior to metabolic work-up was included in the analysis. Urine relative supersaturations of calcium oxalate and brushite (calcium phosphate) were calculated using the EQUIL2 program.<sup>24</sup> Data related to loop and thiazide diuretics and medications that could potentially influence plasma  $25(\text{OH})$  vitamin  $\text{D}_3$  concentration were collected. These included vitamin D supplementation (cholecalciferol, ergocalciferol, calcifediol, and alfacalcidol), carbamazepine, oxcarbazepine, clonazepam, St. John's wort, ritonavir, efavirenz, tenofovir, emtricitabine, glucocorticoids, rifampin, ketoconazole, and calcium antagonists (felodipine, amlodipine, nifedipine, lercanidipine). Noncalcium stones were defined as stone composition containing  $\geq 50\%$  uric acid, struvite, or cystine.

### Statistical Analyses

Continuous variables were reported as medians  $\pm$  interquartile ranges or means  $\pm$  SD. Categorical variables were reported as counts and percentages, as appropriate. All values were first analyzed by descriptive statistics. All statistical tests were 2-sided, and a  $P$ -value  $< 0.05$  was considered statistically significant. Unadjusted and adjusted linear and logistic regression analyses of CYP24A1 activity, measured by the VMDR, were conducted as predictor variables with appropriately transformed outcome variables. Age, sex, body mass index, estimated glomerular filtration rate, and plasma  $25(\text{OH})$  vitamin  $\text{D}_3$  were considered potential confounders and included in the multivariable model. These variables were selected for their known

interaction with the outcome variables and total plasma  $24,25(\text{OH})_2$  vitamin D concentration by affecting  $1,25(\text{OH})_2$  vitamin  $\text{D}_3$  synthesis. To further assess the association between clinical traits of idiopathic hypercalciuria and the VMDR in patients with or without low  $25(\text{OH})$  vitamin  $\text{D}_3$ , we stratified the study population in 2 subgroups ( $< 20$  or  $\geq 20$  ng/ml). In addition, we performed sensitivity analyses for medications (drugs modulating plasma  $25(\text{OH})$  vitamin  $\text{D}_3$  concentration, loop and thiazide diuretics), urinary sodium excretion, which can directly affect urinary calcium excretion and total calcium balance,<sup>25</sup> and for month of VMDR measurement, to account for fluctuations related to seasonality. A fully-adjusted subgroup analysis was conducted specifically on idiopathic calcium stone formers. This excluded patients without available stone composition analysis, those with noncalcium stones, and patients with secondary forms of calcium stones.<sup>26</sup> None of the participants included in this study had an established diagnosis of infantile hypercalcemia. All continuous predictor and outcome variables were scaled to Z-scores. The standardized regression coefficients ( $\beta$ ) were subsequently multiplied by the SD of the corresponding outcome variable. Therefore, the resulting  $\beta$  coefficients are interpreted as the expected change, in absolute values, of the outcome variable per each SD increase in the predictor variable (VMDR), which corresponds to 0.47. All regression models were tested for residuals' normality and homoscedasticity using visual inspection and for highly influential observations by plotting the Cook's distance for each data point. Residual plots were generated and carefully examined for normality through histograms along with Q-Q plots, whereas for homoscedasticity, a scatter plot of the predicted values against the residuals was inspected. All plots displayed expected patterns, suggesting that the assumptions of normality and equal variances were reasonably met for our models. Further, all analyses were checked for incomplete or overcollected 24-hour urine samples using reference values for urinary 24-hour creatinine excretion based on the adult Swiss population.<sup>27</sup> None of the results reported were biased by collection adequacy. Statistical analyses were conducted using the R software, version 4.2.1.<sup>28</sup>

## RESULTS

### Characteristics of the Study Population

A total of 974 kidney stone formers met the predefined eligibility criteria and were included in the analysis. Baseline characteristics of the study population are shown in Table 1. Overall, 69.5% of participants ( $n = 677$ ) were men. Mean age  $\pm$  SD and age at the first

**Table 1.** Baseline characteristics of study population

Characteristics	Study participants (N = 974)
Males	677 (69.5%)
Age, y	47.5 (14.4)
Age at first self-reported stone event, y	36.9 (14.2)
Age at first stone composition analysis, y	45.8 (14.6)
Body mass index, kg/m <sup>2</sup>	27.1 (4.9)
eGFR creatinine Equation CKD-EPI 2009, ml/min per 1.73 m <sup>2</sup> BSA	95.0 (19.7)
Hypertension medication usage	268 (27.6%)
Diabetes	114 (11.7%)
Stone recurrence (≥2 stone events)	793 (81.3%)
Medications affecting plasma 25(OH) vitamin D <sub>3</sub>	71 (7.3%)
Loop diuretics	14 (1.4%)
Thiazide diuretics	127 (13.0%)
DEXA parameters	
Femoral neck BMD, g/cm <sup>2</sup>	0.83 (0.75, 0.93)
Femoral neck T-score, SD	-0.60 (-1.30, 0.00)
Lumbar spine BMD, g/cm <sup>2</sup>	1.01 (0.93, 1.11)
Lumbar spine T-score, SD	-0.60 (-1.40, 0.30)
Blood parameters	
Total plasma calcium, mmol/l	2.34 (2.28, 2.39)
Corrected plasma calcium, mmol/l	2.36 (2.30, 2.42)
Ionized calcium, mmol/l	1.20 (1.17, 1.22)
Phosphate, mmol/l	0.99 (0.87, 1.11)
25(OH) vitamin D <sub>3</sub> , ng/ml	21.00 (14.00, 29.00)
1,25(OH) <sub>2</sub> vitamin D <sub>3</sub> , pmol/l	99.00 (74.79, 127.00)
Total 24,25(OH) <sub>2</sub> vitamin D ng/ml	1.46 (0.73, 2.30)
VMDR	14.78 (11.51, 19.67)
Intact PTH, ng/l	39.00 (30.60, 50.00)
cFGF23, RU/ml	69.50 (54.35, 96.00)
Urinary parameters	
Urinary calcium, mmol/24h	5.14 (3.33, 7.48)
Fractional excretion of calcium, %	2.72 (1.79, 3.57)
RSS for calcium oxalate	4.08 (2.05, 7.62)
RSS for brushite	0.74 (0.27, 2.06)
Relative kidney stone composition ≥50%	
Available stone composition analysis	751 (77.1%)
Total calcium oxalate	595 (79.2%)
Calcium oxalate monohydrate	399 (54.0%)
Calcium oxalate dihydrate	132 (17.9%)
Total calcium phosphate	91 (12.1%)
Apatite	63 (8.4%)
Brushite	15 (2.0%)
Uric acid	60 (8.0%)
Cystine	9 (1.2%)

BKSR, Bern Kidney Stone Registry; BMD, bone mineral density; BSA, body surface area; cFGF23, C-terminal fibroblast growth factor 23; CKD-EPI, Chronic Kidney Disease-Epidemiology Collaboration; DEXA, dual-energy X-ray absorptiometry; eGFR, estimated glomerular filtration rate; PTH, parathyroid hormone; SKSC, Swiss Kidney Stone Cohort; VMDR, Vitamin D metabolite diagnostic ratio.

Characteristics are indicated for all participants enrolled from the BKSR and the SKSC. Categorical variables are described by number of participants *N* (%), continuous variables are described by their mean (SD) or median (25<sup>th</sup>–75<sup>th</sup> percentile).

kidney stone event ± SD were 47.5 ± 14.4 and 36.9 ± 14.2 years, respectively. The majority of participants (*n* = 762, 81%) were recurrent stone formers. A stone composition analysis was available in 77% (*n* = 751) of participants. The most abundant stone type in our cohort (defined as ≥50% of total stone content) was calcium oxalate monohydrate (54%) as reported previously,<sup>29</sup> followed by calcium oxalate

dihydrate (18%), total calcium phosphate (12%), and uric acid (8%).

Median values (interquartile ranges) for plasma 25(OH) vitamin D<sub>3</sub>, total 24,25(OH)<sub>2</sub> vitamin D, and the VMDR were 21.0 (14.0, 29.0) ng/ml, 1.46 (0.73, 2.30) ng/ml, and 14.8 (11.5, 19.7), respectively (Table 1). There was a strong correlation between 25(OH)- and 24,25(OH)<sub>2</sub> vitamin D (Pearson's correlation coefficient = 0.83, *P* < 0.001) (Figure 1a). The VMDR was not normally distributed (Supplementary Figure S1).

A VMDR <25, considered normal (<https://www.mayocliniclabs.com/test-catalog/Overview/63416#Clinical-and-Interpretive>), was present in 87.1% (*n* = 848) of participants.<sup>19,20</sup> A VMDR between 25 and 80, previously reported in monoallelic carriers of pathogenic *CYP24A1* variants, was found in 12.1% (*n* = 118) of participants.<sup>19</sup> A VMDR >80, previously reported in biallelic carriers of pathogenic *CYP24A1* variants, was found in 0.8% (*n* = 8) participants.<sup>19</sup> Baseline characteristics stratified by these VMDR cut-offs are shown in Table 2. A replete 25(OH) vitamin D<sub>3</sub> status (≥20 ng/ml) was found in 56.2% (*n* = 547) of participants. Within this subgroup, 94.9% (*n* = 519) had a VMDR <25, 4.6% (*n* = 25) between 25 and 80, and 0.5% (*n* = 3) >80.


Stone formers with low 25(OH) vitamin D<sub>3</sub> (<20 ng/ml) had significantly higher VMDR compared to the normal range (Figure 1b), in line with previous studies.<sup>19,20</sup> As expected, a negative relationship was observed between 24,25(OH)<sub>2</sub> vitamin D and VMDR (Figure 1c).

The VMDR exhibited seasonal variability, with a median peak value in February, suggesting the lowest *CYP24A1* activity, and with a nadir reached in September, indicating the highest enzyme activity (Supplementary Figure S2, Supplementary Table S1).

## Association Analyses

The unadjusted analysis revealed a significant direct association of VMDR with total and ionized calcium, cFGF23, and fractional excretion of calcium; and an inverse correlation with BMD at the femoral neck (Table 3). After adjusting for multiple confounders, including age, sex, body mass index, estimated glomerular filtration rate, and plasma 25(OH) vitamin D<sub>3</sub> (Table 3), the results remained broadly consistent, confirming the association between VMDR and total (β 0.009 mmol/l; 95% CI: 0.002, 0.016; *P* = 0.02) and ionized calcium (β 0.005 mmol/l; 95% CI: 0.002, 0.008; *P* < 0.01), cFGF23 (β 0.045 RU/ml; 95% CI: 0.004, 0.086; *P* = 0.03), fractional excretion of calcium (β 0.046%; 95% CI: 0.018, 0.074; *P* < 0.01) and BMD at the femoral neck (β -0.005 g/cm<sup>2</sup>; 95% CI: -0.010, -0.001; *P* = 0.04). In addition, absolute





**Figure 1.** Associations between 25(OH) vitamin D<sub>3</sub>, total 24,25(OH)<sub>2</sub> vitamin D and VMDR. Panel (a) shows a scatterplot of the association between 25(OH) vitamin D<sub>3</sub> and total 24,25(OH)<sub>2</sub> vitamin D levels, including linear and natural cubic spline regression lines with green shaded areas representing the 95% confidence bands. The association appears to be approximately linear at 25(OH) vitamin D<sub>3</sub> concentration between 0 to 50 ng/ml, and total 24,25(OH)<sub>2</sub> vitamin D between 0 to 4 ng/ml. Panel (b) shows the association between 25(OH) vitamin D<sub>3</sub> and VMDR restricted to values ≤150 and a natural cubic spline regression line. Panel (c) shows a scatterplot of the association between total 24,25(OH)<sub>2</sub> vitamin D and VMDR restricted to values ≤150 and a natural cubic spline regression line. VMDR, vitamin D metabolite diagnostic ratio

urinary calcium excretion became directly correlated with VMDR ( $\beta$  0.054 mmol/24h; 95% CI: 0.010, 0.097;  $P = 0.02$ ). No association was found between VMDR and plasma 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> or PTH. Only after accounting for cFGF23 as an additional confounder, a trend toward a direct association between 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> and VMDR was observed ( $\beta$  0.131 pmol/l; 95% CI: 0.000, 0.267;  $P = 0.05$ ).

Further, our analysis revealed an inverse relationship between VMDR and the OR to develop calcium oxalate monohydrate-containing stones (Table 4), whereas the likelihood of forming calcium oxalate stones in general was not associated with VMDR. Specifically, a lower CYP24A1 activity, as reflected by a higher VMDR, was associated with a decreased odds of forming stones composed of ≥50% calcium oxalate monohydrate (adjusted OR 0.56; 95% CI: 0.38, 0.79;  $P < 0.01$ ). In contrast, VMDR was associated with an increased odds of stones with a calcium oxalate dihydrate component ≥50% (adjusted OR 1.64; 95% CI: 1.22, 2.35;  $P < 0.01$ ). No correlations between VMDR and relative supersaturations of calcium oxalate or relative supersaturations of brushite were observed.

### Sensitivity Analyses

Sodium intake is an important determinant of urinary calcium excretion,<sup>25</sup> and medications that modulate plasma 25(OH) vitamin D<sub>3</sub> or diuretics may potentially affect the VMDR or urine calcium, respectively. In addition, we found a mild seasonal variability of VMDR. To address these issues, we performed sensitivity analyses that incorporated the month of VMDR measurement, 24-hour urinary sodium excretion, an established proxy of sodium intake, along with the

aforementioned medications. All the associations found remained robust in these sensitivity analyses (Supplementary Tables S2–S7).

Further sensitivity analyses were conducted to examine the persistence of associations in subgroups of participants with varying plasma 25(OH) vitamin D<sub>3</sub> levels. In stone formers with normal plasma 25(OH) vitamin D<sub>3</sub> (≥20 ng/ml), the aforementioned adjusted associations remained significant (Supplementary Table S8). In contrast, in patients with low 25(OH) vitamin D<sub>3</sub> (<20 ng/ml), adjusted associations remained valid only for ionized calcium ( $\beta$  0.01 mmol/l; 95% CI: 0.00, 0.02;  $P = 0.04$ ) and its fractional excretion ( $\beta$  0.08 %; 95% CI: 0.02, 0.14;  $P < 0.01$ ).

### Confirmatory Analysis for Idiopathic Calcium Stone Formers

To confirm these results in the selected population of interest, we conducted a fully-adjusted analysis on idiopathic calcium stone formers (Tables 5 and 6). In this analysis, we excluded noncalcium stone formers, patients without available stone analysis, and those with secondary causes of calcium stones. Consistent with our main analysis, this subgroup exhibited significant direct associations of VMDR with total plasma calcium ( $\beta$  0.014 mmol/l; 95% CI: 0.003, 0.025;  $P = 0.011$ ), ionized calcium ( $\beta$  0.009 mmol/l; 95% CI: 0.001, 0.016;  $P = 0.025$ ), absolute ( $\beta$  0.077 mmol/24h; 95% CI: 0.018, 0.136;  $P = 0.011$ ), and fractional urinary excretion of calcium ( $\beta$  0.051 %; 95% CI: 0.01, 0.093;  $P = 0.016$ ). The inverse association with femoral BMD was also confirmed ( $\beta$  -0.018 g/cm<sup>2</sup>; 95% CI: -0.030, -0.007;  $P = 0.002$ ).

**Table 2.** Blood and urine mineral metabolism parameters and DEXA parameters stratified by subgroups of VMDR

Characteristics	Vitamin D metabolite diagnostic ratio		
	VMDR <25 (N = 848)	VMDR 25–80 (N = 118)	VMDR >80 (N = 8)
Males	604 (71.2%)	68 (57.6%)	5 (62.5%)
Age, years	47.4 (14.2)	47.8 (15.5)	51.1 (15.2)
Age at first self-reported stone event	37.1 (14.1)	36.2 (15.0)	29.3 (9.0)
Age at first stone composition analysis	45.6 (14.3)	46.6 (16.7)	48.3 (17.7)
Body mass index, kg/m <sup>2</sup>	27.0 (4.8)	28.1 (5.4)	24.9 (2.7)
eGFR creatinine Equation CKD-EPI 2009, ml/min per 1.73 m <sup>2</sup> BSA	95.4 (18.7)	93.6 (25.0)	80.2 (25.4)
Hypertension medication usage	232 (27.4%)	34 (28.8%)	2 (25.0%)
Diabetes	94 (11.1%)	20 (16.9%)	0 (0.0%)
Stone recurrence (≥2 stone events)	689 (81.3%)	96 (81.4%)	8 (100.0%)
Medications affecting plasma 25(OH) vitamin D3	55 (6.5%)	15 (12.7%)	1 (12.5%)
Loop diuretics	11 (1.3%)	3 (2.5%)	0 (0.0%)
Thiazide diuretics	115 (13.6%)	11 (9.3%)	1 (12.5%)
DEXA parameters			
Femoral neck BMD, g/cm <sup>2</sup>	0.84 (0.76, 0.93)	0.79 (0.72, 0.88)	0.79 (0.65, 0.84)
Femoral neck T-score, SD	−0.50 (−1.20, 0.10)	−0.80 (−1.48, −0.20)	−1.05 (−2.00, −0.70)
Lumbar spine BMD, g/cm <sup>2</sup>	1.02 (0.94, 1.11)	0.98 (0.85, 1.08)	1.01 (0.97, 1.06)
Lumbar spine T-score, SD	−0.60 (−1.28, 0.30)	−0.70 (−2.00, 0.18)	−0.70 (−0.98, −0.28)
Blood parameters			
Total plasma calcium, mmol/l	2.34 (2.28, 2.39)	2.32 (2.26, 2.39)	2.43 (2.34, 2.53)
Corrected plasma calcium, mmol/l	2.36 (2.30, 2.42)	2.36 (2.29, 2.43)	2.46 (2.34, 2.54)
Ionized calcium, mmol/l	1.20 (1.17, 1.22)	1.20 (1.18, 1.23)	1.24 (1.21, 1.26)
Phosphate, mmol/l	1.00 (0.88, 1.11)	0.98 (0.87, 1.10)	0.96 (0.91, 1.01)
25(OH) vitamin D <sub>3</sub> , ng/ml	22.00 (16.00, 30.00)	12.00 (8.30, 18.00)	12.50 (9.00, 31.30)
1,25(OH) <sub>2</sub> vitamin D <sub>3</sub> , pmol/l	100.00 (76.00, 127.00)	90.00 (67.00, 121.00)	81.50 (72.50, 105.00)
Total 24,25(OH) <sub>2</sub> vitamin D ng/ml	1.65 (0.96, 2.44)	0.37 (0.24, 0.57)	0.12 (0.07, 0.14)
Intact PTH, ng/l	38.60 (30.40, 49.00)	44.00 (32.00, 59.50)	42.00 (21.40, 45.00)
cFGF23, RU/ml	69.30 (54.20, 94.80)	71.10 (55.20, 111.50)	94.80 (65.70, 236.20)
Urine parameters			
Urinary calcium, mmol/24h	5.20 (3.43, 7.48)	4.51 (2.66, 7.15)	9.52 (6.07, 10.30)
Fractional excretion of calcium, %	2.72 (1.84, 3.56)	2.66 (1.55, 3.55)	3.59 (2.95, 5.24)
RSS for calcium oxalate	4.00 (2.00, 7.50)	5.00 (2.70, 8.60)	5.70 (2.60, 7.90)
RSS for brushite	0.70 (0.30, 2.00)	1.00 (0.30, 3.10)	1.30 (0.40, 4.70)
Relative kidney stone composition ≥ 50%			
Total calcium oxalate	525 (79.4%)	65 (77.4%)	5 (83.3%)
Calcium oxalate monohydrate	354 (54.4%)	44 (53.7%)	1 (16.7%)
Calcium oxalate dihydrate	112 (17.2%)	16 (19.5%)	4 (66.7%)
Total calcium phosphate	76 (11.5%)	14 (16.7%)	1 (16.7%)
Apatite	53 (8.0%)	9 (10.7%)	1 (16.7%)
Brushite	12 (1.8%)	3 (3.6%)	0 (0.0%)
Uric acid	57 (8.6%)	3 (3.6%)	0 (0.0%)
Cystine	8 (1.2%)	1 (1.2%)	0 (0.0%)

BMD, bone mineral density; BSA, body surface area; cFGF23, C-terminal fibroblast growth factor 23; CKD-EPI, Chronic Kidney Disease-Epidemiology Collaboration; DEXA, dual-energy X-ray absorptiometry; eGFR, estimated glomerular filtration rate; PTH, parathyroid hormone; VMDR, Vitamin D metabolite diagnostic ratio. Categorical variables are described by number of participants *N* (%), continuous variables are described by their mean (SD) or median (25th–75th percentile).

In terms of stone composition, the VMDR was inversely associated with the odds of calcium oxalate monohydrate stones formation (adjusted OR 0.40; 95% CI: 0.26, 0.63; *P* < 0.001), and positively with calcium oxalate dihydrate stones (adjusted OR 1.62; 95% CI: 1.15, 2.29; *P* = 0.006), in line with our results obtained in the full cohort.

## DISCUSSION

High urinary calcium excretion (“hypercalciuria”) is a risk factor for both kidney stone formation and low

bone mass. However, the underlying mechanisms remain elusive in most patients; thus, the condition is labelled “idiopathic hypercalciuria.”<sup>3,4</sup>

Our study conducted in 2 large and deeply phenotyped Swiss cohorts of kidney stone formers now reveals that the VMDR (and hence the activity of CYP24A1 inversely) is directly associated with urinary calcium, the key prolithogenic abnormality in kidney stone formers. In support of this finding, CYP24A1 activity was associated with kidney stone composition; we observed an direct association of VMDR (and hence inverse association with CYP24A1 activity) with

**Table 3.** Univariable and multivariable linear regression analysis

Outcome variable	Univariable Model				Multivariable Model			
	N	β	95% CI	P-value	N	β	95% CI	P-value
<b>Blood parameters</b>								
Total plasma calcium, mmol/l	960	0.009	0.002, 0.016	0.031	959	0.009	0.002, 0.016	0.016
Ionized calcium, mmol/l	601	0.005	0.002, 0.009	0.002	599	0.005	0.002, 0.008	0.004
Phosphate, mmol/l	962	-0.003	-0.014, 0.008	0.570	961	-0.004	-0.015, 0.007	0.460
<sup>a</sup> 1,25(OH) <sub>2</sub> vitamin D <sub>3</sub> , pmol/l	956	-0.052	-0.174, 0.07	0.400	953	0.029	-0.082, 0.139	0.610
<sup>a</sup> Intact PTH, ng/l	942	0.019	-0.007, 0.045	0.150	939	0.008	-0.018, 0.033	0.560
<sup>b</sup> cFGF23, RU/ml	356	0.065	0.023, 0.108	0.002	352	0.045	0.004, 0.086	0.030
<b>Urinary parameters</b>								
<sup>a</sup> Urinary calcium, mmol/24h	947	0.022	-0.023, 0.067	0.330	944	0.054	0.010, 0.097	0.015
<sup>b</sup> Fractional excretion of calcium, %	931	0.041	0.013, 0.069	0.004	931	0.046	0.018, 0.074	0.001
<sup>a</sup> RSS for calcium oxalate	801	0.042	-0.046, 0.129	0.350	807	0.068	-0.017, 0.154	0.118
<sup>a</sup> RSS for brushite	812	0.077	-0.062, 0.217	0.277	796	0.128	-0.005, 0.261	0.059
<b>DEXA parameters</b>								
<sup>b</sup> Lumbar BMD, g/cm <sup>2</sup>	358	-0.002	-0.007, 0.003	0.470	352	-0.001	-0.006, 0.004	0.650
<sup>b</sup> Femoral BMD, g/cm <sup>2</sup>	359	-0.008	-0.013, -0.002	0.008	353	-0.005	-0.010, -0.001	0.036

BMD, bone mineral density; BMI, body mass index; eGFR, estimated glomerular filtration rate; PTH, parathyroid hormone; RSS, relative supersaturation

<sup>a</sup>natural logarithm transformed

<sup>b</sup>square root transformed

Number of observations (N obs), beta coefficients (β), 95% confidence intervals (95% CI) and P-values are indicated for each comparison.

Association between vitamin D metabolite diagnostic ratio with clinical traits as outcome variables, adjusted for age, sex, BMI, eGFR and 25(OH) vitamin D<sub>3</sub>.

calcium oxalate dihydrate stones, the classical stone type of patients with a high urine calcium-to-oxalate ratio.<sup>29</sup> In contrast, there was an inverse association of VM DR (and thus direct association with CYP24A1 activity) with calcium oxalate monohydrate stones, typically encountered in patients with a high urinary oxalate-to-calcium ratio. Consonant with the function of CYP24A1 as vitamin D-inactivating enzyme, we observed a direct association of VM DR with both total plasma calcium and ionized calcium.

Reduced BMD and an increased fracture risk has been repeatedly observed in kidney stone formers, especially in patients with idiopathic hypercalciuria.<sup>30,31</sup> Intriguingly, we also observed that reduced CYP24A1 activity is associated with a lower BMD at the femoral neck. In contrast, VM DR was not associated with BMD at the lumbar spine, indicating that CYP24A1 activity rather affects cortical than trabecular bone. Given the cross-sectional nature of our analysis, we can only speculate on the underlying

mechanisms. Despite a vitamin D-mediated increase in intestinal calcium absorption in patients with low CYP24A1 activity, urinary calcium losses may prevail, and thus patients are overall in a negative calcium balance, as previously reported in patients with idiopathic hypercalciuria.<sup>32</sup> Indeed, urine calcium excretion was found to be inversely correlated with BMD changes at the femoral neck but not at the lumbar spine in patients with idiopathic hypercalciuria.<sup>32</sup> Alternatively, given the known function of 24,25 (OH)<sub>2</sub> vitamin D in bone remodeling, cell migration, and proliferation,<sup>33-35</sup> low levels of 24,25 (OH)<sub>2</sub> vitamin D may directly be responsible for reduced BMD observed at the femoral neck.<sup>36</sup>

Results of our analyses in kidney stone formers reveal that CYP24A1 activity, estimated by the VM DR, correlates with clinical traits characteristic of idiopathic hypercalciuria after adjustment for multiple confounders, including plasma 25(OH) vitamin D<sub>3</sub> and drugs that modulate its concentration, loop and

**Table 4.** Univariable and multivariable logistic regression analysis

Outcome variables	Univariable Model				Multivariable Model			
	N obs	OR	95% CI	P-value	N obs	OR	95% CI	P-value
<b>Relative kidney stone composition ≥ 50%</b>								
Total calcium oxalate	751	0.912	0.737, 1.142	0.370	745	0.961	0.775, 1.237	0.720
Calcium oxalate monohydrate	739	0.656	0.469, 0.873	0.009	733	0.558	0.377, 0.785	0.002
Calcium oxalate dihydrate	739	1.406	1.103, 1.887	0.015	733	1.636	1.217, 2.353	0.004
Total calcium phosphate	751	1.170	0.924, 1.476	0.150	745	1.152	0.906, 1.493	0.220
Apatite	751	1.158	0.879, 1.463	0.200	745	1.104	0.845, 1.454	0.420
Brushite	751	1.128	0.546, 1.505	0.550	745	1.235	0.479, 1.836	0.500

BMI, body mass index; eGFR, estimated glomerular filtration rate.

Number of observations (N obs), Odds ratios (OR), 95% confidence intervals (95% CI) and P-values are indicated for each comparison.

Univariable and multivariable association between Vitamin D metabolite diagnostic ratio with relative kidney stone composition analysis ≥ 50%. Multivariable model adjusted for age, sex, BMI, eGFR and 25(OH) vitamin D<sub>3</sub>.

**Table 5.** Confirmatory analysis for idiopathic calcium stone formers – linear regression analysis

Outcome variables	N obs	Vitamin D Metabolite Diagnostic Ratio		
		$\beta$	95% CI	P-value
Blood parameters				
Total plasma calcium, mmol/L	606	0.014	0.003, 0.025	0.011
Ionized calcium, mmol/L	379	0.009	0.001, 0.016	0.025
Phosphate, mmol/L	607	-0.007	-0.024, 0.009	0.398
<sup>b</sup> 1,25(OH) <sub>2</sub> Vitamin D <sub>3</sub> , pmol/L	605	0.060	-0.111, 0.23	0.491
<sup>a</sup> Intact PTH, ng/L	593	0.002	-0.039, 0.042	0.938
Urinary parameters				
<sup>a</sup> Urinary calcium, mmol/24h	607	0.077	0.018, 0.136	0.011
<sup>b</sup> Fractional excretion of calcium, %	601	0.051	0.01, 0.093	0.016
<sup>a</sup> RSS for calcium oxalate	504	0.051	-0.045, 0.148	0.298
<sup>a</sup> RSS for brushite	501	0.071	-0.076, 0.217	0.345
DEXA parameters				
<sup>b</sup> Lumbar BMD, g/cm <sup>2</sup>	232	-0.007	-0.018, 0.004	0.189
<sup>b</sup> Femoral BMD, g/cm <sup>2</sup>	233	-0.018	-0.030, -0.007	0.002

BMD, bone mineral density; BMI, body mass index; dRTA, distal renal tubular acidosis; eGFR, estimated glomerular filtration rate; PTH, parathyroid hormone

Multivariable association between Vitamin D metabolite diagnostic ratio with clinical traits as outcome variables, adjusted for age, sex, BMI, eGFR, 25(OH) vitamin D<sub>3</sub>, urinary sodium excretion, month of VMDR measurement, loop and thiazide diuretics, and medications that interfere with plasma 25(OH) vitamin D<sub>3</sub> concentration. Non-calcium stone formers, patients without available stone analysis and with secondary causes for kidney stones were excluded (primary hyperparathyroidism, sarcoidosis, complete dRTA, primary or enteric hyperoxaluria)

<sup>a</sup>natural logarithm transformed

<sup>b</sup>square root transformed

Number of observations (N obs), beta coefficients ( $\beta$ ), 95% confidence intervals (95% CI) and P-values are indicated for each comparison.

thiazide diuretics, and urinary sodium excretion. Notably, the aforementioned associations became even stronger when only idiopathic calcium stone formers were analyzed. However, these observations appear only valid in patients which are 25(OH) vitamin D<sub>3</sub> replete, whereas with 25(OH) vitamin D<sub>3</sub> deficiency (<20 ng/ml), the VMDR will be falsely elevated.

Surprisingly, we observed a significant correlation between CYP24A1 activity and cFGF23, but no association was evident with plasma PTH and 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub>. Of note, only 13 participants exhibited increased serum ionized calcium, and the majority of our cohort did not demonstrate overt hypercalcemia,

**Table 6.** Confirmatory analysis for idiopathic calcium stone formers – logistic regression analysis

Outcome variables	N obs	Vitamin D Metabolite Diagnostic Ratio		
		OR	95% CI	P-value
Total calcium oxalate	595	0.844	0.668, 1.067	0.157
Calcium oxalate monohydrate	588	0.404	0.258, 0.632	<0.001
Calcium oxalate dihydrate	588	1.62	1.147, 2.289	0.006

BMI, body mass index; dRTA, distal renal tubular acidosis; eGFR, estimated glomerular filtration rate

Multivariable logistic regression between Vitamin D metabolite diagnostic ratio with relative kidney stone composition analysis  $\geq 50\%$ . Multivariable model adjusted for age, sex, BMI, eGFR, 25(OH) vitamin D<sub>3</sub>, urinary sodium excretion, loop and thiazide diuretics, and medications that interfere with plasma 25(OH) Vitamin D<sub>3</sub> concentration. Non-calcium stone formers, patients without available stone analysis and with secondary causes for kidney stones were excluded (primary hyperparathyroidism, sarcoidosis, complete dRTA, primary or enteric hyperoxaluria). Number of observations (N obs), Odds ratios (OR), 95% confidence intervals (95% CI) and P-values are indicated for each comparison.

which may have further weakened the association between PTH and reduced CYP24A1 activity. Furthermore, there was only a very weak correlation between 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> and urinary calcium excretion in our study population (Spearman's rho 0.11;  $P < 0.01$ ). FGF23 plays a pivotal role in the regulation of vitamin D metabolism by decreasing circulating 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> through CYP27B1 inhibition and CYP24A1 induction in the kidney.<sup>37</sup> Individuals with infantile hypercalcemia type 1 and very low CYP24A1 activity have high FGF23 levels.<sup>38,39</sup> In our analysis, a trend toward an association between lower CYP24A1 activity and higher 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> was observed after adjusting for FGF23. Thus, we speculate that reduced CYP24A1 activity might lead to higher FGF23, which in turn reduces 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> synthesis through inhibition of CYP27B1. However, 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> is also regulated by PTH as part of a complex interplay between calcium, 25(OH) vitamin D<sub>3</sub>, and PTH. In an observational study, such intricate relationships may be challenging to assess, and reverse causality or alternative explanations, such as tissue-specific nonsystemic effects of CYP24A1 or a 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub>-independent regulation of renal calcium handling by CYP24A1 cannot be excluded. Consonant with this, kidney stone formers were recently shown to exhibit higher CYP24A1 expression in circulating monocytes compared to nonstone formers when individual 1,25(OH)<sub>2</sub> vitamin D<sub>3</sub> concentrations were taken into account (i.e., higher circulating 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> /monocyte CYP24A1 expression ratios).<sup>40</sup>

According to previous reports, a VMDR >80 suggests biallelic pathogenic *CYP24A1* variants, whereas values between 25 and 80 have been reported in heterozygous carriers or in patients with low plasma 25(OH) vitamin D<sub>3</sub>.<sup>10,19</sup> The prevalence of pathogenic variants in *CYP24A1* in the general population has been estimated between 420 and 1960 per 100,000 individuals (0.4 to 2%).<sup>11</sup> The observation that 28 of 547 (5%) of vitamin D-replete kidney stone formers in our cohort had a VMDR >25 indicates that other factors may play an important role in the regulation of CYP24A1 activity. Indeed, CYP24A1 expression is regulated by a myriad of endocrine factors, including PTH, FGF23, 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub>, estrogens, and retinoid receptor ligands as well as inflammatory cytokines such as IL-6 and TNF- $\alpha$ .<sup>41-46</sup> Furthermore, epigenetic modifications at the *CYP24A1* promoter and tissue-specific intronic enhancer modulate transcription of the *CYP24A1* gene.<sup>47,48</sup> Therefore, the VMDR employed in this study integrates both genetic and nongenetic factors influencing CYP24A1 activity. As such, our measurements of the VMDR in both idiopathic calcium-kidney stone formers and unselected



kidney stone formers therefore offer a comprehensive view of biochemical and clinical traits associated with CYP24A1 activity. Additional strengths of our study include the large sample size, the multicenter study design, and a very detailed phenotype including stone analysis and BMD measurements. Our study also has several limitations, including the observational design, the exploratory nature of the analyses, an almost exclusive Caucasian study population, the lack of genetic data, and absence of low risk first-time kidney stone formers in the study population.

Despite these limitations, our analyses significantly extend current knowledge on the role of the vitamin D-inactivating enzyme CYP24A1 in nephrolithiasis. Our results demonstrate that CYP24A1 activity, estimated by VMDR, is directly linked to urine calcium excretion, kidney stone composition, and BMD at the femoral neck. Future studies are now needed to confirm the validity of VMDR as a diagnostic tool in the metabolic work-up of kidney stone formers and to determine if VMDR status prospectively predicts risk of recurrence and is associated with the response to dietary and pharmacologic preventive strategies. If validated, VMDR measurements could offer clinicians valuable insights for tailored treatment recommendations in kidney stone formers.

## DISCLOSURE

CAW reports honoraria from Kyowa Kirin and Advicenne. OB was a consultant for Otsuka, CSL Vifor, Alexion, Bayer, Alnylam, Kyowa Kirin, and Boehringer Ingelheim; and received unrestricted research grants from Otsuka and Alnylam. AR received support for attending meetings and/or travel expenses from Salmon Pharma, Astellas Pharma, and Boehringer Ingelheim and honoraria from CSL Vifor. DGF served as a consultant for Otsuka, Alnylam, Boehringer Ingelheim, and Kyowa Kirin and received unrestricted research grants from Otsuka, Boehringer Ingelheim, and CSL Vifor. All the other authors declared no competing interests.

## ACKNOWLEDGMENTS

We are grateful to the patients who gave their consent to participate in this study, and we thank the study staff at the BKSr and SKSC sites for their support. This study was supported by the Swiss National Science Foundation (grant # 33IC30\_166785/1) and the Swiss National Centre of Competence in Research (NCCR) Kidney.CH.

## DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author.

## SUPPLEMENTARY MATERIAL

Supplementary File (PDF)

**Supplementary Methods.**

**Formulas.**

**Figure S1.** Distribution of VMDR.

**Figure S2.** Seasonal variation of 25(OH) vitamin D<sub>3</sub>, 24,25(OH)<sub>2</sub> vitamin D<sub>3</sub>, and their ratio.

**Table S1.** Seasonal variability of VMDR.

**Table S2.** Sensitivity linear regression analysis for urinary sodium excretion.

**Table S3.** Sensitivity logistic regression analysis for urinary sodium excretion.

**Table S4.** Sensitivity linear regression analysis for medications.

**Table S5.** Sensitivity logistic regression analysis for medications.

**Table S6.** Sensitivity linear regression analysis for seasonality of VMDR measurement.

**Table S7.** Sensitivity logistic regression analysis for seasonality of VMDR measurement.

**Table S8.** Multivariable linear regression analysis stratified by plasma 25(OH) Vitamin D<sub>3</sub>.

**Table S9.** Missing variables report.

**STROBE Statement.**

## REFERENCES

- Ferraro PM, Bargagli M, Trinchieri A, Gambaro G. Risk of kidney stones: influence of dietary factors, dietary patterns, and vegetarian-vegan diets. *Nutrients*. 2020;12:779. <https://doi.org/10.3390/nu12030779>
- Mandel NS, Mandel GS. Urinary tract stone disease in the United States veteran population. II. Geographical analysis of variations in composition. *J Urol*. 1989;142:1516–1521. [https://doi.org/10.1016/s0022-5347\(17\)39145-0](https://doi.org/10.1016/s0022-5347(17)39145-0)
- Daudon M, Bouzidi H, Bazin D. Composition and morphology of phosphate stones and their relation with etiology. *Urol Res*. 2010;38:459–467. <https://doi.org/10.1007/s00240-010-0320-3>
- Pak CY, Britton F, Peterson R, et al. Ambulatory evaluation of nephrolithiasis. Classification, clinical presentation and diagnostic criteria. *Am J Med*. 1980;69:19–30. [https://doi.org/10.1016/0002-9343\(80\)90495-7](https://doi.org/10.1016/0002-9343(80)90495-7)
- Bargagli M, Ferraro PM, Vittori M, Lombardi G, Gambaro G, Somani B. Calcium and vitamin D supplementation and their association with kidney stone disease: a narrative review. *Nutrients*. 2021;13:4363. <https://doi.org/10.3390/nu13124363>
- Maierhofer WJ, Gray RW, Cheung HS, Lemann J. Bone resorption stimulated by elevated serum 1,25-(OH)<sub>2</sub>-vitamin D concentrations in healthy men. *Kidney Int*. 1983;24:555–560. <https://doi.org/10.1038/ki.1983.193>
- Breslau NA, Preminger GM, Adams BV, Otey J, Pak CY. Use of ketoconazole to probe the pathogenetic importance of 1, 25-dihydroxyvitamin D in absorptive hypercalciuria. *J Clin Endocrinol Metab*. 1992;75:1446–1452. <https://doi.org/10.1210/jcem.75.6.1464646>

8. Plum LA, DeLuca HF. Vitamin D, disease and therapeutic opportunities. *Nat Rev Drug Discov*. 2010;9:941–955. <https://doi.org/10.1038/nrd3318>
9. Schlingmann KP, Kaufmann M, Weber S, et al. Mutations in CYP24A1 and idiopathic infantile hypercalcemia. *N Engl J Med*. 2011;365:410–421. <https://doi.org/10.1056/NEJMoa1103864>
10. O’Keeffe DT, Tebben PJ, Kumar R, Singh RJ, Wu Y, Wermers RA. Clinical and biochemical phenotypes of adults with monoallelic and biallelic CYP24A1 mutations: evidence of gene dose effect. *Osteoporos Int*. 2016;27:3121–3125. <https://doi.org/10.1007/s00198-016-3615-6>
11. Nesterova G, Malicdan MC, Yasuda K, et al. 1,25-(OH)2D-24 hydroxylase (CYP24A1) deficiency as a cause of nephrolithiasis. *Clin J Am Soc Nephrol*. 2013;8:649–657. <https://doi.org/10.2215/CJN.05360512>
12. Ferraro PM, Minucci A, Primiano A, et al. A novel CYP24A1 genotype associated to a clinical picture of hypercalcemia, nephrolithiasis and low bone mass. *Urolithiasis*. 2017;45:291–294. <https://doi.org/10.1007/s00240-016-0923-4>
13. Cools M, Goemaere S, Baetens D, et al. Calcium and bone homeostasis in heterozygous carriers of CYP24A1 mutations: a cross-sectional study. *Bone*. 2015;81:89–96. <https://doi.org/10.1016/j.bone.2015.06.018>
14. Bosworth CR, Levin G, Robinson-Cohen C, et al. The serum 24,25-dihydroxyvitamin D concentration, a marker of vitamin D catabolism, is reduced in chronic kidney disease. *Kidney Int*. 2012;82:693–700. <https://doi.org/10.1038/ki.2012.193>
15. Binkley N, Lappe J, Singh RJ, et al. Can vitamin D metabolite measurements facilitate a “treat-to-target” paradigm to guide vitamin D supplementation? *Osteoporos Int*. 2015;26:1655–1660. <https://doi.org/10.1007/s00198-014-3010-0>
16. Ketha H, Singh RJ, Grebe SK, et al. Altered calcium and vitamin D homeostasis in first-time calcium kidney stone-formers. *PLoS One*. 2015;10:e0137350. <https://doi.org/10.1371/journal.pone.0137350>
17. Bonny O, Fuster D, Seeger H, et al. The Swiss Kidney Stone Cohort (SKSC), a longitudinal, multi-centric, observational cohort to study course and causes of kidney stone disease in Switzerland. *Kidney Blood Press Res*. 2023;48:194–201. <https://doi.org/10.1159/000529094>
18. Dhayat NA, Schaller A, Albano G, et al. The vacuolar H<sup>+</sup>-ATPase B1 subunit polymorphism p.E161K associates with impaired urinary acidification in recurrent stone formers. *J Am Soc Nephrol*. 2016;27:1544–1554. <https://doi.org/10.1681/ASN.2015040367>
19. Ketha H, Kumar R, Singh RJ. LC-MS/MS for identifying patients with CYP24A1 mutations. *Clin Chem*. 2016;62:236–242. <https://doi.org/10.1373/clinchem.2015.244459>
20. Kaufmann M, Gallagher JC, Peacock M, et al. Clinical utility of simultaneous quantitation of 25-hydroxyvitamin D and 24,25-dihydroxyvitamin D by LC-MS/MS involving derivatization with DMEQ-TAD. *J Clin Endocrinol Metab*. 2014;99:2567–2574. <https://doi.org/10.1210/jc.2013-4388>
21. Dhayat NA, Ackermann D, Pruijm M, et al. Fibroblast growth factor 23 and markers of mineral metabolism in individuals with preserved renal function. *Kidney Int*. 2016;90:648–657. <https://doi.org/10.1016/j.kint.2016.04.024>
22. Levey AS, Stevens LA, Schmid CH, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150:604–612. <https://doi.org/10.7326/0003-4819-150-9-200905050-00006>
23. Dhayat NA, Schneider L, Popp AW, et al. Predictors of bone mineral density in kidney stone formers. *Kidney Int Rep*. 2022;7:558–567. <https://doi.org/10.1016/j.ekir.2021.12.003>
24. Ackermann D, Brown C, Dunthorn M, Purich DL, Finlayson B. Use of the computer program EQUIL to estimate pH in model solutions and human urine. *Urol Res*. 1989;17:157–161. <https://doi.org/10.1007/BF00256243>
25. Blackwood AM, Sagnella GA, Cook DG, Cappuccio FP. Urinary calcium excretion, sodium intake and blood pressure in a multi-ethnic population: results of the Wandsworth Heart and Stroke Study. *J Hum Hypertens*. 2001;15:229–237. <https://doi.org/10.1038/sj.jhh.1001171>
26. Gambaro G, Croppi E, Coe F, et al. Metabolic diagnosis and medical prevention of calcium nephrolithiasis and its systemic manifestations: a consensus statement. *J Nephrol*. 2016;29:715–734. <https://doi.org/10.1007/s40620-016-0329-y>
27. Forni Ognà V, Ognà A, Vuistiner P, et al. New anthropometry-based age- and sex-specific reference values for urinary 24-hour creatinine excretion based on the adult Swiss population. *BMC Med*. 2015;13:40. <https://doi.org/10.1186/s12916-015-0275-x>
28. R: the R Project for Statistical Computing. Accessed December 16, 2022. <https://www.r-project.org/>
29. Daudon M, Letavernier E, Frochet V, Haymann JP, Bazin D, Jungers P. Respective influence of calcium and oxalate urine concentration on the formation of calcium oxalate monohydrate or dihydrate crystals. *C R Chim*. 2016;19:1504–1513. <https://doi.org/10.1016/j.crci.2016.08.009>
30. Pak CYC, Sakhaee K, Moe OW, et al. Defining hypercalciuria in nephrolithiasis. *Kidney Int*. 2011;80:777–782. <https://doi.org/10.1038/ki.2011.227>
31. Sakhaee K, Maalouf NM, Kumar R, Pasch A, Moe OW. Nephrolithiasis-associated bone disease: pathogenesis and treatment options. *Kidney Int*. 2011;79:393–403. <https://doi.org/10.1038/ki.2010.473>
32. Asplin JR, Donahue S, Kinder J, Coe FL. Urine calcium excretion predicts bone loss in idiopathic hypercalciuria. *Kidney Int*. 2006;70:1463–1467. <https://doi.org/10.1038/sj.ki.5001778>
33. Seo EG, Einhorn TA, Norman AW. 24R,25-dihydroxyvitamin D3: an essential vitamin D3 metabolite for both normal bone integrity and healing of tibial fracture in chicks. *Endocrinology*. 1997;138:3864–3872. <https://doi.org/10.1210/endo.138.9.5398>
34. Ornoy A, Goodwin D, Noff D, Edelstein S. 24, 25-dihydroxyvitamin D is a metabolite of vitamin D essential for bone formation. *Nature*. 1978;276:517–519. <https://doi.org/10.1038/276517a0>
35. Lu X, Chen Z, Mylarapu N, Watsky MA. Effects of 1,25 and 24, 25 vitamin D on corneal epithelial proliferation, migration and vitamin D metabolizing and catabolizing enzymes. *Sci Rep*. 2017;7:16951. <https://doi.org/10.1038/s41598-017-16698-3>
36. Martineau C, Naja RP, Hussein A, et al. Optimal bone fracture repair requires 24R,25-dihydroxyvitamin D3 and its effector molecule FAM57B2. *J Clin Invest*. 2018;128:3546–3557. <https://doi.org/10.1172/JCI98093>
37. Latic N, Erben RG. FGF23 and vitamin D metabolism. *JBM R Plus*. 2021;5:e10558. <https://doi.org/10.1002/jbm4.10558>

38. Molin A, Lemoine S, Kaufmann M, et al. Overlapping phenotypes associated with CYP24A1, SLC34A1, and SLC34A3 mutations: a cohort study of patients with hypersensitivity to vitamin D. *Front Endocrinol.* 2021;12:736240. <https://doi.org/10.3389/fendo.2021.736240>
39. Nizar R, Cantley NWP, Tang JCY, Tang JCY. Infantile hypercalcaemia type 1: a vitamin D-mediated, under-recognised cause of hypercalcaemia. *Endocrinol Diabetes Metab Case Rep.* 2021;21-0058. <https://doi.org/10.1530/EDM-21-0058>
40. Melo TL, Esper PLG, Zambrano LI, Ormanji MS, Rodrigues FG, Heilberg IP. Expression of vitamin D receptor, CYP27B1 and CYP24A1 hydroxylases and 1,25-dihydroxyvitamin D3 levels in stone formers. *Urolithiasis.* 2020;48:19–26. <https://doi.org/10.1007/s00240-019-01163-9>
41. Zierold C, Mings JA, DeLuca HF. Regulation of 25-hydroxyvitamin D3-24-hydroxylase mRNA by 1,25-dihydroxyvitamin D3 and parathyroid hormone. *J Cell Biochem.* 2003;88:234–237. <https://doi.org/10.1002/jcb.10341>
42. Haussler MR, Haussler CA, Whitfield GK, et al. The nuclear vitamin D receptor controls the expression of genes encoding factors which feed the “Fountain of Youth” to mediate healthful aging. *J Steroid Biochem Mol Biol.* 2010;121:88–97. <https://doi.org/10.1016/j.jsbmb.2010.03.019>
43. Dhawan P, Christakos S. Novel regulation of 25-hydroxyvitamin D3 24-hydroxylase (24(OH)ase) transcription by glucocorticoids: cooperative effects of the glucocorticoid receptor, C/EBP beta, and the vitamin D receptor in 24(OH)ase transcription. *J Cell Biochem.* 2010;110:1314–1323. <https://doi.org/10.1002/jcb.22645>
44. Nashold FE, Spach KM, Spanier JA, Hayes CE. Estrogen controls vitamin D3-mediated resistance to experimental autoimmune encephalomyelitis by controlling vitamin D3 metabolism and receptor expression. *J Immunol.* 2009;183:3672–3681. <https://doi.org/10.4049/jimmunol.0901351>
45. Allegretto EA, Shevde N, Zou A, et al. Retinoid X receptor acts as a hormone receptor in vivo to induce a key metabolic enzyme for 1,25-dihydroxyvitamin D3. *J Biol Chem.* 1995;270:23906–23909. <https://doi.org/10.1074/jbc.270.41.23906>
46. Chen XQ, Mao JY, Wang CS, et al. CYP24A1 involvement in inflammatory factor regulation occurs via the Wnt signaling pathway. *Curr Sci.* 2022;42:1022–1032. <https://doi.org/10.1007/s11596-022-2564-x>
47. Luo W, Karpf AR, Deeb KK, et al. Epigenetic regulation of vitamin D 24-hydroxylase/CYP24A1 in human prostate cancer. *Cancer Res.* 2010;70:5953–5962. <https://doi.org/10.1158/0008-5472.CAN-10-0617>
48. Meyer MB, Pike JW. Mechanistic homeostasis of vitamin D metabolism in the kidney through reciprocal modulation of Cyp27b1 and Cyp24a1 expression. *J Steroid Biochem Mol Biol.* 2020;196:105500. <https://doi.org/10.1016/j.jsbmb.2019.105500>