
















RESEARCH

Open Access



Assessment of gene–disease associations and recommendations for genetic testing for somatic variants in vascular anomalies by VASCERN-VASCA

Nicole Revenu^{1†} , Astrid Eijkelenboom^{2†}, Claire Bracquemart^{3†} , Pia Alhopuro⁴ , Judith Armstrong⁵, Eulalia Baselga⁶ , Claudia Cesario⁷ , Maria Lisa Dentici⁸, Melanie Eyries⁹ , Sofia Frisk¹⁰ , Helena Gásdal Karstensen¹¹ , Nagore Gene-Olaciregui¹², Sirpa Kivirikko¹³ , Cinzia Lavarino¹² , Inger-Lise Mero¹⁴, Rodolphe Michiels¹ , Elisa Pisaneschi¹⁵ , Bitten Schönewolf-Greulich¹¹ , Ilse Wieland¹⁶, Martin Zenker^{16†}  and Miikka Vikkula^{17,18,19*†} 

Abstract

Background Vascular anomalies caused by somatic (postzygotic) variants are clinically and genetically heterogeneous diseases with overlapping or distinct entities. The genetic knowledge in this field is rapidly growing, and genetic testing is now part of the diagnostic workup alongside the clinical, radiological and histopathological data. Nonetheless, access to genetic testing is still limited, and there is significant heterogeneity across the approaches used by the diagnostic laboratories, with direct consequences on test sensitivity and accuracy. The clinical utility of genetic testing is expected to increase progressively with improved theragnostics, which will be based on information about the efficacy and safety of the emerging drugs and future molecules. The aim of this study was to make recommendations for optimising and guiding the diagnostic genetic testing for somatic variants in patients with vascular malformations.

Results Physicians and lab specialists from 11 multidisciplinary European centres for vascular anomalies reviewed the genes identified to date as being involved in non-hereditary vascular malformations, evaluated gene–disease associations, and made recommendations about the technical aspects for identification of low-level mosaicism and variant interpretation. A core list of 24 genes were selected based on the current practices in the participating laboratories, the ISSVA classification and the literature. In total 45 gene–phenotype associations were evaluated: 16 were considered definitive, 16 strong, 3 moderate, 7 limited and 3 with no evidence.

Conclusions This work provides a detailed evidence-based view of the gene–disease associations in the field of vascular malformations caused by somatic variants. Knowing both the gene–phenotype relationships and the strength

[†]Nicole Revenu, Astrid Eijkelenboom and Claire Bracquemart contributed equally to this work.

[†]Martin Zenker and Miikka Vikkula contributed equally to this work.

*Correspondence:

Miikka Vikkula

miikka.vikkula@uclouvain.be

Full list of author information is available at the end of the article



of the associations greatly help laboratories in data interpretation and eventually in the clinical diagnosis. This study reflects the state of knowledge as of mid-2023 and will be regularly updated on the VASCERN-VASCA website (VASCERN-VASCA, <https://vascern.eu/groupe/vascular-anomalies/>).

Keywords ERN, Gene curation, ISSVA, Mosaic, Precision medicine, Postzygotic, Somatic, VASCERN-VASCA, Vascular malformation

Background

Vascular anomalies comprise a heterogeneous group of disorders, divided into vascular tumours and vascular malformations [1]. The most widely used classification is that of the International Society for the Study of Vascular Anomalies [2], last updated in 2018. In the majority of cases, vascular malformations occur sporadically, but they may also be observed in several individuals in the same family as part of hereditary diseases or syndromes.

In the last 30 years, significant progress has been made in the elucidation of the genetic causes of vascular malformations. Studies initially focused on the aetiology of the hereditary forms. However, the discovery of somatic *TEK* variants in venous malformation tissues prompted the community to look for such changes in other types of vascular anomalies [3, 4]. The development of deep next-generation sequencing (NGS) techniques, and their increased use due to reduced costs, has facilitated the identification of somatic variants in various types of non-hereditary vascular anomalies. The major pathways involved in hereditary and non-hereditary forms are: 1) the RAS/mitogen-activated protein kinase (MAPK)/extracellular signal-regulated kinase (ERK) pathway, 2) the phosphatidylinositol 3-kinase (PI3K)/protein kinase B (AKT)/mammalian target of rapamycin (mTOR) pathway, and 3) the G-protein coupled receptor signalling pathway [5].

This work focuses on vascular malformations caused by somatic (postzygotic) variants, which represent the majority of patients with vascular malformations. These individuals often manifest unifocal/isolated or segmental vascular malformations. Body asymmetry (overgrowth, less commonly undergrowth) can be associated. The severity varies largely from harmless malformations to complex entities with significant morbidity, such as in CLOVES (congenital lipomatous overgrowth, vascular malformations, epidermal nevi, skeletal/scoliosis and spinal abnormalities), Klippel-Trenaunay syndrome, and Sturge-Weber syndrome. Multiple (multifocal) lesions are sometimes observed, such as in blue rubber bleb nevus syndrome and multifocal sporadic venous malformations, which complicates the diagnosis, as multiple lesions are usually a feature of a hereditary vascular malformation (e.g.,

capillary malformation–arteriovenous malformation, multiple cutaneous and mucosal venous malformations, cerebral cavernous malformation, etc.).

Pathogenic somatic variants in different genes can be associated with similar phenotypes, such as *TEK* and *PIK3CA* in venous malformations [3, 6, 7]. Likewise, somatic variants in the same gene can be associated with a variety of phenotypes. For instance, somatic variants in *PIK3CA* are associated with isolated lymphatic or venous malformation or with combined slow-flow vascular malformations, collectively known as *PIK3CA*-related overgrowth spectrum (PROS) [8].

Many genetic variants that have been identified as somatic changes in vascular malformations are also known as oncogenic variants in various cancer types. This has led to the idea that – pathway-specific pharmacological inhibitors known as cancer drugs could be repurposed for the treatment of patients with vascular malformations [9, 10] and some of them are already in off-label use or ongoing clinical trials. In addition, Alpelisib was granted accelerated approval from the Food and Drug Administration in 2022 for the treatment of patients aged 2 years or older with severe manifestations of PROS who require systemic therapy. Confirmatory evidence is needed for continued approval [11]. For these novel targeted treatment approaches, a precise molecular diagnosis is mandatory and requires genetic investigations alongside the clinical, radiological and histological assessment. Nonetheless, identifying the causative somatic genetic variants can be challenging, as they are only present in a subset of cells. Indeed, the tissue of a vascular malformation contains a mixture of mutated and non-mutated cells [12]. The percentage of cells carrying the pathogenic variant can be quite low, requiring representative test material and ultra-deep sequencing, which may limit the breadth of the gene panel content. In addition, there is a significant heterogeneity across the approaches currently used by the diagnostic laboratories in terms of technology, gene/exon/variant content, type and depth of sequencing, with direct consequences on the test sensitivity.

The aim of this work is to make recommendations to harmonise and optimise somatic genetic testing of patients with vascular malformations, performed in the participating centres, all members of

VASCERN-VASCA or providers of genetic testing for VASCERN-VASCA HCP [13]. The harmonisation mainly focused on a recommended gene content (genes, exons, regions and variants to be covered). Technical aspects for identification of low-level mosaicism were also considered, as well as variant interpretation. Germline genetic testing for hereditary vascular malformations was excluded from this work for two reasons: 1) there is a limited overlap of underlying genetic causes, and 2) the sequencing requirements are different.

Methods

Expert panel

The group of experts comprised physicians and lab specialists from 11 multidisciplinary centres, members of the working group on vascular anomalies (VASCA) of the European Reference Network on Multisystemic Vascular Diseases (VASCERN) or providers of genetic testing for VASCERN-VASCA HCP [13]. The expert group included geneticists (clinical and research), dermatologists and lab specialists in molecular genetics, pathology or oncology. Two task forces were formed: clinical and laboratory. The clinical task force was responsible for gene selection and evaluation of the gene–phenotype associations, and the laboratory task force, for selection of the variants and regions of interest, as well as the recommendations concerning the technical aspects (tissue selection, DNA isolation, sequencing techniques and variant interpretation). Virtual meetings of subgroups or the entire group were organised on a regular basis. The work started in 2020 and finished in mid-2023.

Gene and phenotype selection

Genes of interest in the context of vascular malformations caused by somatic variants were identified through the gene lists of participating expert centres, the ISSVA classification, and a literature review. Only the genes asserted in the literature to be involved in vascular malformations caused by somatic events were considered. Genes involved in hereditary vascular malformations were excluded. If a gene had been reported to be involved both in vascular malformations and vascular tumours (ISSVA classification), the latter association was also assessed (e.g., *GNAQ*). In contrast, genes involved only in vascular tumours were not included (e.g., *GNAI4*, *FOS* and *FOSB*). The ISSVA classification was used to define the list of phenotypes. Phenotypes described after the last update of this classification (e.g., *GJA4*-hepatic haemangioma, and *GJA4*-cutaneous venous malformation with cavernous histology) were added.

Gene–phenotype association evaluation

The clinical validity of gene-phenotype associations was assessed using a modified framework based on the Gene Clinical Validity Curation Process Standard Operating Procedure version 8 [14], developed by the Clinical Genome Resource Gene Curation Working Group (Table S1). This framework is designed to collect, curate and score the evidence for a gene–disease relationship in two categories of evidence: genetic and experimental. Rules for the assessment and scoring of experimental evidence were adopted from the existing standard operating procedure. In contrast, the assessment of genetic evidence on the basis of case-level data required major adaptations, as the existing framework is not applicable for diseases caused by somatic genetic events. Based on the score, gene-phenotype associations were classified in a five-category system: no evidence (0 points), limited (0.5–5.5 points), moderate (6–11.5 points), strong (12–18 points), and definitive (12–18 points and replication over time).

Data reported in the literature were evaluated. Briefly, a basic score of 0.5 was given for a somatic variant in the respective gene in a case with the respective phenotype, for each curated gene-disease pair. Upgrades of +0.5 to +1.5 were given for different levels of evidence for the pathogenicity of the variant: strong prediction of a pathogenic effect by prediction tools: +0.5; proven functional effect of the variant: +0.5–1; variant known in ClinVar as likely pathogenic: +1; and variant known in ClinVar as pathogenic: +1.5. Additional upgrades of 0.5 to 1 were given if the variant detected in one case affected the neighbouring codons (+/– 2) or the same codon, respectively. Thereby, a somatic variant of known pathogenic effect, which affects the same codon as another scored variant reaches a maximum total score of 3. At least five independent observations were necessary to reach the maximum score of 12 for genetic evidence. The evaluation matrix is available as supplemental material (Table S1).

For the genetic evidence, the literature was reviewed either comprehensively or until the maximum score was reached (12 points). Total scores for genetic and functional evidence were calculated according to the published standard operating procedures. Based on the total score, a final classification of the clinical validity of a gene–disease association was proposed (no evidence – limited – moderate – strong – definitive). Classification as “definitive” evidence required replication by > 2 publications over time (> 3 yrs.). The evaluation of each gene–disease pair was done by one member of the clinical task force, followed by presentation and discussion in the panel. The final decision on the classification was taken collectively by the clinical task force. The expert panel

had the option to maintain or change (upgrade or downgrade) the proposed classification that was based on the scoring system. If the panel decided to change, the reason was documented in the curation summary. This work led to a list of genes for which the assessed evidence for a certain phenotype association supported the consideration as a diagnostic gene.

Selection of variants, exons and regions of interest

Based on the gene list constituted by the clinical task force, the laboratory task force searched for all reported somatic variants in these genes using the Leiden Open Variation Database (LOVD: [15]) and ClinVar Miner [16] databases, as well as PubMed and Google Scholar with the keywords “*gene name*+vascular malformation”, “*gene name*+vascular anomaly” and “*gene name*+vascular tumour”. Variants were considered somatic if the observed variant allele fraction (VAF) suggested mosaicism or if a variant was present at different allele fractions among different tissue types from the same individual. After the literature and public database search, all members of the laboratory task force were asked to compare their in-house databases with the collected data and complete the list with unreported variants, if any. Variants were then divided into two groups: 1) variants that have been clearly associated with vascular anomalies in the literature and/or databases; and 2) variants that had emerging yet insufficient evidence, as well as variants reported as somatic hotspots in cancer or germline hotspots in RASopathy genes and *TEK*. Cancer hotspot variants were defined through database queries in Catalogue Of Somatic Mutation In Cancer (COSMIC: [17]) and CancerHotspots [18]. Variants identified in the COSMIC database with ≥ 20 documented instances, or in CancerHotspots with ≥ 10 documented instances, were considered potential oncogenic driver variants, in accordance with previous definitions [19–21].

Technical aspects and variant analysis and interpretation

First, the specific approaches and techniques used by the 11 participating laboratories at the beginning of the work were collected; these included input material, DNA isolation method, sequencing technologies, gene panel content, sequencing platform, bioinformatics pipelines and turn-around time. Based on our experience and a literature review, the laboratory task force drafted analytical workflows that were discussed in the plenary meetings. We provide an expert opinion to improve and standardise laboratory approaches for somatic variant detection and interpretation in the context of vascular malformations caused by postzygotic events.

Results

Gene curation

The formal curation process was applied to 24 genes (Table 1 and Supplementary Material 2). In total, 45 gene–phenotype associations were evaluated. Twelve genes were associated with more than one phenotype (vascular malformation or vascular tumour). Of the 45 evaluated gene–phenotype associations, 16 were considered definitive, 16 strong, 3 moderate 7 limited and 3 with no evidence.

Curation matrices for each gene–disease association evaluation are provided as additional file (Supplementary Material 2). There was at least one “definitive” gene–phenotype association for 10 genes (*AKT1*, *GNAI1*, *GNAQ*, *IDH1*, *KRAS*, *MAP2K1*, *MAP3K3*, *NRAS*, *PIK3CA* and *TEK*). Out of the 24 genes, 11 were published after the last update of the ISSVA classification (May 2018), of which two have at least one “definitive” association (*KRAS* and *NRAS*), five have at least one “strong” association (*AKT3*, *GJA4*, *HRAS*, *PIK3R1* and *PTPN11*), and four have “limited evidence” (*ARAF*, *CBL*, *GNB2* and *PIK3CD*). When the type of vascular malformation was mixed or not specified in the reports, it was considered under the term of “vascular malformation, various types” (*HRAS*, *NRAS* – not linked to KLA).

Diagnostic target regions

Based on the review of reported disease-associated variants per curated gene, we defined two levels of target regions per gene to be considered for diagnostic testing of somatic variants in vascular malformations (Table 2 and Table S2). The first level represents the minimum content that should be included in a diagnostic panel (Table 2). This covers pathogenic and likely pathogenic (P/LP) variants reported in the literature and/or databases [15, 16] in patients with non-hereditary vascular anomalies. No additional unreported variants have been identified in the in-house databases from the participating centres. For the delineation of the extended regions of interest, a broader set of variants was accepted, such as variants with insufficient evidence of pathogenicity, and variants in patients with less well-characterised phenotypes (Table S2). This selection also contains: i) areas defined as hotspots in cancer but not yet observed in vascular malformations, and ii) hotspot germline variants in RASopathy genes and *TEK* [22].

Technical aspects

The detection of somatic variants has special technical requirements. The approaches and techniques, from sample collection to DNA extraction and sequencing

Table 1 Gene–disease associations and strength of association

Gene name	Disease	Disease	Disease	Disease	Disease
AGGF1	Klippel-Trenaunay syndrome NO EVIDENCE				
AKT1	Proteus syndrome DEFINITIVE				
AKT3	CMTC variant (w/o (hemi)megalencephaly) or CM (with LM?) with hypotrophy STRONG				
ARAF	CCLA LIMITED				
BRAF	Arteriovenous malformation STRONG	(Macrocystic) Lymphatic malformation MODERATE	Pyogenic granuloma STRONG		
CBL	Kaposiform lymphangiomatosis LIMITED				
GJA4	Cutaneous venous malformation (cavernous histology) MODERATE	Hepatic hemangioma STRONG			
GNA11	Capillary malformation w/o overgrowth DEFINITIVE	Sturge-Weber syndrome STRONG	Congenital hemangioma STRONG		
GNAQ	Capillary malformation w/o overgrowth DEFINITIVE	Sturge-Weber syndrome DEFINITIVE	Congenital hemangioma DEFINITIVE		
GNAS	Vascular malformation, any type NO EVIDENCE				
GNB2	Sturge-Weber syndrome LIMITED				
HRAS	Vascular malformations, various type STRONG				
IDH1	Maffucci syndrome DEFINITIVE	Spindle cell hemangioma DEFINITIVE			
IDH2	Maffucci syndrome STRONG	Spindle cell hemangioma STRONG			
KRAS	Sporadic arteriovenous malformation DEFINITIVE	Gorham-Stout disease MODERATE	Epithelioid hemangioma LIMITED		
MAP2K1	Arteriovenous malformation DEFINITIVE	Epithelioid hemangioma STRONG			
MAP3K3	Verrucous venous malformation STRONG	Cerebral cavernous malformation DEFINITIVE			
NPM1	Maffucci syndrome NO EVIDENCE				
NRAS	Vascular malformations, low-flow, various type LIMITED	Kaposiform lymphangiomatosis DEFINITIVE	Pyogenic granuloma / tufted angioma LIMITED		
PIK3CA	PIK3CA-related overgrowth spectrum DEFINITIVE	Lymphatic malformation DEFINITIVE	Common venous malformation DEFINITIVE	Megalencephaly-capillary malformation-polymicrogyria DEFINITIVE	Generalised lymphatic anomaly STRONG
PIK3CD	Lymphatic malformation LIMITED				
PIK3R1	Capillary malformation, dilated veins, (mild) overgrowth +/- lymphatic malformation STRONG				
PTPN11	Capillary malformation w/o overgrowth STRONG				
TEK	Common venous malformation DEFINITIVE	Blue Rubber Bleb Nevus syndrome STRONG	Multifocal sporadic Venous malformation STRONG		

technologies, used by the 11 laboratories of the participating centres at the beginning of this study are listed anonymously in Table S3.

Tissue selection and DNA requirements

In participating centres, sequencing analysis for somatic variants was preferably performed on DNA from affected

Table 2 Variants, exons and regions of interest for vascular malformations caused by somatic events

Gene Name	Transcript Reference	Minimal required regions		Optional regions	
		Exons	amino acid residues of interest	Exons	amino acid residues of interest
AKT1	NM_001382430.1	3	E17	3	L52
				4	Q79
AKT3	NM_5465.7	3	E17	None	
ARAF	NM_001654.5	7	S214	9	R255
BRAF	NM_004333.4	15	V600	6	T244 to F247, Q257
				11	G466, G469
				14	N581
				15	N581 to K601
				16	Q636
CBL	NM_005188.4	15	Y774	2	Y102
				7	D359 to K362
				8	Multiple hotspots, splice
				9	C404 to R420, splice
				16	A877
GJA4	NM_002060.3	2	G41	None	
GNA11	NM_002067.5	4	R183	None	
		5	Q209		
GNA14	NM_004297.4	4	Q205	None	
GNAQ	NM_002072.5	4	R183	2	M59, T96
		5	Q209		
GNB2	NM_005273.4	5	K78	None	
HRAS	NM_005343.4	2	G12, G13	3	Q61
		3	T58 to M72	5	F156
IDH1	NM_005896.4	4	R132	None	
IDH2	NM_002168.4	4	R172	4	R140
KRAS	NM_004985.5	2	K5, G12, G13, Q22	2	V14, L19
		3	Q61, Y64	3	T58 to G60
		4	A146	4	K117
				5	D153, F156
MAP2K1	NM_002755.4	2 3	F53 to E62 P105 to 107, C121 to G128	2	L42 to D67
				3	E102 to M146
				5	L177
				6	E203
				7	P264, P294
				8	G301, S331
				15	Y544
				3	G12 to P34
4	T50 to G60				
MAP3K3	NM_002401.5	13	I441	15	Y544
NRAS	NM_002524.5	3	Q61	2	G12 to P34
		4	A146	3	T50 to G60
PIK3CA	NM_006218.4	Full gene		Full gene	
PIK3CD	NM_005026.5	16	L666	13	S520
				24	E1021
PIK3R1	NM_181523.3	11	Y452 to Y467	9	R348
		13	N564 to Y580	10	G376
		14	W583, splice	13	N564, K567, splice
PTPN11	NM_002834.5	3	E76	1_13	Multiple hotspots
		12	V428, A461		
		13	T507		
TEK	NM_00459.5	17	Y897 to A925	15	R849
		22	R1099X	17	L920
		23	T1105 to G1115	18	F960
				23	K1100 to Y1113

This table describes regions of interest (genes, exons, amino acids) that should be studied for somatic exploration of vascular anomalies. The regions of interest are divided into two groups: "Minimal required regions" and "Optional regions". The first one represents what is well described and implicated in the pathogenicity of somatic vascular anomalies, and the second one represents hotspot variants reported in cancer, or germline hotspots in *RASopathy* genes and *TEK*, with insufficient evidence of pathogenicity in non-hereditary vascular anomalies. Reference transcripts are from ensembl.org. Data are from literature and databases (LOVD, Clinvar Miner, Cosmic (v96) and CancerHotspots (v2))

tissue, as the variants are usually absent or present at only very low allele fraction in other sources, such as blood or saliva. Affected tissue samples were obtained by lesional biopsies or from surgical resections. They were further processed in various ways until DNA extraction, mainly as native tissue samples in saline solution, frozen or in RNAlater solution, but some laboratories also used formalin-fixed paraffin embedded (FFPE) tissue (Table S3). The specifics of tissue processing were matched to the laboratory protocols for subsequent analysis (for example, not all the library preparation methods are compatible with genomic DNA (gDNA) from FFPE tissue). For most centres, histological examination was considered in the diagnostic work-up in a pathology lab, especially if the diagnosis was unclear or to ensure that the biopsy is representative (deep malformation). Some laboratories performed tissue dissection (removing unaffected tissue, e. g. epidermidis) to increase the sensitivity of somatic testing (Table S3).

Minimum tissue requirements

The minimum amount of tissue was dependent on the input criteria (quantity and quality) of the laboratory protocols for subsequent analysis. A punch or excisional biopsy size of 4 mm was usually sufficient for subsequent analysis in the laboratory's protocols.

DNA isolation methods

A variety of commercial tissue DNA extraction kits and protocols were used by the participating laboratories to isolate gDNA from the affected tissue. The specifics of the DNA isolation protocols were adapted according to both the tissue preparation and the laboratory protocols for subsequent analysis. The minimum DNA input for analysis ranged from 5 to 200 ng (Table S3). Quality control of gDNA were included in each laboratory protocol depending on the methods used for sequencing part.

Sequencing analysis by NGS

Most centres were using custom targeted panels, while three centres (9, 10 and 11) were using whole-exome sequencing (WES) and applied a virtual in silico panel of genes of interest for the analysis of WES data (Table S3). Two laboratories were using unique molecular identifiers (UMIs) to increase sequencing accuracy. For those centres using WES, the analysed gene panels also included genes for hereditary vascular malformations, to enable germline and somatic testing in the same analysis flow. Higher read depth was used (120X-400X minimal read depth) compared to standard germline WES testing. However, the target regions may have suboptimal coverage compared to custom panels and low-frequency variants can thereby go undetected. Minimal sequencing

depth for custom panels ranged from 100 to 2000X (Table S3).

Read alignment and variant detection

In house pipelines as well as commercial pipelines were used by the participating centres to align and call the variants (Table S3). Bioinformatics pipelines were adapted to the special requirements of somatic variant calling (with a low VAF).

Technical sensitivity and specificity

Pathogenic variants that are detected in vascular malformations caused by somatic events are generally present only in a subset of cells and the mutant allele frequency in DNA from mixed lesional tissue samples may be quite low. In the participating diagnostic routine laboratories, the indicated cut off for the minimum VAF varied from 1 to 3% (Table S3). If depth for detecting low VAF is not reached due to technical limitations, it will reduce the analytical sensitivity of the assay. Notably, approximately 15% ($n=41/262$) of cases with a pathogenic variant in *PIK3CA* detected by the members of this working group had a VAF of < 3% (internal data from the participating centres).

Each laboratory established empirically its sequencing requirements, such as the minimum sequencing depth, a minimum number of variant reads and the minimum VAF to be reported (Table S3). Binomial distribution-based coverage calculation can aid in determining the minimum sequencing depth [23]. UMI-based sequence analysis was successfully used by two of the participating laboratories to increase test accuracy (Table S3). One laboratory used strand-specific amplification, which may help to discriminate genuine C:G>T>A variants from deamination artefacts in FFPE samples. If tissue handling and NGS library flows are combined with the processing of solid tumour specimens, there is a possibility of cross-contamination from solid tumour specimens as the origin of pathogenic variants with a low VAF. One laboratory added common single nucleotide polymorphisms (SNPs) in the library preparation to trace this cross-contamination during analysis of sequences.

Alternative methods of somatic variant detection/validation

Some laboratories used digital PCR (dPCR)/droplet digital PCR (ddPCR) to confirm low frequency variants or detect recurrent hotspots variants in genes such as *GNAQ*, *GNA11*, *PIK3CA*, and *TEK*.

Variant interpretation

Variant filtering

As usually performed in NGS-based analysis, common benign variants and artefacts were removed from the detected list of variants prior to interpretation.

Variant classification and reporting

All centres indicated that their classification of variant pathogenicity was essentially based on ACMG recommendations [24]. The AMP/ACMG/ASCO recommendations were used by the centres to report somatic test results for vascular malformations [25]. All the information required for unambiguous clinical interpretation of the test result was reported according to the guidelines. The Human Genome Variation Society (HGVS) standard nomenclature [26] providing the transcript reference sequence was used to annotate all detected variants. Coverage of the NGS assay, VAF, exons/hotspots that do not meet the quality standards, description of methods, assay performance characteristics, limitations of the assays used and a list of genes covered are the minimal data content indicated in reports [25]. All reported variants were classified as described above, and pathogenic and likely pathogenic alterations were generally disclosed. Unexpected findings or variants of unknown significance were discussed with clinicians prior to reporting. Recommendations for any supplemental testing were made when appropriate.

Discussion

The aim of this work was to propose a performant and accurate genetic testing approach for patients with vascular malformations caused by somatic genetic events. At the beginning of this work, there was a large heterogeneity between the centres in the VASCERN-VASCA regarding the genes and the exonic regions that were screened, as well as in the methodological approaches. Therefore, our first aim was to harmonise the content of the diagnostic gene panel, by proposing a core gene list based on evidence of associations with vascular malformations. Our second aim was to make recommendations for the technical approaches.

Scope of diagnostic somatic testing in vascular malformations

By systematic evaluation of gene-disease associations, we came up with a list of 24 genes, for which there was usable published evidence for their role in vascular malformations driven by somatic variants. For a subset of 17 genes the evidence was rated “strong” or “definitive” for at least one gene-phenotype association. We consider these core genes to be a mandatory content of any diagnostic panel used for a broad spectrum of vascular malformations. We

also recommend the inclusion of a further 4 genes with “limited evidence”, as it can be expected that these genes may be upgraded as new published observations emerge. In contrast, we do not recommend the inclusion of the remaining 3 genes, for which “no evidence” of an association with vascular malformations was found (Table 1).

Since somatic (gain-of-function) variants that are associated with vascular malformations are typically restricted to specific hotspot regions within the respective genes, we also aimed to provide recommendations on variants and regions of interest for each gene. The sites of variants reported in vascular malformations should be covered by a diagnostic test as a minimum requirement (Table 2). As the variants associated with vascular malformations overlap with recurrent oncogenic variants found in cancer in the respective genes, a broader definition of the regions of interest also includes these (Table S2). In general, laboratories using targeted gene panels that only partially cover genes according to their defined hotspots should be aware that novel regions of interest may rapidly emerge.

Notably, the list of genes evaluated in this work is significantly larger than the number of genes mentioned in the last ISSVA classification, illustrating the major progress that has been made since the last update (May 2018). There are 11 new genes (*AKT3*, *ARAF*, *CBL*, *GJA4*, *GNB2*, *HRAS*, *KRAS*, *NRAS*, *PIK3CD*, *PIK3R1* and *PTPN11*) that are not in the ISSVA classification (2018). Of these, seven have at least one “definitive” (*KRAS* and *NRAS*) or “strong” (*AKT3*, *GJA4*, *HRAS*, *PIK3R1* and *PTPN11*) gene-phenotype association. The remaining four genes (*ARAF*, *CBL*, *GNB2* and *PIK3CD*) have been associated with vascular malformations with “limited evidence”. For *ARAF*, two patients have been reported by the same group in 2019 [27] with lymphangiomatosis or central conducting lymphatic anomaly, respectively. Based on the available data, it is not clear whether both variants were somatic or not. The observation of *ARAF* variants in lymphatic malformations has not been replicated since. Case reports have been published for *GNB2*, *CBL* and *PIK3CD*, [28–30] describing patients with Sturge-Weber syndrome (*GNB2*) or lymphatic anomalies (*CBL* and *PIK3CD*). Further evidence is needed to confirm the involvement of these genes in vascular malformations. In the most recent version of the ISSVA classification, the *NPM1* gene was indicated as being associated with Maffucci syndrome. However, a careful review of the literature does not support this association, as the only patient with a variant in this gene reported to date was a 7-year-old girl with Maffucci syndrome and acute myeloid leukaemia [31]. This patient had two somatic variants, both in the leukemic cells and in the vascular lesion: a frameshift variant in the *NPM1* gene and a missense

variant in the *IDH1* gene. As *IDH1* is strongly associated with Maffucci syndrome, the variant identified in this gene can explain this part of the phenotype. Somatic *NPM1* variants are instead associated with acute myeloid leukaemia (OMIM #601,626). Thus, we conclude that the *NPM1* variant is not involved in the pathogenesis of Maffucci syndrome.

There is also increasing evidence that several genes (particularly the core genes of the RAS-MAPK and PI3K-AKT cascades) are involved in a wider range of vascular anomalies than previously recognised, probably reflecting variability in the timing of the mutation, the cell type affected and other factors. For example, the E542K variant in *PIK3CA* has been identified in isolated LM, isolated VM and PROS. New phenotypes were added for a total of five genes: *BRAF* with sporadic AVM (strong evidence) and LM (moderate evidence), *GNA11* with Sturge-Weber syndrome (strong evidence), *MAP2K1* with epithelioid haemangioma (strong evidence), *MAP3K3* with cerebral cavernous malformation (definitive evidence) and *PIK3CA* with generalised lymphatic anomaly (strong evidence).

We evaluated gene–disease associations until variant level, when relevant. As mentioned above, the same pathogenic variants can be associated with different or overlapping phenotypes and with different extension and severity of the disease. However, different variants in the same gene can also be associated with specific phenotypes; for example, the R183Q variant in *GNAQ* has been associated with CM and Sturge-Weber syndrome, but not with non-involuting congenital haemangiomas (NICH) or rapid-involuting congenital haemangiomas (RICH), which are instead associated with the Q209H/L/P variants.

The expanding genetic heterogeneity and phenotypic diversity associated with individual genes challenges the use of narrowly focused diagnostic assays that include only one or a small panel of genes targeted to a defined clinical phenotype. However, the predominance of recurrent variants in some distinctive phenotypes still leaves a possible rationale for the use of targeted methods such as digital PCR, which can be an efficient and cost-effective first-tier approach to search for highly recurrent hotspot variants or to analyse genes with a narrow mutational spectrum, such as *GNAQ*, *GNA11*, *PIK3CA* and *TEK* hotspot variants. However, negative results from such an approach, should be followed by testing of a more comprehensive gene panel.

Genes involved in vascular malformations caused by somatic events may also be analysed in combination with genes involved in hereditary vascular malformations (germline events), although this is not specifically recommended. Instead, the expert panel agreed that the analysis

of genes involved in hereditary vascular malformations is generally not essential in sporadic patients with a type of vascular malformation that is usually caused by somatic events, as the underlying genes are mostly different, and the clinical presentation can usually guide towards somatic or germline testing, respectively. Moreover, for laboratories using custom panels, the inclusion of the genes for hereditary forms would greatly affect the size of the panel, which may conflict with the need for very deep sequencing to detect somatic variants with very low VAFs. The combination of germline and somatic testing in one assay should take into account cost-effectiveness, which also depends on the local laboratory equipment and organisation. Nevertheless, the limitation to a very strictly-defined gene content will become obsolete with the advent (and cost efficiency) of new, very large-scale sequencing technologies. For laboratories using a WES-based approach it is easy to open their analysis to the genes involved in hereditary vascular malformations based on the patient's phenotype (e.g. *KRIT1* for patients with hyperkeratotic cutaneous capillary-venous malformations, and *RASA1*, *EPHB4*, *ENG1*, *ACVRL1*, *SMAD4*, *PTEN* for AVM). However, it has to be noted that WES is usually not performed at a sequencing depth that is sufficient to reach a reliable mosaic detection threshold of 1% VAF. For laboratories using a custom panel, on the other hand, a low threshold for extending the analysis toward the genes involved in hereditary forms can be suggested, if the phenotype is compatible with such a disease. It has to be noted, that multiplicity of vascular lesions or a positive family history are not always present in hereditary vascular malformations. If somatic testing is performed exclusively, potentially pathogenic germline variants should be reported and the patient referred for genetic counselling and germline testing.

Somatic variants in overlapping groups of genes may also be involved in other disorders, for example genes of the PI3K-AKT-mTOR pathway in segmental overgrowth or brain malformations or genes of the RAS-MAPK pathway in neurocutaneous mosaic disorders. Thus, laboratories may find it more useful and efficient to use a combined panel rather than separate panels to test for these groups of conditions.

Technical recommendations for genetic testing of somatic variants

Recommendations for technical aspects of somatic variant testing are summarised in Table 3. The recommendations represent a consensus of expert opinion based on the practice and experience of the reference centres involved in this study, and are intended to assist clinical laboratories in the detection of somatic variants and to ensure a high quality of sequencing results.

Table 3 Technical recommendations for genetic testing of somatic variants

Technical aspect	Recommendation of the VASCERN-VASCA expert group
Tissue sampling	4 mm of affected tissue specimen from debulking/reduction surgery, or skin biopsy of affected tissue containing abnormal vasculature is the minimum tissue size recommended for adequate DNA yield Fresh/frozen and FFPE samples can be considered Blood samples are generally not recommended, as the diagnostic yield is low. Cell-free DNA from blood plasma or lymphatic fluid are a promising alternative source, but need further evaluation
Histologic examination	Recommended for differential diagnostics and/or estimation of the representativeness of the sample. Dissection (removal of epidermis) can be performed to increase the percentage of affected tissue
Tissue culturing	Tissue culturing is not necessary, but culturing endothelial cells from affected tissue may aid subsequent variant detection
DNA extraction	Standard DNA extraction protocols
DNA quality	Low-quality samples can be analysed; however, it reduces sensitivity and may introduce sequencing artefacts. Cost-effectiveness of prior DNA quality control must be determined in each laboratory
Gene-disease validity	Core gene list (Table 1) and updates thereof https://vascern.eu/
Expected VAFs	Mosaic variants can be present at multiple frequencies, expected VAFs are mostly lower than 10%
Target region	Minimum required regions that must be covered for the detection of somatic variants associated with vascular malformation are reported in Table 2. Recommended optional regions are also listed in Table S2
Sequencing analysis by NGS	Various platforms can be used UMI-based sequencing is recommended especially for amplicon-based, but also for hybridisation-based capture technologies. For the latter, if UMIs are not used, read deduplication is recommended, which can be based on read start and end coordinates; Strand-specific amplification or capture can help to discriminate genuine C:G > T > A variants from deamination artefacts in FFPE samples
Technical sensitivity and specificity	Sequencing depth and minimum number of variant reads should be empirically defined. Detection threshold of 1% is recommended. Including common SNPs in the library preparation is advised to detect cross-contamination from tumour samples when handled in parallel
Read alignment, variant detection, and variant filtering	Diagnostically validated pipelines for somatic calling are recommended. Prefiltering is advised to remove benign variants and artefacts
Validation of NGS findings	Recommended for all variants with VAF less than 5% using dPCR or a second NGS analysis
Alternative methods of somatic variant detection	dPCR can be used for the detection of recurrent variants associated with specific phenotypes
Variant classification	ACMG classification with modifications suggested in PMID: 35,997,716 and PMID: 34,040,190 are recommended. Literature/database searches should extend beyond cancer variant data annotation and interpretation
Reporting	AMP/ACMG/ASCO recommendations are largely applicable (PMID: 27,993,330). Variants should be annotated using the HGVS standard nomenclature. Reports should provide VAF, coverage, regions that do not meet quality standards, description of methods, limitations and list of genes analysed. Pathogenic/likely pathogenic variants should be disclosed. Recommendations for supplemental testing should be made, such as germline panel testing, based on the indication and clinical context
Identification of potential germline variants	Potential pathogenic/likely pathogenic germline variants should be included in the report and referral for germline testing and genetic counselling should be advised
Data sharing	Submission of variants with phenotypic, clinical significance, and classification criteria to DNA variant databases, such as ClinVar and LOVD, is recommended to increase knowledge of the genetics of vascular anomalies

Regarding the source of DNA for diagnostic testing, a tissue sample containing abnormal vasculature is the material of choice and gold standard. Since surgical resection tissue is not always available, a skin biopsy with a punch from the site of the lesion can be obtained. Cell culturing from tissue samples is not routinely necessary, but can help increase the VAF if endothelial cells

from the affected tissue are isolated and used for culture [32, 33]. Blood leucocyte DNA is not recommended as the diagnostic yield is relatively low [34], with a few exceptions, such as for some patients with MCAP [35] or MSVM (multiple sporadic venous malformations) [36]. Depending on the phenotype, other non-invasive specimens, such as urine (urinary epithelial cells) [37]

or saliva (buccal cells) [38] may also contain the causative variant, but their use cannot be recommended for the general routine. Cell-free DNA (cfDNA) analysis is emerging as an attractive alternative to tissue-based testing. This would be particularly beneficial when an invasive procedure is not possible and for young patients. Nevertheless, the VAF of disease-causing variants in cfDNA is usually much lower than in lesional tissue DNA (often less than 1%) and requires highly sensitive techniques, such as dPCR [39, 40]. The diagnostic yield of cfDNA in vascular malformations is currently rather low. It may be increased by using cfDNA from an efferent vein [41] or lymphatic fluid [40]. However, the use of cfDNA for diagnostic purposes needs to be further developed and validated.

Low VAFs of the disease-causing somatic variants in mixed tissue specimens or other diagnostic samples from patients with vascular malformations present a major diagnostic challenge. The VAFs of somatic driver variants are generally lower than those typically found in solid tumours. Therefore, routine settings adopted from molecular oncology may be insufficient for vascular malformations. In large patient cohorts, most pathogenic variants are present in lesional tissue samples with a VAF less than 10%, even in patients with severe phenotypes [42, 43]. The expected VAF range depends on the sample type (e.g. [34]) and clinical context (e.g. [44]) and may even fall below 1%. Enrichment of mutant cells in a given sample might be achieved by tissue dissection, microdissection, endothelial cell culturing or cell separation [45, 46]. These approaches may be considered when faced with a negative result. However, the current experience is insufficient to recommend their routine use. In samples of poor DNA quality or quantity, the sensitivity for low-level mosaicism is reduced and the probability of ambiguous results is increased. The expert group recommends to include a quality control of the gDNA starting material to relate the DNA quality assessment with the sequencing results. If tissue handling and NGS library flows are combined with the processing of solid tumour specimens, there is possibility of cross-contamination from solid tumour specimens as the origin of pathogenic variants with a low VAF. Different methods using polymorphic tags in the library preparation and/or post-sequencing bio-informatic analysis can be applied to detect such cross-contamination [47, 48].

Concerning the sensitivity threshold for somatic variants in lesional tissue samples, the expert group recommendation for the minimum VAF is set at 1%. Sequencing depth and minimum number of variant reads needed to reach this threshold should be empirically defined for each work-flow. Standardized set of DNA samples carrying mosaic variants with different

allele fractions are made available by companies helping in panel development. It is also possible to exchange DNA samples between laboratories for genetic tests that are not yet covered by an external quality assessment scheme. Binomial distribution-based coverage calculations can be used to determine the probability to detect a variant at a certain allele ratio [23]. Notably, increasing sequencing depth of libraries, especially those derived from poor quality material, per se has limited effect on sensitivity because of the background level of technical artefacts that are inherent to the technology. Artefacts can arise from different sources, including tissue fixation with formalin, PCR and sequencing; they are therefore dependent on sample type, library preparation and sequencing technique [49]. When FFPE tissue samples are used, common base substitution artefacts, particularly C>T/G>A caused by cytosine deamination, significantly limit the detection threshold [50]. An elegant option to overcome some of these limitations of current NGS technologies and thereby lower the detection threshold is the use of UMI-based sequencing. These are complex molecular barcodes that are added to DNA prior to any PCR amplification steps. This enables the definition of the consensus sequence of PCR multiplicates derived from a single template DNA molecule, thereby eliminating amplification and sequencing artefacts [51]. The use of UMIs is recommended especially for amplicon-based, but also for hybridisation-based capture technologies. For the latter, if UMIs are not used, read deduplication is recommended, which can be based on read start and end coordinates.

Generally, ambiguous findings and results close to the detection threshold of the given test should be interpreted with caution in the context of the clinical indication, and independent confirmation should be sought. This can preferably be achieved by an alternative technique, such as dPCR, or if not available, by repeating the NGS analysis [52]. Digital PCR provides a high depth of genotyping allowing the detection of specific variants at VAFs as low as 0.1%, with a minimal gDNA input requirements (5 ng of gDNA per test) and can also be used if the gDNA quality or quantity is suboptimal (e.g. DNA from FFPE tissue). The use of dPCR as a first-tier test for specific indications is discussed above. The need for specific assays for each change at the nucleotide level and limited multiplexing capabilities constrain the use of dPCR in vascular malformation diagnostics. The use of blocker displacement amplification as an alternative method to confirm variants with very low VAF in vascular malformations has recently been reported [46].

Bioinformatic pipelines for NGS data analysis need to be adapted for the specific needs of low-level mosaic detection. For variant callers the threshold may need

to be adjusted for the desired minimum calling VAF. In addition, for some recurrent hotspot positions, it may be considered to program a systematic variant calling and/or to manually inspect the aligned sequences in a genome viewer such as integrative genomics viewer (IGV: [53]). On the other hand, a more agnostic approach with an exploratory setting based on deep WES may be useful for detecting disease causing variants in novel genes [54].

The interpretation and classification of causal somatic variants in vascular malformations can sometimes be challenging. Although disease-causing variants in several genes largely overlap with oncogenic drivers detected in solid tumours, other variants in specific genes can be found (almost) exclusively in vascular malformations (e.g., variants in the receptor tyrosine kinase *TEK* gene have not been reported in cancer). Furthermore, atypical variants in some oncogenes are (almost) exclusively found in vascular malformations (e.g. atypical *HRAS* and *KRAS* indel variants [55]).

In our experience, the American College of Medical Genetics/Association for Molecular Pathology (ACMG/AMP) general criteria of classification of germline variants [24] were not completely applicable for somatic variant classification. Other classification guidance schemes from oncology, such as the joint consensus recommendations of AMP, ACMG, American Society of Clinical Oncology (ASCO) [25, 56] and European Society for Medical Oncology (ESMO) [57] can be used, but were found to focus on clinical relevance and to lack guidance on variant classification. An exception was the Belgian ComPerMed, which proposed guidelines for biological classification and clinical interpretation of somatic variants [58]. Gene-specific recommendations were recently reported for *PIK3R1* [21], and for *AKT3*, *MTOR*, *PIK3CA* and *PIK3R2* by the ClinGen Brain Malformation Variant Curation Expert Panel [59]. Such gene specific recommendations should be tested and validated for vascular malformations.

Conclusion

Based on the VASCERN-VASCA expertise, our aim was to optimise and guide the diagnostic genetic testing performed for non-hereditary vascular malformations caused by somatic variants. We defined the core gene list that should be tested and evaluated, the level of evidence of the gene–phenotype associations and the variants/regions of interest that should be included in a (virtual) panel. Knowing the gene-phenotype relationship and the strength of this association can greatly help laboratories to interpret the data and contribute to clinical diagnosis. This study reflects the state of knowledge as of mid-2023 and will be regularly updated on the VASCERN-VASCA website [13]. Laboratories around the world should be able to use these data.

Abbreviations

ACMG	American College of Medical Genetics
AMP	Association for Molecular Pathology
ASCO	American Society of Clinical Oncology
cfDNA	Cell-free DNA
ESMO	European Society for Medical Oncology
gDNA	Genomic DNA
COSMIC	Catalogue Of Somatic Mutation In Cancer
CLOVES	Congenital lipomatous overgrowth, vascular malformations, epidermal nevi, skeletal/scoliosis and spinal abnormalities
dPCR	Digital PCR
ddPCR	Droplet digital PCR
HGVS	Human Genome Variation Society
ISSVA	International Society for the Study of Vascular Anomalies
IGV	Integrative genomics viewer
LOVD	Leiden Open Variation Database
MCAP	Megalencephaly-capillary malformation
MSVM	Multiple sporadic venous malformations
NGS	Next-generation sequencing
NICH	Non-involuting congenital haemangiomas
PHTS	<i>PTEN</i> Hamartoma tumour syndrome
PROS	PIK3CA-related overgrowth spectrum
SNP	Single-nucleotide polymorphism
RICH	Rapidly involuting congenital haemangiomas
VAF	Variant allele fraction
UMI	Unique molecular identifier
WES	Whole-exome sequencing

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13023-024-03196-9>.

Supplementary Material 1.

Supplementary Material 2.

Supplementary Material 3.

Supplementary Material 4.

Acknowledgements

The authors of this publication are members of the Vascular Anomaly Working Group (VASCA WG) of the European Reference Network for Rare Multisystemic Vascular Diseases (VASCERN) or providers of genetic testing for VASCERN-VASCA HCP. This project has been supported by the European Reference Network on Rare Multisystemic Vascular Diseases (VASCERN)—Project ID: 101085076, which is funded by the European Union within the framework of the EU for Health Programme. MZ is a member of ERN-ITHACA, funded by the EU4Health programme of the European Union, under GA nr. 101085231. ILM thanks Marte Kirkevold and Mariann Kristensen for their assistance in the acquisition of data. MLD thanks the Dermatology Unit and Genodermatosis Unit of Bambino Gesù Children's Hospital, Rome, Italy. The authors thank Mr. Clovis Lessuisse for excellent secretarial help.

Authors' contributions

Conceptualization: NR, AE, CB, JA, PA, EB, CEL, IW, MZ, MV. Curation matrix adaptation for somatic variants: MZ. Data acquisition: CB, PA, EB, CEL, NGO, SF, ME, HGK, ILM, RM, NR, IW, MV, MZ. Data analysis & interpretation: CB, PA, EB, SF, ME, CEL, ILM, NR, IW, MV, MZ. Data curation: CB, PA, EB, HGK, CEL, ILM, RM, NR, MV, MZ. Writing-original draft (major contributors): NR, AE, CB, MZ, PA, CL. Writing-review & editing: all.

Funding

This project has been supported by the European Reference Network on Rare Multisystemic Vascular Diseases (VASCERN)—Project ID: 101085076, which is funded by the European Union within the framework of the EU for Health Programme. MZ received funding from the German Federal Ministry of Education and Research (BMBF; grant numbers: FKZ 01GM1902A and FKZ 01GM1921A) and is a member of ERN-ITHACA, funded by the EU4Health programme of the European Union, under GA nr. 101085231. PA received special funding for

university-level health research in Finland, University of Helsinki. MV received funding from the Fonds de la Recherche Scientifique—FNRS Grants T.0240.23 & PC005.22, the Fund Genereet managed by the King Baudouin Foundation (Grant 2018-J1810250-211305), and the Walloon Region through the FRFS-WELBIO strategic research programme (WELBIO-CR-2019C-06).

Availability of data and materials

All data analysed during this study are included in the article and its supplementary information files.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

EB: PI for Novartis and Founder of Venthera. All other authors declare no conflicts of interest.

Author details

¹Center for Human Genetics, Cliniques universitaires Saint-Luc, University of Louvain, VASCERN VASCA European Reference Centre, Brussels, Belgium. ²Department of Pathology, Radboud University Medical Center, VASCERN VASCA European Reference Centre, PO Box 9101, 6500, HB, Nijmegen, the Netherlands. ³Normandie Univ, UNICAEN, Service de Génétique, CHU Caen Normandie, BIOTARGEN EA 7450, VASCERN VASCA European Reference Centre, Caen 14000, France. ⁴HUS Diagnostic Center, Laboratory of Genetics, University of Helsinki and Helsinki University Hospital, VASCERN VASCA European Reference Centre, Helsinki, Finland. ⁵Institut de Recerca Sant Joan de Déu, Esplugues de Llobregat, CIBER-ER (Biomedical Network Research Center for Rare Diseases), Instituto de Salud Carlos III (ISCIII), Madrid, and Genomic Unit, Molecular and Genetic Medicine Section, Hospital Sant Joan de Déu, VASCERN VASCA European Reference Centre, Barcelona, Spain. ⁶Department of Dermatology, Hospital Sant Joan de Deu, VASCERN VASCA European Reference Centre, Barcelona, Spain. ⁷Laboratory of Medical Genetics, Translational Cytogenomics Research Unit, Bambino Gesù Children Hospital and Research Institute, IRCCS, VASCERN VASCA European Reference Centre, Rome, Italy. ⁸Medical Genetics Unit, Bambino Gesù Children's Hospital, IRCCS, VASCERN VASCA European Reference Centre, 00165 Rome, Italy. ⁹Sorbonne Université, Département de Génétique, Assistance Publique-Hôpitaux de Paris, Hôpital Pitié-Salpêtrière, VASCERN VASCA European Reference Centre, Paris, France. ¹⁰Department of Molecular Medicine and Surgery, Karolinska Institutet and Department of Clinical Genetics, Karolinska University Hospital, VASCERN VASCA European Reference Centre, Stockholm, Sweden. ¹¹Department of Genetics, Center of Diagnostics, Copenhagen University Hospital - Rigshospitalet, VASCERN VASCA European Reference Centre, Copenhagen, Denmark. ¹²Laboratory of Molecular Oncology, Pediatric Cancer Center Barcelona, Hospital Sant Joan de Déu, VASCERN VASCA European Reference Centre, Barcelona, Spain. ¹³Department of Clinical Genetics, HUS Diagnostic Center, University of Helsinki and Helsinki University Hospital, VASCERN VASCA European Reference Centre, Helsinki, Finland. ¹⁴Department of Medical Genetics, Oslo University Hospital, VASCERN VASCA European Reference Centre, Oslo, Norway. ¹⁵Laboratory of Medical Genetics, Translational Cytogenomics Research Unit, Bambino Gesù Children Hospital and Research Institute, IRCCS, VASCERN VASCA European Reference Centre, Rome, Italy. ¹⁶Institute of Human Genetics, University Hospital Otto-Von-Guericke-University, Magdeburg, Germany. ¹⁷Center for Vascular Anomalies, Cliniques Universitaires Saint-Luc, Brussels, Belgium. ¹⁸Human Molecular Genetics, de Duve Institute, University of Louvain, VASCERN VASCA European Reference Centre, Brussels, Belgium. ¹⁹WELBIO Department, WEL Research Institute, Avenue Pasteur, 6, 1300 Wavre, Belgium.

Received: 12 December 2023 Accepted: 19 April 2024

Published: 22 May 2024

References

- Mulliken JB, Glowacki J. Hemangiomas and vascular malformations in infants and children: a classification based on endothelial characteristics. *Plast Reconstr Surg.* 1982;69(3):412–22.
- International Society for the Study of Vascular Anomalies. ISSVA classification for vascular anomalies. Revised 2018. Available from: <https://www.issva.org/UserFiles/file/ISSVA-Classification-2018.pdf>.
- Limaye N, Wouters V, Uebelhoe M, Tuominen M, Wirkkala R, Mulliken JB, et al. Somatic mutations in angiopoietin receptor gene TEK cause solitary and multiple sporadic venous malformations. *Nat Genet.* 2009;41(1):118–24.
- Limaye N, Boon LM, Vikkula M. From germline towards somatic mutations in the pathophysiology of vascular anomalies. *Hum Mol Genet.* 2009;18(R1):R65–74.
- Queisser A, Seront E, Boon LM, Vikkula M. Genetic Basis and Therapies for Vascular Anomalies. *Circ Res.* 2021;129(1):155–73.
- Limaye N, Kangas J, Mendola A, Godfraind C, Schlogel MJ, Helaers R, et al. Somatic Activating PIK3CA Mutations Cause Venous Malformation. *Am J Hum Genet.* 2015;97(6):914–21.
- Castel P, Carmona FJ, Grego-Bessa J, Berger MF, Viale A, Anderson KV, et al. Somatic PIK3CA mutations as a driver of sporadic venous malformations. *Sci Transl Med.* 2016;8(332):332ra42.
- Canaud G, Hammill AM, Adams D, Vikkula M, Keppler-Noreuil KM. A review of mechanisms of disease across PIK3CA-related disorders with vascular manifestations. *Orphanet J Rare Dis.* 2021;16(1):306.
- Nguyen HL, Boon LM, Vikkula M. Vascular Anomalies Caused by Abnormal Signaling within Endothelial Cells: Targets for Novel Therapies. *Semin Interv Radiol.* 2017;34(3):233–8.
- Van Damme A, Seront E, Dekeuleener V, Boon LM, Vikkula M. New and Emerging Targeted Therapies for Vascular Malformations. *Am J Clin Dermatol.* 2020.
- FDA accelerated approval for Alpelisib. Available from: <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-alpelisib-pik3ca-related-overgrowth-spectrum>.
- Nguyen HL, Boon LM, Vikkula M. Genetics of vascular malformations. *Semin Pediatr Surg.* 2014;23(4):221–6.
- VASCERN-VASCA. Available from: <https://vascern.eu/groupe/vascular-anomalies/>.
- Gene Clinical Validity Curation Process Standard Operating Procedure version 8. Available from: <https://clinicalgenome.org/docs/summary-of-updates-to-the-clingen-gene-clinical-validity-curation-sop-version-8/>.
- LOVD: Leiden Open Variation Database. Available from: <https://www.lovd.nl/>.
- ClinVar Miner database. Available from: <https://clinvarminer.genetics.utah.edu/>.
- COSMIC: Catalogue Of Somatic Mutation In Cancer. Available from: <https://cancer.sanger.ac.uk/cosmic>.
- CancerHotspots database. Available from: <https://www.cancerhotspots.org/#/home>.
- Chang MT, Bhattarai TS, Schram AM, Bielski CM, Donoghue MTA, Jonsson P, et al. Accelerating Discovery of Functional Mutant Alleles in Cancer. *Cancer Discov.* 2018;8(2):174–83.
- Walsh MF, Ritter DI, Kesserwan C, Sonkin D, Chakravarty D, Chao E, et al. Integrating somatic variant data and biomarkers for germline variant classification in cancer predisposition genes. *Hum Mutat.* 2018;39(11):1542–52.
- Cottrell CE, Bender NR, Zimmermann MT, Heusel JW, Corliss M, Evenson MJ, et al. Somatic PIK3R1 variation as a cause of vascular malformations and overgrowth. *Genet Med.* 2021;23(10):1882–8.
- Gelb BD, Cave H, Dillon MW, Gripp KW, Lee JA, Mason-Suares H, et al. ClinGen's RASopathy Expert Panel consensus methods for variant interpretation. *Genet Med.* 2018;20(11):1334–45.
- Eijkelenboom A, Kamping EJ, Kastner-van Raaij AW, Hendriks-Cornelissen SJ, Neveling K, Kuiper RP, et al. Reliable Next-Generation Sequencing of Formalin-Fixed, Paraffin-Embedded Tissue Using Single Molecule Tags. *J Mol Diagn.* 2016;18(6):851–63.

24. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405–24.
25. Li MM, Datto M, Duncavage EJ, Kulkarni S, Lindeman NI, Roy S, et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. *J Mol Diagn*. 2017;19(1):4–23.
26. HGVS standard nomenclature. Available from: <http://www.hgvs.org>.
27. Li D, March ME, Gutierrez-Uzquiza A, Kao C, Seiler C, Pinto E, et al. ARAF recurrent mutation causes central conducting lymphatic anomaly treatable with a MEK inhibitor. *Nat Med*. 2019;25(7):1116–22.
28. Fjaer R, Marciniak K, Sundnes O, Hjorthaug H, Sheng Y, Hammarstrom C, et al. A novel somatic mutation in GNB2 provides new insights to the pathogenesis of Sturge-Weber syndrome. *Hum Mol Genet*. 2021;30(21):1919–31.
29. Foster JB, Li D, March ME, Sheppard SE, Adams DM, Hakonarson H, et al. Kaposiform lymphangiomatosis effectively treated with MEK inhibition. *EMBO Mol Med*. 2020;12(10):e12324.
30. Wang S, Wang W, Zhang X, Gui J, Zhang J, Guo Y, et al. A somatic mutation in PIK3CD unravels a novel candidate gene for lymphatic malformation. *Orphanet J Rare Dis*. 2021;16(1):208.
31. Akiyama M, Yamaoka M, Mikami-Terao Y, Ohyama W, Yokoi K, Arakawa Y, et al. Somatic mosaic mutations of IDH1 and NPM1 associated with cup-like acute myeloid leukemia in a patient with Maffucci syndrome. *Int J Hematol*. 2015;102(6):723–8.
32. Huang L, Couto JA, Pinto A, Alexandrescu S, Madsen JR, Greene AK, et al. Somatic GNAQ Mutation is Enriched in Brain Endothelial Cells in Sturge-Weber Syndrome. *Pediatr Neurol*. 2017;67:59–63.
33. Nikolaev SI, Vetiska S, Bonilla X, Boudreau E, Jauhainen S, Rezai Jahromi B, et al. Somatic Activating KRAS Mutations in Arteriovenous Malformations of the Brain. *N Engl J Med*. 2018;378(3):250–61.
34. Lalonde E, Ebrahimzadeh J, Rafferty K, Richards-Yutz J, Grant R, Toorens E, et al. Molecular diagnosis of somatic overgrowth conditions: A single-center experience. *Mol Genet Genomic Med*. 2019;7(3):e536.
35. Mirzaa G, Timms AE, Conti V, Boyle EA, Girisha KM, Martin B, et al. PIK3CA-associated developmental disorders exhibit distinct classes of mutations with variable expression and tissue distribution. *JCI Insight*. 2016;1(9).
36. Soblet J, Kangas J, Natynki M, Mendola A, Helaers R, Uebelhoer M, et al. Blue Rubber Bleb Nevus (BRBN) Syndrome Is Caused by Somatic TEK (TIE2) Mutations. *J Invest Dermatol*. 2017;137(1):207–16.
37. Michel ME, Konczyk DJ, Yeung KS, Murillo R, Vivero MP, Hall AM, et al. Causal somatic mutations in urine DNA from persons with the CLOVES subgroup of the PIK3CA-related overgrowth spectrum. *Clin Genet*. 2018;93(5):1075–80.
38. Riviere JB, Mirzaa GM, O'Roak BJ, Beddaoui M, Alcantara D, Conway RL, et al. De novo germline and postzygotic mutations in AKT3, PIK3R2 and PIK3CA cause a spectrum of related megalencephaly syndromes. *Nat Genet*. 2012;44(8):934–40.
39. Biderman Waberski M, Lindhurst M, Keppler-Noreuil KM, Sapp JC, Baker L, Gripp KW, et al. Urine cell-free DNA is a biomarker for nephroblastomatosis or Wilms tumor in PIK3CA-related overgrowth spectrum (PROS). *Genet Med*. 2018;20(9):1077–81.
40. Zenner K, Jensen DM, Cook TT, Dmyterko V, Bly RA, Ganti S, et al. Cell-free DNA as a diagnostic analyte for molecular diagnosis of vascular malformations. *Genet Med*. 2021;23(1):123–30.
41. Palmieri M, Pinto AM, di Blasio L, Curro A, Monica V, Sarno LD, et al. A pilot study of next generation sequencing-liquid biopsy on cell-free DNA as a novel non-invasive diagnostic tool for Klippel-Trenaunay syndrome. *Vascular*. 2021;29(1):85–91.
42. Ten Broek RW, Eijkelenboom A, van der Vleuten CJM, Kamping EJ, Kets M, Verhoeven BH, et al. Comprehensive molecular and clinicopathological analysis of vascular malformations: A study of 319 cases. *Genes Chromosomes Cancer*. 2019;58(8):541–50.
43. Keppler-Noreuil KM, Sapp JC, Lindhurst MJ, Parker VE, Blumhorst C, Darling T, et al. Clinical delineation and natural history of the PIK3CA-related overgrowth spectrum. *Am J Med Genet A*. 2014;164(7):1713–33.
44. Brouillard P, Schlögel MJ, Homayun Sepehr N, Helaers R, Queisser A, Fastré E, et al. Non-hotspot PIK3CA mutations are more frequent in CLOVES than in common or combined lymphatic malformations. *Orphanet J Rare Dis*. 2021;16(1):267.
45. Blesinger H, Kaulfuß S, Aung T, Schwoch S, Prantl L, Rößler J, et al. PIK3CA mutations are specifically localized to lymphatic endothelial cells of lymphatic malformations. *PLoS ONE*. 2018;13(7):e0200343.
46. Li D, Sheppard SE, March ME, Battig MR, Surrey LF, Srinivasan AS, et al. Genomic profiling informs diagnoses and treatment in vascular anomalies. *Nat Med*. 2023;29(6):1530–9.
47. Balan J, Koganti T, Basu S, Dina MA, Artymiuk CJ, Barr Fritcher EG, et al. MICon Contamination Detection Workflow for Next-Generation Sequencing Laboratories Using Microhaplotype Loci and Supervised Learning. *J Mol Diagn*. 2023;25(8):602–10.
48. Fiévet A, Bernard V, Tenreiro H, Dehainault C, Girard E, Deshaies V, et al. ART-DeCo: easy tool for detection and characterization of cross-contamination of DNA samples in diagnostic next-generation sequencing analysis. *Eur J Hum Genet*. 2019;27(5):792–800.
49. Singh RR. Next-Generation Sequencing in High-Sensitive Detection of Mutations in Tumors: Challenges, Advances, and Applications. *J Mol Diagn*. 2020;22(8):994–1007.
50. Steiert TA, Parra G, Gut M, Arnold N, Trotta JR, Tonda R, et al. A critical spotlight on the paradigms of FFPE-DNA sequencing. *Nucleic Acids Res*. 2023;51(14):7143–62.
51. Kennedy SR, Schmitt MW, Fox EJ, Kohrn BF, Salk JJ, Ahn EH, et al. Detecting ultralow-frequency mutations by Duplex Sequencing. *Nat Protoc*. 2014;9(11):2586–606.
52. Douzgou S, Rawson M, Baselga E, Danielpour M, Favre L, Kashanian A, et al. A standard of care for individuals with PIK3CA-related disorders: An international expert consensus statement. *Clin Genet*. 2022;101(1):32–47.
53. Integrative genomics viewer (IGV). Available from: <https://igv.org/>.
54. Schönewolf-Greulich B, Karstensen HG, Hjortshøj TD, Jørgensen FS, Harder KM, Frevert S, et al. Early diagnosis enabling precision medicine treatment in a young boy with PIK3R1-related overgrowth. *Eur J Med Genet*. 2022;65(10):104590.
55. Eijkelenboom A, van Schaik FMA, van Es RM, Ten Broek RW, Rinne T, van der Vleuten C, et al. Functional characterisation of a novel class of in-frame insertion variants of KRAS and HRAS. *Sci Rep*. 2019;9(1):8239.
56. Leichsenring J, Horak P, Kreutzfeldt S, Heining C, Christopoulos P, Volckmar AL, et al. Variant classification in precision oncology. *Int J Cancer*. 2019;145(11):2996–3010.
57. Mateo J, Chakravarty D, Dienstmann R, Jezdic S, Gonzalez-Perez A, Lopez-Bigas N, et al. A framework to rank genomic alterations as targets for cancer precision medicine: the ESMO Scale for Clinical Actionability of molecular Targets (ESCAT). *Ann Oncol*. 2018;29(9):1895–902.
58. Froyen G, Le Mercier M, Lierman E, Vandepoele K, Nollet F, Boone E, et al. Standardization of Somatic Variant Classifications in Solid and Haematological Tumours by a Two-Level Approach of Biological and Clinical Classes: An Initiative of the Belgian ComPerMed Expert Panel. *Cancers (Basel)*. 2019;11(12).
59. Lai A, Soucy A, El Achkar CM, Barkovich AJ, Cao Y, DiStefano M, et al. The ClinGen Brain Malformation Variant Curation Expert Panel: Rules for somatic variants in AKT3, MTOR, PIK3CA, and PIK3R2. *Genet Med*. 2022;24(11):2240–8.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.