

Erratum to: Hepatic effects of *Cimicifuga racemosa* extract in vivo and in vitro

S. Lüde · M. Török · S. Dieterle · A. C. Knapp ·
R. Kaeufeler · R. Jäggi · U. Spornitz ·
S. Krähenbühl

Published online: 26 May 2010
© Springer Basel AG 2010

Erratum to: Cell Mol Life Sci (2007) 64:2848–2857
DOI 10.1007/s00018-007-7368-4

In the abstract the correct dose in the sentence “Microvesicular steatosis was found in rats treated with >500 µg/kg body weight cimicifuga extract.” should be 1,000 mg/kg body weight (instead of >500 µg/kg). The corresponding dose in the results part of the article is cited correctly.

The authors apologize for this error and any confusion it may have caused.

The online version of the original article can be found under
doi:[10.1007/s00018-007-7368-4](https://doi.org/10.1007/s00018-007-7368-4).

S. Lüde · M. Török · S. Dieterle · A. C. Knapp ·
S. Krähenbühl (✉)
Division of Clinical Pharmacology & Toxicology
and Department of Research, University Hospital Basel,
4031 Basel, Switzerland
e-mail: kraehenbuehl@uhbs.ch

R. Kaeufeler
Max Zeller Söhne AG, Romanshorn, Switzerland

R. Jäggi
Vitaplant Ltd., Witterswil, Switzerland

U. Spornitz
Institute for Anatomy, Histology and Embryology,
University of Basel, Basel, Switzerland