

## The Membrane-Bound H<sup>+</sup>-ATPase Complex Is Essential for Growth of *Lactococcus lactis*

BRIAN J. KOEBMANN,<sup>1,2</sup> DAN NILSSON,<sup>2</sup> OSCAR P. KUIPERS,<sup>3†</sup> AND PETER R. JENSEN<sup>1\*</sup>

Department of Microbiology, Technical University of Denmark, DK-2800 Lyngby,<sup>1</sup> and Department of Physiology, Chr. Hansen A/S, DK-2970 Hørsholm,<sup>2</sup> Denmark, and Microbial Ingredients Section, NIZO Food Research, NL-6710 BA Ede, The Netherlands<sup>3</sup>

Received 10 January 2000/Accepted 13 June 2000

The eight genes which encode the (F<sub>1</sub>F<sub>o</sub>) H<sup>+</sup>-ATPase in *Lactococcus lactis* subsp. *cremoris* MG1363 were cloned and sequenced. The genes were organized in an operon with the gene order *atpEBFHAGDC*; i.e., the order of *atpE* and *atpB* is reversed with respect to the more typical bacterial organization. The deduced amino acid sequences of the corresponding H<sup>+</sup>-ATPase subunits showed significant homology with the subunits from other organisms. Results of Northern blot analysis showed a transcript at approximately 7 kb, which corresponds to the size of the *atp* operon. The transcription initiation site was mapped by primer extension and coincided with a standard promoter sequence. In order to analyze the importance of the H<sup>+</sup>-ATPase for *L. lactis* physiology, a mutant strain was constructed in which the original *atp* promoter on the chromosome was replaced with an inducible nisin promoter. When grown on GM17 plates the resulting strain was completely dependent on the presence of nisin for growth. These data demonstrate that the H<sup>+</sup>-ATPase is essential for growth of *L. lactis* under these conditions.

The (F<sub>1</sub>F<sub>o</sub>) H<sup>+</sup>-ATPase complex plays an important role in the free energy metabolism of virtually all living cells. The structures of F<sub>1</sub>F<sub>o</sub>-ATPase complexes from different sources are very similar and consist of two parts: a membrane integral part, F<sub>o</sub>, which forms a proton channel, and a soluble part, F<sub>1</sub>, which contains the catalytic site for ATP hydrolysis. In bacteria, the enzyme is located in the cytoplasmic membrane, where it catalyzes the interconversion of ATP and the transmembrane proton gradient. Depending on the particular organism and on the conditions for growth, the enzymes function in the direction of either ATP synthesis or ATP hydrolysis (14). In organisms which contain a respiratory chain, such as *Escherichia coli* and *Bacillus subtilis*, the primary role of the enzyme is to synthesize ATP driven by the proton gradient that results from respiration, when these organisms are supplied with an electron acceptor. In organisms that lack a respiratory chain, or in the absence of electron acceptors, the enzyme generates a transmembrane proton gradient, and this process is then driven by ATP hydrolysis. The anaerobic bacterium *Lactococcus lactis* also possesses an F<sub>1</sub>F<sub>o</sub>-ATPase complex. This bacterium lacks the respiratory chain, and the enzyme here is involved in the extrusion of protons driven by ATP hydrolysis to generate the necessary driving force for solute transport and to maintain an acceptable intracellular pH value (21, 38). The latter function is supported by the fact that the activity of the F<sub>1</sub>F<sub>o</sub>-ATPase in these anaerobic bacteria is enhanced at low external pH (2, 23).

The anaerobic bacteria have an alternative route to generate a proton gradient across the cytoplasmic membrane, namely, through end product excretion. In the so-called energy recycling model, which was first demonstrated by Michels et al.

(27), it was suggested that carrier-mediated excretion of end products can occur in symport with protons, and this contributes to the generation of the transmembrane proton gradient. This mechanism has been thoroughly investigated in *Lactococcus lactis* by Otto et al. (30), and ten Brink et al. (40), who demonstrated that the energy recycling by lactate efflux makes a significant contribution to the generation of the proton gradient in this organism, particularly at high external pH and low external lactate concentrations. An interesting question is then whether this contribution would be sufficient to allow growth of *L. lactis* in the absence of the H<sup>+</sup>-ATPase.

In this paper we report the cloning, sequencing, and characterization of the genes that encode the H<sup>+</sup>-ATPase in *L. lactis* subsp. *cremoris* MG1363. A mutant strain was constructed in which the expression of H<sup>+</sup>-ATPase on the chromosome is under control of the *nisA* promoter. The strain was completely dependent on nisin for growth on GM17 plates, which demonstrates that the H<sup>+</sup>-ATPase is an essential enzyme for growth of *L. lactis*.

### MATERIALS AND METHODS

**Bacterial strains.** The plasmid-free *L. lactis* subsp. *cremoris* strain MG1363 (16) was used to study the *atp* operon in *L. lactis*. *E. coli* K-12 strain BOE270 is highly competent with respect to transformation and was derived from strain MT102, which is an *hsdR* derivative of strain MC1000 [*araD139 (ara-leu)7679 galU galK (lac)174 rpsL thi-1*] (7). BOE270 was used as a host for plasmids in the cloning procedures and for propagation of plasmid DNA in *E. coli*.

**Oligonucleotides and enzymes.** Oligonucleotides were obtained from Hobolth DNA Synthesis (Hillerød, Denmark). Restriction enzymes (Gibco BRL, Pharmacia), *Taq* and *Pfu* DNA polymerases (Pharmacia and AH Diagnostics, respectively), calf intestine alkaline phosphatase (Pharmacia), and T4 DNA ligase (Gibco BRL) were used as recommended by the manufacturers.

**Sequencing and sequence analysis of the H<sup>+</sup>-ATPase operon.** The DNA sequencing was carried out either by the dideoxy nucleotide chain termination method (33) with [ $\alpha$ -<sup>32</sup>P]ddNTP (500 Ci/mmol) (Pharmacia) or by autosequencing by capillary electrophoresis with the Dye Terminator Cycle Sequencing Ready Reaction kit (Perkin-Elmer).

The alignments of DNA and amino acid sequences were performed on the BLAST server at the National Center for Biotechnology Information (NCBI). The numbers given below refer to the numbering used in the GenBank sequence.

**Transformation.** Cells of *E. coli* were made competent by the Ca<sup>2+</sup> method (32). Plasmid DNA was used to transform the cells by a standard transformation procedure (28), and the transformation mixtures were plated at 30°C on Luria-

\* Corresponding author. Mailing address: Department of Microbiology, Technical University of Denmark, Building 301, DK-2800 Lyngby, Denmark. Phone: 45 45252510. Fax: 45 45932809. E-mail: imprj@pop.dtu.dk.

† Present address: Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, NL-9750 AA Haren, The Netherlands.




FIG. 1. Genetic organization of the *L. lactis* *atp* operon and DNA fragments used in this work. The open reading frames are shown as boxes, and the designations of the *atp* genes are shown below the boxes in italic letters. The designations of the H<sup>+</sup>-ATPase subunits are shown above the boxes. The arrow indicates the direction of transcription of the *atp* operon, and the stem loop indicates the putative terminator. The cloned fragments, which are referred to in the text, are indicated in the boxes below the scale.

Bertani agar plates supplemented with either ampicillin (100 µg/ml) or erythromycin (200 µg/ml). Cells of *L. lactis* (16) were made competent by growth in GM17 medium containing 1% glycine and resuspended in 10% glycerol and 0.5 M sucrose as described by Holo and Nes (18). Plasmid DNA was used to transform the cells by electroporation (18), and the cells were allowed to regenerate in SGM17 medium for 2 h and then plated onto Schmidt-Ruppin plates containing the appropriate selective antibiotic.

**Cloning of the *atp* operon from *L. lactis* subsp. *cremoris* MG1363.** Fragments of the *atp* operon were cloned as PCR products or by the plasmid rescue technique (see below). Chromosomal DNA from *L. lactis* MG1363 was used as a template for amplification of DNA. Several primer sets were used to amplify different regions of the *atp* operon (Fig. 1). Here we took advantage of the fact that in an unrelated project, the first part of the *atp* operon from the closely related bacterium *L. lactis* subsp. *lactis* B1014 was accidentally discovered in a clone from a gene library, which allowed us to design primers for the amplification of the genes that encode the F<sub>0</sub> part of the enzyme complex. PCR amplification was carried out in a total volume of 100 µl and in the presence of 0.4 mM concentrations of each deoxynucleoside triphosphate (Boehringer), 3 to 5 µM concentrations of each primer DNA, 0.1 µg of chromosomal DNA, 2.5 U of *Taq* polymerase, and the buffer recommended by the manufacturer (Pharmacia). The reactions were carried out for 25 cycles (1 min at 94°C for denaturing, 1 min at 55°C for annealing, and 2 min at 72°C extension step) by use of a DNA thermal cycler. The resulting PCR products were cloned in pMOSBlue (Amersham) and sequenced. To confirm the correctness of the cloned product, the sequence was also determined directly on the PCR products.

**Cloning of *atpC* by plasmid rescue.** Plasmid pQ1, which harbors the DNA sequence from position 4177 to position 6394 (the C-terminal part *atpG* of the product and the N-terminal part *atpD* of the product) and was obtained by cloning a PCR fragment obtained with primers 3987 (5'-TTGGTGGTGGATC AATGACGGC) and 3991 (5'-TTNCCNTCACGAGTACGNTCC), was inserted into pMOSBlue. This plasmid was used to construct a plasmid for cloning the remaining part of the *atp* operon by the plasmid rescue technique as follows. A 3.2-kb *EcoRI* fragment from pCP12 carrying the *erm* gene and the strong artificial constitutive promoter CP12 (19) was cloned into pQ1 digested with *EcoRI*, resulting in pBK105, in which the *atpGD'* genes (sequence from position 5268 to 6394) had been placed under control of the CP12 promoter. The plasmid pBK105 was then used to transform *L. lactis* MG1363 to erythromycin resistance (2 µg/ml). This plasmid is unable to replicate in *L. lactis*, and only cells with the plasmid integrated into the chromosome will become resistant to erythromycin. If the plasmid integrates into the *atp* operon by a Campbell-type event, the genes *atpDC* will come under control of the CP12 promoter. Chromosomal DNA of some transformants was prepared, and the appropriate integration of pBK105 was verified by PCR techniques. The chromosomal DNA was digested with *SalI*, ligated at a low DNA concentration, and transformed in *E. coli*, which resulted in pRESDC, in which approximately 4.5 kb downstream of the *atp* operon was cloned. Plasmid pRESDC was more extensively characterized and sequenced.

**Primer extension.** Total RNA was extracted from exponentially growing *L. lactis* (30°C, optical density at 600 nm [OD<sub>600</sub>] = 0.5) in GM17 (1% glucose) by the FastRNA kit, BLUE (Bio 101), as recommended by the manufacturer.

Total RNA (10 µg) and <sup>32</sup>P-labeled primer (10 pmol) were heated for 2 min at 80°C in 5 µl of hybridization buffer (100 mM KCl, 50 mM HEPES, pH 7.0), followed by a gradual cooling to 30°C over a 60-min period. Three microliters of a solution containing 250 mM Tris-HCl (pH 8.4), 20 mM MgCl<sub>2</sub>, 20 mM dithiothreitol (DTT), 0.1 mM concentrations of each deoxynucleoside triphosphate, and 0.75 U of avian myeloblastosis virus reverse transcriptase (Life Sciences)/µl was added, and the mixture was incubated at 40°C for 30 min. The extension product was precipitated with ethanol and resuspended in 6 µl of formamide loading buffer, preheated at 85°C for 3 min, and loaded onto a polyacrylamide gel with a set of dideoxy sequencing reactions (33) prepared on a PCR product as a marker. The sequence of the primer used in the 5'-3' direction was 5'-GACCC ATAGCAATTGCTCC-3' (primer 5264).

**Northern blotting.** A single-stranded RNA probe labeled with [<sup>32</sup>P]CTP was derived from a PCR product (primer 5883, 5'-CAACGTGTCCTTCAACGC, and primer T7atpC, 5'-TAATACGACTACTATAGATAAACCACACCAGC AGGGG), which contains *atp'DC'* (position 6918 to 7462) and the T7 promoter, by *in vitro* transcription using T7 RNA polymerase (Promega). A total RNA preparation (12 µg) was dried in a vacuum drier and resuspended in 4.5 µl of H<sub>2</sub>O, 2 µl of 5× formaldehyde gel running (FGR) buffer (0.1 M MOPS [morpholinepropanesulfonic acid] [pH 7], 40 mM sodium acetate, 5 mM EDTA), 3.5 µl of formaldehyde (final concentration, 7% [vol/vol]), and 10 µl of formamide (final concentration, 50% [vol/vol]). The RNA molecules were denatured by incubation for 15 min at 60°C and separated by electrophoresis in a 1.2% (wt/vol) agarose gel containing 2.2% formaldehyde, which was run at 5 V/cm with FGR buffer as the electrophoresis buffer. The gel was then washed in H<sub>2</sub>O for 20 min at room temperature. The RNA was transferred to a Zeta-Probe GT membrane (Bio-Rad) by overnight capillary blotting with 50 mM NaOH as the transfer buffer. The membrane was air dried and prehybridized for 2 h at 42°C in hybridization buffer (1 mM NaCl, 4 mM Na<sub>2</sub>P<sub>2</sub>O<sub>7</sub>, 5× Denhardt's solution, 1% sodium dodecyl sulfate [SDS], 10% [wt/vol] polyethylene glycol 6000, 50 mM Tris-HCl [pH 7.5], 50% [vol/vol] formamide) before the α-<sup>32</sup>P-labeled riboprobe was added. After overnight hybridization at 42°C, the membrane was washed twice for 5 min at room temperature in 2× SSC, twice at 30 min at 65°C in 0.2× SSC-1% SDS, and twice for 30 min at 65°C in 0.1× SSC before being used for autoradiography (1× SSC is 0.15 M NaCl plus 0.015 M sodium citrate). The 0.24- to 9.5-kb RNA ladder from Gibco BRL was used as a molecular size standard.

**Replacement of the chromosomal *atp* promoter in *L. lactis* by the nisin-inducible *nisA* promoter.** A PCR fragment that harbors the DNA sequence from position +998 to position 1850 (the *atpEB'* genes) was amplified using *Taq* polymerase. After polishing the DNA ends with *Pfu* polymerase, the fragment was cloned into the *SfiI* site on the vector pCR-Script Amp SK(+) (Stratagene) (Fig. 2). A plasmid was isolated in which the fragment was inserted in the orientation opposite to that of *lacZ* (pRI13). A 1.5-kb *SalI-PstI* fragment from pNZ8010 (12) that carries the *cat-194* gene and the *nisA* promoter was then cloned into pRI13 digested with *SalI-PstI*, which yielded the plasmid pATP1, in which the *atpEB'* genes had been placed under the control of the (nisin-inducible) *nisA* promoter. A 2.4-kb *ApaI-NotI* fragment from pATP1, which contains the *cat-194* gene, the *nisA* promoter, and the *atpEB'* genes, was cloned into pRC1 digested with *ApaI-NotI*, which gave rise to plasmid pNIS-ATP2. pRC1 is a 3.5-kb derivative of pBluescript II KS in which the *bla* gene has been replaced by the *ermAM* genes to allow for selection of erythromycin resistance in *L. lactis* (25). The strain NZ9000 (12) is a derivative of strain MG1363 (16) in which the *nisR* and *nisK* genes (required for induction of the *nisA* promoter) are integrated into the *pepN* locus on the chromosome. Plasmid pNIS-ATP2 was introduced into strain NZ9000 with selection for erythromycin resistance (2 µg/ml) on plates that contained nisin (5 ng/ml). Since this plasmid is unable to replicate in *L. lactis*, only cells in which the plasmid has integrated into the chromosome should become resistant to erythromycin. If the plasmid integrates into the *atpEB* locus, the transcription of the entire *atp* operon will be placed under the control of the *nisA* promoter. The clones were verified by PCR with primers positioned upstream of the *nisA* promoter and immediately downstream of position 1850.

**Nucleotide sequence accession number.** The sequence of the *lipA* gene, the sequence of the *atp* operon of *L. lactis* subsp. *cremoris* strain MG1363, and the sequence downstream of the *atp* operon (8,912 bp) have been deposited in the NCBI data bank with the accession no. AF059739, and the numbers used in the present paper refer to the numbering used in this sequence.

## RESULTS AND DISCUSSION

**The genes encoding the H<sup>+</sup>-ATPase in *L. lactis*.** The genes encoding the subunits of the H<sup>+</sup>-ATPase were cloned on a series of overlapping fragments, and the complete sequence of




FIG. 2. Cloning strategy used in the replacement of the native *atp* promoter with the nisin-inducible *nisA* promoter. (a) A 1.5-kb *Sall*-*Pst*I fragment from pNZ8010 (12) carrying the *cat-194* gene and the *nisA* promoter was cloned into pRI13 digested with *Sall*-*Pst*I (pATP1). (b) A 2.4-kb *Apa*I-*Nor*I fragment from pATP1, containing the *cat-194* gene, the *nisA* promoter, and the *atpEB'* genes, was then cloned into pRC1 digested with *Apa*I-*Nor*I (pNIS-ATP2). (c) Plasmid pNIS-ATP2 was integrated into the *atp* operon in *L. lactis* strain NZ9000 with selection for erythromycin resistance (2 µg/ml) on plates containing nisin (5 ng/ml), resulting in replacement of the native *atp* promoter with the inducible *nisA* promoter. The designation of the genes is shown above the boxes in italic letters. See Materials and Methods for further details. *S*, *Sall*; *P*, *Pst*I; *A*, *Apa*I; *N*, *Nor*I.

the *atp* operon was determined and analyzed for the presence of open reading frames (Fig. 1). Within a 7-kb region we identified eight open reading frames with putative ribosome binding sites. The deduced amino acid sequences of the eight gene products of the *L. lactis atp* operon were aligned with the corresponding amino acid sequences from other organisms, and the sequences of the *L. lactis* ATPase subunits showed good homology with those of other bacteria (Table 1). The

homologies were particularly high between *L. lactis*, *Streptococcus mutans*, and *Streptococcus bovis*, which confirms the close evolutionary relationships of these bacteria. Among the ATPase subunits, the  $\alpha$ ,  $\beta$ , and  $\gamma$  subunits from the cytoplasmic domain,  $F_1$ , were especially highly conserved. The consensus nucleotide-binding domains, Walker motifs A (GXXXXGKT) and B (L-hydrophobic-hydrophobic-hydrophobic-D) (1, 42), were also conserved in the deduced sequences of the  $\alpha$  and  $\beta$

TABLE 1. Homology between the deduced amino acid sequences of the eight *L. lactis atp* gene products and ATPase subunits from other bacteria

Source of ATPase <sup>a</sup>	% Identity (% similarity) of subunits (gene, size [aa]) <sup>b</sup>							
	c ( <i>atpB</i> , 71)	a ( <i>atpE</i> , 237)	b ( <i>atpF</i> , 168)	$\delta$ ( <i>atpH</i> , 175)	$\alpha$ ( <i>atpA</i> , 500)	$\gamma$ ( <i>atpG</i> , 289)	$\beta$ ( <i>atpD</i> , 469)	$\epsilon$ ( <i>atpC</i> , 141)
<i>B. megaterium</i>	45 (62)	38 (58)	32 (62)	20 (67)	76 (87)	48 (67)	70 (79)	31 (53)
<i>E. coli</i>	34 (72)	27 (61)	33 (55)	22 (48)	52 (85)	35 (59)	64 (90)	26 (55)
<i>S. mutans</i>	56 (76)	47 (65)	46 (71)	31 (57)	84 (92)	59 (75)	79 (85)	53 (69)
<i>Streptococcus faecalis</i>	38 (71)	51 (69)	41 (64)	27 (53)	82 (91)	59 (72)	78 (86)	52 (68)
<i>S. bovis</i>	54 (76)	45 (63)	48 (72)	34 (58)	84 (92)	63 (76)	78 (83)	53 (69)
PS3	40 (65)	35 (57)	32 (59)	24 (44)	74 (86)	46 (66)	72 (81)	34 (52)
<i>Synechococcus</i> sp.	39 (68)	31 (52)	28 (51)	24 (47)	62 (78)	32 (52)	59 (71)	35 (58)

<sup>a</sup> References of bacteria are as follows: *B. megaterium* (6), *E. coli* (43), *S. mutans* (37), *S. faecalis* (36), *S. bovis* DDBJ/EMBL/GenBank database accession no. AB009314), thermophilic bacterium PS3 (29), *Synechococcus* sp. (9).

<sup>b</sup> The genes encode the respective subunits, and the sizes of the subunits are given in amino acids (aa).








FIG. 4. Northern blot analysis. Total RNA was extracted from *L. lactis*, and Northern blot analysis was performed as described in Material and Methods. A ribonucleotide probe labeled with [ $\alpha$ - $^{32}$ P]CTP containing the C-terminus-encoding part of *atpD* and the N-terminus-encoding part of *atpD* (position 7915 to 8459) was used as a probe. The 0.24- to 9.5-kb RNA ladder from Gibco BRL was used as a molecular size standard.

around the transcription start site. The region upstream of the *atp* promoter contains several inverted and direct repeats. Such repeats were also observed in *Enterococcus faecalis* (36), and it was suggested that they may be involved in the regulation of the expression of the *atp* operon at low external pH (2, 22, 23) in order to keep the intracellular pH at an acceptable level.

The size of the *atp* mRNA was determined by Northern blot analysis (Fig. 4), which identified mRNA at approximately 7 kb, which demonstrates that the eight genes are transcribed as a single polycistronic message. Other transcripts could not be identified in the present analysis, in which the 3' end of the *atp* operon was used as a probe. But smaller transcripts might still occur if other probes are employed.

An inverted repeat in the region immediately after *atpC* was recognized, followed by a T-string (7 bp), a structure that resembles a rho-independent terminator (31). The location of

a terminator at this position is also supported by the transcript size found in the Northern analysis.

**The  $F_1F_0$ -ATPase is essential for growth of *L. lactis*.** In the anaerobic bacterium *L. lactis*, the role of the  $H^+$ -ATPase is to maintain the electrochemical proton gradient across the cytoplasmic membrane, and it has been proposed that the  $H^+$ -ATPase functions to regulate the internal pH (4, 11, 21, 24). Is the  $H^+$ -ATPase then essential for growth? In principle, the anaerobic bacteria have the option to generate a proton gradient through carrier-mediated excretion of end products in symport with protons (30).

The electrochemical proton gradient ( $\Delta p$ ) is composed of an electrical component, the transmembrane potential difference ( $\Delta\psi$ ), and a chemical component, the transmembrane pH difference ( $\Delta pH$ ). The magnitude of the energy produced by lactate excretion depends strongly on the  $H^+$ -lactate stoichiometry ( $n$ ) during the excretion process. If  $n$  is 1, the excretion process is electrochemically neutral and only a chemical gradient of protons ( $\Delta pH$ ) can be generated. If  $n$  is 2, the translocation is electrogenic and both a  $\Delta pH$  and a membrane potential ( $\Delta\psi$ ) can be formed. At high pH (6.8) and a low external lactate concentration (<5 mM), ten Brink and Konings determined the  $H^+$ -lactate stoichiometry ( $n$ ) in *L. lactis* to be 1.9 (39). Thus, in principle the contribution of  $H^+$ -lactate efflux may suffice so that the  $H^+$ -ATPase would be dispensable for growth under these conditions.

One way to test how important the  $H^+$ -ATPase is for growth of *L. lactis* would be to replace the chromosomal *atp* promoter with an inducible promoter. In order to replace the original *atp* promoter with an inducible nisin promoter (12), a plasmid, pNIS-ATP2, was constructed, which carries the *atpE* gene and part of the *atpB* gene under the control of the *nisA* promoter. This plasmid, which cannot replicate in *L. lactis*, was integrated into the chromosome of *L. lactis* as described in Materials and Methods (Fig. 2). The resulting strain contained an inducible nisin promoter upstream of the entire chromosomal *atp* operon. When the strain was grown at 30°C on GM17 plates (buffered at pH 7) with different concentrations of nisin, we observed that at very low nisin concentrations the growth of the strain decreased dramatically and in the absence of nisin, growth was completely abolished (Fig. 5). This demonstrates




FIG. 5. Colonies of strain *L. lactis* PJ4700, in which the native *atp* promoter had been replaced by a *nisA* promoter. The strain was streaked on GM17 plus 2  $\mu\text{g}$  of erythromycin/ml at various nisin concentrations (0, 0.25, 0.5, 1, 2, 4, 8, and 16 ng of nisin/ml), and the graph illustrates the average diameter of colonies obtained with the different nisin concentrations.

that the H<sup>+</sup>-ATPase is essential for growth of *L. lactis* under these conditions, presumably because it is essential for maintaining the proton gradient necessary for solute transport and for maintaining the cytoplasmic pH at an acceptable level. This is also in agreement with the observation that the activity of the F<sub>1</sub>F<sub>o</sub>-ATPase in related anaerobic bacteria is enhanced at low external pH (2, 23).

#### ACKNOWLEDGMENTS

We thank Regina Shürmann for excellent technical assistance and Inge Knudsen and Raino K. Hansen for having cloned and sequenced a part of the *atp* operon. We are also grateful to Allan K. Nielsen for his support with the primer extension and Northern blot analysis and to Lene Kragelund for her kind assistance with the autosequencing at Chr. Hansen A/S.

This work was supported by The Danish Academy of Technical Sciences (ATV) and Chr. Hansen A/S.

#### REFERENCES

- Abrahams, J. P., A. G. W. Laslie, R. Lutter, and J. E. Walker. 1994. Structure at 2.8 Å resolution of F<sub>1</sub>-ATPase from bovine heart mitochondria. *Nature* **370**:621–628.
- Abrams, A., and C. Jensen. 1984. Altered expression of the H<sup>+</sup> ATPase in *Streptococcus faecalis* membranes. *Biochem. Biophys. Res. Commun.* **122**:151–157.
- Araya, T., N. Ishinashi, S. Shimamura, K. Tanaka, and H. Takahashi. 1993. Genetic and molecular analysis of the *rpoD* gene from *Lactococcus lactis*. *Biosci. Biotechnol. Biochem.* **57**:88–92.
- Bender, W. A., and R. E. Marquis. 1986. Acid tolerance, proton permeabilities, and membrane ATPases of oral streptococci. *Infect. Immun.* **53**:331–338.
- Bracco, L., D. Kortlarz, A. Kolb, S. Diekmann, and H. Buc. 1989. Synthetic curved DNA sequences can act as transcriptional activators in *Escherichia coli*. *EMBO J.* **8**:4289–4296.
- Brusilow, W. S., M. A. Scarpetta, C. A. Hawthorne, and W. P. Clark. 1989. Organization and sequence of the genes coding for the proton-translocating ATPase of *Bacillus megaterium*. *J. Biol. Chem.* **264**:1528–1533.
- Casabadan, M. J., and S. N. Cohen. 1980. Analysis of gene control signals by DNA fusion and cloning in *Escherichia coli*. *J. Mol. Biol.* **138**:179–207.
- Chiaruttini, C., and M. Millet. 1993. Gene organization, primary structure and RNA processing analysis of a ribosomal RNA operon in *Lactococcus lactis*. *J. Mol. Biol.* **230**:57–76.
- Cozens, A. L., and J. E. Walker. 1987. The organization and sequence of the genes for ATP synthase subunits in the cyanobacterium *Synechococcus* 6301. Support for an endosymbiotic origin of chloroplasts. *J. Mol. Biol.* **194**:359–383.
- Das, A., and L. G. Ljungdahl. 1997. Composition and primary structure of the F<sub>1</sub>F<sub>o</sub> ATP synthase from the obligately anaerobic bacterium *Clostridium thermoaceticum*. *J. Bacteriol.* **179**:3746–3755.
- Dashper, S. G., and E. C. Reynolds. 1992. pH regulation by *Streptococcus mutans*. *J. Dent. Res.* **71**:1159–1165.
- de Ruyter, P. G., O. P. Kuipers, and W. M. de Vos. 1996. Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin. *Appl. Environ. Microbiol.* **62**:3662–3667.
- Fenoll, A., R. Munoz, E. Garcia, and A. D. de la Campa. 1994. Molecular basis of the optochin-sensitive phenotype of pneumococcus: characterization of the genes encoding the F<sub>o</sub> complex of the *Streptococcus pneumoniae* and *Streptococcus oralis* H<sup>+</sup>-ATPases. *Mol. Microbiol.* **12**:587–598.
- Futai, M., and H. Kanazawa. 1983. Structure and function of proton-translocating adenosine triphosphatase (F<sub>1</sub>F<sub>o</sub>): biochemical and molecular biological approaches. *Microbiol. Rev.* **47**:285–312.
- Gartenberg, M. R., and D. M. Crothers. 1991. Synthetic DNA bending sequences increase the rate of *in vitro* transcription initiation at the *Escherichia coli* promoter. *J. Mol. Biol.* **219**:217–230.
- Gasson, M. J. 1983. Plasmid complements of *Streptococcus lactis* NCDO 712 and other lactic streptococci after protoplast-induced curing. *J. Bacteriol.* **154**:1–9.
- Gay, N. J. 1984. Construction and characterization of an *Escherichia coli* strain with a *uncI* mutation. *J. Bacteriol.* **158**:820–825.
- Holo, H., and I. F. Nes. 1989. High-frequency transformation, by electroporation, of *Lactococcus lactis* subsp. *cremoris* grown with glycine in osmotically stabilized media. *Appl. Environ. Microbiol.* **55**:3119–3123.
- Jensen, P. R., and K. Hammer. 1998. The sequence of spacers between the consensus sequences modulates the strength of prokaryotic promoters. *Appl. Environ. Microbiol.* **64**:82–87.
- Jensen, P. R., and O. Michelsen. 1992. Carbon and energy metabolism of *atp* mutants of *Escherichia coli*. *J. Bacteriol.* **174**:7635–7641.
- Kobayashi, H. 1985. A proton-translocating ATPase regulates pH of the bacterial cytoplasm. *J. Biol. Chem.* **260**:72–76.
- Kobayashi, H., N. Murakami, and T. Unemoto. 1982. Regulation of the cytoplasmic pH in *Streptococcus faecalis*. *J. Biol. Chem.* **257**:13246–13252.
- Kobayashi, H., T. Suzuki, N. Kinoshita, and T. Unemoto. 1984. Amplification of the *Streptococcus faecalis* proton-translocating ATPase by a decrease in cytoplasmic pH. *J. Bacteriol.* **158**:1157–1160.
- Kobayashi, H., T. Suzuki, and T. Unemoto. 1986. Streptococcal cytoplasmic pH is regulated by changes in amount and activity of a proton-translocating ATPase. *J. Biol. Chem.* **261**:627–630.
- Le Bourgeois, P., M. Lautier, M. Mata, and P. Ritzenthaler. 1992. New tools for the physical and genetic mapping of *Lactococcus* strains. *Gene* **111**:109–114.
- McCarn, D. F., R. A. Whitaker, J. Alam, J. M. Vrba, and S. E. Curtis. 1988. Genes encoding the alpha, gamma, delta, and four F<sub>o</sub> subunits of ATP synthase constitute an operon in the cyanobacterium *Anabaena* sp. strain PCC 7120. *J. Bacteriol.* **170**:3448–3458.
- Michels, P. A., J. P. Michels, J. Boonstra, and W. N. Konings. 1979. Generation of an electrochemical proton gradient in bacteria by the excretion of metabolic end products. *FEMS Microbiol. Lett.* **5**:357–364.
- Miller, J. H. 1972. Experiments in molecular genetics, p. 352–355. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Ohta, S., M. Yohda, M. Ishizuka, H. Hirata, T. Hamamoto, Y. Otawara-Hamamoto, K. Matsuda, and Y. Kagawa. 1988. Sequence and overexpression of subunits of adenosine triphosphate synthase in thermophilic bacterium PS3. *Biochim. Biophys. Acta* **933**:141–155.
- Otto, R., A. S. Sonnenberg, H. Veldkamp, and W. N. Konings. 1980. Generation of an electrochemical proton gradient in *Streptococcus cremoris* by lactate efflux. *Proc. Natl. Acad. Sci. USA* **77**:5502–5506.
- Platt, T. 1986. Transcription termination and the regulation of gene expression. *Annu. Rev. Biochem.* **55**:339–372.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
- Santana, M., M. S. Ionescu, A. Vertes, R. Longin, F. Kunst, A. Danchin, and P. Glaser. 1994. *Bacillus subtilis* F<sub>1</sub>F<sub>o</sub> ATPase: DNA sequence of the *atp* operon and characterization of *atp* mutants. *J. Bacteriol.* **176**:6802–6811.
- Saraste, M., N. J. Gay, A. Eberle, M. J. Runswick, and J. E. Walker. 1981. The *atp* operon: nucleotide sequence of the genes for the γ, β, and ε subunits of *Escherichia coli* ATP synthase. *Nucleic Acids Res.* **9**:5287–5296.
- Shibata, C., T. Ehara, K. Tomura, K. Igarashi, and H. Kobayashi. 1992. Gene structure of *Enterococcus hirae* (*Streptococcus faecalis*) F<sub>1</sub>F<sub>o</sub>-ATPase, which functions as a regulator of cytoplasmic pH. *J. Bacteriol.* **174**:6117–6124.
- Smith, A. J., R. G. Quivey, Jr., and R. C. Faustoferri. 1996. Cloning and nucleotide sequence analysis of the *Streptococcus mutans* membrane-bound, proton-translocating ATPase. *Gene* **183**:87–96.
- Suzuki, T., and H. Kobayashi. 1989. Regulation of the cytoplasmic pH by a proton-translocating ATPase in *Streptococcus faecalis* (*faecium*). A computer simulation. *Eur. J. Biochem.* **180**:467–471.
- ten Brink, B., and W. N. Konings. 1982. The electrochemical proton gradient and lactate concentration gradient in *Streptococcus cremoris* grown in batch culture. *J. Bacteriol.* **152**:682–686.
- ten Brink, B., R. Otto, U. P. Hansen, and W. N. Konings. 1985. Energy recycling by lactate efflux in growing and nongrowing cells of *Streptococcus cremoris*. *J. Bacteriol.* **162**:383–390.
- van de Guchte, M., J. Kok, and G. Venema. 1992. Gene expression in *Lactococcus lactis*. *FEMS Microbiol. Rev.* **88**:73–92.
- Walker, J. E., M. Saraste, J. Runswick, and N. J. Gay. 1982. Distantly related sequences in the α- and β-subunits of ATP synthase, myosin, kinases and other ATP requiring enzymes and a common nucleotide binding fold. *EMBO J.* **1**:945–951.
- Walker, J. E., M. Saraste, and J. N. Gay. 1984. The *unc* operon: nucleotide sequence, regulation and structure of ATP-synthase. *Biochim. Biophys. Acta* **768**:164–200.