

# Photo-cycloaddition reactions of vinyl diazo compounds

Received: 18 October 2023

Accepted: 26 April 2024

Published online: 29 May 2024

Check for updates

Ming Bao<sup>1</sup>, Klaudia Łuczak<sup>2</sup>, Wojciech Chaładaj<sup>2</sup>✉, Marriah Baird<sup>1</sup>,  
Dorota Gryko<sup>2</sup>✉ & Michael P. Doyle<sup>1</sup>✉


Heterocyclic rings are important structural scaffolds encountered in both natural and synthetic compounds, and their biological activity often depends on these motifs. They are predominantly accessible via cycloaddition reactions, realized by either thermal, photochemical, or catalytic means. Various starting materials are utilized for this purpose, and, among them, diazo compounds are often encountered, especially vinyl diazo compounds that give access to donor-acceptor cyclopropenes which engage in  $[2+n]$  cycloaddition reactions. Herein, we describe the development of photochemical processes that produce diverse heterocyclic scaffolds from multisubstituted oximido-vinyl diazo compounds. High chemoselectivity, good functional group tolerance, and excellent scalability characterize this methodology, thus predisposing it for broader applications. Experimental and computational studies reveal that under light irradiation these diazo reagents selectively transform into cyclopropenes which engage in cycloaddition reactions with various dipoles, while under thermal conditions the formation of pyrazole from vinyl diazo compounds is favored.

The development of effective, sustainable methodologies giving access to valuable heterocyclic molecules is of paramount importance to modern synthetic chemistry. Among approaches to their synthesis, cycloaddition reactions stand at the forefront, and the use of visible light as the only source of energy for these transformations is highly appealing. Recently, diazo compounds that produce various reactive intermediates have attracted considerable interest<sup>1–7</sup>. In particular, vinyl diazo compounds have proven to be valued reactants for the synthesis of diverse carbo- and heterocyclic compounds<sup>8–10</sup>, especially through cycloaddition reactions<sup>11,12</sup>. Their catalytic applications include highly enantioselective  $[3+3]$ -,  $[3+2]$ -, and  $[3+1]$ -cycloaddition<sup>13</sup>. Furthermore, silyl-protected enol diazoacetates are known to form stable donor-acceptor cyclopropenes that provide a resting state for incipient metallovinyl carbenes in cycloaddition reactions<sup>14</sup>. Their photochemically induced dinitrogen extrusion to form vinylcarbene intermediates has been investigated<sup>15,16</sup>, and the resultant formation of cyclopropene products with alkyl and aryl substituents is well known<sup>17–19</sup>. Cyclopropenes are highly strained


alkenes that are, themselves, highly reactive in cycloaddition reactions<sup>20,21</sup>; they are activated by strain to undergo “click”-like cycloaddition reactions and although few reports have documented these transformations<sup>13,22–25</sup>, they have already been employed for site-specific protein conjugation<sup>26,27</sup>, cell labeling<sup>28,29</sup> and other materials development functions<sup>30,31</sup>.

The linkage between vinyl diazo compounds and cyclopropenes provides versatility in product formation. Donor-acceptor cyclopropenes are produced catalytically from silyl-protected enoldiazo compounds via metallovinyl carbene intermediates<sup>32</sup>, as well as thermally<sup>33</sup> in nearly quantitative yield (Fig. 1A).  $\beta$ -Aryl/alkyl vinyl diazo carbonyl compounds undergo non-reversible catalytic formation of unstable cyclopropenes via metallovinyl carbenes<sup>34</sup>, but thermally they form 3*H*-pyrazoles that transform into 1*H*-pyrazoles by 1,5-H transfer<sup>35</sup>. In particular, styryldiazoacetates undergo the thermal formation of 1*H*-pyrazoles, but they do not form cyclopropenes in catalytic reactions<sup>36</sup>. In catalytic reactions with vinyl diazo compounds, dipolar species can intercept metallo-vinylcarbenes prior to cyclopropene

<sup>1</sup>Department of Chemistry, The University of Texas at San Antonio, San Antonio, Texas, USA. <sup>2</sup>Institute of Organic Chemistry Polish Academy of Sciences, Kasprzaka 44/52, Warsaw, Poland. ✉ e-mail: [wojciech.chaladaj@icho.edu.pl](mailto:wojciech.chaladaj@icho.edu.pl); [dorota.gryko@icho.edu.pl](mailto:dorota.gryko@icho.edu.pl); [michael.doyle@utsa.edu](mailto:michael.doyle@utsa.edu)



**Fig. 1 | Transformations of vinyl diazo compounds.** **A** Catalytic and thermal formation of cyclopropenes from vinyl diazo compounds; **B** Cyclopropene cycloaddition reactions; **C** Tandem cycloaddition reactions of vinyl diazo compounds.



**Fig. 2 | Photoreactivity of vinyl diazo compounds.** **A** Transformation of 3-phenyl-2-diazo butanoate. **B** Calculated reactivity of methyl 2-diazo-3-butenate in the ground and excited states.


formation but with suitable donor-acceptor substituents; the cyclopropene can be returned to the metallo-vinylcarbene to produce [3+n]-cycloaddition products<sup>14</sup>. Because of their inherent reactivity, the double bond in cyclopropenes undergoes [2+n]-cycloaddition with viable dipolar species (Fig. 1B)<sup>20,21,37,38</sup>. However, although vinyl diazo compounds are precursors to cyclopropenes, they have not been used as direct precursors for [2+n]-cycloaddition products that can be achieved with cyclopropenes; instead, the [2+n]-cycloaddition products are formed from cyclopropenes, which, in turn, are separately prepared from vinyl diazo compounds or by alternative methods. We now report a general, efficient methodology for the conversion of vinyl diazo compounds to [2+n]-cycloaddition products through the photochemical production of intermediate cyclopropenes (Fig. 1C).

## Results and Discussion


### Generation of cyclopropenes from vinyl diazo compounds

Vinyl diazo compounds form either cyclopropenes (dinitrogen extrusion)<sup>33,39</sup> or 1H-pyrazoles (cycloaddition/rearrangement)<sup>34,36</sup>. This dichotomy is found in the photochemical and thermal reactions of  $\beta$ -aryl-/alkyl-vinyl diazoacetates. The thermal reaction of 3-phenyl-2-diazo-3-butenate (**1a**) produces corresponding 1H-pyrazole **2a** in high yield, whereas dinitrogen is lost in the photochemical reaction to give cyclopropene dimer **4a** resulting from an ene-reaction<sup>40</sup> of the unstable cyclopropene intermediate **3a** (Fig. 2A). The same **4a** is obtained in the rhodium acetate catalyzed reaction of **1a**<sup>34</sup>.

To understand the basis for the divergent outcomes from thermal and photochemical reactions, the chemistry of model methyl



**Fig. 3** | [3+2]-Cycloaddition reaction of diazo compounds. **A** Model reaction of oximidovinyldiazo acetate **5a** with nitron **6**. **B** DFT calculations on the cycloaddition [3+2] of model cyclopropene with nitron **6**.




**Fig. 4** | Cycloaddition reactions of oximidovinyldiazo compounds with various dipoles. **A** Model reaction of oximidovinyldiazo acetate **5** with nitron **6**. **B** Model reaction of oximidovinyldiazo acetate **5** with *N,N*-cyclic azamethine ylide **8**. **C** Model reaction of oximidovinyldiazo acetate **5** with nitrile oxide **10**.

2-diazobut-3-enoate **I** was investigated computationally on S0 and S1 potential energy surfaces (Fig. 2B)<sup>41,42</sup>. In the ground state (S0 PES), cyclization of the model vinyl diazoacetate **I** towards pyrazole **II** is preferred by 22.5 kJ/mol over extrusion of nitrogen ( $\Delta G^\ddagger = 115.6$  and 138.1 kJ/mol, respectively, Fig. 2B), although both are hardly accessible at ambient temperature. The latter of the discussed pathways provides carbene **III**, which easily ( $\Delta G^\ddagger = 13.4$  kJ/mol) cyclizes, yielding cyclopropane **IV**. The dimerization of **IV** via an ene-reaction must overcome a barrier of  $\Delta G^\ddagger = 110.8$  kJ/mol. A similar reactivity pattern was observed, i.e., preference for cyclization over extrusion of N<sub>2</sub> under thermal conditions, for vinyl diazo compounds substituted with simple alkyl and aryl groups (see, SI). Conversely, the calculated mechanistic scenario on the S1 PES is considerably different. The cycloisomerization of methyl 2-diazobut-3-enoate **VIII** to pyrazole system **X** not only features a considerable barrier ( $\Delta G^\ddagger = 118.1$  kJ/mol) but is also highly endergonic ( $\Delta G = 92.3$  kJ/mol). In contrast, the excited vinyl diazoacetate **VIII** loses dinitrogen in a practically barrierless event ( $\Delta G^\ddagger = 3.4$  kJ/mol). However, the subsequent cyclization of

resulting carbene **IX** into a cyclopropane ring cannot occur at S1 PES (high-energy cyclopropane in the excited state spontaneously opens towards the vinyl carbene). Thus, it presumably first undergoes internal conversion to ground state **III**, at which it then converts into cyclopropane system **IV**. In contrast, cyclization of carbene **VI** in the triplet state, which is a ground state<sup>15,16</sup>, through **TSS** is associated with a very high barrier of  $\Delta G^\ddagger = 188.0$  kJ/mol, which is not accessible under the reaction conditions.

### Cycloaddition reactions of vinyl diazo compounds


Based on these theoretical studies, we expected that cyclopropenes could be generated from vinyl diazo compounds under light irradiation and, as such, engage in reactions with dipoles. In this regard, we previously observed traces of [3+2]-cycloaddition product formed via cyclopropane intermediates, accompanying the intended metallovinylcarbene [3+*n*]-cycloaddition product, in the Rh-catalyzed reaction of enol diazoacetates with isoquinolinium dicyanomethylides<sup>32,34</sup>. To evaluate the possibility of developing photochemical tandem reaction,



**Fig. 5 | Photocatalytic formal [3 + 2]-cycloaddition of diverse oximidovinyl diazo esters with nitrones. A** The general transformation; **B** Reaction scope of oximidovinyl diazoacetates; **C** Reaction scope of nitrones; **D** Reaction of other types of vinyl diazo compounds with nitrones; **E** Reaction of silyl-protected enol diazoacetate **48**.

generation of the cyclopropene from vinyl diazo compounds followed by its reaction with dipoles, leading to heterocycles, oximidovinyl diazo acetate **5a** was chosen as a model diazo compound (Fig. 3). Such vinyl diazo reagents are conveniently prepared from 1,2,3-triazine 1-oxides<sup>43</sup>. Since cyclopropene cycloaddition with nitrones is known to occur in reactions involving metallo-vinylcarbene intermediates<sup>44–48</sup>,

we commenced our studies with nitrones **6a** as a reliable model dipole<sup>49</sup>. Photolysis with blue light gave the expected isooxazolidine cycloaddition product **7** as a mixture of diastereoisomers (7:1 ratio) in 91% yield. Control experiments revealed that the reaction is photochemical in nature. Under light irradiation cyclopropene forms, as confirmed by the <sup>1</sup>H NMR analysis, and after the addition of the



**Fig. 6 | Photocatalytic formal [3 + 2]-cycloaddition of diverse oximodivinyldiazo esters with azomethine imines. A** The general transformation; **B** Reaction scope of oximodivinyldiazoacetates and azomethine imines; **C** Reaction of other types of vinyl diazo compounds with azomethine imines.

nitron, thermal cycloaddition occurs. Because cyclopropene can undergo rapid dimerization via an ene reaction<sup>40</sup>, intermediate **3b** was not isolated in a pure form. According to DFT calculations, cycloaddition of nitron **6** to the model cyclopropene (TMS protected) proceeds with a Gibbs free energy of activation of 85.2 and 90.0 kJ/mol for major and minor isomers, respectively, which matches the observed diastereoselectivity for the reaction with cyclopropene **3b**. Expectedly, for the reaction of simple acrylate (not a strained analog of cyclopropene) with nitron **6** a barrier of 109.1 kJ/mol was calculated, corroborating the higher reactivity of strained cyclopropene over acrylate.

To test the generality of the developed tandem transformation, we also explored azomethine imines and nitrile oxides as other dipole-type substrates. Both reactions gave [3 + 2] products; for azomethine ylide **8** cycloadduct **9** was observed, while an unexpected pyridine *N*-oxide **11** formed from the cycloadduct produced in reactions of some oximodivinyldiazo esters with 2,4,6-trimethylbenzotriole oxide (**10**) (Fig. 4).


Realizing the generality of the proposed strategy and the importance of heterocyclic scaffolds, we next optimized the reaction conditions and evaluated the substrate scope for the newly developed [2 + 3]-cycloaddition reactions.

**Cycloaddition of oximodivinyldiazo acetates with nitrones.** The model reaction of oximodivinyldiazo acetate **5a** with diphenylmethanimine oxide (**6**) was performed in different solvents, and the highest *dr* ratio (7:1) and yield (91% isolated) was obtained in acetone (see Table S1 in SI). Reactions in chlorocarbon solvents (dichloromethane, chloroform, 1,2-dichloroethane) also occurred in high yields, but their *dr* were significantly diminished to only 1:2 to 1:3. Using LED light sources (40 W) at 400 nm produced the cycloaddition product **7** in only 30% yield. The dominant stereoisomer in all cases is the one in which the phenyl group is *trans* to the carboethoxy group. The stereochemistry of the two diastereomers was determined spectroscopically based on the X-ray structure of the dominant isomer **30** after desilylation (Fig. 5).

The scope of the reaction was determined with a broad spectrum of oximodivinyldiazo compounds **5** and a representative selection of nitrones **6** (Fig. 5).

Reactions with alkyl and aryl substituted oximodivinyldiazo compounds **5** under the optimal conditions gave isooxazolidine products **7**, **12–24** in high yields and modest *dr* values (Fig. 5B). Steric factors appear to play a major role in determining diastereoselectivity. The highest *dr* was achieved with the vinyl diazo ester without a






**Fig. 7 | Photocatalytic formal [3 + 2]-cycloaddition of oximidovinyl diazo esters with nitrile oxide compounds.** **A** The general transformation; **B** Formation of pyridine-*N*-oxides from photolytic cyclization/silica induced rearrangement. X-ray

structure of **56** is of the hydrogen-bonded dimer; **C** Reaction scope for [3 + 2]-cycloaddition of oximidovinyl diazoacetates **5** with mesitylnitrile oxide (**10**); **D** Reaction scope of oximidovinyl diazo acetate **5a** with various aryl nitrile oxides.

substituent at the gamma position (**22**), and the lowest *dr* intriguingly comes from the reaction of the vinyl diazo ester with a bulky menthyl ester of oximidovinyl diazoacetate **24**. In all these reactions, the competing ene reaction of intermediate cyclopropane **3b** was only a minor component of the reaction products, amounting to less than 10% yield; the only exception being the reaction of vinyl diazo ester with no substituent at the gamma position leading to product **22**.

Similarly, the size and the nature of substituents on the nitron influence the diastereoselectivity of the reaction (Fig. 5C). For Ar and alkyl substituted nitrones, yields remained at the same level (**25–33**) while for glyoxylic acid derived nitron reactions were less efficient (**34, 36**). The yields, however, improved when deprotected oximidovinyl diazoacetate was used as a starting material, suggesting that the free hydroxyl group, by forming hydrogen bonds influences the reaction's efficacy (products **35, 37**). The biggest impact on the stereoselectivity of the reaction had the replacement of *N*-Bn with bulky *t*Bu group for which reactions produced only one diastereoisomer (compare **34** with **36**).

Oximidovinyl diazo compounds **5** are stable reagents in contrast to vinyl diazo and aryl vinyl diazo derivatives. However, when tested in the developed method, gratifyingly, desired cycloaddition products **38–47** were also formed from these less stable derivatives, though with slightly diminished yields compared to reactions with oximidovinyl diazoacetates (Fig. 5D). The outcome of the photolytic reaction with aryl-substituted vinyl diazoacetates show contrasting behavior; with  $\beta$ -phenyl vinyl diazoacetate the yields and *dr* of isooxazolidine cycloaddition products **38–40** were comparable to those with similarly substituted oximidovinyl diazoacetates, whereas with  $\gamma$ -phenyl vinyl diazoacetate (styryl diazoacetate) competition with intramolecular pyrazole **2a** formation reduced the yield of the isooxazolidine cycloaddition product **41** which was formed with high diastereoselectivity. In general, reactions of vinyl diazo compounds with nitrones derived from glyoxylic acid ester were less efficient. Though diastereoselectivity remained at the same level for *N*-Bn derivatives, *N*-*t*Bu analogs proved superior in furnishing products **42** and **44** as single diastereoisomers.



**Fig. 8 | Reaction of other vinyl diazo compounds with nitrile oxide. A** Cycloaddition of other types of vinyl diazo compounds with aryl nitrile oxides. **B** Photocatalytic formal [3 + 2]-cycloaddition of 3-substituted vinyl diazo compounds with nitrile oxide compounds.

Furthermore, preliminary experiments confirmed that silyl-protected enol diazoacetate **48**, which forms a stable donor-acceptor cyclopropene, is also a suitable substrate in the developed tandem transformation (Fig. 5E). In this case, the bicyclic product was, however, not isolated. During the purification, the TIPS protecting group cleaves inducing the subsequent rearrangement giving product **50** in 39% yield.

**Cycloaddition of oximido vinyl diazo acetates with *N,N*-cyclic azamethine imines.** *N,N*-Cyclic azamethine imines are also suitable dipoles in cycloaddition reactions that have been reported to undergo [3 + 2]-cycloaddition reactions with propargylic and  $\alpha,\beta$ -unsaturated carbonyl compounds<sup>50,51</sup>, and cyclic enamines<sup>52</sup>. With vinyl diazo compounds, these azamethine imines are unreactive unless a transition metal catalyst converts the vinyl diazo compound to a metalvinylcarbene which then undergoes a N-N cleavage reaction to form a diimide with azomethine rather than [3 + 3]-cycloaddition<sup>53,54</sup>. In our case, photochemical tandem reaction of *N,N*-cyclic azamethine ylides **8** with vinyl diazo compound **5a** gave tricyclic pyrazolone **9** without the use of any catalyst (Fig. 6B).


Yields of cycloaddition reactions are comparable to those obtained for nitrones without the need for fine tuning of the reaction conditions. The competing ene dimer **4b** was obtained at most in less than 10% yield. Diastereoselectivities are also on a similar level with the *trans*-R/Ar isomer dominant (configuration certified by X-ray analysis). Interestingly, the reaction with styryldiazoacetate gave the [3 + 2]-cycloaddition product **59** in high yield but with no diastereoselectivity and without the evidence of pyrrole formation (Fig. 6C). In contrast, only one diastereoisomer **60** is produced from the reaction with the unsubstituted ethyl 2-diazo-3-butenoate. With ethyl 3-phenyl-2-diazo-3-butenoate, only the ene dimer **4a** was isolated, and there was no evidence for the formation of the [3 + 2]-cycloaddition product, whereas reaction with ethyl 3-methyl-2-diazo-3-butenoate resulted in a

high yield of the cycloaddition product **61** although with low diastereoselectivity (Fig. 6C).


**Cycloaddition of oximido vinyl diazo acetates with nitrile oxides.** Nitrile oxides are relatively reactive dipolar species that are well known to undergo [3 + 2]-cycloaddition reactions with alkenes and alkynes to form diverse heterocyclic compounds<sup>55–57</sup>, that have been used in the synthesis of natural products<sup>58</sup>. We selected the relatively stable aromatic nitrile oxide **10a** with the mesityl aromatic ring that does not dimerize and combined this nitrile oxide with vinyl diazoesters. As anticipated, under photolysis with blue light, in the presence of nitrile oxide **10a**, cyclopropane intermediate **3a** undergoes rapid [3 + 2]-cycloaddition with high diastereoselectivity (Fig. 7).

Intriguingly, some of the cycloaddition products converted to new chemical structures (**11**, **62–65**) during chromatography on silica gel, of which one was identified both spectroscopically and by X-ray crystallography to be pyridine-*N*-oxides **11** (Fig. 7B). However, [3 + 2]-cycloaddition products **66–72** did not undergo rearrangement on silica gel even over long periods of time, or in the presence of Lewis acids (Fig. 7C). Products **73–80**, obtained from other aryl nitrile oxides in high yield, were stable and, intriguingly, did not rearrange regardless of the position of substituents and the electronic character of the phenyl ring of nitrile oxide (Fig. 7D). Reactions with vinyl diazo esters other than the oximido vinyl diazo esters **5** also formed stable cycloaddition products (**81** and **82**) but with low stereoselectivity (Fig. 8A). As acidic conditions seemed to promote conversion of isoxazole products to pyridine-*N*-oxides, we wondered whether Sc(OTf)<sub>3</sub> treatment of the products from [3 + 2]-cycloaddition of 3-substituted vinyl diazo esters with mesityl nitrile oxide (**10a**) would lead to pyridine derivatives (Fig. 8B). In this case however, the reaction resulted in ring opening of the cyclopropane ring and formation of 2*H*-1,2-oxazine **83–85** in the absence of water or afforded cyclic product **86** in the presence of water. On the other hand, ketenimine products (**87** and

## A) Mechanistic experiments



## B) Proposed mechanism



**Fig. 9** | Mechanistic studies for rearrangement of the [3 + 2]-cycloaddition product from ketenime intermediate to pyridine-*N*-oxide, furan, and ester/amide products. **A** Control experiments. **B** Proposed mechanism.

**88**) were obtained in the two-step reaction from ethyl 2-diazo-3-phenylbut-3-enoate with aryl nitrile oxides (Fig. 8B).

To determine how the pyridine-*N*-oxide was formed, we subjected several of the [3 + 2]-cycloaddition products to heating at reflux in chloroform. Only spirocyclopentyl derivative **72** underwent reaction and the product of this thermal reaction was ketenimine **89**, anticipated to be formed by the Lossen rearrangement, which was previously reported to occur in a dirhodium(II)-catalyzed process involving formation of an intermediate [3 + 2]-cycloaddition product similar to **89** (Fig. 9A, a)<sup>59</sup>. Indeed, when cycloaddition product **90** from cycloaddition of 4-fluorophenyl oximidodiazooacetates with mesitylnitrile oxide **10a** was treated with AcOH in chloroform, the mixed ester/amide **91** as formed in 90% yield, which is similar to the product from the dirhodium(II) catalyzed reaction, thus also pointing to a ketenimine intermediate (Fig. 9A, b). The same treatment of the

benzyl and cyclohexyl analogs of **68** produced the corresponding ester/amides in 77% (2:1 *dr*) and 83% (3:1 *dr*) yields, respectively. Consistent with this intermediate, treatment of **90** with the less acidic 2,2,2-trifluoroethanol or hexafluoroisopropyl alcohol (HFIP) formed furan **92** in 57% and 78% yield, respectively, that further confirms the ketenimine intermediate (Fig. 9A, c).

The formation of ketenimines intermediates from the isoxazole-related products obtained from nitrile oxides explains the formation of pyridine-*N*-oxides, ester/amides, and furan products. In the proposed mechanism, ketenimine **A** is formed by the Lossen rearrangement<sup>60,61</sup>, although the reactivity of the [3 + 2]-cycloaddition products towards this rearrangement is not evident. This multi-substituted intermediate has two basic centers (the imine and oxime nitrogens) whose protonation influences subsequent reactions (Fig. 9B). Although protonation may not be required for the formation



of the pyridine-*N*-oxide **B** (path a), protonation of the imine nitrogen of intermediate, **A** activates the central ketenimine carbon for nucleophilic attack by the ketone oxygen to form furan derivative **C** (path b). If both basic nitrogens are protonated, neither intramolecular reaction occurs, and ketenimine hydrolysis occurs to produce **D** (path c).

## Conclusions

In conclusion, we have found that direct photolysis of vinyl diazo compounds selectively leads to cyclopropenes, which contrasts with the same reactions under thermal conditions that favors the formation of pyrazoles. Once formed, these reactive cyclopropene intermediates undergo [3+2]-cycloaddition with diverse dipolar species to yield heterocyclic scaffolds, highly prized by the pharmaceutical industry. Reactions of vinyl diazo compounds with nitrones, *N,N*-cyclic azamethine ylides, and nitrile oxides afforded heterocyclic products in high yields with mostly good stereoselectivities. The most diastereoselective reactions with nitrones were those derived from *N*-*t*Bu-glyoxylic acid, for which single isomers were observed. *N,N*-cyclic azamethine ylides react with the photochemically-generated cyclopropenes in modest to good yields and diastereoselectivities, and the transformation is general for vinyl diazoacetates. Nitrile oxides show similar generality in its cycloaddition reactions with photochemically-generated cyclopropenes, but its bicyclic isoxazole products exhibit a diversity of reactions and reactivities, some of which, especially the formation of pyridine-*N*-oxides, were unexpected. A great variety of heterocyclic scaffolds accessible by the developed method emphasizes the importance of cyclopropene generated from diazo compounds in a photochemical manner. Further studies of its unique reactivity are currently underway in our laboratory.

## Methods

### General procedure for [3 + 2]-cycloadditions with nitrones

To a 10-mL oven-dried vial with a magnetic stirring bar, vinyl diazo compound (0.1 mmol) in 1.0 mL acetone was added over 1 min to a solution of nitrones (0.12 mmol, 1.2 equiv.) in the same solvent (1.0 mL) at room temperature with irradiation by 440 nm blue LED (40 W), and the reaction mixture was stirred for 5–15 min under these conditions. When the reaction was complete (monitored by TLC), the reaction mixture was purified by flash column chromatography on silica gel without additional treatment (hexanes: EtOAc = 20:1 to 15:1) to give the pure [3 + 2]-cycloaddition products in good yields.

### General procedure for [3 + 2]-cycloaddition with azamethine imines

To a 10-mL oven-dried vial with a magnetic stirring bar, vinyl diazo compound (0.12 mmol, 1.2 equiv.) in 1.0 mL acetone was added to a solution of azamethine imine (0.1 mmol) in the same solvent (1.0 mL) via a syringe pump over 2 h at room temperature with irradiation by 440 nm blue LED (40 W). When the reaction was complete (monitored by TLC), the reaction mixture was purified by flash column chromatography on silica gel without additional treatment (hexanes: EtOAc = 20:1 to 15:1) to give the pure [3 + 2]-cycloaddition products in good yields.

### General procedure for [3 + 2]-cycloaddition with nitrile oxides

To a 10-mL oven-dried vial with a magnetic stirring bar, vinyl diazo compound (0.12 mmol, 1.2 equiv.) in 1.0 mL DCM was added to a solution of the nitrile oxide (0.1 mmol) in the same solvent (1.0 mL) via a syringe pump over 1 h at room temperature with irradiation by 440 nm blue LED. When the reaction was complete (monitored by TLC), the reaction mixture was purified by flash column chromatography on silica gel without additional treatment (hexanes: EtOAc = 20:1 to 1:1) to give the pure [3 + 2]-cycloaddition and cycloaddition/rearrangement products in good yields.

## Data availability

The Authors declare that all relevant data generated and analyzed during this study, which include experimental, spectroscopic, crystallographic and computational data, are included in this article and its supplementary information. Should any raw data files be needed in another format, they are available from the corresponding author upon request. Crystallographic data for the structures reported in this article have been deposited at the Cambridge Crystallographic Data Center under deposition numbers CCDC 1) 2282181, 2) 2295973, 3) 2279914, 4) 2279928, 5) 2279917, 6) 2279931. Coordinates of the optimized structures are present as source data. Copies of the data can be obtained free of charge via <https://www.ccdc.cam.ac.uk/structures/>. Source data are provided in this paper.

## References

1. Wang, J. & Qiu, D. *Recent developments of diazo compounds in organic synthesis*. (World Scientific: London, 2020).
2. Wang, J., Che, C. M. & Doyle, M. P. (eds) *Transition metal-catalyzed carbene transformations*. (Wiley-VCH GmbH, Weinheim, Germany, 2022).
3. Ford, A. et al. Modern organic synthesis with  $\alpha$ -diazocarbonyl compounds. *Chem. Rev.* **115**, 9981–10080 (2015).
4. Jurgberg, I. D. & Davies, H. M. L. Blue light-promoted photolysis of aryl diazoacetates. *Chem. Sci.* **9**, 5112–5118 (2018).
5. Wang, H., Wang, S., George, V., Galder, L. & König, B. Photo-induced homoligation of carbonyl compounds for iterative syntheses. *Angew. Chem. Int. Ed.* **61**, e202211578 (2022).
6. Durka, J., Turkowska, J. & Gryko, D. Lightning diazo compounds? *ACS Sustain. Chem. Eng.* **9**, 8895–8918 (2021).
7. Yang, Z., Stivanin, M. L., Jurberg, I. D. & Koenigs, R. M. *Chem. Soc. Rev.* **49**, 6833–6847 (2020).
8. López, E., González-Pelayo, S. & López, L. A. Recent developments in coinage metal catalyzed transformations of stabilized vinyl diazo compounds: beyond carbenic pathways. *Chem. Rec.* **17**, 312–325 (2017).
9. Cheng, Q.-Q., Yu, Y., Yedoyan, J. & Doyle, M. P. Vinyl diazo reagents and metal catalysts: a versatile toolkit for heterocycle and carbocycle construction. *ChemCatChem* **10**, 488–496 (2018).
10. López, E., Bernardo, O. & López, L. A. Coinage metal-catalyzed carbonyl and heterocyclizations involving alkenyl carbene intermediates as C3 synthons. *Tetrahedron Lett.* **109**, 154156 (2022).
11. Marichev, K. O. & Doyle, M. P. Catalytic asymmetric cycloaddition reactions of enoldiazo compounds. *Org. Biomol. Chem.* **17**, 4183–4195 (2019).
12. Liu, L. & Zhang, J. Gold-catalyzed transformations of  $\alpha$ -diazocarbonyl compounds: Selectivity and diversity. *Chem. Soc. Rev.* **45**, 506–516 (2016).
13. Cheng, Q.-Q., Deng, Y., Lankelma, M. & Doyle, M. P. Cycloaddition reactions of enoldiazo compounds. *Chem. Soc. Rev.* **46**, 5425–5443 (2017).
14. Dong, K., Marichev, K. O. & Doyle, M. P. The role of donor-acceptor cyclopropenes in metal carbene reactions. conversion of *E*-substituted enoldiazoacetates to *Z*-substituted metallo-enolcarbenes. *Organometallics* **38**, 4043–4050 (2019).
15. Ciszewski, L. W., Rybicka-Jasinska, K. & Gryko, D. Recent developments in photochemical reactions of diazo compounds. *Org. Biomol. Chem.* **17**, 432–448 (2019).
16. Zhang, Y., Kubicvki, J. & Platz, M. S. Ultrafast UV-visible and infrared spectroscopic observation of a singlet vinyl carbene and the intramolecular cyclopropanation reaction. *J. Am. Chem. Soc.* **131**, 13602–13603 (2009).
17. Closs, G. L. & Closs, L. E. Alkenylcarbenes as precursors of cyclopropenes. *J. Am. Chem. Soc.* **83**, 2015–2016 (1961).
18. Padwa, A., Blacklock, T. J., Getman, D., Hatanaka, N. & Loza, R. On the problem of regioselectivity in the photochemical ring-opening

- reaction of 3-phenyl- and 3-vinyl-substituted cyclopropenes to indenes and 1,3-cyclopentadienes. *J. Org. Chem.* **43**, 1481–1492 (1978).
19. Zhu, Z.-B., Wei, Y. & Shi, M. Recent developments of cyclopropene chemistry. *Chem. Soc. Rev.* **40**, 5534–5563 (2011).
20. Molchanov, A. P., Efremova, M. M. & Kuznetsov, M. A. Cyclopropenes and methylenecyclopropanes in 1,3-dipolar cycloaddition reactions. *Russ. Chem. Bull.* **71**, 620–650 (2022).
21. Ravasco, J. M. J., Monteiro, C. M. & Trindade, A. F. Cyclopropenes: a new tool for the study of biological systems. *Org. Chem. Front.* **4**, 1167–1198 (2017).
22. Gahatory, D., Sen, R., Kuzmyn, A. R., Escorihuela, J. & Zuilhof, H. Strain-promoted cycloaddition of cyclopropenes with *o*-quinones: a rapid click reaction. *Angew. Chem. Int. Ed.* **57**, 10118–10122 (2018).
23. Yang, J., Liang, Y., Šečkutė, J., Houk, K. N. & Devaraj, N. K. Synthesis and reactivity comparisons of 1-methyl-3-substituted cyclopropene mini-tags for tetrazine bioorthogonal reactions. *Chem. Eur. J.* **20**, 3365–3375 (2014).
24. Oblak, E. Z., VanHeyst, M. D., Li, J., Weimer, A. J. & Wright, D. L. Cyclopropene cycloadditions with annulated furans: Total synthesis of (+)- and (–)-frondosin B and (+)-frondosin A. *J. Am. Chem. Soc.* **136**, 4309–4315 (2014).
25. Kumar, G. S. & Lin, Q. Light-triggered click chemistry. *Chem. Rev.* **121**, 6991–7031 (2021).
26. Borrmann, A. et al. *Bioconjugate Chem.* **26**, 257–261 (2015).
27. Bruins, J. J. et al. Inducible, site-specific protein labeling by tyrosine oxidation-strain-promoted (4 + 2) cycloaddition. *Bioconjugate Chem.* **28**, 1189–1193 (2017).
28. George, A. et al. Accelerated strain-promoted and oxidation-controlled cyclooctyne-quinone cycloaddition for cell labeling. *ChemistrySelect* **2**, 7117–7122 (2017).
29. Patterson, D. M., Nazarova, L. A., Xie, B., Kamber, D. N. & Prescher, J. A. Functionalized cyclopropenes as bioorthogonal chemical reporters. *J. Am. Chem. Soc.* **134**, 18638–18643 (2012).
30. Jonker, A. M. et al. A fast and activatable cross-linking strategy for hydrogel formation. *Adv. Mater.* **27**, 1235–1240 (2015).
31. Kaur, G., Singh, G. & Singh, J. Photochemical tuning of materials: a click chemistry perspective. *Mater. Today Chem.* **8**, 56–84 (2018).
32. Xu, X., Zavalij, P. Y. & Doyle, M. P. Catalytic asymmetric syntheses of quinolizidines by dirhodium-catalyzed dearomatization of isoquinolinium/pyridinium methylides—the role of catalyst and carbene source. *J. Am. Chem. Soc.* **135**, 12439–12447 (2013).
33. Deng, Y., Jing, C. & Doyle, M. P. Dinitrogen extrusion from enoldiazo compounds under thermal conditions: synthesis of donor-acceptor cyclopropenes. *Chem. Commun.* **51**, 12924–12927 (2015).
34. Zheng, H., Faghihi, I. & Doyle, M. P. Copper(I)-catalyzed highly enantioselective [3+3]-cycloaddition of  $\beta$ -aryl/Alkyl vinyl diazoacetates with nitrones. *Helv. Chim. Acta* **104**, e2100081 (2021).
35. Kardile, R. D. & Liu, R. S. Gold(I)-catalyzed reactions between 2-(1-alkynyl)-2-alken-1-ones and vinyl diazo ketones for divergent synthesis of nonsymmetric heteroaryl-substituted triarylmethanes: N-versus C-attack paths. *Org. Lett.* **22**, 8229–8233 (2020).
36. Padwa, A., Kulkarni, Y. S. & Zhang, Z. Reaction of carbonyl compounds with ethyl lithio-diazoacetate. studies dealing with the rhodium(II)-catalyzed behavior of the resulting adducts. *J. Org. Chem.* **55**, 4144–4153 (1990).
37. Liu, F., Liang, Y. & Houk, K. N. Bioorthogonal cycloadditions: computational analysis with the distortion/interaction model and predictions of reactivities. *Acc. Chem. Res.* **50**, 2297–2308 (2017).
38. Vicente, R. Recent progresses towards the strengthening of cyclopropene chemistry. *Synthesis* **48**, 2343–2360 (2016).
39. Marichev, K. O. et al. Rhodium(II)-catalysed generation of cycloprop-1-en-1-yl ketones and their rearrangement to 5-aryl-2-siloxyfurans. *Chem. Commun.* **54**, 9513–9516 (2018).
40. Deng, Q., Thomas, B. E., Houk, K. N. & Dowd, P. Transition structures of the ene reactions of cyclopropene. *J. Am. Chem. Soc.* **119**, 6902–6908 (1997).
41. Frisch, M. J. et al. Gaussian 16, Revision C.01, (Gaussian, Inc., Wallingford CT, 2019).
42. Marenich, A. V., Cramer, C. J. & Truhlar, D. G. Universal solvation model based on solute electron density and on a continuum model of the solvent defined by the bulk dielectric constant and atomic surface tensions. *J. Phys. Chem. B* **113**, 6378–6396 (2009).
43. Bao, M. et al. Expanded access to vinyl diazo compounds and their catalytic enantioselective application. *Chem Catal.* **3**, (in press 2023).
44. Wang, X., Xu, X., Zavalij, P. Y. & Doyle, M. P. Asymmetric formal [3 + 3]-cycloaddition reactions of nitrones with electrophilic vinylcarbene intermediates. *J. Am. Chem. Soc.* **133**, 16402–16405 (2011).
45. Wang, X., Abrahams, Q. M., Zavalij, P. Y. & Doyle, M. P. Highly regio- and stereoselective dirhodium vinylcarbene induced Nitrono cycloaddition with subsequent cascade carbenoid aromatic cycloaddition/N-O cleavage and rearrangement. *Angew. Chem. Int. Ed.* **51**, 5907–5910 (2012).
46. Qian, Y. et al. Rhodium(II)- and copper(II)-catalyzed reactions of enol diazoacetates with nitrones: metal carbene versus Lewis acid directed pathways. *Angew. Chem. Int. Ed.* **51**, 5900–5903 (2012).
47. Cheng, Q.-Q., Yedoyan, J., Arman, H. & Doyle, M. P. Copper-catalyzed divergent addition reactions of enoldiazoacetamides with nitrones. *J. Am. Chem. Soc.* **138**, 44–47 (2016).
48. Adly, F. G., Marichev, K. O., Jensen, J. A., Arman, H. & Doyle, M. P. Enoldiazoacetates for highly enantioselective [3 + 3]-cycloaddition with nitrones catalyzed by copper(I) with chiral BOX ligands. *Org. Lett.* **21**, 40–44 (2019).
49. Diev, V. V. et al. Nitrono cycloadditions to 1,2-diphenylcyclopropenes and subsequent transformations of the isoxazolidine cycloadducts. *J. Org. Chem.* **73**, 2396–2399 (2008).
50. Deepthi, A., Thomas, N. V. & Sruthi, S. L. An overview of the reactions involving azomethine imines over half a decade. *N. J. Chem.* **45**, 8847–8873 (2021).
51. Borah, B. & Chowhan, L. R. Recent updates on the stereoselective synthesis of structurally functionalized spiro-oxindoles mediated by isatin *N,N'*-cyclic azomethine imine 1, 3-dipoles. *Tetrahedron Lett.* **104**, 154014 (2022).
52. Cao, V. D., Kim, K., Kwak, J. & Joung, S. [3 + 2] Cycloaddition reaction of the endocyclic *N*-silyl enamine and *N,N'*-cyclic azomethine imine. *Org. Lett.* **24**, 1974–1978 (2022).
53. Qian, Y., Zavalij, P. Y., Hu, W. & Doyle, M. P. Bicyclic pyrazolidinone derivatives from diastereoselective catalytic [3+3]-cycloaddition reactions of enoldiazoacetates with azomethine imines. *Org. Lett.* **15**, 1564–1567 (2013).
54. Filatov, A. S. et al. Stereo- and regioselective 1,3-dipolar cycloaddition of the stable ninhydrin-derived azomethine ylide to cyclopropenes: Trapping of unstable cyclopropene dipolarophiles. *J. Org. Chem.* **84**, 7017–7036 (2019).
55. Rane, D. & Sibi, M. Recent advances in nitrile oxide cycloadditions. synthesis of isoxazolines. *Curr. Org. Synth.* **8**, 616–627 (2011).
56. Easton, C. J., Hughes, C. M. M., Savage, G. P. & Simpson, G. W. Cycloaddition reactions of nitrile oxides with alkenes. *Adv. Heterocycl. Chem.* **60**, 261–327 (1994).
57. Heaney, F. Nitrile oxide/alkyne cycloadditions—a credible platform for synthesis of bioinspired molecules by metal-free molecular clicking. *Eur. J. Org. Chem.* **16**, 3043–3048 (2012).
58. Bode, J. W. & Carreira, E. M. Stereoselective syntheses of ephothilones A and B via nitrile oxide cycloadditions and related studies. *J. Org. Chem.* **66**, 6410–6424 (2001).
59. Xu, X., Shabashov, D., Zavalij, D. & Doyle, M. P. Unexpected catalytic reactions of silyl-protected enol diazoacetates with nitrile oxides

- that form 5-arylamino-furan-2(3H)-one-4-carboxylates. *Org. Lett.* **14**, 800–803 (2012).
60. Cui, Y., Zhang, G. & Ding, C. Recent advances in Lossen rearrangement. *Chin. J. Org. Chem.* **42**, 2015–2017 (2022).
61. Thomas, M. et al. The lossen rearrangement from free hydroxamic acids. *Org. Biomol. Chem.* **17**, 5420–5427 (2019).

## Acknowledgements

Financial support for this work was supported by the U.S. National Science Foundation (CHE 2054845) for M.P.D. and M. B. and by NCN Poland OPUS UMO-2019/35/B/ST4/03435 for D.G. and K.Ł. Calculations have been carried out using resources provided by Wrocław Center for Networking and Supercomputing (<http://wcss.pl>), grant No. 518. for W.CH.

## Author contributions

M. B. and K. Ł. performed photo-catalytic studies of the vinyl diazo compounds and characterization of products. M. B. and M. Baird prepared the vinyl diazo compounds. M. B., D. G., and M. P. D. conceived and designed the experiments, W. CH. prepared DFT calculations, and M. B., K. Ł., D. G., and M. P. D. prepared the manuscript. All authors contributed to discussions and commented on the manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41467-024-48274-5>.

**Correspondence** and requests for materials should be addressed to Wojciech Chatadaj, Dorota Gryko or Michael P. Doyle.

**Peer review information** *Nature Communications* thanks shengbiao tang and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. A peer review file is available.

**Reprints and permissions information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024