

CORRECTION

Open Access



# Correction: The microbiota-gut-brain axis participates in chronic cerebral hypoperfusion by disrupting the metabolism of short-chain fatty acids

Weiping Xiao<sup>1,2†</sup>, Jiabin Su<sup>1,2†</sup>, Xinjie Gao<sup>1,2</sup>, Heng Yang<sup>1,2</sup>, Ruiyuan Weng<sup>1,2</sup>, Wei Ni<sup>1,3\*</sup> and Yuxiang Gu<sup>1,2\*</sup>

**Correction: *Microbiome* 10, 62 (2022)**

**<https://doi.org/10.1186/s40168-022-01255-6>**

Following publication of the original article [1], the author reported that in Figure 9D, the first image is a duplicate of the second image. The correct image is below.

---

<sup>†</sup>Weiping Xiao and Jiabin Su contributed equally to this work.

The original article can be found online at <https://doi.org/10.1186/s40168-022-01255-6>.

\*Correspondence:

Wei Ni

[hsniwei@fudan.edu.cn](mailto:hsniwei@fudan.edu.cn)

Yuxiang Gu


[guyuxiang1972@126.com](mailto:guyuxiang1972@126.com)

<sup>1</sup> Department of Neurosurgery, Huashan Hospital, Fudan University, Shanghai 200040, China

<sup>2</sup> Institute of Neurosurgery, Fudan University, Shanghai 200052, China

<sup>3</sup> Shanghai Key Laboratory of Brain Function and Restoration and Neural Regeneration, Shanghai 200052, China





#### Reference

1. Xiao W, Su J, Gao X, et al. The microbiota-gut-brain axis participates in chronic cerebral hypoperfusion by disrupting the metabolism of short-chain fatty acids. *Microbiome*. 2022;10:62. <https://doi.org/10.1186/s40168-022-01255-6>.