

Lysophosphatidic acid stimulates the proliferation and motility of malignant pleural mesothelioma cells through lysophosphatidic acid receptors, LPA₁ and LPA₂

Tadaaki Yamada,¹ Seiji Yano,^{1,2} Hirokazu Ogino,¹ Kenji Ikuta,¹ Soji Kakiuchi,¹ Masaki Hanibuchi,¹ Takanori Kanematsu,¹ Tetsuo Taniguchi,³ Yoshitaka Sekido³ and Saburo Sone^{1,4}

¹Department of Internal Medicine and Molecular Therapeutics, Institute of Health Biosciences, University of Tokushima Graduate School, 3-18-15 Kuramoto-cho, Tokushima 770-8503; ²Division of Medical Oncology, Cancer Research Institute, Kanazawa University, 13-1 Takaramachi, Kanazawa, Ishikawa 920-0934; ³Division of Molecular Oncology, Aichi Cancer Center Research Institute, Aichi, 1-1 Kanokoden, Chikusa-ku, Nagoya 464-0021, Japan

(Received January 18, 2008/Revised April 12, 2008/Accepted April 13, 2008/Online publication July 29, 2008)

Lysophosphatidic acid (LPA) is one of the simplest natural phospholipids. This phospholipid is recognized as an extracellular potent lipid mediator with diverse effects on various cells. Although LPA is shown to stimulate proliferation and motility via LPA receptors, LPA₁ and LPA₂, in several cancer cell lines, the role of LPA and LPA receptors for malignant pleural mesothelioma (MPM) has been unknown. MPM is an aggressive malignancy with a poor prognosis and the incidence is increasing and is expected to increase further for another 10–20 years worldwide. Therefore, the development of novel effective therapies is needed urgently. In this study, we investigated the effect of LPA on the proliferation and motility of MPM cells. We found that all 12 cell lines and four clinical samples of MPM expressed LPA₁, and some of them expressed LPA₂, LPA₃, LPA₄ and LPA₅. LPA stimulated the proliferation and motility of MPM cells in a dose-dependent manner. Moreover, LPA-induced proliferation was inhibited by Ki16425, an inhibitor of LPA₁, and small interfering RNA against LPA₁, but not LPA₂. Interestingly, LPA-induced motility was inhibited by small interfering RNA against LPA₂, but not LPA₁, unlike a number of previous reports. These results indicate that LPA is a critical factor on proliferation through LPA₁, and on motility through LPA₂ in MPM cells. Therefore, LPA and LPA receptors, LPA₂ as well as LPA₁, represent potential therapeutic targets for patients with MPM. (*Cancer Sci* 2008; 99: 1603–1610)

Malignant pleural mesothelioma (MPM) is an aggressive tumor arising from the mesothelial cells in the pleural cavity. It is frequently diagnosed at a locally advanced stage and is refractory to conventional treatment regimens, such as chemotherapy and radiation therapy, and so its prognosis is extremely poor. Several etiological factors, including asbestos,^(1,2) iron,⁽³⁾ the simian virus 40 (SV40), radiation and thorotrast,⁽⁴⁾ have been reported to be involved in the development of MPM. Of these factors, exposure to asbestos is most closely related to the development of MPM. Prior to the recognition of its adverse health effects, large amounts of asbestos were used worldwide. As the latent period between the first exposure to asbestos and the occurrence of MPM tends to be 30–40 years, it is expected that there will be rapid increases in the number of MPM patients in Europe and Australia in 2015–2020⁽⁵⁾ and in Japan in 2010–2040.⁽²⁾ Thus, the incidence of MPM is expected to increase further for another 10–20 years worldwide, although the disease has already reached its peak incidence in the USA.⁽⁵⁾ The major reasons for the poor prognosis of MPM are: (i) the highly proliferating and invasive characteristics of the disease; and (ii) its resistance to conventional chemotherapy and radiotherapy. Therefore, novel effective therapeutic strategies based on the

molecular mechanisms of proliferation and motility/invasion of MPM are required to improve the prognosis of this disease. While several growth factors, including platelet-derived growth factors A and B, epidermal growth factor, transforming growth factor- β and vascular endothelial growth factor, were reported to be involved in the progression of MPM,⁽⁵⁾ the molecular mechanisms of the proliferation and motility/invasion of MPM are not fully understood.

Lysophosphatidic acid (LPA) is one of the simplest natural phospholipids. It is present in serum at concentrations of 2–20 μM ⁽⁶⁾ and is derived from various enzymatic mechanisms, such as activated platelets, fibroblasts, adipocytes and ovarian tumors.^(6–8) Recently, this phospholipid has attracted a great deal of attention as a potent lipid mediator responsible for multiple cellular processes⁽⁹⁾ such as cell proliferation, platelet aggregation, smooth muscle contraction and cytoskeletal reorganization. In fact, extracellular LPA has been shown to be associated with various diseases, including atherosclerosis⁽¹⁰⁾ and cancer.^(11–13) LPA exerts its multiple biological functions via three types of G protein-coupled receptor (GPCR), LPA₁, LPA₂ and LPA₃. These receptors share 50–54% amino acid homology and transmit their signals via three families of heterotrimeric G proteins, G_i, G_q and G_{12/13}.⁽¹⁴⁾ Furthermore, more recent studies have identified orphan glutamyl transpeptidase binding protein-coupled receptors (GPCR), LPA₄⁽¹⁵⁾ and LPA₅.⁽¹⁶⁾

The mechanisms of action of LPA receptors differ among subtypes, and they are expressed at different levels in both normal and malignant tissues. In normal tissues, LPA₁ is highly expressed in the brain and heart, LPA₂ is highly expressed in the testis and leukocytes, LPA₃ is highly expressed in the kidney and prostate^(17,18) and LPA₄ is expressed highly in the ovary.⁽¹⁵⁾ In malignant tumors, LPA₁ is dominant in brain tumors,⁽¹⁹⁾ LPA₂ is predominantly expressed in colon, stomach, thyroid and breast cancers^(20–22) and LPA₃ is expressed at relatively high levels in ovarian and prostate cancers.^(19,23) Another study showed that the expression pattern of LPA receptors is associated with cancer differentiation in gastric cancer cells.⁽²⁴⁾ Moreover, LPA₁ has recently been reported to control tumor cell proliferation in prostate⁽²⁵⁾ and breast cancer cells⁽²⁶⁾ and LPA₂ was shown to mediate the growth of ovarian,⁽²⁷⁾ thyroid⁽²¹⁾ and colon cancer cells.⁽²⁸⁾ Interestingly, LPA₁ also stimulates the motility of various types of cancer, such as pancreas, colon, glioblastoma and gastric

⁴To whom correspondence should be addressed.
E-mail: ssone@clin.med.tokushima-u.ac.jp

cancer.^(19,24,28,29) Thus, LPA receptors play crucial roles at least in cancer proliferation and motility/invasion. However, the roles of LPA receptors and their downstream signaling pathways in MPM are unknown.

The present study was performed to explore the expression of LPA receptors in human MPM cell lines and further elucidate the roles of five LPA receptors in the malignant properties of MPM cells in terms of proliferation and motility.

Materials and Methods

Cell lines and culture conditions. In this study, we used 12 human MPM cell lines: EHMES-1, EHMES-10, MSTO-211H, Y-MESO-8A, NCI-H28, NCI-H290, NCI-H513, NCI-H2052, NCI-H2373, NCI-H2452, ACC-MESO-1 and ACC-MESO-4. EHMES-1 and EHMES-10^(30,31) were kindly provided by Dr Hironobu Hamada (Ehime University, Matsuyama, Japan). NCI-H290 and NCI-H513 were kindly provided by Dr Adi F. Gazdar (University of Texas South-western Medical Center, Dallas, TX, USA). Y-MESO-8A, ACC-MESO-1 and ACC-MESO-4⁽³²⁾ were established in Aichi Cancer Center Research Institute. MSTO-211H, NCI-H28, NCI-H2052, NCI-H2373 and NCI-H2452 were purchased from the American Type Culture Collection (Manassas, VA, USA).

All human tumor cell lines were maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum (FBS), penicillin (100 U/mL) and streptomycin (50 µg/mL); this was designated as RPMI-1640 in this study. Cells were cultured in a humidified CO₂ incubator at 37°C.

Clinical specimens. All human MPM samples were obtained from the Department of Internal Medicine and Molecular Therapeutics, University of Tokushima Graduate School. Written informed consent was obtained from all subjects prior to tissue sampling.

Reagents. 1-Oleoyl-LPA (18:1) and Ki16425 were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Expression of LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅. LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅ mRNA expression were determined by reverse transcription polymerase chain reaction (RT-PCR). Total RNA samples were isolated using an RNeasy Mini kit and RNase-free DNase kits (Qiagen, Valencia, CA, USA) in accordance with the manufacturer's recommendations. Total RNA were reverse transcribed using an Omniscript RT kit (Qiagen). The primers for LPA₁, LPA₂, LPA₃, LPA₄, LPA₅ and β-actin were as follows: LPA₁, 5'-TGGCTGCCATCTCTACTTCC-3' and 5'-AACCAATCCAGGAGTCCAGC-3'; LPA₂, 5'-CATCATGCTTCCCGAC AACG-3' and 5'-GGGCTTACCAAGGATACGCAG-3'; LPA₃, 5'-AGTGTCACTATGACAAGC-3' and 5'-GAGATGTTGCAGAGGC-3'; LPA₄, 5'-TGAAGGCTTCCAAACGTGTCTG-3' and 5'-GTTCAAGTTGCAAGGCACAAGGT-3'; LPA₅, 5'-CTGGATCTAAACCGCCACAG-3' and 5'-GCCTGGAAGGGGATGT-3'; and β-actin, 5'-AAGAGAGGCATCCTCACCT-3' and 5'-TACATGGCTGGGGTGTGAA-3'.

Polymerase chain reaction was performed with Ex Taq Hot Start Version (Takara Bio, Shiga, Japan). The bands were visualized by ethidium bromide staining. Data shown are representative of three independent experiments.

Cell proliferation assay. Cell proliferation was measured by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide dye reduction method.⁽³³⁾ Briefly, tumor cells seeded in 96-well plates (5 × 10³/well) were incubated in RPMI-1640 for 24 h. The cells were starved for 24 h by replacing the medium with serum-free RPMI-1640 medium containing 0.1% fatty acid-free bovine serum albumin (BSA), with or without LPA and/or Ki16425, an inhibitor of LPA₁ (and LPA₃ at higher doses).⁽³⁴⁾ The cells were cultured for 48 h. Then, an aliquot of 50 µL stock 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide solution (2 mg/mL; Sigma, St. Louis, MO, USA) was added to each plate and the cells were incubated for 2 h at

37°C. The media containing 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide solution were removed and the dark-blue crystals were dissolved by adding 100 µL dimethylsulfoxide. Absorbance was measured with an MTP-120 microplate reader (Corona Electric, Ibaraki, Japan) at test and reference wavelengths of 550 and 630 nm, respectively. Data shown are representative of three independent experiments.

Cell migration assay. Cell migration was quantified in Falcon Cell Culture Inserts (8-µm pore size; BD Biosciences, San Jose, CA, USA). Cells suspended in serum-free RPMI-1640 containing 0.1% fatty acid-free BSA, with or without LPA and/or Ki16425, were added to the upper chamber (1 × 10⁵ cells in 100 µL/well). Serum-free RPMI-1640 containing 0.1% fatty acid-free BSA, containing human fibronectin (5 µg/mL; BD Biosciences) as a chemoattractant was placed in the lower chamber. In the case of treatment with Ki16425, the cells were pre-incubated with 10 µM Ki16425 for 30 min. The cells were allowed to migrate for 3 h at 37°C. Non-migratory cells were removed from the top filter surface with a cotton swab. Migrated cells, attached to the bottom surface, were fixed with methanol, stained with Giemsa, mounted and then counted. Data shown are representative of three independent experiments.

RNA interference. Three different duplexed Stealth RNAi (Invitrogen, Carlsbad, CA, USA) against LPA₁ and LPA₂, and Stealth RNAi Negative Control Kit (Invitrogen) were used for RNA interference assay. Briefly, aliquots of 1 × 10⁵ 211H, H28 or H2052 cells in 2 mL antibiotic-free medium were plated on 6-well plates and incubated at 37°C for 24 h, and the cells were then transfected with small interfering RNA (siRNA) (250 pmol) or scramble RNA (siSCR) using Lipofectamine 2000 (5 µL; Invitrogen, Carlsbad, CA, USA) in accordance with the manufacturer's instructions. After 24-h incubation, the cells were washed twice with phosphate-buffered saline (PBS), and incubated in antibiotic-containing medium. These cells were then used for proliferation or migration assay as described above. LPA₁ and LPA₂ knockdown were confirmed by RT-PCR analysis. The sense sequences of siRNA were as follows: siLPA1#1, 5'-AACCAAUCCAGGAGUCCAGCAGAUG-3' and 5'-CAUCUGCUGGACUCCUGGAUUGGUU-3'; siLPA1#2, 5'-AUUACAGGGAUGGAAGUAGAGAUGG-3' and 5'-CCAUCUCUACUCCAUCCUGUAAU-3'; siLPA1#3, 5'-AUAGAUUGCCACCAUGACCAAUAGG-3' and 5'-CCUAUUGGUAUGGUGGCAUCUAU-3'; siLPA2#1, 5'-UACACAGCAGCAUUGACCAGUAGU-3' and 5'-ACUCACUGGUAUUGCUGUGUA-3'; siLPA2#2, 5'-AUGUAUAGUGGACAGACUCGCGGGU-3' and 5'-ACCCGCGAGUCUGUCCACUAUACAU-3'; and siLPA2#3, 5'-UGUGGAACAUGAGGAAGAGGUAGGC-3' and 5'-GCCUACUCUCCUCAUGUCCACA-3'.

Western blotting. Tumor cells were washed twice with PBS, harvested in cell lysis buffer (20 mM Tris, pH 7.4, 150 mM NaCl, 1 mM ethylenediaminetetraacetic acid, 1 mM ethyleneglycoltetraacetic acid, 1% Triton X-100, 2.5 mM sodium pyrophosphate, 1 mM β-glycerolphosphate, 1 mM Na₃VO₄, 1 µg/mL leupeptin and 1 mM phenylmethylsulfonyl fluoride), and flash-frozen on dry ice. After allowing the cells to thaw, the cell lysates were collected with a rubber scraper, sonicated and centrifuged at 14 000g (4°C for 20 min). The total protein concentration was measured using a Pierce BCA Protein Assay Kit (Pierce, Rockland, IL, USA). Cell lysates were subjected to sodium dodecylsulfate polyacrylamide gel electrophoresis, and the proteins were then transferred onto polyvinylidene fluoride (PVDF) membranes (Atto, Tokyo, Japan). The membranes were blocked with Blocking One (Nacalai Tesque, Kyoto, Japan) for 1 h at room temperature, and then the blots were incubated at 4°C overnight with anti-phospho-p44/42 mitogen-activated protein kinase (MAPK) (Thr202/Tyr204), anti-p44/42 MAPK (1:1000 dilution; Cell Signaling Technology, Beverly, MA, USA), or anti-β-actin antibody (1:5000 dilution; Sigma), followed by incubation for




Fig. 1. Expression of lysophosphatidic acid (LPA) receptors, LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅ in human malignant pleural mesothelioma cell lines, and clinical specimens. Reverse transcription polymerase chain reaction (RT-PCR) was performed to determine expression of the LPA receptors, LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅ in human MPM cell lines (a) and clinical specimens (b). RT-PCR products were electrophoresed on a 2% agarose gel and stained with ethidium bromide. Data shown are representative of three independent experiments with similar results.

2 h at room temperature with secondary antibodies (horseradish peroxidase-conjugated species-specific Ab). Immunoreactive bands were visualized using enhanced chemiluminescent substrate (Pierce).

RhoA activity assay. We determined RhoA activation as the amount of RhoA-glutamyl transpeptidase (GTP) binding using a G-LISA RhoA Activation Assay Biochem Kit (Cytoskeleton, Denver, CO, USA), in accordance with the manufacturer's recommendations. Briefly, tumor cells (2×10^5 /well) seeded in 6-well plates were incubated in RPMI-1640 containing 10% FBS for 24 h. The cells were starved for 24 h by replacing the medium with serum-free RPMI-1640 medium containing 0.1% BSA. The dishes were then treated with LPA (10 μ M) for 0, 1, 3, 6, 12 and 30 min. The tumor cells were washed twice with PBS and harvested in cell lysate with a cell scraper, sonicated and centrifuged at $10\,000 \times g$ (4°C for 2 min). The total protein concentration was measured using Precision Red Advanced Protein Assay Reagent. Equal amounts of cell lysate protein were incubated in Rho-GTP affinity plates for 30 min at 4°C. To the plates was added anti-RhoA primary antibody, followed by secondary horseradish peroxidase (HRP)-labeled antibody and HRP detection reagent, and incubated at room temperature for the time recommended by the manufacturer. Absorbance was measured with an MTP-120 microplate reader at a wavelength of 490 nm. Data shown are representative of three independent experiments with similar results.

Statistical analysis. All data, expressed as means \pm standard error (SE), were analyzed by one-way ANOVA. The statistical significance of differences was assessed by Fisher's protected least-significant difference test. $P < 0.05$ was considered statistically significant. Statistical analyses were performed using StatView ver. 5.0 (SAS Institute, Cary, NC, USA).

Results

Human MPM cell lines expressed LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅ at various levels. In the first set of experiments, we examined the mRNA expression of five LPA receptors (LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅) in 12 human MPM cell lines by RT-PCR (Fig. 1a). All of the MPM cell lines examined expressed LPA₁ mRNA at various levels. Of these cell lines, 211H, H28, H290, H2052 and H2373 cells showed higher levels of expression as compared with the others. On the other hand, some of the MPM

cell lines expressed LPA₂. Y-MESO-8 A, H290, H2052 and ACC-MESO-4 cells showed high levels of LPA₂ expression as compared with the other cell lines. Similarly, some of the MPM cell lines expressed LPA₃, LPA₄ and LPA₅. In parallel experiments, we examined LPA₁, LPA₂, LPA₃, LPA₄ and LPA₅ mRNA expression in tumor specimens obtained from four patients with MPM (Fig. 1b). Consistent with the results in the cell lines examined, all samples expressed LPA₁ and some expressed LPA₂, LPA₃, LPA₄ and LPA₅.

LPA stimulated the proliferation of MPM cells via LPA₁. We determined the effects of LPA on proliferation of MPM cells. Exogenous LPA at physiological concentrations (up to 20 μ M) significantly stimulated the proliferation of five of the 11 MPM cell lines examined (data not shown). Of these cell lines, 211H and H28 cells responded well to LPA in a dose-dependent manner (Fig. 2a). Therefore, these two cell lines were used to investigate whether LPA receptors are involved in LPA-induced proliferation, when the effects of Ki16425, an inhibitor of LPA₁ (and LPA₃ at higher doses), on proliferation were examined. Ki16425 inhibited LPA-induced proliferation of both 211H and H28 cells in a dose-dependent manner (Fig. 2b). In a parallel experiment, we examined the effects of siRNA against LPA₁ and LPA₂. Knockdown of LPA₁ resulted in significant inhibition of the LPA-induced proliferation of both 211H and H28 cells. These results were confirmed using serum-starved media with LPA, as well as media containing 10% FBS (data not shown). However, knockdown of LPA₂ had little impact on LPA-induced proliferation of 211H or H28 cells (Fig. 2c). These results indicated that LPA stimulated the proliferation of these two MPM cell lines via LPA₁ receptors.

LPA activates p44/42 MAPK through LPA₁. To determine the mechanism by which LPA₁ mediated proliferation of MPM cells, we next examined the phosphorylation of MAPK, which is the downstream signal of GPCR. MPM cells were stimulated with LPA (10 μ M) for 0, 2 or 6 h, and the phosphorylation of MAPK was evaluated by Western blotting. While LPA did not affect MAPK protein expression, it induced the phosphorylation of MAPK (phospho-p44/42 MAPK) in both 211H and H28 cells. The level of phosphorylation became highest after 2 h and the effect was still substantial after 6 h (Fig. 3a). Therefore, we evaluated the phosphorylation of MAPK after 2-h incubation with LPA. In a parallel experiment, siRNA knockdown of LPA₁, but not of LPA₂, markedly reduced the phosphorylation of




Fig. 2. Effects of lysophosphatidic acid (LPA) on cell proliferation of human malignant pleural mesothelioma cells, and inhibition by antagonist and knockdown of LPA₁. Tumor cells (5×10^3 /well) plated in triplicate in 96-well plates were incubated overnight in RPMI-1640 containing 10% fetal bovine serum. The cells were starved for 24 h by replacing the media with serum-free RPMI-1640 containing 0.1% fatty acid-free bovine serum albumin (BSA). Different doses of LPA and Ki16425, an inhibitor of LPA₁, were then added, and the cells were cultured for 48 h. Then, an aliquot of 50 μ L of stock 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide solution (2 mg/mL) was added to each plate and the cells were incubated for 2 h at 37°C. LPA stimulated proliferation of 211H and H28 cells in a dose-dependent manner (a). LPA-induced proliferation of 211H and H28 cells was inhibited by Ki16425 in a dose-dependent manner (b). After transfection with small interfering RNA (siRNA) or scramble RNA (siSCR) using Lipofectamine 2000 in accordance with the manufacturer's instructions, the cells were starved for 24 h by replacing the media with serum-free RPMI-1640 containing 0.1% fatty acid-free BSA. LPA (10 μ M) was then added and cells were cultured for 48 h. The cells were then subjected to 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay. Knockdown of LPA₁ (siLPA1#1, siLPA1#2 and siLPA1#3) significantly inhibited LPA-induced proliferation of 211H and H28 cells, while knockdown of LPA₂ (siLPA2#1, siLPA2#2 and siLPA2#3) had little impact on LPA-induced proliferation of 211H or H28 cells (c). Data shown are representative of three independent experiments with similar results. *Significantly different from control ($P < 0.001$). **Significantly different from siSCR ($P < 0.005$).

p44/42 MAPK in these two MPM cell lines (Fig. 3b), indicating that LPA activates the p44/42 MAPK pathway via LPA₁.

LPA stimulates cell motility via LPA₂. We next examined the effect of LPA on the motility of MPM cells using a cell migration assay. LPA stimulated migration of six of 10 MPM cell lines (data not shown). Of these MPM cell lines, LPA induced a significant increase in migration of H2052, H2373 and ACC-MESO-4 cells in a dose-dependent manner (Fig. 4a). However, it did not affect the proliferation of these MPM cells (data not shown). These observations exclude the possibility that the effect of cell motility was attributable to differences in proliferation. In turn, these cell lines were used to examine whether LPA receptors were involved in LPA-induced motility, when the effect of Ki16425, an inhibitor of LPA₁ (and LPA₃ at higher doses), on the motility was examined. Ki16425 had no effect on LPA-induced motility in these MPM cells (Fig. 4b). To clarify the involvement of LPA₁ and LPA₂, we further knocked down LPA₁ and LPA₂ expression, respectively, with siRNA using H2052 cells, because LPA induced the motility of H2052 cells most efficiently. Knockdown of LPA₂ markedly inhibited LPA-induced motility of H2052 cells, while knockdown of LPA₁

did not (Fig. 4c). These results strongly suggest that LPA induces motility via LPA₂ in H2052 cells.

LPA activates RhoA through LPA₂. We next investigated the effects of LPA on the activation of RhoA, which is a downstream signal of GPCR and one of the key molecules for cell motility. H2052 cells were stimulated with LPA (10 μ M) for 0, 1, 3, 6, 12 or 30 min, and activation of RhoA was evaluated by G-LISA RhoA Activation Assay (Biochem Kit). LPA induced activation of RhoA, the degree of which became strongest after 1 min, and the effect decreased within 30 min (Fig. 5a). Therefore, we evaluated RhoA after 1-min incubation with LPA. Knockdown of LPA₂ with siRNA, but not of LPA₁, significantly reduced RhoA activation in comparison with the control level (Fig. 5b). These results suggest that LPA stimulates cell motility through an LPA₂-mediated RhoA pathway.

Discussion

In the present study, we demonstrated that all of the 12 cell lines and four clinical samples of MPM examined expressed LPA₁, while other types of LPA receptors (LPA₂, LPA₃, LPA₄ and




Fig. 3. Effects of lysophosphatidic acid (LPA) on phospho-p44/42 mitogen-activated protein kinase (MAPK), and inhibition by knockdown of LPA₁. Tumor cells were incubated with LPA (10 μM) for 0, 2 and 6 h. Then, the cell lysates were harvested and the protein level and phosphorylation of p44/42 MAPK were determined by immunoblotting analysis. LPA-induced phosphorylation in 211H and H28 cells occurred after 2 h, and the effect was still substantial after 6 h (a). Moreover, knockdown of LPA₁ with small interfering RNA markedly reduced the activation of p44/42 MAPK. However, knockdown of LPA₂ had little impact (b). Data shown are representative of three independent experiments with similar results. siSCR, scramble RNA.

LPA₂) were expressed only in some of the cell lines and clinical samples, and LPA stimulated the proliferation and motility of several MPM cell lines in a dose-dependent manner. There was no apparent correlation between the response to LPA and the expression level of LPA receptors, although MPM cells with a high level of LPA₁ tend to stimulate proliferation by LPA (data not shown). Neither were the histological subtypes, such as epithelial, biphasic and sarcomatoid, particularly correlated with the response to LPA (data not shown). We also examined the role of LPA and its receptors using a non-cancerous mesothelial cell line, Met-5A. Met-5A cells expressed LPA₁ and LPA₂, and LPA induced the proliferation, but not migration (data not shown). These observations suggest that MPM is biologically heterogeneous and uses various factors and signaling pathways for cell proliferation and motility. Our results also clearly showed that particular populations of MPM cells essentially utilized the LPA/LPA receptor pathway for both cell proliferation and motility. As the biomarker for MPM, Lindholm *et al.* reported that gene copy number losses are a major mechanism for carcinogenesis and copy number changes are associated with a recurrent pattern.⁽³⁵⁾ Pass *et al.* reported to predict survival and progression of MPM patients, using a 27-gene expression profile.⁽³⁶⁾ However, to our knowledge, this is the first report of the involvement of LPA/LPA receptors in the malignant phenotype of MPM.

Lysophosphatidic acid is known to act on specific GPCR to elicit a wide range of cellular responses, ranging from cell proliferation and motility of various types of solid tumor.⁽³⁷⁾ MPM tends to advance locally with pleural effusion, rather than distant metastasis, and the extent of the disease is directly associated with the prognosis of MPM patients.⁽⁵⁾ It is important to

understand the molecular mechanism of MPM pathogenesis in terms of proliferation and motility, because molecules responsible for MPM progression seem to be ideal targets for therapy of MPM.

Recently, the differential roles of five GPCR as LPA receptors have attracted a great deal of attention. Of these receptors, LPA₁ and LPA₂ are thought to play important roles in the signaling pathways of cancer cells. Several studies indicated that LPA₁ controls cell proliferation in prostate cancer⁽²⁵⁾ and breast cancer cells.⁽²⁶⁾ Similarly, LPA₂ was found to mediate LPA-induced growth of ovarian,⁽²⁷⁾ thyroid⁽²¹⁾ and colon cancer cells.⁽²⁹⁾ Thus, the role of LPA receptors in cell proliferation triggered by LPA appears to depend on the type of cancer cell line examined. In this study, we demonstrated that LPA₁, but not LPA₂, is a key receptor in LPA-stimulated proliferation of MPM cells using silencing of LPA receptors.

G protein-coupled receptors, including LPA receptors, have multiple intracellular signaling mechanisms by small GTPase proteins, Ras, Rho, Rac and their downstream cascades.^(37,38) The activation of Ras-MAPK responsible for cell proliferation can lead to the nuclear translocation of MAPK followed by gene transcription and cell cycle progression, and this signaling plays a pivotal role in cancer cell proliferation and differentiation. In the present study, we showed that LPA activated the p44/42 MAPK pathway and siRNA for LPA₁, but not LPA₂, caused marked inhibition of cell proliferation and phosphorylation of p44/42 MAPK. These results strongly suggest that MAPK is involved in the LPA₁-mediated proliferation of MPM cells induced by LPA.

Malignant pleural mesothelioma cells have high motility, facilitate free movement in the pleural cavity during respiration by extending lubricating glycoproteins, and show disordered




Fig. 4. Effects of lysophosphatidic acid (LPA) on migration of human malignant pleural mesothelioma (MPM) cells, and inhibition by knockdown of LPA₂. Cell migration was measured in Falcon Cell Culture Inserts (8- μm pore size). Cells suspended in serum-free RPMI-1640 containing 0.1% fatty acid-free bovine serum albumin (BSA) with or without LPA and/or Ki16425 were added to the upper chamber (1×10^5 cells at 100 μL /well). Serum-free RPMI-1640 containing 0.1% fatty acid-free BSA and human fibronectin (5 $\mu\text{g}/\text{mL}$) as a chemoattractant was added to the lower chamber. When the cells treated with Ki16425, the cells were pre-incubated with 10 μM Ki16425 for 30 min. The cells were allowed to migrate for 3 h at 37°C. LPA stimulated migration of MPM cells, H2052, H2373 and ACC-MESO-4 in a dose-dependent manner (a). Ki16425 had no effect on LPA-induced motility in these MPM cells (b). After transfection with small interfering RNA (siRNA) or scramble RNA (siSCR), the cells were starved for 24 h by replacing the media with serum-free RPMI-1640 containing 0.1% fatty acid-free BSA. Knockdown of LPA₂ by siRNA significantly inhibited the LPA-induced motility of H2052 cells, while knockdown of LPA₁ did not (c). Data shown are representative of three independent experiments. *Significantly different from control ($P < 0.005$). **Significantly different from siSCR ($P < 0.005$).




Fig. 5. Effects of lysophosphatidic acid (LPA) on RhoA activity, and inhibition by knockdown of LPA₂. To assess RhoA activation, we determined RhoA activation as the amount of RhoA-glutamyl transpeptidase (GTP) binding using a G-LISA RhoA Activation Assay Biochem Kit in accordance with the manufacturer's recommendations. H2052 cells were treated with LPA (10 μM) for 0, 1, 3, 6, 12 and 30 min, and the cell lysates were then prepared for the assay. LPA stimulation of RhoA activity in H2052 cell was highest after 1 min and the effect decreased within 30 min (a). Knockdown of LPA₂ with small interfering RNA significantly reduced the activation of RhoA, while knockdown of LPA₁ had little effect (b). Data shown are representative of three independent experiments. *Significantly different from control ($P < 0.05$). **Significantly different from scramble RNA (siSCR; $P < 0.005$).

invasion into the thoracic cavity.⁽⁵⁾ LPA is already well known to stimulate cell motility in many cell types, such as fibroblasts, gliomas and T-cell lymphomas.⁽³⁹⁾ LPA-stimulated motility of cancer cells was recently reported to require signaling via LPA₁ in many types of solid tumor, including pancreas, colon, glioblastoma and gastric cancer.^(19,24,28,29) However, this was not the case in our studies that indicated that the motility of MPM cells was augmented by LPA through LPA₂, but not LPA₁. This was confirmed in the present study using two different methods. First, treatment with Ki16425, an inhibitor of LPA₁, did not inhibit cell migration. Second, knockdown of LPA₂, but not LPA₁, effectively suppressed LPA-induced motility of MPM (H2052) cells. These results demonstrated that LPA-induced motility of MPM cells is mediated partly, if not completely, through its interaction with LPA₂.

RhoA is a small GTPase protein that mediates actin microfilament organization necessary for adhesion, motility and changes in cell shape.⁽⁴⁰⁾ Recent studies have implicated the pathway of RhoA activation, which acts downstream of LPA receptors, in tumor invasion.^(41,42) We demonstrated that LPA activated RhoA and that the effect was significantly, but not completely, diminished by knockdown of LPA₂, suggesting that LPA stimulated migration of H2052 cells was mediated by LPA₂ through the RhoA activation pathway. Further studies are warranted to determine the involvement of other signaling pathways in LPA₂-mediated cell-motility of MPM cells.

To explore antitumor activity by inhibiting LPA receptors, we used an inhibitor of LPA₁, Ki16425, because it is only one commercially available LPA inhibitor at present. The severe combined immunodeficiency (SCID) mice bearing s.c. inoculated MSTO-211H cells were treated daily from days 19–30 with s.c.

injection of Ki16425 at 20 mg/kg/day. The tumor growth was measured every 3 days. Under these experimental conditions, treatment with Ki16425 did not significantly inhibit the tumor growth of MSTO-211 cells (data not shown). However, we can not make conclusions based on this experiment, because Ki16425 is an inhibitor of LPA₁, but not LPA₂. In addition, because a limited amount of Ki16425 was available, only one dose of Ki16425 was used and treatment was given at the late stage of 12 days. Therefore, further experiments with higher doses of Ki16425 for longer duration or dual inhibitors of LPA₁ and LPA₂, if available, are warranted to elucidate the real role of LPA receptors in mesothelioma cells *in vivo*.

In summary, we demonstrated that MPM are heterogeneous in terms of the response to LPA and the expression of LPA receptors. However, at least in particular populations of MPM, LPA stimulated proliferation via LPA₁ through MAPK and motility via LPA₂ through RhoA. These novel findings suggest that LPA₁ and LPA₂ may be one of the therapeutic targets for controlling the proliferation and motility of MPM cells. As MPM develops into locally advanced disease, novel strategies targeting both LPA₁ and LPA₂ might be beneficial for suppressing the progression of MPM. Further *in vivo* experiments with dual inhibitors of LPA₁ and LPA₂ are warranted.

Acknowledgments

This work was supported by Grants-in-Aid 17016051 for Cancer Research from the Ministry of Education, Science, Sports and Culture of Japan (S. Sone). We thank Dr Hironobu Hamada for the EHMES-1 and EHMES-10 cell lines and Dr Adi F. Gazdar for the NCI-H290 and NCI-H513 cell lines.

References

- 1 Broaddus VC. Asbestos, the mesothelial cell and malignancy: a matter of life or death. *Am J Respir Cell Mol Biol* 1997; **17**: 657–9.
- 2 Morinaga K, Kishimoto T, Sakatani M, Akira M, Yokoyama K, Sera Y. Asbestos-related lung cancer and mesothelioma in Japan. *Industrial Health* 2001; **39**: 65–74.
- 3 Dufresne A, Bégin R, Churg A, Massé S. Mineral fiber content of lungs in patients with mesothelioma seeking compensation in Quebec. *Am J Respir Crit Care Med* 1996; **153**: 711–18.
- 4 Carbone M, Kratzke RA, Testa JR. The pathogenesis of mesothelioma. *Semin Oncol* 2002; **29**: 2–17.
- 5 Robinson BW, Lake RA. Advances in malignant mesothelioma. *N Engl J Med* 2005; **353**: 1591–603.
- 6 Eichholtz T, Jalink K, Fahrenfort I, Moolenaar WH. The bioactive phospholipid lysophosphatidic acid is released from activated platelets. *Biochem J* 1993; **291**: 677–80.
- 7 Gerrard JM, Robinson P. Identification of the molecular species of lysophosphatidic acid produced when platelets are stimulated by thrombin. *Biochim Biophys Acta* 1989; **1001**: 282–5.
- 8 Shen Z, Belinson J, Morton RE, Xu Y, Xu Y. Phorbol 12-myristate 13-acetate stimulates lysophosphatidic acid secretion from ovarian and cervical cancer cells but not from breast or leukemia cells. *Gynecol Oncol* 1998; **71**: 364–8.
- 9 Aoki J, Taira A, Takanezawa Y *et al*. Serum lysophosphatidic acid is produced through diverse phospholipase pathways. *J Biol Chem* 2002; **277**: 48737–44.
- 10 Siess W, Zangl KJ, Essler M *et al*. Lysophosphatidic acid mediates the rapid activation of platelets and endothelial cells by mildly oxidized low density lipoprotein and accumulates in human atherosclerotic lesions. *Proc Natl Acad Sci USA* 1999; **96**: 6931–6.
- 11 Fang X, Schummer M, Mao M *et al*. Lysophosphatidic acid is a bioactive mediator in ovarian cancer. *Biochim Biophys Acta* 2002; **1582**: 257–64.
- 12 Mills GB, May C, Hill M, Campbell S, Shaw P, Marks A. Ascitic fluid from human ovarian cancer patients contains growth factors necessary for intraperitoneal growth of human ovarian adenocarcinoma cells. *J Clin Invest* 1990; **86**: 851–5.
- 13 Xu Y, Gaudette DC, Boynton JD *et al*. Characterization of an ovarian cancer activating factor in ascites from ovarian cancer patients. *Clin Cancer Res* 1995; **1**: 1223–32.
- 14 Contos JJ, Ishii I, Chun J. Lysophosphatidic acid receptors. *Mol Pharmacol* 2000; **58**: 1188–96.
- 15 Noguchi K, Ishii S, Shimizu T. Identification of p2y9/GPR23 as a novel G protein-coupled receptor for lysophosphatidic acid, structurally distant from the Edg family. *J Biol Chem* 2003; **278**: 25 600–6.
- 16 Lee CW, Rivera R, Gardell S, Dubin AE, Chun J. GPR92 as a new G12/13- and Gq-coupled lysophosphatidic acid receptor that increases cAMP, LPA5. *J Biol Chem* 2006; **281**: 23 589–97.
- 17 An S, Bleu T, Hallmark OG, Goetzl EJ. Characterization of a novel subtype of human G protein-coupled receptor for lysophosphatidic acid. *J Biol Chem* 1998; **273**: 7906–10.
- 18 Bandoh K, Aoki J, Hosono H *et al*. Molecular cloning and characterization of a novel human G-protein-coupled receptor, EDG7, for lysophosphatidic acid. *J Biol Chem* 1999; **274**: 27776–85.
- 19 Kishi Y, Okudaira S, Tanaka M *et al*. Autotaxin is overexpressed in glioblastoma multiforme and contributes to cell motility of glioblastoma by converting lysophosphatidylcholine to lysophosphatidic acid. *J Biol Chem* 2006; **281**: 17492–500.
- 20 Shida D, Watanabe T, Aoki J *et al*. Aberrant expression of lysophosphatidic acid (LPA) receptors in human colorectal cancer. *Lab Invest* 2004; **84**: 1352–62.
- 21 Schulte KM, Beyer A, Köhrer K, Oberhäuser S, Röher HD. Lysophosphatidic acid, a novel lipid growth factor for human thyroid cells: over-expression of the high-affinity receptor edg4 in differentiated thyroid cancer. *Int J Cancer* 2001; **92**: 249–56.
- 22 Kitayama J, Shida D, Sako A *et al*. Over-expression of lysophosphatidic acid receptor-2 in human invasive ductal carcinoma. *Breast Cancer Res* 2004; **6**: R640–6.
- 23 Mills GB, Eder A, Fang X *et al*. Critical role of lysophospholipids in the pathophysiology, diagnosis, and management of ovarian cancer. *Cancer Treat Res* 2002; **107**: 259–83.
- 24 Shida D, Kitayama J, Yamaguchi H *et al*. Dual mode regulation of migration by lysophosphatidic acid in human gastric cancer cells. *Exp Cell Res* 2004; **301**: 168–78.
- 25 Guo R, Kasbohm EA, Arora P *et al*. Expression and function of lysophosphatidic acid LPA1 receptor in prostate cancer cells. *Endocrinology* 2006; **147**: 4883–92.
- 26 Boucharaba A, Serre CM, Guglielmi J, Bordet JC, Clezardin P, Peyruchaud O. The type 1 lysophosphatidic acid receptor is a target for therapy in bone metastases. *Proc Natl Acad Sci USA* 2006; **103**: 9643–8.
- 27 Goetzl EJ, Dolezalova H, Kong Y *et al*. Distinctive expression and functions of the type 4 endothelial differentiation gene-encoded G protein-coupled receptor for lysophosphatidic acid in ovarian cancer. *Cancer Res* 1999; **59**: 5370–5.

- 28 Shida D, Kitayama J, Yamaguchi H *et al.* Lysophosphatidic acid (LPA) enhances the metastatic potential of human colon carcinoma DLD1 cells through LPA1. *Cancer Res* 2003; **63**: 1706–11.
- 29 Yamada T, Sato K, Komachi M *et al.* Lysophosphatidic acid (LPA) in malignant ascites stimulates motility of human pancreatic cancer cells through LPA1. *J Biol Chem* 2004; **279**: 6595–605.
- 30 Yokoyama A, Kohno N, Fujino S *et al.* Origin of heterogeneity of interleukin-6 (IL-6) levels in malignant pleural effusions. *Oncol Rep* 1994; **1**: 507–11.
- 31 Nakataki E, Yano S, Matsumori Y *et al.* Novel orthotopic implantation model of human malignant pleural mesothelioma (EHMES-10 cells) highly expressing vascular endothelial growth factor and its receptor. *Cancer Sci* 2006; **97**: 183–91.
- 32 Usami N, Fukui T, Kondo M *et al.* Establishment and characterization of four malignant pleural mesothelioma cell lines from Japanese patients. *Cancer Sci* 2006; **97**: 387–94.
- 33 Yano S, Muguruma H, Matsumori Y *et al.* Antitumor vascular strategy for controlling experimental metastatic spread of human small-cell lung cancer cells with ZD6474 in natural killer cell-depleted severe combined immunodeficient mice. *Clin Cancer Res* 2005; **11**: 8789–98.
- 34 Ohta H, Sato K, Murata N *et al.* Ki16425, a subtype-selective antagonist for EDG-family lysophosphatidic acid receptors. *Mol Pharmacol* 2003; **64**: 994–1005.
- 35 Lindholm PM, Salmenkivi K, Vauhkonen H *et al.* Gene copy number analysis in malignant pleural mesothelioma using oligonucleotide array CGH. *Cytogenet Genome Res* 2007; **119**: 46–52.
- 36 Harvey I, Pass Zhandong L, Anil W *et al.* Gene expression profiles predict survival and progression of pleural mesothelioma. *Clin Cancer Res* 2004; **10**: 849–59.
- 37 Mills GB, Mooleenaar WH. The emerging role of lysophosphatidic acid in cancer. *Nat Rev Cancer* 2003; **3**: 582–91.
- 38 Stähle M, Veit C, Bachfischer U *et al.* Mechanisms in LPA-induced tumor cell migration: critical role of phosphorylated ERK. *J Cell Sci* 2003; **116**: 3835–46.
- 39 Hama K, Aoki J, Fukaya M *et al.* Lysophosphatidic acid and autotaxin stimulate cell motility of neoplastic and non-neoplastic cells through LPA1. *J Biol Chem* 2004; **279**: 17634–9.
- 40 Do TV, Symowicz JC, Berman DM *et al.* Lysophosphatidic acid down-regulates stress fibers and up-regulates pro-matrix metalloproteinase-2 activation in ovarian cancer cells. *Mol Cancer Res* 2007; **5**: 121–31.
- 41 Michiels F, Habets GG, Stam JC, van der Kammen RA, Collard JG. A role for Rac in Tiam1-induced membrane ruffling and invasion. *Nature* 1995; **375**: 338–40.
- 42 Itoh K, Yoshioka K, Akedo H, Uehata M, Ishizaki T, Narumiya S. An essential part for Rho-associated kinase in the transcellular invasion of tumor cells. *Nat Med* 1999; **5**: 221–5.