

B-cell linker protein prevents aneuploidy by inhibiting cytokinesis

Hiroki Kamino,¹ Manabu Futamura,¹ Yasuyuki Nakamura,¹ Noriaki Kitamura, Koki Kabu and Hirofumi Arakawa²

Cancer Medicine and Biophysics Division, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

(Received July 21, 2008/Revised August 14, 2008/Accepted August 15, 2008/Online publication November 19, 2008)

Aneuploidy is a hallmark of human cancers. Although the maintenance of genomic integrity by p53 is important in preventing aneuploidy, its mechanism remains to be elucidated. Here we report evidence that B-cell linker protein (BLNK) mediates the inhibition of cytokinesis, which generates tetraploidy but prevents aneuploidy. We identified BLNK as a transcriptional target of p53. Surprisingly, ectopic expression of exogenous BLNK inhibited cytokinesis, resulting in the formation of tetraploid cells. Indeed, BLNK was involved in the generation of spontaneously arising binucleate tetraploid cells. Interestingly, cytokinesis after DNA damage was inhibited in p21^{-/-} and p53^{+/+} cells, but not in p53^{-/-} cells. BLNK knockdown in p53^{+/+} and p21^{-/-} cells enhanced cytokinesis after DNA damage, leading to the generation of aneuploid cells. In addition, a BLNK-downregulated human pre-B leukemia cell line showed increased cytokinesis and aneuploidy after DNA damage compared with two other pre-B leukemia cell lines expressing higher levels of BLNK. These results suggest that BLNK acts as a mediator of p53 in the inhibition of cytokinesis, which prevents aneuploidy. We propose that the inhibition of cytokinesis is crucial for the maintenance of genomic integrity. (Cancer Sci 2008; 99: 2444–2454)

The tumor suppressor p53 prevents malignant transformation when cells suffer stresses including severe DNA damage.⁽¹⁾ Hence, p53 is known as 'a guardian of the human genome'.⁽²⁾ Indeed, p53 is mutated in more than 50% of all human cancers, emphasizing its essential role in tumorigenesis in any of the human cancers. The p53 gene encodes a transcription factor that binds to a specific sequence of its downstream target gene. Therefore, p53 exerts its functions via transcriptional activation of various target genes. Although four major genes, including those functioning in cell-cycle arrest, apoptosis, DNA repair, and antiangiogenesis, are considered to be involved in the core mechanism of p53-regulated tumor suppression, a considerable number of target genes have been reported and their functions display great diversity, implying that many p53-target genes still remains unknown.^(3,4) Because the maintenance of genomic integrity by p53 is critical in preventing aneuploidy,^(5,6) identification of the p53-target genes involved in this function is important.

Aneuploidy is a major characteristic of various human cancers.⁽⁷⁾ Therefore, aneuploidy likely plays a significant role in cancer initiation and progression.⁽⁸⁾ There are several checkpoint systems to prevent aneuploidy in normal cells, which monitor proper chromosome segregation, cytokinesis, and cell-cycle progression. Recently, the p53-regulated tetraploidy checkpoint was reported to prevent the progression of aneuploidy.^(9–12) At this checkpoint, most cells that suffer from DNA damage progress to mitosis without cytokinesis and arrest at G₁ phase of the cell cycle.⁽¹²⁾ Therefore, the arrested cells with DNA damage become tetraploid at G₁ phase. If the function of p53 is impaired, the tetraploid cells at G₁ begin to progress through the cell cycle, replicate DNA, and continue cell division, resulting in the progression of aneuploidy. Although p21/WAF1 has been shown to play a

critical role in this checkpoint pathway,⁽¹²⁾ the precise mechanism remains to be elucidated.

B-cell linker protein (BLNK),⁽¹³⁾ also called SLP-65⁽¹⁴⁾ or B-cell adaptor containing SH2 domain protein (BASH),⁽¹⁵⁾ is a B-cell adaptor molecule and plays an essential role in signal transduction from pre-B-cell receptor and B-cell antigen receptor. As BLNK encodes no intrinsic enzymatic activity, its function is to serve as a scaffold for assembling molecular complexes.⁽¹⁶⁾ After phosphorylation by the kinase Syk, BLNK couples enzymes (phospholipase C, γ 2, and Vav) and additional linker proteins (Grb2 and Nck), and regulates downstream signaling pathways.^(13–15) The important function of BLNK has been demonstrated by the phenotype of BLNK-deficient mice, which show an incomplete block in B-cell development at the pre-B-cell and immature B-cell stage, suggesting that BLNK is essential for mature development of B cells.⁽¹⁷⁾ More importantly, the BLNK-deficient mice have a high incidence of spontaneous pre-B-cell lymphoma, which likely results from the enhanced proliferative capacity of BLNK-deficient pre-B cells.⁽¹⁸⁾ Consistent with this finding, approximately 50% of human childhood pre-B acute lymphoblastic leukemias show complete loss or drastic reduction of BLNK expression.⁽¹⁹⁾ Those results provide evidence that BLNK acts as a tumor suppressor.^(18–20)

Cytokinesis is the physical process by which a cell divides after the completion of mitosis.^(21,22) Cell division is completed through the sequential coordination of chromosome segregation and cytokinesis. Mitosis without cytokinesis can produce multinucleate polyploid cells. Cytokinesis at the wrong location and time can lead to the generation of aneuploid cells or mitotic catastrophe. However, despite its importance in cell growth and division, little attention has been paid to the role of the inhibition of cytokinesis in tumorigenesis. Here we report on a cytokinesis-blocking role of BLNK, which is a candidate tumor-suppressor gene in pre-B-cell leukemia. In addition, BLNK-regulated cytokinesis inhibition appears to be involved in the p53-regulated maintenance of genomic integrity, which prevents aneuploidy. We therefore propose that cytokinesis block is an essential safeguard system that prevents aneuploidy arising from cell division with chromosome missegregation and breakage or DNA damage.

Materials and Methods

Detection of binucleate cells. Various HDK1 and HMEC4 cells, including parental, GFPsi, BLNKsi-1 (si-1), BLNKsi-2 (si-2), and BLNKsi-3 (si-3) cells, were seeded in six-well plates at a density of 1.5×10^5 or 2×10^5 cells/plate, respectively. After 72

¹These authors contributed equally to this work.

²To whom correspondence should be addressed. E-mail: harakawa@ncc.go.jp

(HDK1) or 24 h (HMEC4), we randomly captured up to 50 fields of vision, each of which included approximately 20 cells, as photographic images using an Olympus IX71 microscope and DP70-BSW software (Olympus, Tokyo, Japan). The numbers of mononucleate, binucleate, and multinucleate cells were counted in approximately 500–600 independent cells. The ratio of the binucleate cells (including approximately 10% multinucleate cells) to the total number of cells was calculated and evaluated statistically.

Detection of micronucleate cells. Various HCT116-p53^{+/+} or p21^{-/-} cells, including parental, GFPsi, si-1, si-2, and si-3 cells, were seeded in six-well plates at a density of 3×10^5 cells/plate. After 24 h, the cells were treated with 1 μ g/mL adriamycin for 2 h. At the indicated times, 25–30 fields of vision, each of which included approximately 20 cells, were captured as photographic images using an Olympus IX71 microscope with DP70-BSW software. The numbers of mononucleate, binucleate, and micronucleate cells were counted in approximately 500 independent cells. The ratio of the mononucleate, binucleate, or micronucleate cells to the total number of cells was calculated and evaluated statistically.

Cytokinesis index. HCT116-p21^{-/-} or HCT116-p21^{-/-}-BLNKsi-1 cells were treated with 1 μ g/mL adriamycin for 2 h, and then 250–350 independent cells were monitored until 24 h after treatment by time-lapse microscopy. The number of cells that had undergone cytokinesis was counted, and the ratio of the cells that had undergone cytokinesis to the total number of cells was calculated and evaluated statistically.

Fluorescence *in situ* hybridization analysis. To examine aneuploidy with probes specific for chromosomes 8 and 12, CEP8 (30-160008; Abbott Molecular, Des Plaines, IL, USA) and CEP12 (D12Z3, 32-112012; Abbott Molecular) DNA probe kits were used. The cells were fixed in Carnoy's liquid (acetic acid : methanol = 3:1) at room temperature for 30 min. Slide glasses were allowed to dry, and were incubated sequentially in 2 \times saline-sodium citrate (SSC) and 0.1% Nonidet P-40 (NP-40) at 37°C for 30 min. The slides were then incubated in increasing concentrations of ethanol (70, 85, and 100%) for 1 min each. Then, the slides were incubated in 70% formamide and 2 \times SSC at 73°C for 5 min, and incubated in increasing concentrations of ethanol (70, 85, and 100%) for 1 min each. The slides were allowed to dry completely, and then 10 μ L hybridization solution containing labeled probes at a concentration of 1 ng/ μ L per probe was added to each slide glass, which was incubated at 42°C for 16 h. The slides were washed three times in 50% formamide and 2 \times SSC at 45°C for 10 min, and then incubated in 0.1% NP-40 and 2 \times SSC at 45°C for 5 min. The nuclei on the slides were stained with 4', 6-diamidino-2-phenylindole (DAPI) II. The chromosome 8 (green) and 12 (red) signals were examined using an Olympus IX71 microscope and DP70-BSW software in 300 independent nuclei to detect aneuploid cells, which were defined as the number cells with zero, one, or three signals of either chromosome 8 or chromosome 12. The ratio of aneuploid cells to the total number of cells was calculated and evaluated statistically.

Results

Identification of BLNK as a novel p53-inducible gene. In order to identify new p53-target genes, we screened p53-inducible genes using microarray analysis, as reported previously.⁽²³⁾ The expression level of BLNK was notably elevated in the hepatoblastoma cell line HepG2 infected with Ad-p53-wt (data not shown). BLNK is an adaptor protein that is involved in B-cell receptor signaling,^(13–15) and its role in B-cell tumorigenesis has previously been reported in both human and mouse pre-B leukemias.^(18–20) As the expression of BLNK can be induced by p53, we speculated that it might be involved in the

tumorigenesis of various cell types in addition to B cells. We therefore further analyzed the activities of BLNK. As indicated in Figure 1a,b, BLNK expression was strongly induced in response to DNA damage in p53 wild-type colorectal cancer cell lines (HCT116-p53^{+/+} and LS174T-control), but the induction of BLNK was severely impaired in the corresponding isogenic p53-deficient cell lines (HCT116-p53^{-/-} and LS174T-p53si). Similar results were obtained in other cancer cell lines, including HepG2 and the breast cancer cell line MCF7 (data not shown). These results suggest that endogenous p53 activates the transcription of BLNK in response to DNA damage in various cell types in addition to B cells.

We identified a possible p53-binding sequence (p53BS) in intron 1 of the BLNK gene (Fig. 1c). A chromatin-immunoprecipitation assay indicated that the DNA fragment containing p53BS was immunoprecipitated with an anti-p53 antibody, and was polymerase chain reaction amplified with the precipitated genomic DNA; this suggested that p53 interacts with p53BS *in vivo* (Fig. 1d). Moreover, a heterologous reporter assay showed that the transcriptional activities of luciferase reporter plasmids containing p53BS (pGL-pro-BS, pGL-promoter 1xBS, and pGL-promoter 2xBS) were strongly enhanced when they were cotransfected with wild-type (WT)-p53 but not mutant (MT)-p53, indicating that p53BS is a p53-responsive sequence of BLNK (Fig. 1d). Based on these findings, we concluded that BLNK is a bona fide target of the tumor suppressor p53.

B-cell linker protein inhibits cytokinesis and induces binucleate tetraploid cells. To explore the function of BLNK as a mediator of p53, we prepared an adenovirus vector containing full-length BLNK complementary DNA (cDNA; Ad-BLNK), which was designed to express the full-length BLNK protein. As indicated in Figure 2a,b, when p53-deficient cells (such as HCT116-p53^{-/-} and HepG2-p53si) were infected with Ad-BLNK, the number of tetraploid cells with 4 N DNA content increased by 24 h. This phenomenon was also observed in the p53-mutated glioblastoma cell line U373MG (Fig. 2c). All of the cell lines infected with Ad-BLNK experienced cell death 48–72 h after infection. These results suggest that the overexpression of BLNK preferentially generates tetraploidy.

To investigate the mechanism of the BLNK-induced tetraploidy, U373MG cells infected with Ad-BLNK were monitored by time-lapse microscopy. Surprisingly, most of the cells infected with Ad-BLNK showed inhibition of cytokinesis, resulting in the generation of tetraploid cells (Fig. 2d). These results suggest that BLNK might be involved in the inhibition of cytokinesis and the generation of tetraploid cells.

The role of BLNK is closely related to the generation of tetraploid cells via cytokinesis block. A previous report showed that chromosome non-disjunction accelerated tetraploid cell formation in order to prevent aneuploidy.⁽²⁴⁾ According to this model, tetraploid cells could be actively generated by an unknown mechanism in order to stop cell cycle progression of cells with chromosome non-disjunction.⁽²⁴⁾ We therefore speculated that BLNK might generate tetraploid cells in order to prevent aneuploidy. To validate this model, we inhibited the expression of BLNK with short interfering RNA (siRNA) against the BLNK sequence in normal human dermal keratinocyte (HDK1) cells, which were used previously,⁽²⁴⁾ as well as in normal human mammary epithelial cells (HMEC4). As shown in Figure 3a,b, the expression of endogenous BLNK mRNA was significantly downregulated in HDK1-si-1, HDK1-si-2, HMEC4-si-1, and HMEC4-si-2 cells. Consistent with these expression levels, BLNK knockdown significantly decreased the number of spontaneously arising binucleate tetraploid cells in these lines (Fig. 3a,b). Our results suggest that BLNK is important in the generation of spontaneously arising binucleate tetraploid cells.

B-cell linker protein is involved in the p53-regulated mononucleate tetraploid checkpoint. Contrary to previous observations,⁽²⁴⁾ DNA




Fig. 1. Identification of B-cell linker protein (BLNK) as a direct target gene of p53. Endogenous p53-dependent induction of BLNK in (a) HCT116-p53^{+/+} and p53^{-/-} cells, or (b) LS174T-control-si and p53-si cells after DNA damage by treatment with 1 μg/mL adriamycin. Expression levels are shown by (a) reverse transcription-polymerase chain reaction, and (b) northern and western blot analysis. Beta 2-microglobulin expression or coomassie brilliant blue (CBB) staining was used as a loading control. p21^{WAF1} was used as a positive control. (c) Identification of p53-binding sequence (p53BS) in the genomic DNA of BLNK. Black boxes indicate the locations and relative sizes of 17 exons; the arrow indicates the potential p53BS in intron 1 of BLNK. (d) Binding of p53 with p53BS of BLNK and p53-dependent transcriptional activity of p53BS. A chromatin-immunoprecipitation assay was carried out on the DNA-protein complex, which was immunoprecipitated with anti-p53 antibody from HepG2 cells infected with Ad-p53 or Ad-EGFP at a multiplicity of infection of 30. The heterologous luciferase reporter plasmid containing p53BS (258 bp), 1 × BS (20 bp) or 2 × BS (40 bp) was cotransfected with the plasmid designed to express wild-type p53 (wt), mutant p53 (mt), or no p53 (mock) into H1299 cells. The luciferase activity 24 h after transfection is shown in relation to the activity of the pGL3-promoter vectors without p53BS, 1 × BS, or 2 × BS.

damage did not enhance the generation of binucleate tetraploid cells in our current system (Fig. 3c,d). G₁ arrest of the mononucleate tetraploid cells after DNA damage was previously suggested to be involved in a p53-regulated DNA-damage checkpoint that induces long-term cell cycle arrest and subsequent cellular senescence.⁽²⁵⁾ We therefore proposed that BLNK-induced cytokinesis block might be involved in the DNA-damage checkpoint. Indeed, after DNA damage, the HCT116-p53^{+/+} cells moved from cell cycle phase G₂ to G₁ without cytokinesis, as indicated by decreased cyclin B1 and

CDK1 expression, and increased expression of cyclin E, cyclin D, and p27 (Fig. 4a,b). These cells eventually arrested in G₁ phase in the mononucleate tetraploid state until at least 72 h after the DNA damage when some cell death was observed (Fig. 4a,c). By contrast, HCT116-p53^{-/-} cells had a tetraploid DNA content 24 h after DNA damage (Fig. 4a) and cell cycle arrest was not observed, as indicated by increased levels of cyclin B1 and CDK1 expression, and decreased expression of cyclin E, cyclin D, and p27 (Fig. 4b). Eventually, the tetraploid DNA content decreased as a consequence of cell division




Fig. 2. B-cell linker protein (BLNK) induces binucleate tetraploid cells by inhibiting cytokinesis. (a–c) BLNK induces tetraploidy. p53-wild-type and p53-deficient cancer cell lines, including (a) HCT116-p53^{+/±} and p53^{-/-}, (b) HepG2-control and p53si, and (c) U373MG cells were infected with Ad-BLNK at a multiplicity of infection of 10. The DNA contents were examined by fluorescence-activated cell sorting (FACS) analysis at the indicated times. The results are shown as FACS images. (d) BLNK inhibits cytokinesis. A time-lapse microscopic analysis was carried out to monitor the cytokinesis of U373MG cells infected with Ad-BLNK at a multiplicity of infection of 10. Arrows (black, red, blue) indicate the cells that have undergone inhibition of cytokinesis.




Fig. 3. B-cell linker protein (BLNK) is involved in the generation of spontaneously arising binucleate tetraploid cells. (a,b) Spontaneously arising binucleate tetraploid cells. Downregulation of endogenous BLNK expression by short interfering RNA (siRNA) in (a) HDK1 or (b) HMEC4 cells. BLNK mRNA was downregulated with three different siRNA: BLNKsi-1 (si-1), BLNKsi-2 (si-2), and BLNKsi-3 (si-3). Parental HDK1 and HMEC4 cells (parent), or HDK1 and HMEC4 cells infected with retrovirus containing the siRNA of green fluorescence protein-si (GFPsi), were used as negative controls. Endogenous BLNK mRNA expression was examined by real-time polymerase chain reaction analysis. BLNK was involved in spontaneously arising binucleate tetraploidy. The numbers of binucleate cells in (a) HDK or (b) HMEC cells, in which BLNK was downregulated with the siRNA of si-1, si-2, and si-3, were counted. Parental HDK and HMEC cells, or HDK and HMEC cells infected with GFPsi siRNA, were used as negative controls. Error bars \pm SD ($n = 4$); $*P < 0.01$, Student's *t*-test. (c,d) Binucleate tetraploid cells after DNA damage. Endogenous expression levels of BLNK mRNA in various (c) HDK1 or (d) HMEC4 cells after DNA damage. Endogenous BLNK mRNA expression was examined by real-time polymerase chain reaction analysis in parental, control (GFPsi), and BLNK-knockdown (si-1, si-2, and si-3) HDK1 or HMEC4 cells after DNA damage with 1.0 μ g/mL adriamycin. DNA damage did not induce the generation of binucleate tetraploid cells. The numbers of binucleate cells in (c) HDK or (d) HMEC cells, in which BLNK was downregulated with si-1, si-2, and si-3 siRNA, were counted after DNA damage by 1.0 μ g/mL adriamycin (ADR). Parental HDK and HMEC cells, or HDK and HMEC cells infected with GFPsi siRNA, were used as negative controls. Error bars \pm SD ($n = 4$); $*P < 0.01$, Student's *t*-test.

leading to aneuploidy or cell death (Fig. 4a). In addition, the cells had abnormal multilobulated nuclei (referred to hereafter as micronucleate cells) as described previously⁽²⁶⁾ (Fig. 4c). However, the phenotype of the HCT116-p21^{-/-} cells appeared to differ from that of the HCT116-p53^{-/-} cells. In the early phase, similar to the HCT116-p53^{-/-} cells, the HCT116-p21^{-/-} cells became tetraploid by 24 h, and the cell cycle subsequently proceeded with a significant fraction of dead cells (Fig. 4a). However, in contrast to the p53^{-/-} cells, many HCT116-p21^{-/-} cells were not only micronucleate but also relatively large in size, and had a polyploid (>4N) phenotype 72 h after DNA damage (Fig. 4a,c), implying a failure of cytokinesis. Time-lapse microscopy confirmed that HCT116-p21^{-/-} cells had entered M phase, but many failed to complete cytokinesis, thereby resulting in endomitosis and a polyploid phenotype (data not shown). We hypothesized that BLNK might contribute to the failure of cytokinesis and the polyploid phenotype in the HCT116-p21^{-/-} cells.

We thus examined the expression levels of p53, p21, and BLNK in these cell lines. As expected, p53 and p21 expression

was not detected in HCT116-p53^{-/-} and p21^{-/-} cells, respectively (Fig. 4b). Interestingly, the expression level of BLNK in the HCT116-p21^{-/-} cells was as much as twice that in the HCT116-p53^{+/+} cells (Fig. 4b). Therefore, in order to examine the role of BLNK in cytokinesis inhibition after DNA damage, its expression was downregulated with siRNA against the BLNK sequence (Supporting Fig. S1a,b). As shown in Figure 4d, BLNK knockdown significantly enhanced the generation of micronucleate cells after DNA damage in HCT116-p53^{+/+} and p21^{-/-} cells.

B-cell linker protein prevents aneuploidy by inhibiting cytokinesis.

To determine whether the elevation of micronucleate cells was caused by enhanced cytokinesis, we monitored the HCT116-p21^{-/-} parent cells and the BLNK-si-1 cells until 24 h after DNA damage using time-lapse microscopy, and counted the number of cells that completed cytokinesis during this period. As indicated in Figure 5a, BLNK knockdown in the BLNK-si-1 cells enhanced cytokinesis by as much as three times compared with the parental cells, implying that BLNK was involved in the failure of cytokinesis in the HCT116-p21^{-/-} cells. We proposed that the failure of cytokinesis induced by BLNK might prevent




Fig. 4. B-cell linker protein (BLNK) is involved in the p53-regulated mononucleate tetraploid G₁ checkpoint after DNA damage. (a) The DNA contents of HCT116-p53^{+/+}, p53^{-/-}, and p21^{-/-} cells after DNA damage. HCT116-p53^{+/+}, p53^{-/-}, and p21^{-/-} cells were treated with 1.0 μg/mL adriamycin, and the DNA contents were analyzed at 0, 24, 48, and 72 h after the treatment, by green fluorescence activated cell sorting (FACS) analysis. The polyploidy phenotype after DNA damage was observed only in HCT116-p21^{-/-} cells: M1 ≤ 2 N; M2 = 2 N; M3 = 2 N–4 N; M4 = 4 N; M5 ≥ 4 N. The results were shown by FACS images (left) and bar graphs (right). (b) Expression levels of various cell cycle regulators and BLNK in HCT116-p53^{+/+}, p53^{-/-}, and p21^{-/-} cells after DNA damage. HCT116-p53^{+/+}, p53^{-/-}, and p21^{-/-} cells were treated with 1.0 μg/mL adriamycin, and the expression levels of p53, p21/waf1, cyclin B1, CDK1/cdc2, cyclin E, cyclin D1, and p27 were examined by western blot analysis with specific antibodies at the times indicated. β-Actin was used as a loading control. BLNK was highly expressed in HCT116-p21^{-/-} cells after DNA damage. BLNK mRNA expression was examined by reverse transcription–polymerase chain reaction. Each expression level is shown relative to the level of BLNK at 0 h in HCT116-p53^{+/+} cells. (c) Nuclear and cellular appearance of HCT116-p53^{+/+} and p21^{-/-}, and large polyploid cells in HCT116-p21^{-/-}. Images are shown 48 h after DNA damage. (d) BLNK inhibited the generation of micronucleate cells after DNA damage. HCT116-p53^{+/+} or p21^{-/-} cells, in which BLNK expression was downregulated with the short interfering RNA (siRNA) of BLNKsi-1 (si-1), BLNKsi-2 (si-2), and BLNKsi-3 (si-3), were treated with 1.0 μg/mL adriamycin, and the numbers of mononucleate, binucleate, and micronucleate cells were counted 24 h after DNA damage. Parental HCT116-p53^{+/+} and p21^{-/-} cells (parent) or these cells infected with green fluorescence protein siRNA (GFPsi) were used as negative controls. Error bars ± SD (n = 4); *P < 0.01, **P < 0.001, Student's *t*-test between GFPsi and BLNK-si cells (si-1, si-2 and si-3). White, gray, and black bars indicate cells with mononucleate, binucleate, and multilobulated nuclei, respectively.




Fig. 5 B-cell linker protein (BLNK) prevents aneuploidy by inhibiting cytokinesis after DNA damage. (a) Cytokinesis enhancement by BLNK knockdown. HCT116-p21^{+/+} and p21^{-/-}-BLNKsi-1 cells were treated with 1.0 μ g/mL adriamycin, and then monitored for 24 h by time-lapse microscopy. The numbers of cells that completed cytokinesis were counted during this period. The cytokinesis index indicates the ratio of the cells that underwent cytokinesis to the total number of cells. Error bars \pm SD ($n = 5$); $**P < 0.001$, Student's t -test. (b) Impairment of BLNK induces aneuploidy after DNA damage. HCT116-p21^{+/+} and p21^{-/-}-BLNKsi-1 cells were treated with 1.0 μ g/mL adriamycin, and 24 h after treatment, the numbers of chromosomes 8 and 12 in 300 independent cells were examined by fluorescence *in situ* hybridization (FISH) analysis. Signals of 0, 1, and 3 were judged to indicate aneuploidy. The experiments were repeated four times. Error bars \pm SD ($n = 4$); $*P < 0.01$, $**P < 0.001$, Student's t -test. (c) Representative results of aneuploid HCT116-p21^{+/+} cells. HCT116-p21^{+/+} cells were treated with 1.0 μ g/mL adriamycin, and 24 h after the treatment the cells were subjected to FISH analysis, in which probes of chromosome 8 (green) and chromosome 12 (red) were used to detect aneuploidy. Signals of 0, 1, and 3 in either chromosome 8 or 9 were judged to indicate aneuploidy: 1 = diploidy (chromosome 8, 2; chromosome 12, 2); 2 = tetraploidy (chromosome 8, 4; chromosome 12, 4); 3 = aneuploidy (arrow, chromosome 8, 2; chromosome 12, 3); 4 = aneuploidy (arrow, chromosome 8, 1; chromosome 12, 2); 5 = aneuploidy (arrow, chromosome 8, 2; chromosome 12, 1); 6 = aneuploidy (chromosome 8, 1; chromosome 12, 1); 7 = aneuploidy (chromosome 8, 2; chromosome 12, 1); 8 = aneuploidy (chromosome 8, 2; chromosome 12, 0); 9 = aneuploidy (chromosome 8, 3; chromosome 12, 2); 10 = aneuploidy (chromosome 8, 3; chromosome 12, 3); 11 = aneuploidy (chromosome 8, 3; chromosome 12, 1); 12 = aneuploidy (chromosome 8, 4; chromosome 12, 3); 13 = aneuploidy (arrow, chromosome 8, 1; chromosome 12, 2); 14 = aneuploidy (chromosome 8, 3; chromosome 12, 4); 15 = aneuploidy (chromosome 8, 10; chromosome 12, 9). (d) Impairment of BLNK-induced aneuploidy after DNA damage. HCT116-p53^{+/+}-parent, p53^{+/+}-BLNKsi-1, and p53^{-/-}-parent cells were treated with 1.0 μ g/mL adriamycin, and 24 h after the treatment the numbers of chromosomes 8 and 12 in 300 independent cells were examined by FISH analysis. Signals of 0, 1, and 3 were judged to indicate aneuploidy. Error bars \pm SD ($n = 2$); $*P < 0.05$, $**P < 0.01$, Student's t -test.

aneuploidy derived from the division of cells with DNA damage. Therefore, cytokinesis enhancement after DNA damage would cause an increase in the number of aneuploid cells. To validate this notion, we examined aneuploid cells in these lines by fluorescence *in situ* hybridization analysis with probes for chromosomes 8 and 12, as described previously.⁽²⁴⁾ Importantly, the enhancement of cytokinesis caused by the inhibition of BLNK significantly increased the number of aneuploid cells in the HCT116-p21^{-/-} and HCT116-p53^{+/+} cell lines (Fig. 5b,d). Representative signals of diploidy and aneuploidy are shown in Figure 5c. Interestingly, the frequency of aneuploid cells in the HCT116-p53^{+/+}-BLNKsi-1 cell line was similar to that in the HCT116-p53^{-/-} cell line (Fig. 5d). These results strongly suggest that BLNK is involved in a p53-regulated DNA-damage checkpoint via cytokinesis inhibition, that it plays a pivotal role in the generation of mononucleate tetraploid cells in G₁ phase, and that impairment of BLNK function induces cytokinesis after DNA damage, thereby leading to aneuploidy.

Potential role of BLNK as a cytokinesis inhibitor in pre-B-cell leukemia. Finally, we evaluated the role of BLNK as a cytokinesis inhibitor in human pre-B-cell leukemia cells. BLNK was originally identified as a mediator of B-cell receptor signaling,⁽¹³⁻¹⁵⁾ and its inactivation is closely related to the initiation of human and mouse pre-B-cell leukemia,⁽¹⁸⁻²⁰⁾ implying that it is a tumor-suppressor gene for B cells. As shown in Figure 6a, BLNK expression was severely impaired in HPB-null pre-B leukemia cells, whereas high expression levels were detected in REH and LAZ-221 pre-B leukemia cells. Interestingly, an accumulation of endogenous BLNK in the REH cell line was observed at the spindle midzone in late anaphase and at the midbody in telophase, especially overlapping the signals of Aurora-B at the midbody in late telophase (Supporting Fig. S2a). BLNK signals were also detected at the midbody in cytokinesis, and were colocalized between the two signals of Aurora-B (Supporting Fig. S2b). In addition, binucleate cells (~10%) were observed in the REH cell line, in which the BLNK signals tended to accumulate in the region between the two nuclei (Supporting Fig. 2a). Using these pre-B leukemia cell lines, we examined the status of cell division reflecting cytokinesis under various conditions, including floating or attached, and with or without DNA damage. Compared with the REH and LAZ-221 cells, the number of HPB-null cells increased more rapidly under all of the conditions tested, implying a higher frequency of cell division (Fig. 6b). In addition, the REH and LAZ-221 cells underwent cell cycle arrest with 2N and 4N DNA contents after DNA damage, whereas the NPB-null cells showed continued cell division and had an aneuploid DNA content (2-4N; Fig. 6b). To validate this result, we carried out fluorescence *in situ* hybridization analysis of these cell lines with the same probes as indicated in Figure 5c. Representative signals of diploidy and aneuploidy are shown in Supporting Figure S3. As indicated in Figure 6c, we detected aneuploid cell levels of approximately 14% (chromosome 8 or 12) or 23% (both chromosome 8 and 12) after DNA damage in the HPB-null cells; these levels were approximately four times higher than those in two other cell lines expressing high levels of BLNK. Taken together, these results suggest that BLNK plays an important role in cytokinesis inhibition, in order to prevent aneuploidy after DNA damage, even in B-cell lines.

Discussion

In the present paper, we demonstrated that BLNK functions as a cytokinesis inhibitor generating tetraploidy in order to prevent aneuploidy. Based on our findings and previous reports^(12,24,25) it appears likely that BLNK-induced tetraploidy plays a critical role in blocking cytokinesis of cells with chromosome mis-segregations, chromosome breakages, and DNA damage, thereby

preventing the occurrence of aneuploidy. We also demonstrated that there are two major states of tetraploidy, binucleate and mononucleate, in G₁-arrested cells: the former is probably generated from spontaneous chromosome mis-segregations,⁽²⁴⁾ whereas the latter is likely to be a G₁-arrest DNA-damage checkpoint that is regulated by p53.^(12,25) We consider the primary role of BLNK in the generation of tetraploidy to be the inhibition of cytokinesis (Figs 2,3; Supporting Fig. S2). However, the BLNK-knockdown experiment suggested that BLNK is also involved in keeping cells that have incurred DNA damage in a state of mononucleate tetraploidy at the G₁ phase of the cell cycle (Fig. 4). If this is the case, what is the regulatory mechanism for BLNK-induced mononucleate tetraploidy? Mitosis, as well as cytokinesis, did not occur during the movement of the mononucleate tetraploid cells from cell cycle phase G₂ to G₁ after DNA damage. If BLNK regulates only cytokinesis, how can the inhibition of BLNK promote mitosis? We speculate that, in addition to the inhibition of cytokinesis, BLNK might also regulate factors that trigger the start of mitosis, which determines the timing of the cell entering M phase. In fact, it is generally thought that M phase can be completed through sequential coordination of mitosis and cytokinesis, implying that the factor that regulates cytokinesis is also involved in mitosis. Further careful investigation of this process will be required to clarify the mechanism in detail.

A tetraploid-checkpoint hypothesis has been suggested to explain the p53-regulated checkpoint system.^(9-12,25) In this model, p53 recognizes tetraploid cells as abnormalities, such that cell cycle is arrested at G₁ phase in a p53-dependent manner, followed by senescence or cell death. However, in our model, p53 induces tetraploidy through the activation of BLNK, in order to prevent aneuploidy. Therefore, we consider p53-induced tetraploidy, regardless of whether it is mononucleate or binucleate, to be the 'last bastion' that blocks the generation of aneuploid cells, rather than being an abnormal state (Fig. 6d). It is likely that, in response to DNA damage, p53 stops the cell cycle at G₁ phase via p21/WAF1, and blocks cytokinesis via BLNK, leading to the G₁ arrest of tetraploid cells (Fig. 6d). Consistent with previous observations,^(12,25) DNA damage-induced tetraploid cells moved from cell cycle phase G₂ to G₁ without mitosis and cytokinesis in our current study. Thus, we speculate that BLNK might prevent accidental division of cells with DNA insult during their movement from cell cycle phase G₂ to G₁. We propose that p53-regulated tetraploid G₁ arrest mediated by the cooperation of p21/WAF1 and BLNK is an important safeguard system to maintain genomic integrity (Fig. 6d).

Recently, Takahashi *et al.* reported that sustained activation of reactive oxygen species-protein kinase C (ROS-PKC) σ signaling blocks cytokinesis irreversibly.⁽²⁶⁾ This irreversible cytokinetic block is likely to act as a second barrier to cellular immortalization, ensuring stable cell cycle arrest in human senescent cells.⁽²⁶⁾ Interestingly, we observed that human normal cell lines including HMEC4 and HDK1 sustained high-level BLNK protein expression at least until 7 days after DNA damage, when a senescence-like cell cycle arrest was induced in more than 50% of the cells (M. Futamura and H. Arakawa, unpublished data, 2008). Therefore, the irreversible cytokinetic block in senescent cells may be mediated by BLNK. It will be very interesting to further explore the role of BLNK in this novel tumor-suppressive mechanism.

Accumulating evidence suggests that the molecules promoting cytokinesis are strongly expressed in human cancers⁽²⁷⁻³⁰⁾ and that their activation or inactivation enhances cancer cell growth or death, respectively.⁽²⁷⁻²⁹⁾ Therefore, the molecules that are involved in cytokinesis progression have potential for use as molecular targets for cancer therapy.^(31,32) We demonstrated that BLNK was localized at the midzone and the midbody in late anaphase, telophase, and cytokinesis (Supporting Fig. S2a), and in the middle region of the two nuclei of binucleate cells, even




Fig. 6. Potential role of B-cell linker protein (BLNK) as an inhibitor of cytokinesis in pre-B-cell leukemia. (a) BLNK expression in human pre-B-cell leukemia. Three human pre-B-cell leukemia cell lines (HPB-null, REH, and LAZ-221) were irradiated at the indicated doses, and BLNK expression at 24 h after DNA damage was examined by western blot analysis. β -Actin was used as a loading control. (b) HPB-null cells lacking BLNK showed enhanced cell division and impaired tetraploid cell cycle arrest. Three human pre-B-cell leukemia cells (HPB-null, REH, and LAZ-221) were irradiated at 2.5 Gy, and the numbers of live cells were examined at the indicated times in the floating (gray bar) or attached (black bar) condition. The numbers of live cells without γ -irradiation were also examined at the indicated times in the floating (white bar) or attached (striped bar) condition. The DNA contents of each cell 24 h after DNA damage were determined by fluorescence activated cell sorting analysis. (c) Aneuploidy was generated preferentially from HPB-null cells lacking BLNK after DNA damage. HPB-null, REH, and LAZ-221 cells were irradiated at 2.5 Gy, and 24 h after treatment, the numbers of signals for chromosomes 8 and 12 in 300 independent cells were examined by fluorescence *in situ* hybridization analysis. Signals of 0, 1, and 3 were judged to indicate aneuploidy. The experiments were repeated twice. Error bars \pm SD ($n = 2$); $*P < 0.01$, Student's *t*-test. (d) Hypothetical model of BLNK-regulated cytokinesis.

in interphase (Supporting Fig. S2a). In addition, the signal was colocalized with Aurora-B in cytokinesis (Supporting Fig. S2b). This result suggests a number of possible ways in which BLNK might inhibit cytokinesis. At the midzone and the midbody, Aurora-B interacts with and regulates several mediators that are involved in cytokinesis promotion, including RhoA,⁽³³⁾ PRC1,⁽³⁴⁾ MgcRacGAP,⁽³⁵⁾ and Plk.⁽³⁶⁾ All of these molecules are thought to promote cytokinesis, and their dysfunction causes failure of cytokinesis and multinucleate polyploidy in cells. BLNK probably interacts with at least one of these molecules to prevent cytokinesis.

On the other hand, we actually observed that the enforced expression of BLNK in U373MG cells strongly induced activation of RhoA, but not Rac1 and Cdc42 (K. Kabu and H. Arakawa, unpublished data, 2008). RhoA activation is really critical for cytokinesis progression.⁽²¹⁾ However, at the late stage of cell division, the downregulation of RhoA activity at the midbody plays a crucial role in completion of cytokinesis probably through MgcRacGAP activation by Aurora B.^(21,35) Thus, we speculate that BLNK may prevent this downregulation of RhoA activity at the midbody, resulting in the failure of cytokinesis and generation of tetraploid cells.

Although a number of cytokinesis promoters have been reported thus far, a physiological inhibitor of cytokinesis has not previously been identified. We believe that BLNK is the first

candidate for a physiological cytokinesis inhibitor. Our findings thus shed light on the importance of the inhibition of cytokinesis in the maintenance of genomic integrity. In addition, our findings highlight the need for further investigations of the precise mechanism of this physiological cytokinesis inhibition.

Acknowledgments

We thank B. Vogelstein for the gift of the p53-knockout colorectal cancer cell lines, M. Kibara for the HPB-null, REH, and LAZ-221 cell lines, T. Kiyono and D.A. Galloway for the HDK1 and HMEC4 cell lines, and D. Kitamura for helpful comments on our experiments. We also thank S. Usuda, A. Ishizaka, and I. Hyo for technical assistance. This work was supported, in part, by grants from the Ministry of Health, Labour, and Welfare, Japan, and the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

Author contributions

H.K., M.F., Y.N., N.K., K.K., and H.A. carried out biochemical and cell biological experiments and analyzed the data; H.K., M.F., Y.N., and H.A. designed experiments; H.K. and H.A. wrote the paper. All authors discussed the results and commented on the manuscript.

References

- Vogelstein B, Lane D, Levine AJ. Surfing the p53 network. *Nature* 2000; **408**: 307–10.
- Vousden KH. Live or let die: The cell's response to p53. *Nat Rev Cancer* 2002; **2**: 594–604.
- Nakamura Y. Isolation of p53-target genes and their functional analysis. *Cancer Sci* 2004; **95**: 7–11.
- Arakawa H. p53, apoptosis and axon-guidance molecules. *Cell Death Differ* 2005; **12**: 1057–65.
- Liu G, Parant JM, Lang G *et al*. Chromosome stability, in the absence of apoptosis, is critical for suppression of tumorigenesis in Trp53 mutant mice. *Nat Genet* 2004; **36**: 63–8.
- Barboza JA, Liu G, Ju Z, El-Naggar AK, Lozano G. p21 delays tumor onset by preservation of chromosomal stability. *Proc Natl Acad Sci USA* 2006; **103**: 19 842–7.
- Kops GJ, Weaver BA, Cleveland DW. On the road to cancer: aneuploidy and the mitotic checkpoint. *Nat Rev Cancer* 2005; **5**: 773–85.
- Rajagopalan H, Lengauer C. Aneuploidy and cancer. *Nature* 2004; **432**: 338–41.
- Margolis RL. Tetraploidy and tumor development. *Cancer Cell* 2005; **8**: 353–4.
- Margolis RL, Lohez OD, Andreassen PR. G1 tetraploidy checkpoint and the suppression of tumorigenesis. *J Cell Biochem* 2003; **88**: 673–83.
- Andreassen PR, Lohez OD, Lacroix FB, Margolis RL. Tetraploid state induces p53-dependent arrest of nontransformed mammalian cells in G1. *Mol Biol Cell* 2001; **12**: 1315–28.
- Andreassen PR, Lacroix FB, Lohez OD, Margolis RL. Neither p21^{WAF1} nor 14-3-3 σ prevents G2 progression to mitotic catastrophe in human colon carcinoma cells after DNA damage, but p21^{WAF1} induces stable G1 arrest in resting tetraploid cells. *Cancer Res* 2001; **61**: 7660–8.
- Fu C, Turck CW, Kurosaki T, Chan AC. BLNK: a central linker protein in B cell activation. *Immunity* 1998; **9**: 93–103.
- Wienands J, Schweikert J, Wollscheid B *et al*. SLP-65: a new signaling component in B lymphocytes which requires expression of the antigen receptor for phosphorylation. *J Exp Med* 1998; **188**: 791–5.
- Goitsuka R, Fujimura Y, Mamada H *et al*. BASH, a novel signaling molecule preferentially expressed in B cells of the brusa of Fabricius. *J Immunol* 1998; **161**: 5804–8.
- Kurosaki T. Genetic analysis of B cell antigen receptor signaling. *Annu Rev Immunol* 1999; **17**: 555–92.
- Ishiai M, Kurosaki M, Pappu R *et al*. BLNK required for coupling Syk to PLC gamma 2 and Rac1-JNK in B cells. *Immunity* 1999; **10**: 117–25.
- Flemming A, Brummer T, Reth M, Jumaa H. The adaptor protein SLP-65 acts as a tumor suppressor that limits pre-B cell expansion. *Nat Immunol* 2003; **4**: 13–5.
- Jumaa H, Bossaller L, Portugal K *et al*. Deficiency of the adaptor SLP-65 in pre-B-cell acute lymphoblastic leukemia. *Nature* 2003; **423**: 452–6.
- Hayashi K, Yamamoto M, Nojima T, Goitsuka R, Kitamura D. Distinct signaling requirements for Dmu selection, IgH allelic exclusion, pre-B cell transition, and tumor suppression in B cell progenitors. *Immunity* 2003; **18**: 825–36.
- Piekny A, Werner M, Glotzer M. Cytokinesis: welcome to the Rho zone. *Trends Cell Biol* 2005; **15**: 651–8.
- Straight AF, Field CM. Microtubules, membrane and cytokinesis. *Curr Biol* 2000; **10**: R760–70.
- Masuda Y, Futamura M, Kamino H *et al*. The potential role of DFNA5, a hearing impairment gene, in p53-mediated cellular response to DNA damage. *J Hum Genet* 2006; **51**: 652–64.
- Shi Q, King RW. Chromosome nondisjunction yields tetraploid rather than aneuploid cells in human cell lines. *Nature* 2005; **437**: 1038–42.
- Bhonde MR, Hanski ML, Notter M *et al*. Equivalent effect of DNA damage-induced apoptotic cell death or long-term cell cycle arrest on colon carcinoma cell proliferation and tumour growth. *Oncogene* 2006; **25**: 165–75.
- Takahashi A, Ohtani N, Yamakoshi K *et al*. Mitogenic signalling and the p16^{INK4a}-Rb pathway cooperate to enforce irreversible cellular senescence. *Nat Cell Biol* 2006; **8**: 1291–7.
- Shimo A, Nishidate T, Ohta T *et al*. Elevated expression of protein regulator of cytokinesis 1, involved in the growth of breast cancer cells. *Cancer Sci* 2007; **98**: 174–81.
- Kanehira M, Katagiri T, Shimo A *et al*. Oncogenic role of MPHOSPH1, a candidate-testis antigen specific to human bladder cancer. *Cancer Res* 2007; **67**: 3276–85.
- Hayama S, Daigo Y, Yamabuki T *et al*. Phosphorylation and activation of cell division associated 8 by aurora kinase B plays a significant role in human lung carcinogenesis. *Cancer Res* 2007; **67**: 4113–22.
- Vishioni B, Oudejans JJ, Vos W, Rodriguez JA, Giaccone G. Frequent overexpression of aurora B kinase, a novel drug target, in non-small lung carcinoma patients. *Mol Cancer Ther* 2006; **5**: 2905–13.
- Keen N, Taylor S. Aurora-kinase inhibitors as anticancer agents. *Nat Rev Cancer* 2004; **4**: 927–36.
- Carvajal RD, Tse A, Schwartz GK. Aurora kinases: new targets for cancer therapy. *Clin Cancer Res* 2006; **12**: 6869–75.
- Yoshizaki H, Ohba Y, Kurokawa K *et al*. Activity of Rho-family GTPase during cell division as visualized with FRET-based probes. *J Cell Biol* 2003; **162**: 223–32.
- Jiang W, Jimenez G, Wells NJ *et al*. PRC1: a human mitotic spindle-associated CDK substrate protein required for cytokinesis. *Mol Cell* 1998; **2**: 877–85.
- Minoshima Y, Kawashima T, Hirose K *et al*. Phosphorylation by Aurora B converts MgcRacGAP to a RhoGAP during cytokinesis. *Dev Cell* 2003; **4**: 549–60.
- Nigg EA. Polo-like kinases: positive regulators of cell division from start to finish. *Curr Opin Cell Biol* 1998; **10**: 776–83.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Fig. S1. Inhibition of B-cell linker protein (BLNK) expression by short interfering RNA in HCT116-p53^{+/+} and HCT116-p21^{-/-} cells. (a) Downregulation of BLNK mRNA expression in (a) HCT116-p53^{+/+} and (b) HCT116-p21^{-/-} cells. Various cells, including parental (P), GFPsi (G), BLNKsi-1 (si-1), BLNKsi-2 (si-2), and BLNKsi-3 (si-3), were treated with 1.0 µg/mL adriamycin, and the BLNK mRNA expression levels were examined by real-time polymerase chain reaction 24 and 48 h after DNA damage. The expression levels are shown relative to the level (= 1) of (a) HCT116-p53^{+/+} and (b) HCT116-p21^{-/-} parental cells without DNA damage.

Fig. S2. Localization of B-cell linker protein (BLNK) at the mid-zone and mid-body during late anaphase, telophase, and cytokinesis. BLNK and Aurora B were colocalized at the mid-body in telophase. In REH cells, BLNK (green) and Aurora B (red) proteins were stained with anti-BLNK and anti-Aurora B antibodies, respectively. DNA (blue) was stained with Topro 3.B, BLNK and Aurora B were localized at the midbody in cytokinesis. BLNK (green) was localized at the center of the midbody between the two signals (red) of Aurora B in cytokinesis.

Fig. S3. Aneuploidy was preferentially generated from HPB-null cells lacking B-cell linker protein (BLNK) after DNA damage. Representative fluorescence *in situ* hybridization images are shown. The signals indicate the number of chromosomes 8 (green) and 12 (red): 1 = diploidy (chromosome 8, 2; chromosome 12, 2); 2 = aneuploidy (chromosome 8, 1; chromosome 12, 1); 3 = aneuploidy (chromosome 8, 2; chromosome 12, 0); 4 = aneuploidy (chromosome 8, 1; chromosome 12, 2); 5 = aneuploidy (chromosome 8, 2; chromosome 12, 1); 6 = aneuploidy (arrow, chromosome 8, 3; chromosome 12, 2). The individual (8 or 12) and combined (8 and 12) are shown.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.