




DATA NOTE

# The genome sequence of the John Dory, *Zeus faber* Linnaeus, 1758 [version 1; peer review: 2 approved, 1 not approved]

Patrick Adkins<sup>1</sup>, Joanna Harley<sup>1</sup>, Rachel Brittain<sup>1</sup>, Kesella Scott-Somme<sup>1</sup>, Freja Azzopardi <sup>1</sup>, Marine Biological Association Genome Acquisition Lab, Darwin Tree of Life Barcoding collective, Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team,

Wellcome Sanger Institute Scientific Operations: Sequencing Operations, Wellcome Sanger Institute Tree of Life Core Informatics team, Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

<sup>1</sup>The Marine Biological Association, Plymouth, England, UK

---

**V1** First published: 19 Mar 2024, 9:150  
<https://doi.org/10.12688/wellcomeopenres.21140.1>  
Latest published: 19 Mar 2024, 9:150  
<https://doi.org/10.12688/wellcomeopenres.21140.1>

---

## Abstract




We present a genome assembly from an individual *Zeus faber* (the John Dory; Chordata; Actinopteri; Zeiformes; Zeidae). The genome sequence is 804.7 megabases in span. Most of the assembly is scaffolded into 22 chromosomal pseudomolecules. The mitochondrial genome has also been assembled and is 16.72 kilobases in length.



## Keywords

Zeus faber, John Dory, genome sequence, chromosomal, Zeiformes

## Open Peer Review

Approval Status   

	1	2	3
<b>version 1</b> 19 Mar 2024	 <a href="#">view</a>	 <a href="#">view</a>	 <a href="#">view</a>

1. **Shawn M. Burgess**, National Human Genome Research Institute, Bethesda, USA
2. **Naiara Rodriguez-Ezpeleta** , AZTI Basque Research and Technology Alliance, Bizkaia, Spain
3. **Nastaran Mazloumi** , University of Tasmania Institute for Marine and Antarctic Studies, Hobart, Australia

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding author:** Darwin Tree of Life Consortium ([mark.blaxter@sanger.ac.uk](mailto:mark.blaxter@sanger.ac.uk))

**Author roles:** **Adkins P:** Investigation, Resources, Writing – Review & Editing; **Harley J:** Investigation, Resources; **Brittain R:** Investigation, Resources; **Scott-Somme K:** Investigation, Resources; **Azzopardi F:** Writing – Original Draft Preparation;

**Competing interests:** No competing interests were disclosed.

**Grant information:** This work was supported by Wellcome through core funding to the Wellcome Sanger Institute [206194, <https://doi.org/10.35802/206194>] and the Darwin Tree of Life Discretionary Award [218328, <https://doi.org/10.35802/218328>]. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2024 Adkins P *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Adkins P, Harley J, Brittain R *et al.* **The genome sequence of the John Dory, *Zeus faber* Linnaeus, 1758 [version 1; peer review: 2 approved, 1 not approved]** Wellcome Open Research 2024, 9:150 <https://doi.org/10.12688/wellcomeopenres.21140.1>

**First published:** 19 Mar 2024, 9:150 <https://doi.org/10.12688/wellcomeopenres.21140.1>

## Species taxonomy

Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Paracanthopterygii; Zeiogadaria; Zeariae; Zeiformes; Zeidae; *Zeus*; *Zeus faber* Linnaeus, 1758 (NCBI:txid64108).

## Background

John Dory, *Zeus faber* Linnaeus, 1758, also known as St Peter's fish, is a solitary, demersal marine fish characterised by a laterally compressed body of golden-brown colouration with a black spot on either side and long dorsal spines (Wheeler, 1978). It is widely distributed in the eastern Atlantic, Mediterranean, Pacific and Indian Oceans, and along the entire West African coast, typically occupying depths of 0–200 m (Iwamoto, 2015; Maravelias *et al.*, 2007). Recently, it has been recorded for the first time in the Black Sea (Aydm & Karadurmuş, 2023). John Dory has a large, highly protrusible mouth allowing it to predate on relatively large fish which they target using well-developed eyes (Kim *et al.*, 2020; Stergiou & Fourtouni, 1991).

In the eastern Mediterranean, John Dory begins its life feeding on small zooplankton, such as mysids, until it reaches a total length (TL) of approximately 80 mm (Stergiou & Fourtouni, 1991). It gradually switches to small benthopelagic fishes then, as it grows, it predate on larger schooling pelagic teleosts (Kim *et al.*, 2020; Stergiou & Fourtouni, 1991). In Korean coastal waters there is also varying diet composition with size and age (Kim *et al.*, 2020). Off the Portuguese coast however, there is no prey switching from juvenile to adult life stages (Silva, 1999). Predominantly, evidence suggests it is an opportunistic feeder, switching prey depending on food availability and abundance which can vary seasonally and with life stage (Kim *et al.*, 2020). Recently, meso- and microplastics have been found to occur in the gastrointestinal tracts of *Z. faber* in the Mediterranean (Bottari *et al.*, 2019).

*Z. faber* is known to make 'croaking' or 'barking' noises upon capture onboard (Radford *et al.*, 2018). These vocalisations have since been documented *in-situ* in Australia and were found to induce an escape response in conspecifics and heterospecifics such as the Australian Snapper (*Pagurus auratus*), suggesting they make sounds as a territorial display against competitors (Radford *et al.*, 2018).

John Dory is a commercially important species valued for human consumption, meal, and fish oil, as well as being important as a gamefish and in the aquarium trade (Iwamoto, 2015). It has been a prominent species in mixed species trawl fisheries in the British Isles and is a notable by-catch species for trawl gears globally (Dunn, 2001; Iwamoto, 2015). Despite being a highly regarded food fish, the only stock assessment for the British Isles, to our knowledge, was carried out between April 1994 and March 1996 (Dunn, 2001). Dunn describes

landings from commercial fisheries in England and Wales alongside biological samples and catch data from the English Channel. John Dory was most abundant in the south and southwest of the British Isles (Dunn, 2001; Wheeler, 1969). Evidence suggests that the English Channel is a seasonal nursery ground for *Z. faber*: seasonal peaks in landings coincided with the period of recruitment, in this case during quarters three and four, when individuals were just over 1 year old at the transition between juvenile and adult life stages species (Dunn, 2001). Recruitment total length of the species was approximately 23 cm TL. Most commercial landings were in the range 23–29 cm TL with a maximum observed TL of 59 cm. Mean TL of first maturity was approximately 26 cm for males and 34.5 cm for females. The global conservation status of this species was last assessed in 2013, deemed 'Data Deficient' on the IUCN Red List (Iwamoto, 2015), which reflects the lack of historical data and biological information for this species (Dunn, 2001). Because of this, stock status and fishing pressure is uncertain for *Z. faber*.

In the Northeast Atlantic (FAO area 27), consuming John Dory may prove harmful to human health, as mercury levels of  $0.68 \pm 0.07 \mu\text{g g}^{-1}$  exceed the maximum limit deemed safe for human consumption ( $0.5 \mu\text{g g}^{-1}$ ) (Vieira *et al.*, 2021). John Dory imported from Senegal and sold in Turkish fish markets, as well as individuals caught around Turkey, have also been found to carry larvae of the nematode *Anisakis pegreffii*, with potential implications for human health if consumed (Pekmezci, 2019; Yardimci *et al.*, 2014).

Molecular investigation of this species has shown significant genetic differentiation (7.44%) between clades in the North Atlantic/Mediterranean region and Australasia, indicating the possibility that they have speciated (Ward *et al.*, 2008). The first genome of *Z. faber* was generated in 2016 for a study that suggests immune-related genes play an important role in teleost evolution and speciation (Malmström *et al.*, 2016). Here we present the second published genome of John Dory, collected and sequenced as part of the Darwin Tree of Life project (Blaxter *et al.*, 2022). This dataset will be important for furthering our understanding of teleost pathology, immunology, evolution and phylogenetics (Malmström *et al.*, 2016; Ward *et al.*, 2008).

## Genome sequence report

The genome was sequenced from an individual *Zeus faber* (Figure 1) collected from Bigbury Bay, UK (50.27, -3.97). A total of 43-fold coverage in Pacific Biosciences single-molecule HiFi long reads was generated. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data. Manual assembly curation corrected 9 missing joins or mis-joins, reducing the scaffold number by 1.04%.

The final assembly has a total length of 804.7 Mb in 190 sequence scaffolds with a scaffold N50 of 34.5 Mb (Table 1). The snail plot in Figure 2 provides a summary of the assembly statistics, while the distribution of assembly scaffolds on GC proportion and coverage is shown in Figure 3. The cumulative



**Figure 1.** Photograph of the *Zeus faber* (fZeuFab8) specimen used for genome sequencing.

assembly plot in Figure 4 shows curves for subsets of scaffolds assigned to different phyla. Most (97.08%) of the assembly sequence was assigned to 22 chromosomal-level scaffolds. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size (Figure 5; Table 2). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.

The estimated Quality Value (QV) of the final assembly is 52.6 with  $k$ -mer completeness of 99.98%, and the assembly has a BUSCO v5.3.2 completeness of 96.4% (single = 94.9%, duplicated = 1.4%), using the actinopterygii\_odb10 reference set ( $n = 3,640$ ).


**Table 1.** Genome data for *Zeus faber*, fZeuFab8.1.

Project accession data		
Assembly identifier	fZeuFab8.1	
Species	<i>Zeus faber</i>	
Specimen	fZeuFab8	
NCBI taxonomy ID	64108	
BioProject	PRJEB63619	
BioSample ID	SAMEA111562156	
Isolate information	fZeuFab8 (DNA, Hi-C and RNA sequencing)	
Assembly metrics*		Benchmark
Consensus quality (QV)	52.6	$\geq 50$
$k$ -mer completeness	99.98%	$\geq 95\%$
BUSCO**	C:96.4%[S:94.9%,D:1.4%], F:1.4%,M:2.2%,n:3,640	$C \geq 95\%$
Percentage of assembly mapped to chromosomes	97.08%	$\geq 95\%$
Sex chromosomes	None	localised homologous pairs
Organelles	Mitochondrial genome: 16.72 kb	complete single alleles
Raw data accessions		
PacificBiosciences SEQUEL II	ERR11641070, ERR11641069	
Hi-C Illumina	ERR11641144, ERR11641145	
PolyA RNA-Seq Illumina	ERR11641143	
Genome assembly		
Assembly accession	GCA_960531495.1	


Accession of alternate haplotype	GCA_960530785.1
Span (Mb)	804.7
Number of contigs	1,078
Contig N50 length (Mb)	1.4
Number of scaffolds	190
Scaffold N50 length (Mb)	34.5
Longest scaffold (Mb)	65.76

\* Assembly metric benchmarks are adapted from column VGP-2020 of "Table 1: Proposed standards and metrics for defining genome assembly quality" from [Rhie et al. \(2021\)](#).

\*\* BUSCO scores based on the actinopterygii\_odb10 BUSCO set using version 5.3.2. C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at [https://blobtoolkit.genomehubs.org/view/fZeuFab8\\_1/dataset/fZeuFab8\\_1/busco](https://blobtoolkit.genomehubs.org/view/fZeuFab8_1/dataset/fZeuFab8_1/busco).



**Figure 2. Genome assembly of *Zeus faber*, fZeuFab8.1: metrics.** The BlobToolKit snail plot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 804,731,948 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (65,762,550 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (34,476,449 and 28,869,016 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the actinopterygii\_odb10 set is shown in the top right. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/fZeuFab8\\_1/dataset/fZeuFab8\\_1/snail](https://blobtoolkit.genomehubs.org/view/fZeuFab8_1/dataset/fZeuFab8_1/snail).



**Figure 3. Genome assembly of *Zeus faber*, fZeuFab8.1: BlobToolKit GC-coverage plot.** Sequences are coloured by phylum. Circles are sized in proportion to sequence length. Histograms show the distribution of sequence length sum along each axis. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/fZeuFab8\\_1/dataset/fZeuFab8\\_1/blob](https://blobtoolkit.genomehubs.org/view/fZeuFab8_1/dataset/fZeuFab8_1/blob).

Metadata for specimens, barcode results, spectra estimates, sequencing runs, contaminants and pre-curation assembly statistics are given at <https://links.tol.sanger.ac.uk/species/64108>.


## Methods

### Sample acquisition and nucleic acid extraction

A *Zeus faber* specimen (specimen ID MBA-211116-004A, ToLID fZeuFab8) was collected from Bigbury Bay, UK (latitude 50.27, longitude -3.97) on 2021-11-16 using an otter trawl deployed from the RV Sepia. The collectors were Patrick Adkins, Joanna Harley, Rachel Brittain, Kesella Scott-Somme (all Marine Biological Association) and identified by Rachel Brittain, and then preserved in liquid nitrogen. The fish died as part of a trawl attached to another project and

was opportunistically taken and dissected by the DToL team who were also on board the Sepia that day.

The workflow for high molecular weight (HMW) DNA extraction at the Wellcome Sanger Institute (WSI) includes a sequence of core procedures: sample preparation; sample homogenisation, DNA extraction, fragmentation, and clean-up. In sample preparation, the fZeuFab8 sample was weighed and dissected on dry ice (Jay *et al.*, 2023). Tissue was homogenised using a PowerMasher II tissue disruptor (Denton *et al.*, 2023a). HMW DNA was extracted in the WSI Scientific Operations core using the Automated MagAttract v2 protocol (Oatley *et al.*, 2023). The DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor



**Figure 4. Genome assembly of *Zeus faber*, fZeuFab8.1: BlobToolKit cumulative sequence plot.** The grey line shows cumulative length for all sequences. Coloured lines show cumulative lengths of sequences assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/fZeuFab8\\_1/dataset/fZeuFab8\\_1/cumulative](https://blobtoolkit.genomehubs.org/view/fZeuFab8_1/dataset/fZeuFab8_1/cumulative).

3 system with speed setting 31 (Bates *et al.*, 2023). Sheared DNA was purified by solid-phase reversible immobilisation (Strickland *et al.*, 2023): in brief, the method employs a 1.8X ratio of AMPure PB beads to sample to eliminate shorter fragments and concentrate the DNA. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system.


RNA was extracted from tissue of fZeuFab8 in the Tree of Life Laboratory at the WSI using the RNA Extraction: Automated MagMax™ mirVana protocol (do Amaral *et al.*, 2023). The RNA concentration was assessed using a Nanodrop spectrophotometer and a Qubit Fluorometer using the Qubit RNA Broad-Range Assay kit. Analysis of the

integrity of the RNA was done using the Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay.

Protocols developed by the WSI Tree of Life laboratory are publicly available on protocols.io (Denton *et al.*, 2023b).

### Sequencing

Pacific Biosciences HiFi circular consensus DNA sequencing libraries were constructed according to the manufacturers' instructions. Poly(A) RNA-Seq libraries were constructed using the NEB Ultra II RNA Library Prep kit. DNA and RNA sequencing was performed by the Scientific Operations core at the WSI on Pacific Biosciences SEQUEL II (HiFi) and Illumina NovaSeq 6000 (RNA-Seq) instruments. Hi-C data were also generated from tissue of fZeuFab8 using the Arima2 kit and sequenced on the Illumina NovaSeq 6000, Illumina NovaSeq 6000 instrument.



**Figure 5. Genome assembly of *Zeus faber*, fZeuFab8.1: Hi-C contact map of the fZeuFab8.1 assembly, visualised using HiGlass.** Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/l/?d=ODHdK-fnRfy3tLmo69JGwQ>.

**Table 2. Chromosomal pseudomolecules in the genome assembly of *Zeus faber*, fZeuFab8.**

INSDC accession	Chromosome	Length (Mb)	GC%
OY482845.1	1	65.76	43.5
OY482846.1	2	52.95	43.5
OY482847.1	3	40.01	43.5
OY482848.1	4	38.03	43.5
OY482849.1	5	37.98	44.0
OY482850.1	6	36.47	44.5
OY482851.1	7	36.1	44.0
OY482852.1	8	35.01	44.0
OY482853.1	9	34.68	44.5
OY482854.1	10	34.48	44.0
OY482855.1	11	33.96	43.5
OY482856.1	12	33.1	44.0
OY482857.1	13	32.89	44.5
OY482858.1	14	32.3	43.5
OY482859.1	15	32.12	44.5
OY482860.1	16	31.67	44.5

INSDC accession	Chromosome	Length (Mb)	GC%
OY482861.1	17	31.65	42.5
OY482862.1	18	31.31	44.5
OY482863.1	19	29.26	44.0
OY482864.1	20	28.87	44.5
OY482865.1	21	27.29	45.0
OY482866.1	22	25.41	44.0
OY482867.1	MT	0.02	42.5

#### Genome assembly, curation and evaluation

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021) and haplotypic duplication was identified and removed with purge\_dups (Guan *et al.*, 2020). The assembly was then scaffolded with Hi-C data (Rao *et al.*, 2014) using YaHS (Zhou *et al.*, 2023). The assembly was checked for contamination and corrected as described previously (Howe *et al.*, 2021). Manual curation was performed using HiGlass (Kerpedjiev *et al.*, 2018) and PretextView (Harry, 2022). The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2023), which runs MitoFinder (Allio *et al.*, 2020) or MITOS (Bernt *et al.*, 2013) and uses these annotations to select the final



**Table 3. Software tools: versions and sources.**

Software tool	Version	Source
BlobToolKit	4.1.7	<a href="https://github.com/blobtoolkit/blobtoolkit">https://github.com/blobtoolkit/blobtoolkit</a>
BUSCO	5.3.2	<a href="https://gitlab.com/ezlab/busco">https://gitlab.com/ezlab/busco</a>
Hifiasm	0.19.5-r587	<a href="https://github.com/chhylp123/hifiasm">https://github.com/chhylp123/hifiasm</a>
HiGlass	1.11.6	<a href="https://github.com/higlass/higlass">https://github.com/higlass/higlass</a>
Merqury	MerquryFK	<a href="https://github.com/thegenemyers/MERQURY.FK">https://github.com/thegenemyers/MERQURY.FK</a>
MitoHiFi	3	<a href="https://github.com/marcelauliano/MitoHiFi">https://github.com/marcelauliano/MitoHiFi</a>
PretextView	0.2	<a href="https://github.com/wtsi-hpag/PretextView">https://github.com/wtsi-hpag/PretextView</a>
purge_dups	1.2.5	<a href="https://github.com/dfguan/purge_dups">https://github.com/dfguan/purge_dups</a>
sanger-tol/genomenote	v1.0	<a href="https://github.com/sanger-tol/genomenote">https://github.com/sanger-tol/genomenote</a>
sanger-tol/readmapping	1.1.0	<a href="https://github.com/sanger-tol/readmapping/tree/1.1.0">https://github.com/sanger-tol/readmapping/tree/1.1.0</a>
YaHS	1.2a.2	<a href="https://github.com/c-zhou/yahs">https://github.com/c-zhou/yahs</a>

mitochondrial contig and to ensure the general quality of the sequence.

A Hi-C map for the final assembly was produced using bwa-mem2 (Vasimuddin *et al.*, 2019) in the Cooler file format (Abdennur & Mirny, 2020). To assess the assembly metrics, the *k*-mer completeness and QV consensus quality values were calculated in Merqury (Rhie *et al.*, 2020). This work was done using Nextflow (Di Tommaso *et al.*, 2017) DSL2 pipelines “sanger-tol/readmapping” (Surana *et al.*, 2023a) and “sanger-tol/genomenote” (Surana *et al.*, 2023b). The genome was analysed within the BlobToolKit environment (Challis *et al.*, 2020) and BUSCO scores (Manni *et al.*, 2021; Simão *et al.*, 2015) were calculated.

Table 3 contains a list of relevant software tool versions and sources.

### Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature

of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

### Data availability

European Nucleotide Archive: *Zeus faber* (John dory). Accession number PRJEB63619; <https://identifiers.org/ena.embl/PRJEB63619> (Wellcome Sanger Institute, 2023). The genome sequence is released openly for reuse. The *Zeus faber* genome sequencing initiative is part of the Darwin Tree of Life (DTOL) project. All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the Ensembl pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in Table 1.

## Author information

Members of the Marine Biological Association Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.8382513>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.4893703>.

Members of the Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team are listed here: <https://doi.org/10.5281/zenodo.10066175>.

Members of Wellcome Sanger Institute Scientific Operations: Sequencing Operations are listed here: <https://doi.org/10.5281/zenodo.10043364>.

Members of the Wellcome Sanger Institute Tree of Life Core Informatics team are listed here: <https://doi.org/10.5281/zenodo.10066637>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.5013541>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

## References

- Abdennur N, Mirny LA: **Cooler: Scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics.* 2020; **36**(1): 311–316. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, et al.: **MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour.* 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Aydin M, Karadurmus U: **First record of the benthopelagic fish John dory *Zeus faber* (Linnaeus, 1758) in the Black Sea coasts of Türkiye.** *Aquat Res.* 2023; **6**(2): 159–165. [Publisher Full Text](#)
- Bates A, Clayton-Lucey I, Howard C: **Sanger Tree of Life HMW DNA Fragmentation: Diagenode Megaruptor<sup>®</sup>3 for LI PacBio.** *protocols.io.* 2023. [Publisher Full Text](#)
- Bernt M, Donath A, Jühling F, et al.: **MITOS: Improved *de novo* metazoan mitochondrial genome annotation.** *Mol Phylogenet Evol.* 2013; **69**(2): 313–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Blaxter M, Mieszekowska N, Di Palma F, et al.: **Sequence locally, think globally: The Darwin Tree of Life Project.** *Proc Natl Acad Sci U S A.* 2022; **119**(4): e2115642118. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bottari T, Savoca S, Mancuso M, et al.: **Plastics occurrence in the gastrointestinal tract of *Zeus faber* and *Lepidopus caudatus* from the Tyrrhenian Sea.** *Mar Pollut Bull.* 2019; **146**: 408–416. [PubMed Abstract](#) | [Publisher Full Text](#)
- Challis R, Richards E, Rajan J, et al.: **BlobToolKit - interactive quality assessment of genome assemblies.** *G3 (Bethesda).* 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, et al.: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm.** *Nat Methods.* 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Denton A, Oatley G, Cornwell C, et al.: **Sanger Tree of Life Sample Homogenisation: PowerMash.** *protocols.io.* 2023a. [Publisher Full Text](#)
- Denton A, Yatsenko H, Jay J, et al.: **Sanger Tree of Life Wet Laboratory Protocol Collection V.1.** *protocols.io.* 2023b. [Publisher Full Text](#)
- Di Tommaso P, Chatzou M, Floden EW, et al.: **Nextflow enables reproducible computational workflows.** *Nat Biotechnol.* 2017; **35**(4): 316–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- do Amaral RJV, Bates A, Denton A, et al.: **Sanger Tree of Life RNA Extraction: Automated MagMax<sup>™</sup> mirVana.** *protocols.io.* 2023. [Publisher Full Text](#)
- Dunn M: **The biology and exploitation of John dory, *Zeus faber* (Linnaeus, 1758) in the waters of England and Wales.** *ICES J Mar Sci.* 2001; **58**(1): 96–105. [Publisher Full Text](#)
- Guan D, McCarthy SA, Wood J, et al.: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics.* 2020; **36**(9): 2896–2898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harry E: **PretextView (Paired REad TEXTure Viewer): A desktop application for viewing pretext contact maps.** 2022; [Accessed 19 October 2022]. [Reference Source](#)
- Howe K, Chow W, Collins J, et al.: **Significantly improving the quality of genome assemblies through curation.** *GigaScience.* Oxford University Press, 2021; **10**(1): g1aa153. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Iwamoto T: ***Zeus faber*, The IUCN Red List of Threatened Species.** 2015; [Accessed 2 February 2024]. [Reference Source](#)
- Jay J, Yatsenko H, Narváez-Gómez JP, et al.: **Sanger Tree of Life Sample Preparation: Triage and Dissection.** *protocols.io.* 2023. [Publisher Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, et al.: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol.* 2018; **19**(1): 125. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kim HJ, Kim HG, Oh CW: **Diet composition and feeding strategy of John Dory, *Zeus faber*, in the coastal waters of Korea.** *J Ecol Environ.* 2020; **44**(1): 8. [Publisher Full Text](#)
- Malmstrøm M, Matschiner M, Tørresen OK, et al.: **Evolution of the immune system influences speciation rates in teleost fishes.** *Nat Genet.* 2016; **48**(10): 1204–1210. [PubMed Abstract](#) | [Publisher Full Text](#)
- Manni M, Berkeley MR, Seppay M, et al.: **BUSCO update: Novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol.* 2021; **38**(10): 4647–4654. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Maravelias CD, Tsitsika EV, Papaconstantinou C: **Seasonal dynamics, environmental preferences and habitat selection of John Dory (*Zeus faber*).** *Estuar Coast Shelf S.* 2007; **72**(4): 703–710. [Publisher Full Text](#)
- Oatley G, Denton A, Howard C: **Sanger Tree of Life HMW DNA Extraction: Automated MagAttract v.2.** *protocols.io.* 2023. [Publisher Full Text](#)
- Pekmezci GZ: **Occurrence of *Anisakis pegreffii* (Nematoda: Anisakidae) Larvae in Imported John Dory (*Zeus faber*) from Senegalese Coast Sold in Turkish Supermarkets.** *Acta Parasitol.* 2019; **64**(3): 582–586. [PubMed Abstract](#) | [Publisher Full Text](#)
- Radford CA, Putland RL, Mensinger AF: **Barking mad: The vocalisation of the John Dory, *Zeus faber*.** *PLoS One.* 2018; **13**(10): e0204647. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rao SSP, Huntley MH, Durand NC, et al.: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell.* 2014; **159**(7): 1665–1680. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, et al.: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature.* 2021; **592**(7856): 737–746. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Rhie A, Walenz BP, Koren S, *et al.*: **Mercury: Reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol.* 2020; **21**(1): 245.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Silva A: **Feeding habits of John Dory, *Zeus faber*, off the Portuguese continental coast.** *J Mar Biol Assoc U K.* 1999; **79**(2): 333–340.

[Publisher Full Text](#)

Simão FA, Waterhouse RM, Ioannidis P, *et al.*: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.**

*Bioinformatics.* 2015; **31**(19): 3210–3212.

[PubMed Abstract](#) | [Publisher Full Text](#)

Stergiou KI, Fourtouni H: **Food habits, ontogenetic diet shift and selectivity in *Zeus faber* Linnaeus, 1758.** *J Fish Biol.* 1991; **39**(4): 589–603.

[Publisher Full Text](#)

Strickland M, Cornwell C, Howard C: **Sanger Tree of Life Fragmented DNA clean up: Manual SPRI.** *protocols.io.* 2023.

[Publisher Full Text](#)

Surana P, Muffato M, Qi G: **sanger-tol/readmapping: sanger-tol/readmapping v1.1.0 - Hebridean Black (1.1.0).** *Zenodo.* 2023a.

[Publisher Full Text](#)

Surana P, Muffato M, Sadasivan Baby C: **sanger-tol/genomenote (v1.0.dev).** *Zenodo.* 2023b.

[Publisher Full Text](#)

Uliano-Silva M, Ferreira JGRN, Krashennikova K, *et al.*: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads.** *BMC Bioinformatics.* 2023; **24**(1): 288.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vasimuddin M, Misra S, Li H, *et al.*: **Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems.** In: *2019 IEEE International Parallel and*

*Distributed Processing Symposium (IPDPS).* IEEE, 2019; 314–324.

[Publisher Full Text](#)

Vieira HC, Ramirez MMB, Bordalo MD, *et al.*: **Total and Organic Mercury in Fish from Different Geographical Areas in the North Atlantic Ocean and Health Risk Assessment.** *Expos Health.* 2021; **13**(3): 361–373.

[Publisher Full Text](#)

Ward R, Costa F, Holmes B, *et al.*: **DNA barcoding of shared fish species from the North Atlantic and Australasia: minimal divergence for most taxa, but *Zeus faber* and *Lepidopus caudatus* each probably constitute two species.** *Aquat Biol.* 2008; **3**(1): 71–78.

[Publisher Full Text](#)

Wellcome Sanger Institute: **The genome sequence of the John Dory, *Zeus faber* Linnaeus, 1758.** European Nucleotide Archive, [dataset], accession number PRJEB63619, 2023.

Wheeler A: **The Fishes of the British Isles and North-West Europe.** London, UK: Macmillan, 1969.

[Reference Source](#)

Wheeler A: **Key to the Fishes of Northern Europe: a guide to the identification of more than 350 species.** London: Frederick Warne (Publishers) Ltd, 1978.

[Reference Source](#)

Yardimci B, Pekmezci GZ, Onuk EE: **Pathology and molecular identification of *Anisakis pegreffii* (Nematoda: Anisakidae) infection in the John Dory, *Zeus faber* (Linnaeus, 1758) caught in Mediterranean Sea.** *Ankara Univ Vet Fak.* 2014; **61**(3): 233–236.

[Publisher Full Text](#)

Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool.** *Bioinformatics.* 2023; **39**(1): btac808.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

# Open Peer Review

Current Peer Review Status:   

---

## Version 1

Reviewer Report 08 August 2024

<https://doi.org/10.21956/wellcomeopenres.23384.r83999>

© 2024 Mazloumi N. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Nastaran Mazloumi** 

University of Tasmania Institute for Marine and Antarctic Studies, Hobart, Tasmania, Australia

### Title:

The chosen title “The genome sequence of the John Dory, *Zeus faber* Linnaeus, 1758” lacks clarity and intention. You need to choose a better title to define why you have done this study and what the genome sequestration is going to do!

### Background:

In the background section, this paragraph below is disjointed and very hard to follow:

“John Dory is a commercially important species valued for human consumption, meal, and fish oil, as well as being important as a gamefish and in the aquarium trade (Iwamoto, 2015). It has been a prominent species in mixed species trawl fisheries in the British Isles and is a notable by-catch species for trawl gears globally (Dunn, 2001; Iwamoto, 2015). Despite being a highly regarded food fish, the only stock assessment for the British Isles, to our knowledge, was carried out between April 1994 and March 1996 (Dunn, 2001). Dunn describes landings from commercial fisheries in England and Wales alongside biological samples and catch data from the English Channel. John Dory was most abundant in the south and southwest of the British Isles (Dunn, 2001; Wheeler, 1969). Evidence suggests that the English Channel is a seasonal nursery ground for *Z. faber*: seasonal peaks in landings coincided with the period of recruitment, in this case during quarters three and four, when individuals were just over 1 year old at the transition between juvenile and adult life stages species (Dunn, 2001). Recruitment total length of the species was approximately 23 cm TL. Most commercial landings were in the range 23–29 cm TL with a maximum observed TL of 59 cm. Mean TL of first maturity was approximately 26 cm for males and 34.5 cm for females. The global conservation status of this species was last assessed in 2013, deemed ‘Data Deficient’ on the IUCN Red List (Iwamoto, 2015), which reflects the lack of historical data and biological information for this species (Dunn, 2001). Because of this, stock status and fishing pressure is uncertain for *Z. faber*”.

This paragraph provides valuable information for reader but it significantly lacks clarity and confinement. For instance, human health issues with over the threshold elements in the flesh is one argument, data deficient stock assessment is another argument. Similarly, the genetically distinguished populations in different parts of the world that is discussed in the following paragraph, is another statement: "Molecular investigation of this species has shown significant genetic differentiation (7.44%) between clades in the North Atlantic/Mediterranean region and Australasia, indicating the possibility that they have speciated (Ward et al., 2008). The first genome of *Z. faber* was generated in 2016 for a study that suggests immune-related genes play an important role in teleost evolution and speciation (Malmstrøm et al., 2016). Here we present the second published genome of John Dory, collected and sequenced as part of the Darwin Tree of Life project (Blaxter et al., 2022). This dataset will be important for furthering our understanding of teleost pathology, immunology, evolution and phylogenetics (Malmstrøm et al., 2016; Ward et al., 2008)" - this confuse the reader as to why this study is important and why you made efforts to sequence genomes for John Dory.

Background section, requires substantial revision in term of clarity and confinement and more importantly the message the author want to communicate with the reader as to why they have done this study and what gaps this is going to fill and how it serves the intention of the study. It needs to be specifically clarified as to whether this is for seafood industry and human health matter, or conservation matter? Or even management of the stock? What is the main intention here. without this valuable bit, it is not recommended to jump into the methodology and analysis.

**Summary:**

Despite my encouragement for scientists to publish their work, I must admit that this particular research lacks clarity and conciseness. Therefore, I have no choice but to recommend rejection and suggest resubmission after thorough revisions, including expanding the research and writing in a more concise manner.

**Is the rationale for creating the dataset(s) clearly described?**

No

**Are the protocols appropriate and is the work technically sound?**

Partly

**Are sufficient details of methods and materials provided to allow replication by others?**

No

**Are the datasets clearly presented in a useable and accessible format?**

No

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Fisheries sciences, Biological Sciences, Climate Change, Fish Stock Assessment, influences of environmental and anthropogenic impacts on marine resources

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for**

reasons outlined above.

Reviewer Report 14 June 2024

<https://doi.org/10.21956/wellcomeopenres.23384.r80568>

© 2024 Rodriguez-Ezpeleta N. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Naiara Rodriguez-Ezpeleta** 

AZTI Basque Research and Technology Alliance, Bizkaia, Spain

This manuscript presents the genome sequence of a fish species (John Dory) that is in general poorly studied and for which conservation strategies can not be appropriately defined. Considering that understanding the genetic connectivity is critical for defining management units, and that full genomes facilitate the analysis and interpretation of population genomic data, this study is considered important for improving the conservation of the John Dory. The methods are well described and the results are pertinent. I have only one comment that I think the authors can address easily:

- In the introduction they should better explain what is the knowledge of this species in terms of how many populations exist and potential subspecies. The data of "significant genetic differentiation (7.44%) between clades in the North Atlantic/Mediterranean region and Australasia" seem to be taken from the abstract of a cited paper, but there is not much information here to understand what is the main issue that could be resolved with the help of a full genome sequence.

**Is the rationale for creating the dataset(s) clearly described?**

Yes

**Are the protocols appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and materials provided to allow replication by others?**

Yes

**Are the datasets clearly presented in a useable and accessible format?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Population genomics, molecular evolution, high-throughput sequencing data analysis

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 10 May 2024

<https://doi.org/10.21956/wellcomeopenres.23384.r79083>

© 2024 Burgess S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



### Shawn M. Burgess

National Human Genome Research Institute, Bethesda, USA

Adkins and colleagues have submitted a manuscript describing the genomic assembly of a John Dory fish (*Zeus faber*). The sequencing was performed using PacBio HiFi and Hi-C data and the assembly using hifiasm and manual curation. The resulting assembly exceeds general benchmarks for high-quality assemblies including scoring for BUSCO, k-mer completeness, and percentage of sequence mapped to chromosomes. All raw and analyzed datasets are deposited in publicly available repositories. The data are clear, the approaches adequately documented. I have no substantial objections to any of the data presented. It should be a very useful assembly for researchers who are interested in John Dory biology.

#### Is the rationale for creating the dataset(s) clearly described?

Yes

#### Are the protocols appropriate and is the work technically sound?

Yes

#### Are sufficient details of methods and materials provided to allow replication by others?

Yes

#### Are the datasets clearly presented in a useable and accessible format?

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Zebrafish genetics and genomics. Fish genome assembly.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---