

EDITORIAL

The hydrocarbon pollution crisis: Harnessing the earth hydrocarbon-degrading microbiome

As part of the battle against climate change, the decarbonization of human activities has been acted in many countries worldwide. Thus, in order to limit the planet warming, it is expected to reduce the combustion of fossil fuels for decreasing drastically the production of greenhouse gas emissions. Although beneficial for reducing carbon dioxide (CO₂) production to fight against climate change, this countermeasure unfortunately does not mean that hydrocarbon pollution is behind us because hydrocarbon pollution has many sources (Duran & Cravo-Laureau, 2016) that will remain. It is estimated that the industrial and petroleum activities, which have already left behind a multitude of hydrocarbon-contaminated sites that still need to be restored, release accidentally between 1.7 and 8.8 million tonnes of oil into the environment each year (Ambaye et al., 2022). The decarbonization is also expected to have a beneficial impact on decreasing the industrial release of hydrocarbons into the environment by reducing oil spill frequency and consequences (Little et al., 2021). In addition to the direct contribution of hydrocarbons resulting from the continued use of fossil fuels, the human activities also generate indirect inputs such as wildfires introducing polycyclic aromatic hydrocarbon (PAH) into the environment (Campos et al., 2019; Paul et al., 2023). Particularly “mega fires,” which burn large forest areas, are becoming more frequent as a consequence of climate change (Bracewell et al., 2023; van Oldenborgh et al., 2021). Of course, the natural sources of hydrocarbon contamination, such as volcanic activities and marine oil seeps, as well as biogenic sources (Duran & Cravo-Laureau, 2016), continuously emit hydrocarbons into the environment. Thus, hydrocarbons last to be of concern for the environment in the future. In order to mitigate the impact of hydrocarbons on the environment, exploiting the hydrocarbon degradation potential that microorganisms have is a challenge to meet for scientists and engineers concerned about hydrocarbon pollution.

Important key knowledge has been gained on microbial hydrocarbon degradation as well as on the ecology of microbial communities inhabiting hydrocarbon-contaminated sites. The hydrocarbon

degradation capacity has been described for a large number of microorganisms from diverse terrestrial and aquatic environments. Several specialist hydrocarbon-degrading microbial taxa have been described and isolated, as for example, the marine obligate hydrocarbonoclastic bacteria (OHCB) observed to bloom during marine oil spills (Yakimov et al., 2007), hydrocarbon-tolerant fungi found in petroleum-contaminated sediment (Álvarez-Barragán et al., 2021), and alkane-degrading specialist populations in soil (Hamamura et al., 2013). The characterization of specialist hydrocarbon-degrading microbial taxa has unveiled the degradation pathways, both aerobic and anaerobic, for a wide range of hydrocarbon compounds from alkanes to PAHs (Chunyan et al., 2023). While gaining knowledge, various bioremediation strategies have been proposed to mitigate the impact of hydrocarbon in contaminated sites (Goñi-Urriza et al., 2013). However, the different existing bioremediation processes for the in situ treatment of hydrocarbon-contaminated sites have to content several technical, environmental, and regulatory hurdles before reaching optimal efficiency and societal acceptability (Boopathy, 2000). Particularly, the release of microorganisms, genetically modified or not, into the environment following a bio-augmentation approach is controversial at both technical and ethical point of views (Lensch et al., 2024). Apart from the regulatory and ethical issues associated with the use of microorganisms for in situ bioremediation, which are not addressed here, there is no clear evidence that the added microorganisms colonize the contaminated site and effectively degrade the hydrocarbons (Wu et al., 2019). The difficulty for the added microorganisms to incorporate well-established microbial community, known as colonization resistance in ecological theory, depends on several factors controlling the microbial community mixing, i.e., microbial community coalescence (Châtillon, Duran, et al., 2023; Rillig et al., 2015). These factors include the microbial communities legacy based on the history of contamination (Hafez et al., 2022), the size of the added community (Rillig et al., 2015), and priority effects (Tucker

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Microbial Biotechnology* published by Applied Microbiology International and John Wiley & Sons Ltd.

& Fukami, 2014). Nevertheless, the current knowledge on hydrocarbon microbial ecology provides crucial information opening the way for the implementation of bioremediation processes based on synthetic biology approaches (de Lorenzo, 2008).

The ecology of microbial communities living in hydrocarbon-contaminated sites has been conceptualized on the basis of experimental and in situ observations. According to the disturbance theory (Allison & Martiny, 2008), the resistance, resilience, and functional redundancy of microbial communities in response to hydrocarbon contamination have been illustrated in several studies (Châtillon, Cébron, et al., 2023; Chronopoulou et al., 2013; Stauffert et al., 2014; Zerebecki et al., 2022). However, the microbial communities have been shown extremely dynamic and interactive in response to hydrocarbon contamination (Head et al., 2006; McGenity et al., 2012), requiring to switch beyond the concepts of resistance, resilience, and redundancy in order to better understand the structure or function relationships (Bissett et al., 2013). The ecological succession, microbial taxa replacement over time, has been observed in many cases after hydrocarbon contamination experimentally (Bordenave, Fourçans, et al., 2004; Bordenave, Jézéquel, et al., 2004; Cerqueda-García et al., 2020) and in situ after an oil spill (Kimes et al., 2014; Péquin et al., 2022). Such ecological succession relies on the fact that hydrocarbon degradation is effected by a consortium of microorganisms, not all of which are directly involved in the degradation process. The microbial interactions have an extremely important ecological role in hydrocarbon degradation; hydrocarbon consumers receiving beneficial products (vitamins, EPS, metals, etc.) from associated microbes in return for their detoxifying activity (Louati et al., 2013; McGenity, 2014); even some associated microbes can also provide biosurfactants increasing hydrocarbon bioavailability that facilitates their degradation (Brito et al., 2009; McKew et al., 2007; Schweitzer et al., 2022). The microbial interactions also involve co-metabolism and synergistic processes in which the degradation of hydrocarbons is enhanced by the use of a co-substrate (Chen & Aitken, 1999; García-Rivero & Peralta-Pérez, 2008) or terminal electron acceptors (oxygen, nitrogen, sulphur) produced by other (micro) organisms (McGenity, 2014). Also, the microbial interactions allow microorganisms to colonize novel niches as illustrated by the inter-kingdom interaction between fungi and bacteria (Álvarez-Barragán et al., 2022), in which bacteria use the hyphosphere as 'fungal highway' for dispersal (Álvarez-Barragán et al., 2023). Understanding the network of processes within microbial assemblages and the underlying mechanisms from which they arise is of paramount importance to achieve efficient bioremediation practices for

oil-polluted sites. It is accepted that trade-offs drive microbial assemblages (Østman et al., 2014), particularly niche specialization has been recognized as key process in hydrocarbon degradation, each degradation step being performed by different microbial functional groups (Dalby et al., 2008). The establishment of ecotypes, step in niche specialization (Gushgari-Doyle et al., 2022), has been demonstrated by the emergence of ecotypes adapted to either hydrocarbon structure (Kleindienst et al., 2015; McKew et al., 2007), temperature (Bargiela et al., 2015) or oxygen conditions (Terrisse et al., 2017). The genomic evolution conducting to niche specialization involves several adaptation mechanisms such as switch of integron gene cassettes (Abella et al., 2015; Huang et al., 2009) and horizontal gene transfer (Shahi et al., 2017); mechanisms that have been proposed to spread key degradation genes to harness microbial communities for hydrocarbon degradation in a synthetic biology approach (French et al., 2020).

In recent years, the concept of synthetic biology has been extended to microbial community (Borchert et al., 2021; De Roy et al., 2014) and even to the ecosystem (Hammond et al., 2023). The design of engineered microbial community, crucial step in the synthetic biology approaches, relies on microbial ecology principles and rules governing microbial assemblage processes such as community coalescence, habitat filtering, taxa replacement and turnover, and priority effects (Bernstein, 2019) for improving not only the biodegradation capacity but also favouring the colonization (Rocca et al., 2021; Ruan et al., 2024). Several strategies have been proposed to design synthetic microbial communities (SynComs) either by enrichment techniques resulting in simplified communities, defined as top-down approaches, or reconstruction of microbial consortia by assembling bacterial strains, known as bottom-up approaches (Bernstein, 2019; De Roy et al., 2014; Hu et al., 2022). Anent the top-down approach, the current momentum for obtaining engineered microbiomes owning the desired functions can implement successive sub-culturing under defined conditions (Li et al., 2023) applying as well high-throughput techniques (Duran et al., 2022), or artificial selection applying sequential breeding (Swenson et al., 2000), or directed evolution subjecting community to perturbation cycles (Sánchez et al., 2021). The construction of SynComs, synthetic microbial assemblages, following a bottom-up approach combines isolated microorganisms, genetically modified or not. The current efforts to build SynComs are based on the metabolic division of labour (MDOL) concept, the different steps of a metabolic pathway being performed by distinct populations in order to reduce the burden for each population (Tsoi et al., 2018). The design of SynComs also consider functional traits and potential niche

occupancy (Jing et al., 2024), ecological coexistence (Chen et al., 2023), as well as microbial species interactions within the microbiome and outside the microbiome with organisms in its environment (Leggieri et al., 2021), that will benefit hydrocarbon bioremediation treatments. In order to direct the construction of microbial consortia, guide their transfer into the environment and anticipate their effects once released, SynComs draws largely on the recent developments of powerful computational genomic analyses (Jing et al., 2024), mathematical models (Tsoi et al., 2018), machine learning (Ghannam & Techtmann, 2021), and artificial intelligence (Patowary et al., 2023). The engineered microbial consortia have not only promising potential for developing technologies to reclaim hydrocarbon-contaminated ecosystems (Yang et al., 2021), but they have also been proposed as tool for microbiome rescue to restore ecological stability in damaged ecosystems (de Lorenzo, 2018; Shade, 2023). Several stratagems and formulations have been proposed to deliver microorganisms and microbial consortia into contaminated sites to ensure their colonization and maintenance overcoming colonization resistance (Das & Chandran, 2011), which have been conceptualized as 'Environmental Galenics' (de Lorenzo, 2022). Most of them involve (bio)surfactants, foams, encapsulation, and immobilization of cells using supports such as biochar or nanoparticles (Mrozik & Piotrowska-Seget, 2010; Patowary et al., 2023). Even, engineered horizontal gene transfer has been proposed especially for the treatment of large contaminated areas (de Lorenzo, 2022). Albeit the progress made in building artificial microbial communities, whether synthetic or not, to improve both degradation and colonization capacities for efficient bioremediation, some technical hurdles still need to be overcome to achieve effective bioremediation. Particularly, it is still required to improve the stability of microbial community by strengthening microbial interactions (Xu & Yu, 2021), as well as establish rules for a safe introduction of engineered microorganism into the environment (Wang & Zhang, 2019). The most glaring shortcoming is the lack of effective control of the artificial microbial communities in time, space, and composition for which promising strategies have been proposed relying, respectively, on signalling and adhesive systems, and symbiotic relationships (Grandel et al., 2021). Likewise, although it has been suggested that gene spread via horizontal transfer is self-limited until fitness advantages over selection pressure persist (de Lorenzo, 2010), control systems should also consider the resilience of the local microbial community with respect to ecosystem biogeochemical cycling functions (French et al., 2020).

The evaluation of the performance of the bioremediation treatment requires robust monitoring methods

to determine the efficiency of hydrocarbon degradation and removal, the effectiveness in reducing the toxicity, as well as the innocuousness of the applied technologies for human health and the environment. Thus, it is not enough to just assess hydrocarbon content in the contaminated environment by analytical chemistry methods but the monitoring toolbox must also include a battery of (eco)-toxicity tests in order to appraise the ecological status (Schuijt et al., 2021). In the recent years, the development of microbial biomarkers for reporting the ecological status has gained momentum leading to the elaboration of microbial indexes such as microgAMBI (Aylagas et al., 2017), which adds the panoply of test. However, holistic approaches in line with the 'One Health' concept are recommended (Burke et al., 2017) for a comprehensive evaluation of the treatments' performances. In a world driven by economics, the implementation of bioremediation treatment for the rehabilitation of hydrocarbon-contaminated sites, like any damaged site whatever the cause, must include collaboration with socio-economic players within translational ecology framework (Enquist et al., 2017) and promote circular economy (Jagaba et al., 2022). The development of such initiatives will clearly demonstrate the financial benefits generated by taking care of environmental health, thus paving the way for the upturn of environmental engineering harnessing the Earth hydrocarbon-degrading microbiome for reclaiming hydrocarbon-contaminated sites.

AUTHOR CONTRIBUTIONS

Robert Duran: Conceptualization; investigation; writing – original draft; writing – review and editing.

Cristiana Cravo-Laureau: Conceptualization; writing – review and editing.

FUNDING INFORMATION

The research work of the authors was funded by the MAEWA project (H2020-EU-PRIMA programme, PRIMA-P012-0004-01) supported by the French National Research Agency (ANR-23-P012-0004-01), the BIOMIC project (Interreg Sudoe Programme, European Regional Development Fund SOE4/P1/F0993), and the AQUASALT project (H2020-EU-ERANET-MED programme, NMED-0003-01) supported by the French National Research Agency (ANR-17-NMED-0003-01).

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests.

Robert Duran 

Cristiana Cravo-Laureau 

*Universite de Pau et Des Pays de l'Adour, E2S
UPPA, CNRS, IPREM, Pau, France*

Correspondence

Robert Duran, Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Pau, France.
Email: robert.duran@univ-pau.fr

ORCID

Robert Duran  <https://orcid.org/0000-0002-2351-8553>

Cristiana Cravo-Laureau  <https://orcid.org/0000-0003-2011-813X>

REFERENCES

- Abella, J., Bielen, A., Huang, L., Delmont, T.O., Vujaklija, D., Duran, R. et al. (2015) Integron diversity in marine environments. *Environmental Science and Pollution Research*, 22, 15360–15369.
- Allison, S.D. & Martiny, J.B.H. (2008) Resistance, resilience, and redundancy in microbial communities. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 11512–11519.
- Álvarez-Barragán, J., Cravo-Laureau, C. & Duran, R. (2022) Fungal-bacterial network in PAH-contaminated coastal marine sediment. *Environmental Science and Pollution Research*, 29, 72718–72728.
- Álvarez-Barragán, J., Cravo-Laureau, C., Wick, L.Y. & Duran, R. (2021) Fungi in PAH-contaminated marine sediments: cultivable diversity and tolerance capacity towards PAH. *Marine Pollution Bulletin*, 164, 112082.
- Álvarez-Barragán, J., Cravo-Laureau, C., Xiong, B., Wick, L.Y. & Duran, R. (2023) Marine fungi select and transport aerobic and anaerobic bacterial populations from polycyclic aromatic hydrocarbon-contaminated sediments. *MBio*, 14, e0276122.
- Ambaye, T.G., Chebbi, A., Formicola, F., Prasad, S., Gomez, F.H., Franzetti, A. et al. (2022) Remediation of soil polluted with petroleum hydrocarbons and its reuse for agriculture: recent progress, challenges, and perspectives. *Chemosphere*, 293, 133572.
- Aylagas, E., Borja, Á., Tangherlini, M., Dell'Anno, A., Corinaldesi, C., Michell, C.T. et al. (2017) A bacterial community-based index to assess the ecological status of estuarine and coastal environments. *Marine Pollution Bulletin*, 114, 679–688.
- Bargiela, R., Mapelli, F., Rojo, D., Chouaia, B., Tornés, J., Borin, S. et al. (2015) Bacterial population and biodegradation potential in chronically crude oil-contaminated marine sediments are strongly linked to temperature. *Scientific Reports*, 5, 11651.
- Bernstein, H.C. (2019) Reconciling ecological and engineering design principles for building microbiomes. *mSystems*, 4, e00106-19. Available from: <https://doi.org/10.1128/msystems.00106-19>
- Bissett, A., Brown, M.V., Siciliano, S.D. & Thrall, P.H. (2013) Microbial community responses to anthropogenically induced environmental change: towards a systems approach. *Ecology Letters*, 16, 128–139.
- Boopathy, R. (2000) Factors limiting bioremediation technologies. *Bioresource Technology*, 74, 63–67.
- Borchert, E., Hammerschmidt, K., Hentschel, U. & Deines, P. (2021) Enhancing microbial pollutant degradation by integrating evolutionary principles with environmental biotechnology. *Trends in Microbiology*, 29, 908–918.
- Bordenave, S., Fourçans, A., Blanchard, S., Goñi, M.S., Caumette, P. & Duran, R. (2004) Structure and functional analyses of bacterial communities changes in microbial mats following petroleum exposure. *Ophelia*, 58, 195–203.
- Bordenave, S., Jézéquel, R., Fourçans, A., Budzinski, H., Merlin, F.X., Fourel, T. et al. (2004) Degradation of the "Erika" oil. *Aquatic Living Resources*, 17, 261–267.
- Bracewell, S.A., Barros, T.L., Mayer-Pinto, M., Dafforn, K.A., Simpson, S.L. & Johnston, E.L. (2023) Contaminant pulse following wildfire is associated with shifts in estuarine benthic communities. *Environmental Pollution*, 316, 120533.
- Brito, E.M.S., Duran, R., Guyoneaud, R., Goni-Urriza, M., Garcia de Oteyza, T., Crapez, M.A.C. et al. (2009) A case study of in situ oil contamination in a mangrove swamp (Rio De Janeiro, Brazil). *Marine Pollution Bulletin*, 58, 418–423.
- Burke, T.A., Cascio, W.E., Costa, D.L., Deener, K., Fontaine, T.D., Fulk, F.A. et al. (2017) Rethinking environmental protection: meeting the challenges of a changing world. *Environmental Health Perspectives*, 125, A43–A49.
- Campos, I., Abrantes, N., Pereira, P., Micaelo, A.C., Vale, C. & Keizer, J.J. (2019) Forest fires as potential triggers for production and mobilization of polycyclic aromatic hydrocarbons to the terrestrial ecosystem. *Land Degradation & Development*, 30, 2360–2370.
- Cerqueda-García, D., García-Maldonado, J.Q., Aguirre-Macedo, L. & García-Cruz, U. (2020) A succession of marine bacterial communities in batch reactor experiments during the degradation of five different petroleum types. *Marine Pollution Bulletin*, 150, 110775.
- Châtillon, E., Cébron, A., Rigal, F., Cagnon, C., Lorgeoux, C., Faure, P. et al. (2023) Functional redundancy in response to runoff input upholds microbial community in hydrocarbon-contaminated land-sea continuum. *Environmental Pollution*, 335, 122330.
- Châtillon, E., Duran, R., Rigal, F., Cagnon, C., Cébron, A. & Cravo-Laureau, C. (2023) New insights into microbial community coalescence in the land-sea continuum. *Microbiological Research*, 267, 127259.
- Chen, S.-H. & Aitken, M.D. (1999) Salicylate stimulates the degradation of high-molecular weight polycyclic aromatic hydrocarbons by *Pseudomonas saccharophila* P15. *Environmental Science & Technology*, 33, 435–439.
- Chen, X., Zhou, X., Geng, P., Zeng, Y., Hu, F., Sun, P. et al. (2023) Advancing biodegradation of petroleum contaminants by indigenous microbial consortia through assembly strategy innovations. *Chemical Engineering Journal*, 475, 146142.
- Chronopoulou, P.-M., Fahy, A., Coulon, F., Paisse, S., Goni-Urriza, M., Peperzak, L. et al. (2013) Impact of a simulated oil spill on benthic phototrophs and nitrogen-fixing bacteria in mudflat mesocosms. *Environmental Microbiology*, 15, 242–252.
- Chunyan, X., Qaria, M.A., Qi, X. & Daochen, Z. (2023) The role of microorganisms in petroleum degradation: current development and prospects. *Science of the Total Environment*, 865, 161112.
- Dalby, A.P., Kormas, K.A., Christaki, U. & Karayanni, H. (2008) Cosmopolitan heterotrophic microeukaryotes are active bacterial grazers in experimental oil-polluted systems. *Environmental Microbiology*, 10, 47–56.
- Das N, Chandran P (2011) Microbial degradation of petroleum hydrocarbon contaminants: an overview. *Biotechnology Research International*, 2011, 941810.
- de Lorenzo, V. (2008) Systems biology approaches to bioremediation. *Current Opinion in Biotechnology*, 19, 579–589.
- de Lorenzo, V. (2010) Environmental biosafety in the age of synthetic biology: do we really need a radical new approach? Environmental fates of microorganisms bearing synthetic genomes could be predicted from previous data on traditionally engineered bacteria for in situ bioremediation. *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology*, 32, 926–931.
- de Lorenzo, V. (2018) Environmental microbiology to the rescue of planet earth. *Environmental Microbiology*, 20, 1910–1916.

- de Lorenzo, V. (2022) Environmental Galenics: large-scale fortification of extant microbiomes with engineered bioremediation agents. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 377, 20210395.
- De Roy, K., Marzorati, M., Van den Abbeele, P., Van de Wiele, T. & Boon, N. (2014) Synthetic microbial ecosystems: an exciting tool to understand and apply microbial communities. *Environmental Microbiology*, 16, 1472–1481.
- Duran, C., Zhang, S., Yang, C., Falco, M.L., Cravo-Laureau, C., Suzuki-Minakuchi, C. et al. (2022) Low-cost gel-filled microwell array device for screening marine microbial consortium. *Frontiers in Microbiology*, 13, 1031439.
- Duran, R. & Cravo-Laureau, C. (2016) Role of environmental factors and microorganisms in determining the fate of polycyclic aromatic hydrocarbons in the marine environment. *FEMS Microbiology Reviews*, 40, 814–830.
- Enquist, C.A., Jackson, S.T., Garfin, G.M., Davis, F.W., Gerber, L.R., Littell, J.A. et al. (2017) Foundations of translational ecology. *Frontiers in Ecology and the Environment*, 15, 541–550.
- French, K.E., Zhou, Z. & Terry, N. (2020) Horizontal 'gene drives' harness indigenous bacteria for bioremediation. *Scientific Reports*, 10, 15091.
- García-Rivero, M. & Peralta-Pérez, M. (2008) Cometabolism in the biodegradation of hydrocarbons. *Revista Mexicana de Ingeniería Química*, 7, 1–12.
- Ghannam, R.B. & Techtman, S.M. (2021) Machine learning applications in microbial ecology, human microbiome studies, and environmental monitoring. *Computational and Structural Biotechnology Journal*, 19, 1092–1107.
- Goñi-Urriza, M.S., Cravo-Laureau, C. & Duran, R. (2013) Microbial bioremediation of aquatic environments. In: Féraud, J.-F. & Blaise, C. (Eds.) *Encyclopedia of aquatic ecotoxicology*. Netherlands, Dordrecht: Springer, pp. 709–720.
- Grandel, N.E., Reyes Gamas, K. & Bennett, M.R. (2021) Control of synthetic microbial consortia in time, space, and composition. *Trends in Microbiology*, 29, 1095–1105.
- Gushgari-Doyle, S., Lui, L.M., Nielsen, T.N., Wu, X., Malana, R.G., Hendrickson, A.J. et al. (2022) Genotype to ecotype in niche environments: adaptation of *Arthrobacter* to carbon availability and environmental conditions. *ISME Communications*, 2, 32.
- Hafez, T., Ortiz-Zarragoitia, M., Cagnon, C., Cravo-Laureau, C. & Duran, R. (2022) Legacy and dispersant influence microbial community dynamics in cold seawater contaminated by crude oil water accommodated fractions. *Environmental Research*, 212, 113467.
- Hamamura, N., Ward, D.M. & Inskeep, W.P. (2013) Effects of petroleum mixture types on soil bacterial population dynamics associated with the biodegradation of hydrocarbons in soil environments. *FEMS Microbiology Ecology*, 85, 168–178.
- Hammond, M.P., Kolasa, J. & Fung, P. (2023) Synthetic ecosystems: an emerging opportunity for science and society? *Oikos*, 2023, e09816.
- Head, I.M., Jones, D.M. & Roling, W.F.M. (2006) Marine microorganisms make a meal of oil. *Nature Reviews Microbiology*, 4, 173–182.
- Hu, H., Wang, M., Huang, Y., Xu, Z., Xu, P., Nie, Y. et al. (2022) Guided by the principles of microbiome engineering: accomplishments and perspectives for environmental use. *mLife*, 1, 382–398.
- Huang, L., Cagnon, C., Caumette, P. & Duran, R. (2009) First gene cassettes of Integrins as targets in finding adaptive genes in metagenomes. *Applied and Environmental Microbiology*, 75, 3823–3825.
- Jagaba, A.H., Kutty, S.R.M., Lawal, I.M., Birniwa, A.H., Affam, A.C., Yaro, N.S.A. et al. (2022) Circular economy potential and contributions of petroleum industry sludge utilization to environmental sustainability through engineered processes – a review. *Cleaner and Circular Bioeconomy*, 3, 100029.
- Jing, J., Garbeva, P., Raaijmakers, J.M. & Medema, M.H. (2024) Strategies for tailoring functional microbial synthetic communities. *The ISME Journal*, 18, wrae049.
- Kimes, N.E., Callaghan, A.V., Suflita, J.M. & Morris, P.J. (2014) Microbial transformation of the Deepwater horizon oil spill—past, present, and future perspectives. *Frontiers in Microbiology*, 5, 603.
- Kleindienst, S., Grim, S., Sogin, M., Bracco, A., Crespo-Medina, M. & Joye, S.B. (2015) Diverse, rare microbial taxa responded to the Deepwater horizon deep-sea hydrocarbon plume. *The ISME Journal*, 10, 400–415.
- Leggieri, P.A., Liu, Y., Hayes, M., Connors, B., Seppälä, S., O'Malley, M.A. et al. (2021) Integrating systems and synthetic biology to understand and engineer microbiomes. *Annual Review of Biomedical Engineering*, 23, 169–201.
- Lensch, A., Lindfors, H.A., Duwenig, E., Fleischmann, T., Hjort, C., Kärenlampi, S.O. et al. (2024) Safety aspects of microorganisms deliberately released into the environment. *EFB Bioeconomy Journal*, 4, 100061.
- Li, P., Liang, X., Shi, R., Wang, Y., Han, S. & Zhang, Y. (2023) Unraveling the functional instability of bacterial consortia in crude oil degradation via integrated co-occurrence networks. *Frontiers in Microbiology*, 14, 1270916.
- Little, D.I., Sheppard, S.R.J. & Hulme, D. (2021) A perspective on oil spills: what we should have learned about global warming. *Ocean and Coastal Management*, 202, 105509.
- Louati, H., Ben Said, O., Soltani, A., Got, P., Mahmoudi, E., Cravo-Laureau, C. et al. (2013) The roles of biological interactions and pollutant contamination in shaping microbial benthic community structure. *Chemosphere*, 93, 2535–2546.
- McGenity, T.J. (2014) Hydrocarbon biodegradation in intertidal wetland sediments. *Current Opinion in Biotechnology*, 27, 46–54.
- McGenity, T.J., Folwell, B.D., McKew, B.A. & Sanni, G.O. (2012) Marine crude-oil biodegradation: a central role for interspecies interactions. *Aquatic Biosystems*, 8, 10.
- McKew, B.A., Coulon, F., Yakimov, M.M., Denaro, R., Genovese, M., Smith, C.J. et al. (2007) Efficacy of intervention strategies for bioremediation of crude oil in marine systems and effects on indigenous hydrocarbonoclastic bacteria. *Environmental Microbiology*, 9, 1562–1571.
- Mrozik, A. & Piotrowska-Seget, Z. (2010) Bioaugmentation as a strategy for cleaning up of soils contaminated with aromatic compounds. *Microbiological Research*, 165, 363–375.
- Østman, B., Lin, R. & Adami, C. (2014) Trade-offs drive resource specialization and the gradual establishment of ecotypes. *BMC Evolutionary Biology*, 14, 113.
- Patowary, R., Devi, A. & Mukherjee, A.K. (2023) Advanced bioremediation by an amalgamation of nanotechnology and modern artificial intelligence for efficient restoration of crude petroleum oil-contaminated sites: a prospective study. *Environmental Science and Pollution Research International*, 30, 74459–74484.
- Paul, M.J., LeDuc, S.D., Boaggio, K., Herrick, J.D., Kaylor, S.D., Lassiter, M.G. et al. (2023) Effects of air pollutants from wildfires on downwind ecosystems: observations, knowledge gaps, and questions for assessing risk. *Environmental Science & Technology*, 57, 14787–14796.
- Péquin, B., Cai, Q., Lee, K. & Greer, C.W. (2022) Natural attenuation of oil in marine environments: a review. *Marine Pollution Bulletin*, 176, 113464.
- Rillig, M.C., Antonovics, J., Caruso, T., Lehmann, A., Powell, J.R., Veresoglou, S.D. et al. (2015) Interchange of entire communities: microbial community coalescence. *Trends in Ecology & Evolution*, 30, 470–476.
- Rocca, J.D., Muscarella, M.E., Peralta, A.L., Izabel-Shen, D. & Simonin, M. (2021) Guided by microbes: applying community coalescence principles for predictive microbiome Engineering.

- mSystems*, 6, e00538-21. Available from: <https://doi.org/10.1128/msystems.00538-21>
- Ruan, Z., Chen, K., Cao, W., Meng, L., Yang, B., Xu, M. et al. (2024) Engineering natural microbiomes toward enhanced bioremediation by microbiome modeling. *Nature Communications*, 15, 4694.
- Sánchez, Á., Vila, J.C.C., Chang, C.Y., Diaz-Colunga, J., Estrela, S. & Rebolleda-Gomez, M. (2021) Directed evolution of microbial communities. *Annual Review of Biophysics*, 50, 323–341.
- Schuijt, L.M., Peng, F.-J., van den Berg, S.J.P., Dingemans, M.M.L. & Van den Brink, P.J. (2021) (eco)toxicological tests for assessing impacts of chemical stress to aquatic ecosystems: facts, challenges, and future. *Science of the Total Environment*, 795, 148776.
- Schweitzer, H.D., Smith, H.J., Barnhart, E.P., McKay, L.J., Gerlach, R., Cunningham, A.B. et al. (2022) Subsurface hydrocarbon degradation strategies in low- and high-sulfate coal seam communities identified with activity-based metagenomics. *npj Biofilms and Microbiomes*, 8, 7.
- Shade, A. (2023) Microbiome rescue: directing resilience of environmental microbial communities. *Current Opinion in Microbiology*, 72, 102263.
- Shahi, A., Ince, B., Aydin, S. & Ince, O. (2017) Assessment of the horizontal transfer of functional genes as a suitable approach for evaluation of the bioremediation potential of petroleum-contaminated sites: a mini-review. *Applied Microbiology and Biotechnology*, 101, 4341–4348.
- Stauffert, M., Duran, R., Gassie, C. & Cravo-Laureau, C. (2014) Response of archaeal communities to oil spill in bioturbated mudflat sediments. *Microbial Ecology*, 67, 108–119.
- Swenson, W., Arendt, J. & Wilson, D.S. (2000) Artificial selection of microbial ecosystems for 3-chloroaniline biodegradation. *Environmental Microbiology*, 2, 564–571.
- Terrisse, F., Cravo-Laureau, C., Noël, C., Cagnon, C., Dumbrell, A.J., McGenity, T.J. et al. (2017) Variation of oxygenation conditions on a Hydrocarbonoclastic microbial community reveals Alcanivorax and Cycloclasticus ecotypes. *Frontiers in Microbiology*, 8, 1549.
- Tsoi, R., Wu, F., Zhang, C., Bewick, S., Karig, D. & You, L. (2018) Metabolic division of labor in microbial systems. *Proceedings of the National Academy of Sciences of the United States of America*, 115, 2526–2531.
- Tucker, C.M. & Fukami, T. (2014) Environmental variability counteracts priority effects to facilitate species coexistence: evidence from nectar microbes. *Proceedings of the Royal Society B: Biological Sciences*, 281, 20132637.
- van Oldenborgh, G.J., Krikken, F., Lewis, S., Leach, N.J., Lehner, F., Saunders, K.R. et al. (2021) Attribution of the Australian bush-fire risk to anthropogenic climate change. *Natural Hazards and Earth System Sciences*, 21, 941–960.
- Wang, F. & Zhang, W. (2019) Synthetic biology: recent progress, biosafety and biosecurity concerns, and possible solutions. *Journal of Biosafety and Biosecurity*, 1, 22–30.
- Wu, M., Wu, J., Zhang, X. & Ye, X. (2019) Effect of bioaugmentation and biostimulation on hydrocarbon degradation and microbial community composition in petroleum-contaminated loessal soil. *Chemosphere*, 237, 124456.
- Xu, C. & Yu, H. (2021) Insights into constructing a stable and efficient microbial consortium. *Chinese Journal of Chemical Engineering*, 30, 112–120.
- Yakimov, M.M., Timmis, K.N. & Golyshin, P.N. (2007) Obligate oil-degrading marine bacteria. *Current Opinion in Biotechnology*, 18, 257–266.
- Yang, Y., Zhang, Z.-W., Liu, R.-X., Ju, H.-Y., Bian, X.-K., Zhang, W.-Z. et al. (2021) Research progress in bioremediation of petroleum pollution. *Environmental Science and Pollution Research*, 28, 46877–46893.
- Zerebecki, R.A., Heck, K.L., Jr. & Valentine, J.F. (2022) Biodiversity influences the effects of oil disturbance on coastal ecosystems. *Ecology and Evolution*, 12, e8532.