

# Quality of clinical guidelines: It matters as it impacts patient care

Clinical guidelines are of vital importance in providing healthcare professionals with updated evidence-based recommendations eventually aiming to improve patient care and outcomes. Nonetheless, guidelines often lack evidence or transparency in how evidence was interpreted and graded. Historically, the development of guidelines was done in GOBSAT - ('Good old men sitting around the table') style by collating data from literature mixed with experience-based opinions. These guidelines often fail to address some of the essential and pivotal factors that are fundamental to high-quality guidelines. In addition, they often lack applicability, practicality, and the outcomes that really matter to patients.

In recent decades, there has been a shift to making the guideline development process more transparent, robust, and evidence-based, whilst involving multiple stakeholders (e.g., patients, general practitioners, or paramedics). Within the editorial board of United European Gastroenterology (UEG) Journal, a Guidelines Taskforce was established that aims at fostering a heightened awareness among readership and contributing authors on the quality of guidelines and the critical importance of transparency throughout the guideline development process.

In order to scrutinize the adherence to the standards of guidelines within the field of gastroenterology, the Taskforce collected and evaluated all 16 guidelines<sup>1-17</sup> published in the UEG Journal between January 2020 and March 2022. The assessment was conducted by four reviewers, who were equipped with the necessary expertise through participation in the International Guideline Training and Certification (INGUIDE) program (<https://inguide.org/>).

The four reviewers used the Appraisal of Guidelines for Research Evaluation II (AGREE II) tool,<sup>18</sup> which is a widely acknowledged instrument that provides both authors and (external) reviewers with a standardized methodology for assessing the critical quality metrics of guidelines. It features a detailed checklist designed to improve the reporting of guidelines, structured around six key domains:

1. Scope and purpose
2. Stakeholder involvement
3. Rigor of development
4. Clarity of presentation

5. Applicability
6. Editorial independence

These domains collectively address the essential components that underpin the reliability and validity of guidelines. The scoring is performed using a 7-point scale (ranging from 1: Strongly Disagree to 7: Strongly Agree) based on specific items within the domain itself. Each domain is scored by summing up all the scores of the items within the domain and scaling the total as a percentage of the maximum possible score. This results in one score per guideline per domain.

Overall, the reviewed guidelines were of good quality, mostly achieving above 70% in each domain (Figure 1). The highest scores were reached in **Domain 1—scope and purpose** (median 80%), as the scope and purpose of each guideline was generally explicit. Strikingly, the objectives of guidelines were not always specific to the clinical topic described in the guideline. For instance, a more specific objective could be 'to lower the risk of involuntary loss of stool' rather than 'to improve patient outcomes'.


The lowest scores were observed in **Domain 5—applicability** (median 53%). The domain includes factors, such as, whether the guideline provides recommendations or a tool on how to implement guidelines in practice, how to overcome facilitators/barriers and their cost implications. Most of the assessed guidelines lacked evaluation of barriers leading to reduced applicability of the guidelines. Performing pilot testing of the guideline or seeking feedback from key stakeholders or a health economist may highlight challenges in applicability and facilitate early adjustments before widespread implementation.

**Domain 3—rigor of development** (median 70%) is often considered as the most critical domain. Multiple frameworks are being used to appraise the quality of available evidence. To be relevant and practical, guidelines must follow a solid framework, involving a transparent systematic review process and a clear presentation of the evidence strength and recommendation rationale. The most frequently used framework is Grading of Recommendations Assessment, Development and Evaluation (GRADE).<sup>19</sup> It provides a systematic approach for developing and presenting evidence, has gained

---

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. United European Gastroenterology Journal published by Wiley Periodicals LLC. on behalf of United European Gastroenterology.



**FIGURE 1** Boxplot showing median and ranges of AGREE II scores per domain for the 16 included guidelines.

widespread acceptance and is currently endorsed by over 100 organizations, including the World Health Organization and the National Institute for Health and Care Excellence. Most of the reviewed guidelines indeed attempted to use GRADE, but did not apply it thoroughly. There is a scope to promote further training and education, such as INGUIDE (<https://inguide.org/>) or the UEG guideline development course (<https://ueg.eu/p/269>), prior to taking part in guideline development.

To further increase transparency, we would recommend the use of an Evidence-to-Decision framework, which provides the reader background, research evidence, considerations from the panel members, and drawn conclusions. To ensure the methodological rigor, (preferably multiple) external experts with different expertise (including a methodologist) could be involved.

Overall, guidelines generally had a multidisciplinary approach involving multiple stakeholders (covered in **Domain 2—stakeholder involvement**; median 67%), yet roles of the stakeholders within the guideline panel were frequently not clear, as these were not always explicitly stated in the published articles. The most important health questions can be missed if representatives of one or more disciplines are missing.

**Domain 4—clarity of presentation** (median 72%) and **Domain 6—editorial independence** (median 71%) both had medium to high scores. **Domain 4** appraises the quality of recommendations, which should be specific and unambiguous. Guidelines often lacked specification of the intent or purpose of a recommendation (e.g., ‘to improve quality of life’).

In conclusion, this editorial underscores the importance of standardized guideline development in order to improve transparency, rigor and applicability. The Quality of Care Committee of UEG is actively promoting educational events (<https://ueg.eu/p/269>), tools and frameworks to facilitate this process,<sup>20</sup> ultimately aiming at a more uniform landscape of high-quality clinical guidance.

## KEYWORDS

AGREE II, assessment, gastroenterology, GRADE, guideline, quality

## CONFLICT OF INTEREST STATEMENT

Speaking fee from Galapagos and Abbvie. Haluk Tarik Kani received speaking fee from Janssen and Abbvie.

Iris J. M. Levink<sup>1,2</sup>

Alberto Balduzzi<sup>3</sup>

Irene Marafini<sup>4</sup>

Haluk Tarik Kani<sup>5,6</sup>

Yasuko Maeda<sup>7</sup>

on behalf of the UEGJ Guideline Taskforce

<sup>1</sup>Department of Gastroenterology and Hepatology, Erasmus MC University Medical Centre, Rotterdam, The Netherlands

<sup>2</sup>Department of Internal Medicine, Reinier de Graaf Gasthuis, Delft, The Netherlands

<sup>3</sup>Department of Surgery, Dentistry, Paediatrics and Gynaecology, Unit of General and Pancreatic Surgery, The Pancreas Institute Verona, University of Verona, Verona, Italy

<sup>4</sup>Policlinico Universitario Tor Vergata, Gastroenterology Unit, Rome, Italy

<sup>5</sup>Department of Gastroenterology, Marmara University, School of Medicine, Istanbul, Turkey

<sup>6</sup>Marmara University Institute of Gastroenterology, Istanbul, Turkey

<sup>7</sup>Department of Surgery, Queen Elizabeth University Hospital, Glasgow, UK

## Correspondence

Iris J. M. Levink.

Email: [i.levink@erasmusmc.nl](mailto:i.levink@erasmusmc.nl)

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

## ORCID

Iris J. M. Levink <https://orcid.org/0000-0001-9825-8728>

Irene Marafini <https://orcid.org/0000-0001-8258-3235>

Yasuko Maeda <https://orcid.org/0000-0002-4081-4741>

## REFERENCES

- Al-Toma A, Volta U, Auricchio R, Castillejo G, Sanders DS, Cellier C, et al. European Society for the Study of Coeliac Disease (ESsCD) guideline for coeliac disease and other gluten-related disorders. *UEG J.* 2019;7(5):583–613. <https://doi.org/10.1177/2050640619844125>
- Assmann SL, Keszhelyi D, Kleijnen J, Anastasiou F, Bradshaw E, Brannigan AE, et al. Guideline for the diagnosis and treatment of Faecal Incontinence—a UEG/ESCP/ESNM/ESPCG collaboration. *United Eur Gastroenterol J.* 2022;10(3):251–86. <https://doi.org/10.1002/ueg2.12213>
- Bischoff SC, Barazzoni R, Busetto L, Campmans-Kuijpers M, Cardinale V, Chermesh I, et al. European guideline on obesity care in patients with gastrointestinal and liver diseases – joint European Society for Clinical Nutrition and Metabolism/United European Gastroenterology guideline. *United Eur Gastroenterol J.* 2022;10(7):665–722. <https://doi.org/10.1002/ueg2.12280>

4. D'Amico F, Rubin DT, Kotze PG, Magro F, Siegmund B, Kobayashi T, et al. International consensus on methodological issues in standardization of fecal calprotectin measurement in inflammatory bowel diseases. *United Eur Gastroenterol J.* 2021;9(4):451–60. <https://doi.org/10.1002/ueg2.12069>
5. Hammer HF, Fox MR, Keller J, Salvatore S, Basilisco G, Hammer J, et al. European guideline on indications, performance, and clinical impact of hydrogen and methane breath tests in adult and pediatric patients: European Association for Gastroenterology, Endoscopy and Nutrition, European Society of Neurogastroenterology and Motility, and European Society for Paediatric Gastroenterology Hepatology and Nutrition consensus. *United Eur Gastroenterol J.* 2022;10(1):15–40. <https://doi.org/10.1002/ueg2.12133>
6. Keller J, Hammer HF, Afolabi PR, Benninga M, Borrelli O, Dominguez-Munoz E, et al. European guideline on indications, performance and clinical impact of <sup>13</sup>C-breath tests in adult and pediatric patients: an EAGEN, ESNM, and ESPGHAN consensus, supported by EPC. *United Eur Gastroenterol J.* 2021;9(5):598–625. <https://doi.org/10.1002/ueg2.12099>
7. Kruis W, Germer C.-T, Böhm S, Dumoulin FL, Frieling T, Hampe J, et al. German guideline diverticular disease/diverticulitis: PART I. *United Eur Gastroenterol J.* 2022;10(9):923–39. <https://doi.org/10.1002/ueg2.12309>
8. Kruis W, Germer C.-T, Böhm S, Dumoulin FL, Frieling T, Hampe J, et al. German guideline diverticular disease/diverticulitis: PART II. *United Eur Gastroenterol J.* 2022;10(9):940–57. <https://doi.org/10.1002/ueg2.12313>
9. Löhr J-M, Beuers U, Vujasinovic M, Alvaro D, Frøkjær JB, Buttgerit F, et al. European Guideline on IgG4-related digestive disease – UEG and SGF evidence-based recommendations. *UEG J.* 2020;8(6):637–66. <https://doi.org/10.1177/2050640620934911>
10. Markar S, Andreou A, Bonavina L, Florez ID, Huo B, Kontouli K.-M, et al. UEG and EAES rapid guideline: update systematic review, network meta-analysis, CINeMA and GRADE assessment, and evidence-informed European recommendations on surgical management of GERD. *United Eur Gastroenterol J.* 2022;10(9):983–98. <https://doi.org/10.1002/ueg2.12318>
11. Miehke S, Guagnozzi D, Zabana Y, Tontini GE, Kanstrup Fiehn A-M, Wildt S, et al. European guidelines on microscopic colitis: United European Gastroenterology and European Microscopic Colitis Group statements and recommendations. *United Eur Gastroenterol J.* 2021;9(1):13–37. <https://doi.org/10.1177/2050640620951905>
12. Oude Nijhuis R, Zaninotto G, Roman S, Boeckstaens G, Fockens P, Langendam M, et al. European guidelines on achalasia: United European Gastroenterology and European Society of Neurogastroenterology and Motility recommendations. *UEG J.* 2020;8(1):13–33. <https://doi.org/10.1177/2050640620903213>
13. Savarino E, Zingone F, Barberio B, Marasco G, Akyuz F, Akpinar H, et al. Functional bowel disorders with diarrhoea: clinical guidelines of the United European Gastroenterology and European Society for Neurogastroenterology and Motility. *United Eur Gastroenterol J.* 2022;10(6):556–84. <https://doi.org/10.1002/ueg2.12259>
14. Schol J, Wauters L, Dickman R, Drug V, Mulak A, Serra J, et al. United European Gastroenterology (UEG) and European Society for Neurogastroenterology and Motility (ESNM) consensus on gastroparesis. *United Eur Gastroenterol J.* 2021;9(8):287–306. <https://doi.org/10.1111/nmo.14237>
15. Terlouw LG, Moelker A, Abrahamsen J, Acosta S, Bakker OJ, Baumgartner I, et al. European guidelines on chronic mesenteric ischaemia – joint United European Gastroenterology, European Association for Gastroenterology, Endoscopy and Nutrition, European Society of Gastrointestinal and Abdominal Radiology, Netherlands Association of Hepatogastroenterologists, Hellenic Society of Gastroenterology, Cardiovascular and Interventional Radiological Society of Europe, and Dutch Mesenteric Ischemia Study group clinical guidelines on the diagnosis and treatment of patients with chronic mesenteric ischaemia. *UEG J.* 2020;8(4):371–95. <https://doi.org/10.1177/2050640620916681>
16. Valori R, Cortas G, de Lange T, Salem Balfaqih O, de Pater M, Eisendrath P, et al. Performance measures for endoscopy services: a European Society of Gastrointestinal Endoscopy (ESGE) quality improvement initiative. *UEG J.* 2019;7(1):21–44. <https://doi.org/10.1177/2050640618810242>
17. Wauters L, Dickman R, Drug V, Mulak A, Serra J, Enck P, et al. United European Gastroenterology (UEG) and European Society for Neurogastroenterology and Motility (ESNM) consensus on functional dyspepsia. *United Eur Gastroenterol J.* 2021;9(3):307–31. <https://doi.org/10.1002/ueg2.12061>
18. Brouwers MC, Kerkvliet K, Spithoff K. The AGREE Reporting Checklist: a tool to improve reporting of clinical practice guidelines. *BMJ.* 2016;352:i1152. <https://doi.org/10.1136/bmj.i1152>
19. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ.* 2008;336(7650):924–6. <https://doi.org/10.1136/bmj.39489.470347.ad>
20. Boltin D, Lambregts DM, Jones F, Siterman M, Bonovas S, Cornberg M, et al. UEG framework for the development of high-quality clinical guidelines. *UEG J.* 2020;8:851–64. <https://doi.org/10.1177/2050640620950854>