

CORRECTION

Correction: Gender differences in quality of life in coronary artery disease patients with comorbidities undergoing coronary revascularization


Tom H. Oreeel, Pythia T. Nieuwkerk, Iris D. Hartog, Justine E. Netjes, Alexander B. A. Vonk, Jorrit Lemkes, Hanneke W. M. van Laarhoven, Michael Scherer-Rath, Mirjam A. G. Sprangers, José P. S. Henriques

The columns of the parameters beta (β) and standard errors (SE) in Table 2 are incorrect. It should have been chi-square statistics “ χ^2 ” and “df” respectively. Please see the correct Table 2 here.

Table 2. Wald type II chi-square statistics of the linear mixed models for each HRQOL measure.

Variables	χ^2	df	P value
Physical Health (PCS)			
Time	95.024	1	< .001*
Gender	15.049	1	< .001*
Time \times gender	4.843	1	.028*
Comorbidity burden:			
Total comorbidity conditions	16.819	2	< .001*
Charlson comorbidity categories	12.798	2	.002*
Comorbidity burden \times gender:			
Total comorbidity conditions	9.448	2	.009*
Charlson comorbidity categories	3.715	2	.156
Comorbidity burden \times time:			
Total comorbidity conditions	0.348	2	.840
Charlson comorbidity categories	0.990	2	.610
Mental health (MCS)			
Time	44.794	1	< .001*
Gender	1.351	1	.245
Time \times gender	0.206	1	.650
Comorbidity burden:			
Total comorbidity conditions	2.143	2	.343
Charlson comorbidity categories	4.787	2	.091
Comorbidity burden \times gender:			
Total comorbidity conditions	2.775	2	.250
Charlson comorbidity categories	5.135	2	.077
Comorbidity burden \times time:			
Total comorbidity conditions	4.823	2	.090
Charlson comorbidity categories	2.344	2	.310
Disease-specific HRQoL (SAQ)			

(Continued)



OPEN ACCESS

Citation: Oreeel TH, Nieuwkerk PT, Hartog ID, Netjes JE, Vonk ABA, Lemkes J, et al. (2024) Correction: Gender differences in quality of life in coronary artery disease patients with comorbidities undergoing coronary revascularization. PLoS ONE 19(7): e0307984. <https://doi.org/10.1371/journal.pone.0307984>

Published: July 24, 2024

Copyright: © 2024 Oreeel et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 2. (Continued)

Variables	χ^2	df	P value
Time	248.309	1	< .001*
Gender	15.787	1	< .001*
Time \times gender	1.564	1	.211
Comorbidity burden:			
Total comorbidity conditions	2.035	2	.362
Charlson comorbidity categories	7.115	2	.029*
Comorbidity burden \times gender:			
Total comorbidity conditions	13.761	2	.001*
Charlson comorbidity categories	3.132	2	.209
Comorbidity burden \times time			
Total comorbidity conditions	6.877	2	.032*
Charlson comorbidity categories	0.906	2	.636

*P values < 0.05 were considered significant.

<https://doi.org/10.1371/journal.pone.0307984.t001>

Reference

1. Oreel TH, Nieuwkerk PT, Hartog ID, Netjes JE, Vonk ABA, Lemkes J, et al. (2020) Gender differences in quality of life in coronary artery disease patients with comorbidities undergoing coronary revascularization. PLOS ONE 15(6): e0234543. <https://doi.org/10.1371/journal.pone.0234543> PMID: 32555617