



HHS Public Access

Author manuscript

Am J Kidney Dis. Author manuscript; available in PMC 2024 August 06.

Published in final edited form as:

Am J Kidney Dis. 2024 June ; 83(6): 840. doi:10.1053/j.ajkd.2024.04.002.

Erratum regarding “Serum β -Trace Protein and β 2-Microglobulin as Predictors of ESRD, Mortality, and Cardiovascular Disease in Adults With CKD in the Chronic Renal Insufficiency Cohort (CRIC) Study” (*Am J Kidney Dis.* 2016;68(1):68–76)

In the Original Investigation entitled “Serum β -Trace Protein and β 2-Microglobulin as Predictors of ESRD, Mortality, and Cardiovascular Disease in Adults With CKD in the Chronic Renal Insufficiency Cohort (CRIC) Study” that appeared in the July 2016 issue of *AJKD*,¹ there was an error regarding the source of the β -Trace Protein (BTP) and β 2-Microglobulin (B2M) samples. The samples were reported as being measured in serum; however, the samples were measured in EDTA plasma.

Reference

1. Foster MC, Coresh J, Hsu C, et al. Serum β -Trace Protein and β 2-Microglobulin as Predictors of ESRD, Mortality, and Cardiovascular Disease in Adults With CKD in the Chronic Renal Insufficiency Cohort (CRIC) Study. *Am J Kidney Dis.* 2016;68(1):68–76. doi:10.1053/j.ajkd.2016.01.015 [PubMed: 26948990]