

Draft genome sequence of multidrug-resistant *Kurthia gibsonii* strain Hakim RU_BHWE isolated from sewage water in Bangladesh

M. Romance,¹ Md. Arif-Uz-Zaman Polash,¹ Nusrat Zahan,¹ Jafor Raihan,¹ Md. Sumon Ali,¹ Muhib Ullah Khan,¹ Subir Sarker,² Md. Hakimul Haque^{1,2}

AUTHOR AFFILIATIONS See affiliation list on p. 3.

ABSTRACT We have sequenced the genome of *Kurthia gibsonii* strain Hakim RU_BHWE, isolated from sewage water. The assembled genome consists of 2.891 Mb with 58.6883× coverage, presenting an average GC content of 36.60%. This genome includes 8 CRISPR arrays, 3 prophages, 3 antibiotic resistance genes, and 12 virulence factor genes.

KEYWORDS whole genome, sewage water, multidrug-resistant, *Kurthia gibsonii*, Bangladesh

Since its discovery in 1883 by Hermann Kurth, *Kurthia* spp. has been known for its wide environmental distribution and its potential to cause opportunistic infections (1–5). Genome sequencing is crucial for understanding the survival, adaptation, and role of *Kurthia* spp. in antimicrobial resistance (6). Reports of multidrug-resistant *Kurthia* spp. in humans, animals, food, and the environment underscore the importance of ongoing surveillance through One Health approaches to understand its molecular epidemiology and implement effective public health strategies (1–3, 5, 7–9).

The research techniques and protocols for this study were approved by the Institute of Biological Science (IBSc) at the University of Rajshahi, Bangladesh, under Memo No. 56/321/IAMEBBC/IBSc. In September 2023, we collected samples of sewage water at the University of Rajshahi (24.3733°N, 88.6049°E), following standard procedures. The water samples were mixed thoroughly, transferred to sterile tubes, and transported to the laboratory. We then inoculated these samples on urinary tract infection agar (HiMedia, India) and incubated them aerobically at 37°C for 18–24 hours (10). *Kurthia gibsonii* was isolated by streaking the cultures on tryptic soy agar (HiMedia), followed by staining and biochemical tests (11). Antibigram study of the isolates was performed using the disk diffusion method (12), following Clinical and Laboratory Standards Institute guidelines (13). The strain exhibited resistance to penicillin, amoxicillin, tetracycline, and doxycycline. We cultured the isolated strain in nutrient broth (HiMedia) overnight at 37°C and then extracted its genomic DNA using the Qiagen DNA Mini Kit (QIAGEN, Hilden, Germany). The genomic DNA was enzymatically fragmented using the NEBNext dsDNA Fragmentase Kit (NEB, Massachusetts, USA), and size selection was carried out with solid-phase reversible immobilization beads (14). A sequencing library was prepared using the Nextera DNA Flex Library Preparation Kit (Illumina, San Diego, CA, USA), and the library was sequenced with 2 × 150 paired-end reads on the Illumina NextSeq 2000 platform. Quality checks were performed using FastQC v.0.11.7 (15). Raw paired-end reads ($n = 2,450,168$) were trimmed using Trimmomatic v.0.39 (16), and genome assembly was conducted using Unicycler v.0.4.9 (17). The annotation of the genome was carried out using PGAP v.3.0 (18). The assembled genome was analyzed for antibiotic resistance genes (ARGs) using CARD v.3.2.4 with RGI v.6.0.2 (19) and ResFinder v.4.1

Editor John J. Dennehy, Department of Biology, Queens College, Queens, New York, USA

Address correspondence to Md. Hakimul Haque, hakim.ahvs@ru.ac.bd.

The authors declare no conflicts of interest.

Received 22 May 2024

Accepted 26 June 2024

Published 22 July 2024

Copyright © 2024 Romance et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

TABLE 1 Genomic traits of the *Kurthia* strain Hakim RU_BHWE

Elements	Values
Genome size	2,891,399 bp
Genome coverage	58.6883x
G + C content	36.60%
Number of contigs	104
Contig L50	10
Contig N50	101,097 bp
Total genes	2,920
Coding sequences	2,868
Coding genes	2,836
RNA genes	52
tRNA genes	44
rRNAs genes	3
ncRNAs genes	5
Pseudo genes	32
Genes assigned to SEED subsystems	2,943
Number of subsystems	261

(20), mobile genetic elements (MGEs) using mobileOG-db (21), virulence factor genes using VFDB with VFAnalyzer v.4.0 (22), pathogenicity index using PathogenFinder v.1.1 (23), sequence type using MLST v.2.0 (24), CRISPR arrays using CRISPRimmunity (25), prophages using PHASTER (26), and metabolic functional features using RAST v.2.0 (27). We used default parameters for all tools, unless noted otherwise.

The traits of the draft genomes are documented in Table 1. Notably, 3 ARGs, 12 virulence genes, and 95 MGEs were predicted. MLST classified the genome as sequence type unknown. The genome exhibited eight CRISPR arrays with signature genes (*Cas14j*, *WYL*, *csa3*, *cas1*, *cas2*, *cas4*, *cas5*, *cas7*, *DEDDh*, and *cas8c*) and three prophages. RAST analysis uncovered 261 subsystems comprising 2,943 genes with 27% coverage (Fig. 1).




FIG 1 Metabolic functional features in the assembled genome of the *Kurthia gibsonii* strain Hakim RU_BHWE in SEED viewer. The 27% coverage indicates the completeness of functional roles within a specific subsystem across different genomes.

ACKNOWLEDGMENTS

Rajshahi University funded this study under Bangladesh's University Grants Commission (financial year 2023–2024).

M.R.: conceptualization, data curation, formal analysis, investigation, methodology, project administration, and writing (original draft); A.U.-Z.P.: conceptualization, data curation, formal analysis, methodology, software, and writing (original draft); N.Z.: conceptualization, data curation, investigation, and methodology; J.R.: formal analysis, methodology, and software; M.S.A.: formal analysis, methodology, and software; M.U.K.: formal analysis, methodology, and software; S.S.: formal analysis, writing (review and editing), and software; M.H.H.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, resources, supervision, validation, writing (original draft, review, and editing).

AUTHOR AFFILIATIONS

¹Department of Veterinary and Animal Sciences, University of Rajshahi, Rajshahi, Bangladesh

²Biomedical Sciences and Molecular Biology, College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, Australia

AUTHOR ORCIDS

Md. Hakimul Haque  <http://orcid.org/0000-0002-3856-6478>

AUTHOR CONTRIBUTIONS

M. Romance, Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration | Md. Arif-Uz-Zaman Polash, Conceptualization, Data curation, Formal analysis, Methodology, Software | Nusrat Zahan, Conceptualization, Data curation, Investigation, Methodology | Jafor Raihan, Formal analysis, Methodology, Software | Md. Sumon Ali, Formal analysis, Methodology, Software | Muhib Ullah Khan, Formal analysis, Methodology, Software | Subir Sarker, Formal analysis, Software, Writing – review and editing | Md. Hakimul Haque, Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Resources, Supervision, Validation, Writing – original draft, Writing – review and editing

DATA AVAILABILITY

The study on *Kurthia gibsonii* strain Hakim RU_BHWE, conducted using the whole genome sequencing shotgun approach, was submitted to National Center for Biotechnology Information/GenBank, and the assembly was deposited under accession number [JBCHWB000000000](https://ncbi.nlm.nih.gov/assembly/JBCHWB000000000). The pertinent data, including the original readings, were stored with BioProject accession number [PRJNA1102855](https://ncbi.nlm.nih.gov/bioproject/PRJNA1102855), BioSample accession number [SAMN41030973](https://ncbi.nlm.nih.gov/biosample/SAMN41030973), and Sequence Read Archive accession number [SRR28762083](https://ncbi.nlm.nih.gov/sra/SRR28762083). The specific version mentioned in this document is labeled as [JBCHWB000000000.1](https://ncbi.nlm.nih.gov/assembly/JBCHWB000000000.1).

REFERENCES

1. Kurth H. 1883. Ueber Bacterium zopfii, eine neue Bakterienart. *Berichte der Deutschen Botanischen Gesellschaft* 1:97–100.
2. Ongrádi J, Stercz B, Kövesdi V, Nagy K, Chatlyne L. 2014. Isolation of *Kurthia gibsonii* from non-gonorrhoeal urethritis: implications for the pathomechanism upon surveying the literature. *Acta Microbiol Immunol Hung* 61:79–87. <https://doi.org/10.1556/AMicr.61.2014.1.8>
3. Dourou D, Spyrelli ED, Doulgeraki AI, Argyri AA, Grounta A, Nychas G-JE, Chorianopoulos NG, Tassou CC. 2021. Microbiota of chicken breast and thigh fillets stored under different refrigeration temperatures assessed by next-generation sequencing. *Foods* 10:765. <https://doi.org/10.3390/foods10040765>
4. Shaw S, Keddle RM. 1983. A numerical taxonomic study of the genus *Kurthia* with a revised description of *Kurthia zopfii* and a description of *Kurthia gibsonii* sp. nov. *Syst Appl Microbiol* 4:253–276. [https://doi.org/10.1016/S0723-2020\(83\)80054-X](https://doi.org/10.1016/S0723-2020(83)80054-X)
5. Mukhopadhyay BC, Mitra S, Kazi TA, Mandal S, Biswas SR. 2019. Draft genome sequence of cold-tolerant *Kurthia gibsonii* B83, isolated from spinach leaf. *Microbiol Resour Announc* 8:e01480-18. <https://doi.org/10.1128/MRA.01480-18>
6. Baker S, Thomson N, Weill FX, Holt KE. 2018. Genomic insights into the emergence and spread of antimicrobial-resistant bacterial pathogens. *Science* 360:733–738. <https://doi.org/10.1126/science.aar3777>

7. Gardner GA. 1969. Physiological and morphological characteristics of *Kurthia zopfii* isolated from meat products. *J Appl Bacteriol* 32:371–380. <https://doi.org/10.1111/j.1365-2672.1969.tb00986.x>
8. Sharma BC, Subba R, Saha A. 2012. *Kurthia* sp, a novel member of phosphate solubilising bacteria from rhizospheric tea soil of Darjeeling hills, India. *J of Pharm Bioallied Sci* 2:36–39. <https://doi.org/10.9790/3008-0233639>
9. Lozica L, Maurić Maljković M, Mazić M, Gottstein Ž. 2022. *Kurthia gibsonii*, a novel opportunistic pathogen in poultry. *Avian Pathol* 51:26–33. <https://doi.org/10.1080/03079457.2021.1993132>
10. Romance M, Khan MU, Islam MS, Islam MF, Haque MH. 2024. Draft genome sequence of multidrug-resistant *Klebsiella pneumoniae* Hakim-RU strain isolated from a patient with urinary tract infections in Bangladesh. *Microbiol Resour Announc* 13:e0008924. <https://doi.org/10.1128/mra.00089-24>
11. Bergey DH, Buchanan RE, Gibbons NE. 1974. *Bergey's Manual of Determinative Bacteriology*. 8th ed, p 966–1097. American Society for Microbiology, Washington, DC, USA.
12. Bauer AW, Kirby WMM, Sherris JC, Turck M. 1966. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 45:493–496. https://doi.org/10.1093/ajcp/45.4_ts.493
13. M100-S32. 2022. Performance standards for antimicrobial susceptibility testing. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
14. Liu D, Li Q, Luo J, Huang Q, Zhang Y. 2023. An SPRI beads-based DNA purification strategy for flexibility and cost-effectiveness. *BMC Genomics* 24:125. <https://doi.org/10.1186/s12864-023-09211-w>
15. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Available from: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
16. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
17. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>
18. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>
19. Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A, Huynh W, Nguyen A-LV, Cheng AA, Liu S, et al. 2020. CARD 2020: antibiotic resistance surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res* 48:D517–D525. <https://doi.org/10.1093/nar/gkz935>
20. Florensa AF, Kaas RS, Clausen P, Aytan-Aktug D, Aarestrup FM. 2022. ResFinder - an open online resource for identification of antimicrobial resistance genes in next-generation sequencing data and prediction of phenotypes from genotypes. *Microb Genom* 8:000748. <https://doi.org/10.1099/mgen.0.000748>
21. Brown CL, Mullet J, Hindi F, Stoll JE, Gupta S, Choi M, Keenum I, Vikesland P, Pruden A, Zhang L. 2022. mobileOG-dB: a manually curated database of protein families mediating the life cycle of bacterial mobile genetic elements. *Appl Environ Microbiol* 88:e0099122. <https://doi.org/10.1128/aem.00991-22>
22. Liu B, Zheng D, Zhou S, Chen L, Yang J. 2022. VFDB 2022: a general classification scheme for bacterial virulence factors. *Nucleic Acids Res* 50:D912–D917. <https://doi.org/10.1093/nar/gkab1107>
23. Cosentino S, Voldby Larsen M, Møller Aarestrup F, Lund O. 2013. Pathogenfinder - distinguishing friend from foe using bacterial whole genome sequence data. *PLoS ONE* 8:e77302. <https://doi.org/10.1371/journal.pone.0077302>
24. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>
25. Zhou F, Yu X, Gan R, Ren K, Chen C, Ren C, Cui M, Liu Y, Gao Y, Wang S, Yin M, Huang T, Huang Z, Zhang F. 2023. CRISPRimmunity: an interactive web server for CRISPR-associated important molecular events and modulators used in geNome editing tool identifying. *Nucleic Acids Res* 51:W93–W107. <https://doi.org/10.1093/nar/gkad425>
26. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–21. <https://doi.org/10.1093/nar/gkw387>
27. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, et al. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>