

A scoping review of evidence-based guidance and guidelines published by general practice professional organizations

Emer O'Brien^{1,*}, Seamus Duffy¹, Velma Harkins², Susan M. Smith^{1,3}, Noirin O'Herlihy², Aisling Walsh⁴, Barbara Clyne¹, Emma Wallace^{1,5}

¹Department of General Practice, RCSI University of Medicine and Health Sciences, Dublin, Ireland

²Irish College of General Practitioners, Dublin, Ireland

³Department of Public Health and Primary Care, Trinity College Dublin, Dublin Ireland

⁴Department of Public Health and Epidemiology, RCSI University of Medicine and Health Sciences, Dublin, Ireland

⁵Department of General Practice, University College Cork, Cork, Ireland

*Corresponding author: Department of General Practice, RCSI University of Medicine and Health Sciences, 123 St Stephens Green, Dublin 2, Ireland. E-mail: emerobrien@rcsi.ie

Background: General practitioners (GPs) need robust, up-to-date evidence to deliver high-quality patient care. There is limited literature regarding the role of international GP professional organizations in developing and publishing clinical guidelines to support GPs clinical decision making.

Objective: To identify evidence-based guidance and clinical guidelines produced by GP professional organizations and summarize their content, structure, and methods of development and dissemination.

Methods: Scoping review of GP professional organizations following Joanna Briggs Institute guidance. Four databases were searched and a grey literature search was conducted. Studies were included if they were: (i) evidence-based guidance documents or clinical guidelines produced de novo by a national GP professional organization, (ii) developed to support GPs clinical care, and (iii) published in the last 10 years. GP professional organizations were contacted to provide supplementary information. A narrative synthesis was performed.

Results: Six GP professional organizations and 60 guidelines were included. The most common de novo guideline topics were mental health, cardiovascular disease, neurology, pregnancy and women's health and preventive care. All guidelines were developed using a standard evidence-synthesis method. All included documents were disseminated through downloadable pdfs and peer review publications. GP professional organizations indicated that they generally collaborate with or endorse guidelines developed by national or international guideline producing bodies.

Conclusion: The findings of this scoping review provide an overview of de novo guideline development by GP professional organizations and can support collaboration between GP organizations worldwide thus reducing duplication of effort, facilitating reproducibility, and identifying areas of standardization.

Protocol registration: Open Science Framework: <https://doi.org/10.17605/OSF.IO/JXQ26>.

Key words: clinical practice guidelines, evidence-based practice, family practitioner, general practice, primary healthcare

Introduction

General practitioners (GPs) practice medicine in the setting of the community and the family,¹ and are responsible for providing comprehensive and ongoing care to every individual seeking medical support irrespective of their illness, sex, or age.² Internationally there are variations in healthcare structures and the cultural settings that GPs practice in, but there are also similar components. For example in many countries GPs act as gatekeepers, facilitating access to hospital and speciality care and certain diagnostic tests.³

GPs endeavour to use a patient centred approach to achieve shared decision making, through the integration of clinical evidence, clinical judgement, and patient priorities.^{4,5} In this setting, accessible, succinct, evidence-based guidance is required by GPs to support patient care.^{6–10} Clinical practice guidelines, being systematically developed statements, based on a comprehensive evaluation of evidence, aim to address

this need and support practitioners' to make healthcare decisions.¹¹ However, a review of 45 UK clinical practice guidelines reported a significant number of guideline recommendations were based on studies with little relevance to primary care.¹² With up to 80% of GP consultations involving the management of patients with multiple chronic medical conditions, GPs require adequate decision support to deliver such complex care in the primary care setting.^{13–16} High workload and time pressure are significant barriers to utilization of clinical guidelines,¹⁷ however, despite these barriers, GPs are more likely to use guidelines that involved GP contributors during the development stage and where the evidence is applicable to primary care.¹⁸

Availability of resources and national guideline development agencies influence how GP professional organizations develop and disseminate clinical guidelines.¹⁹ National guideline agencies may approach GP organizations to endorse

Key messages

- GP professional organizations produce de novo clinical guidelines.
- Mental health is the most common de novo guideline category.
- Dissemination occurs via downloadable pdfs and peer review publication.
- GP professional organizations also collaborate on and endorse other guidelines.

their guidelines.²⁰ GP organizations may also adapt or adopt existing national and/or international guidelines and disseminate the findings to their members in the context of primary care.^{21–23} Some GP organizations play a central role in developing de novo guidelines for GPs. For example, the Dutch College of GPs/Nederlands Huisartsen Genootschap (NHG) develop clinical guidelines which cover a range of primary care presentations.²⁴ Assimilating how GP professional organizations develop and disseminate clinical guidelines for their members, would facilitate collaboration between GP organizations thus reducing duplication of effort and promoting a standardization of processes to support GPs in their clinical decision making.²⁵

The aim of this scoping review was to identify what evidence-based guidance is published by GP professional organizations internationally to support GPs in their clinical decision making. The objectives were: (i) to identify the topics covered, (ii) to review the methods used to develop evidence-based guidance and/or clinical guidelines and how these guidance documents are structured, and (iii) to explore how evidence is disseminated to GPs.

Methods

This scoping review was preregistered on Open Science Framework and a study protocol has been published.²⁵

Given that many GP guidelines may not be published as peer reviewed publications and the evident heterogeneity of nomenclature (e.g. guides versus guidance versus clinical guidelines), a combination of a bibliographic database search, grey literature search, and GP organization author contacts was conducted.

Scoping review study design

The scoping review was conducted in accordance with JBI methodology for scoping reviews^{26–28} and is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis extension for scoping reviews (PRISMA-ScR).²⁹ A scoping review was selected as it is designed to address a broad research question by mapping a body of literature in that area.^{30,31} The research team comprised researchers and knowledge users (GPs and Irish College of General Practitioners) and through regular meetings identified the specific research question, designed the search strategy, and synthesized the evidence.

Eligibility Articles were included where they were an evidence-based guidance document or clinical guideline (henceforth “guideline”) produced by a national GP professional organization, either de novo (new and updated versions) or through adaptation (Table 1). These guidelines had to support GPs clinical decision making and patient clinical care and be published in the last 10 years for currency. For the purposes of this scoping review “published” refers to guidelines that

are made freely available by GP professional organizations on their website or through peer reviewed publication. It does not include guidelines that are collaborations with or endorsed by other, non GP organizations. No language restrictions were applied. English versions were sought on the organizations websites or translated via Google translate.

Search strategy The search strategy (Appendix 1) was developed in consultation with an information specialist (PM), and applied to 4 bibliographic databases (Medline, Embase, Cochrane Library, and Scopus). The search was conducted on 12 April 2021.

Grey literature search

In addition to NHS evidence and Guideline central, websites of included organizations were searched for clarification regarding the production method of de novo guidelines. This search was not exhaustive, e.g. if additional de novo guidelines were discovered from the organization websites these were not charted as per the database search.

Screening and data extraction Two reviewers (SD and EOB) screened titles for eligibility, a third reviewer (EW) resolved conflicts. One reviewer (EOB) completed the data charting, with 20% checked by the second reviewer (SD).

Quality appraisal and analysis Consistent with established scoping review methodology,²⁶ we did not appraise the risk of bias of included studies, nor did we summarize the data quantitatively (meta-analysis). Findings were synthesized narratively with descriptive statistics (e.g. frequencies and percentages).

GP professional organizations were contacted to provide supplementary information. A total of 39 organizations were contacted through professional links via the Irish College of General Practitioners ($n = 6$) and links with the European Society for Quality and Safety in Practice ($n = 33$). GP professional organizations were invited to provide additional information regarding their published guidelines (see Appendix 2) which was based on the inclusion criteria for the review (Table 1). Responses were collected using Survey Monkey, transferred to Excel, and data analysed using descriptive statistics. The purpose was to clarify methods of guideline production and publication within GP organizations included in the scoping review and identify any other relevant publications.

Results

Search results

A total of 14,142 titles and abstracts were identified after duplicates removed. Of these, 125 articles full texts were assessed for eligibility with 60 full texts guidelines being included (Appendix 3 and Fig. 1) from 6 organizations. Findings

Table 1. Summary of the eligibility criteria for the review.

Inclusion criteria	Excluded
National GP professional organization	Regional organizations
Evidence based guidance or guideline development	Commentary
(i) Explicit method of development	Position statement
(ii) Includes literature review	
(iii) Reviewed by committee or experts	
(iv) Recommendation formation not necessary once distinction made between guidance and guideline	
Published by GP professional organizations	Collaborations with other non GP specialties or other guideline producing bodies
(i) Produced de novo or by adaptation	Endorsements by other national or international guideline producing bodies
(ii) Open access on GP professional organization website and/or	GP organizations with members only access
(iii) Peer review publication	
Published in the last 10 years	Publications >10 years old
All languages	
Patient clinical care	Governance document
	Policy document
World Organization of National Colleges, Academies and Academic Associations of General Practitioners/Family Physicians (WONCA) definition of GP ²	

from the websites of included organizations revealed that additional de novo guidelines not retrieved from the database search were also available. The 60 identified guidelines from the database search are therefore a subset of de novo guidelines from the included organizations.

Of the 39 GP professional organizations contacted to provide supplementary information, 13 responded, representing input from 12 unique GP professional organizations (Appendix 4). Four of these 12 organizations met the inclusion criteria; being national organizations and producing evidence-based de novo guidelines. Eight organizations were excluded based on production method, members only access or being a regional organization (Table 2). Guideline topics covered in the questionnaire included all methods of production. These findings were documented for the purposes of comparison and context, although they were outside the inclusion criteria of de novo production. Of the 6 organizations identified in the database search, 4 responded to the contact for supplementary information and 2 did not.

Characteristics of included guidelines

The 6 organizations included were from the Netherlands (The Dutch College of General Practitioners [NHG]), Germany (the German College of General Practitioners and Family Practitioners [DEGAM]), Belgium (Belgian Society for General Practitioners/Family Physicians [Domus Medica]), the United States (the American Association of Family Practitioners [AAFP]), Canada (the College of Family Physicians of Canada [CFPC]), and Australia (the Royal Australian College of GPs [RACGP]). From these organizations, 60 guidelines were included, 12 of which were in English and all others were translated (from German or Dutch to English) or accessed in English from the organizations website. Figure 2 shows the number of guidelines retrieved from the search for each organization, the majority were from NHG ($n = 38$).

Guideline topics

Across the 6 organizations and 60 included guidelines, topics were distributed across 19 clinical categories (Table

3). Mental health comprised the most guidelines ($n = 7$), addressing broad categories such as anxiety,³² depression,³³ sleep problems,³⁴ and addiction management.^{35,36} Two organizations covered very specific guidelines such as managing medically unexplained physical symptoms and somatoform disorder³⁷ and depression following coronary syndrome.³⁸ In the cardiovascular disease category, chest pain,³⁹ atrial fibrillation,^{40,41} venous leg ulcers,⁴² and venous thromboembolism⁴³ were identified. Atrial fibrillation is one of only 3 de novo guidelines retrieved from the AAFP⁴¹; other AAFP guidelines such as cholesterol management and management of hypertension are produced in collaboration with other specialties or by endorsing external organizations such as the US preventive services task force. Chronic disease management, pain management, and preventive care were the top 3 categories (all methods of production) identified from GP organizations (Appendix 4). All organizations identified in the scoping review produce guidelines by collaboration with external organizations and the RACGP, AAFP, and CFPC all endorse guidelines developed by external organizations.

Methods of development of de novo guidelines

All 60 included guidelines contained explicit descriptions of the evidence-based methods involved in their development. A total of 57 guidelines from the 6 organizations followed a similar process of development, outlined in Table 4. In summary, this process involves topic identification and formation of a working group comprising of GPs and other medical specialists, allied health professionals, scientific experts, and patients as appropriate. Once conflicts of interest are declared, guideline-specific questions are formulated. A systematic literature search is conducted, by either an evidence team or member of the working group, depending on the organization and funds available. For example, the RACGP utilize the PEER team while the NHG utilize a methodologist employed by the college. Following the search, recommendations are developed using an approach such as GRADE, utilizing the evidence synthesized,⁹² however use of the GRADE process




Fig. 1. PRISMA flow diagram of articles retrieved for scoping review on evidence-based guidance documents or clinical guidelines produced by general practice professional organizations, 2010–2021.


may depend on external funding.⁷⁴ The final steps of consensus, review and guideline publication is carried out by all 6 organizations.

There was 1 guide from the RACGP⁹¹ which was distinguished from a guideline, as there was no formulation of weighted recommendations. This was explicitly stated

as part of the method of production. Two guidelines were developed using the ADAPTE process,^{66,93} examining currently available international and/or national guidelines as a basis for recommendations, with the remainder of the process following the organizations de novo production method.

Table 2. Summary of excluded organizations from those contacted.

GP professional organization	Reason for exclusion
Swedish Association of General Practice	Produce guidelines with other specialities
Royal College of General Practitioners	Produce guidelines in association with other national bodies
Hong Kong College of Family Physicians	Do not produce de novo guidelines & regional
Austrian Society of General and Family Physicians	Do not produce de novo guidelines
Greek Association of General Practitioners	Do not produce de novo guidelines & regional
Serbian Medical Society	Do not produce de novo guidelines
Irish College of General Practitioners	Members only access to guidelines
Slovenian Family Medicine Association	Members only access to guidelines



* NHG The Dutch College of General Practitioners; DEGAM The German College of General Practitioners and Family Practitioners; Domus Medica Belgian Society for General Practitioners/Family Physicians; AAFP The American Association of Family Practitioners; CFPC The College of Family Physicians of Canada; RACGP The Royal Australian College of General Practitioners.

Fig. 2. Number of de novo guidelines retrieved from the database search for each general practice professional organization. The number of de novo guidelines found on each organizations website shown for comparison.

Guideline updating processes varied across the 6 organizations. Five of the organizations updated guidelines after a pre-defined time period, ranging from 3–5 years (RACGP, AAFP, and Domus Medica) to 5–10 years (NHG and the CFPC). The guidelines and web resources from DEGAM did not contain this information.^{39,47,49,55,71,76,79,81,83} In addition to the timeframes specified, the 5 organizations state that guideline updating can be prompted by a change in the evidence and/or national guideline updating on a particular topic. The method is a modified version of the original evidence-based approach (Table 4).

Structure and dissemination of guidelines

All 6 organizations make guidelines available in downloadable portable document format (Pdf) directly from their organizations website. The length of guideline Pdf's varied from 30 to 150 pages depending on the topic. Five organizations also provided a summary version of the guideline to accompany the Pdf. DEGAM, NHG, and CFPC provide a summary

infographic.^{39,47,49,55,71,76,79,81,83} Websites of NHG, RACGP, and AAFP provide summary versions, but the full version can only be downloaded as a Pdf. Five organizations provide patient information material. For Modus Medica, only the guideline was available via open access, all other material was accessible only to members. Patient information leaflets differ between organizations, e.g. NHG support a dedicated website for patients (<https://www.gpinfo.nl/>), while the CFPC refers to patient information within their guidelines and also signposts to external patient support groups.

In terms of dissemination, in addition to publishing the full guideline on the organizations website, 3 organizations (CFPC, AAFP, and DEGAM) publish full guidelines, and 1 (NHG) publishes a summary, in their organizations journal. The included Belgium guideline was published in a peer reviewed journal.⁶⁶ Continuing medical education meetings are used for dissemination by CFPC, AAP, NHG, and RACGP. The NHG, RACGP, and CFPC use webinars and e-learning modules, while the NHG and RACGP use periodicals and workshops (Table 5). In the DEGAM guideline on Dementia the strategy for dissemination is described and includes GPs involved in developing the guideline and the publication media drive on completion of a guideline.⁷⁶ Although beyond the scope of this review, of note, clinical decision support systems are used by the CFPC and NHG as a form of implementation, while NHG also use financial incentives and continuing medical education credits.

Discussion

Summary of principal findings

This scoping review included 6 GP professional organizations that publish a range of guidelines for frontline GPs with explicit methods of development and recommendations. Topics cover a wide range of clinical areas including mental health, cardiovascular care, neurology, pregnancy and women's health and preventive care. These guidelines are available on organizations websites as downloadable Pdfs, with summary documents and patient information. Dissemination strategies include peer reviewed publications, webinars, and continuing medical education meetings.

Comparison with previous literature

To our knowledge, this review is the first to map guidelines published by GP professional organizations internationally to support GPs in their clinical decision making. Mental health topics were the top clinical category for de novo guideline production. Mental health conditions are a common presenting condition to general practice⁹⁶ and a more significant challenge now due to the COVID-19 pandemic.⁹⁷ Mental health is a leading cause of significant morbidity⁹⁸ and clinical importance is one of the criteria supporting the need for guideline development.⁹⁹ While, GPs dealing with mental health issues may require access to psychologists and psychiatrists,¹⁰⁰ they remain the first point of care and need diagnostic and therapeutic tools to deal with such presentations. In this context, the updated NHG guidelines on Depression, e.g. now includes these tools as well as website links to online learning modules for GP members to upskill in the delivery of such care.³³ Chronic health conditions were the main clinical category from GP organizations and

Table 3. Summary of guidelines per clinical category.

Mental health (<i>n</i> = 7)	Problem drinking ³⁵	Anxiety ³²	Depression ³³	Medically unexplained physical symptoms ³⁷	Sleep problems ³⁴	Depression post coronary syndrome ³⁸	Opioid use disorder ³⁶
Cardiovascular disease (<i>n</i> = 6)	Atrial fibrillation ^{40,41,44}	Chest pain ³⁹	Venous leg ulcers ⁴²	Venous thromboembolism ⁴³			
Neurology (<i>n</i> = 5)	Facial paralysis ⁴⁵	Head injury ⁴⁶	Stroke ^{47,48}	Acute dizziness ⁴⁹			
Pregnancy and women's health (<i>n</i> = 5)	Preconception care ⁵⁰	Contraception ⁵¹	Menopause ⁵²	Vaginal bleeding ⁵³	Labour and delivery after CS ⁵⁴		
Preventive care (<i>n</i> = 5)	Cardiovascular prevention ⁵⁵	Lipid guidelines ⁵⁶	Risk prevention (SNAP) ⁵⁷	Smoking cessation ⁵⁸	Preventive activities in GP ⁵⁹		
Sexual health and GU (<i>n</i> = 4)	STD consultation ⁶⁰	Female urinary incontinence ⁶¹	Sexual problems ⁶²	Male micturition problems ⁶³			
Paediatrics (<i>n</i> = 4)	ADHD ⁶⁴	Obesity ⁶⁵	Prolonged cough ⁶⁶	Asthma ⁶⁷			
Renal and GIT (<i>n</i> = 4)	Acute diarrhoea ⁶⁸	Gastric symptoms ⁶⁹	Diverticulitis ⁷⁰	Chronic kidney disease ⁷¹			
Musculoskeletal (<i>n</i> = 3)	Lumbosacral radicular syndrome ⁷²	Hand and wrist symptoms ⁷³	Osteoarthritis ⁷⁴				
Care of the elderly (<i>n</i> = 3)	Delirium ⁷⁵	Dementia ^{76,77}					
Respiratory (<i>n</i> = 3)	COPD ⁷⁸	Acute cough ⁷⁹	Asthma ⁸⁰				
Endocrine (<i>n</i> = 2)	Thyroid disorders ^{81,82}						
ENT (<i>n</i> = 2)	Sore throat ^{83,84}						
Rheumatology and allergy (<i>n</i> = 2)	Food allergy ⁸⁵	Polymyalgia rheumatic ⁸⁶					
Dermatology and immunology (<i>n</i> = 1)	Eczema ⁸⁷						
Infectious diseases (<i>n</i> = 1)	Influenza pandemic ⁸⁸						
Ophthalmology (<i>n</i> = 1)	Visual symptoms ⁸⁹						
Cannabinoids	⁹⁰						
Genomics	⁹¹						

topics such as COPD,⁷⁸ asthma,⁸⁰ depression,³³ obesity,⁶⁵ and chronic kidney disease⁷¹ were all included guidelines from the review. The complexity of this care is a considerable challenge for frontline GPs, especially as these patients may suffer from multiple chronic conditions.¹⁴ GPs play a central role in the coordination of this care and for this GP-specific guidance is required.¹⁵ DEGAM and RACGP publish multimorbidity guidelines on their websites. In addition to guidelines, the literature emphasizes the importance of GPs unique relationships with their patients, the need for policies and “models of practice” that allow nurturing of this relationship in order to manage such complex care.¹⁰¹

The findings of this review are consistent with existing literature on guideline development, structure, and updating.^{99,102,103} Each organization has a transparent production method and the de novo process aligns with the Institute of Medicine standards for the development of trustworthy guidelines and the Guideline International Network key components for guideline development.^{104,105} Governance structures and access to government and other national

guideline producing bodies,¹⁹ as well as resources, a need for information sharing and a drive to reduce duplication of effort, all influence the choice of the guideline development process.¹⁰⁶ The review findings reflect this, as in addition to producing de novo guidelines, all included organizations, produce guidelines by collaborating with other specialties and 50% endorse guidelines from other non GP guideline producing organizations. Guidelines are also developed by adaptation of other available guidelines,¹⁰⁷ 2 such guidelines are included in this review.^{59,66} The review findings also support other recent literature from Belgium on the quality of evidence-based medicine resources in primary care,¹⁰⁸ which suggests that being concise, of direct clinical relevance and adapted to the local situation is essential for evidence-based medicine resources.¹⁰⁸

Recent literature on guideline dissemination focussing on the format and language of guidelines outlines the importance of document structure in helping improve their use in practice. Being user friendly, especially relating to the length of the document and how information is visualized is important

Table 4. Summary of methods of guideline development and updating.

Methods	Topic identification	Working group and conflict of interest develop initial draft	Systematic search	Recommendation (level of evidence and strength of recommendations)	Consensus and draft formation	External review	Final draft	Publication	Citations	Update	Update citations
Dutch College of General Practitioners	✓	✓	✓	✓	✓	✓	✓	✓	37,46,50,64,70,73	5–10 years/ evidence	32–35,40,42–45,48,51–53,60–63,65,67–69,72,75,77,78,80,82,84–87,89
College of Family Physicians of Canada	✓	✓	✓	✓	✓	✓	✓	✓	36,56,90	6 years/ evidence	
German College of General Practitioners and Family Physicians	✓	✓	✓	✓	✓	✓	✓	✓	39,49,55,71,76,81,83		47,79
Domus Medica (Belgian College)	✓	✓	✓	✓	✓		✓	✓		5 years	66
American Academy of Family Physicians	✓	✓	✓	✓	✓	✓	✓	✓		5 years/ evidence	47,55,71
Royal Australian College of General Practitioners	✓	✓	✓	✓	✓	✓	✓	✓	91	3–5 years	57–59,74,88

for end-users.¹⁰⁹ Wide variation in the length of guideline pdf documents is evident in this review. Although summary pdfs and summarized website versions of guidelines are available, quick access to pertinent information such as using visual aids and hyperlinks may help to improve their implementation in practice.¹¹⁰ The findings of the review show that multiple dissemination strategies are used, including education tools, infographics, patient information, and a publicity drive at the time of guideline publication. The importance of these strategies is highlighted by the findings of a recent Cochrane review of tools to promote uptake of guidelines, where provision of education materials likely improves adherence to guidelines.¹¹¹

Strengths and limitations

The challenge in this review was to map the landscape of information being produced given the fact that not all GP professional organizations publish their guidelines in peer reviewed journals and also the heterogeneity of the nomenclature for this search. Contacting GP organizations and conducting an extensive grey literature search which included website searches of key organizations helped address this. However, completing a more exhaustive website search (including those organizations whose guidelines were not freely available) and mapping a complete topic list of de novo guidelines from each

organization would allow a more detailed comparison of specific guideline recommendations, although that was outside the scope of this review.

This scoping review was limited to de novo production of guidelines, this excluded certain national GP organizations that develop guidelines in association with national bodies. This decision was based on the varying number of GPs that are members of these national bodies and if those GP members represent their organization. Furthermore, in choosing to map the de novo production of guidelines, the full breadth of activity undertaken by GP organizations in terms of guidelines for use in clinical practice was not captured in this review. A good example of this are the AAFP guidelines on Cholesterol Management and Management of Hypertension, both excluded but very relevant for use in GP. The authors believe it is important to map what guidelines GP professional organizations are producing, given the specific challenges for GPs identifying guidelines that are relevant to general practice, as most are single disease focussed, may not account for the patient perspective, and may not consider the challenges relating to cost and resources for general practice.¹¹²

As this was a scoping review, we did not quality appraise included guidelines so we were not in a position to comment on the overall quality of the included guidelines.

Table 5. Summary of guideline structures and modes of dissemination.

Structure	Pdf	Summary	Patient information leaflet	Education	Audit	Infographics	Citations	Podcast	E-learning module	Periodicals	CME meeting	Faculty study days	Webinars	Workshops	Social media	Newsletter	Peer review publication	
NHG	✓	✓	✓	✓	✓	✓	32-35,37,40,42-46,48,50-53,60-65,67-70,72,73,77,78,80,82,84-87,89,94		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
GFPC	✓	✓	✓	✓	✗	✓	36,56,95	✓	✓	✓	✓		✓				✓	✓
DEGAM	✓	✓	✓	✓	✗	✓	39,47,49,53,71,76,79,81,83										✓	✓
Domus Medica	✓						66										✓	✓
AAFP	✓	✓	✗	✗	✗	✗	38,41,54				✓				✓	✓	✓	✓
RACGP	✓	✓	✗	✗	✗	✗	57-59,74,88,91		✓	✓	✓		✓	✓			✓	✓

Research implications

There is scope for further research to map guidelines being published by GP organizations worldwide regardless of the production method. This would facilitate a broader understanding of the complex nature of guidelines necessary for GPs working on the frontline in any given healthcare system.

Implementation strategies were beyond the scope of this review but we did note that in general these were not identified as part of the guidelines identified. There is a gap in the evidence on which type of implementation strategies are effective,¹¹³⁻¹¹⁵ and although there is some literature in general practice on these strategies,¹¹⁶ there is a need for further research on the type and effectiveness of implementation strategies, e.g. use of financial or other incentives to encourage uptake. International GP organizations would be well placed to collaborate on such research.

Clinical and policy implications

This review shows that international GP organizations, despite working in different healthcare systems, with different governance structures, produce guidelines with similar topics, methods, and dissemination strategies for use by their members. The standard transparent approach to production methods can facilitate a standardization across GP organizations thus promoting sharing of resources, but this needs to be balanced with the need for local relevance to improve the success of guideline implementation.¹¹⁷ Emerging evidence on the effectiveness of guideline implementation in terms of improving patient outcomes, may influence policy around the use of alternative strategies to support the use of guidelines in practice.

Conclusion

This scoping review has highlighted specific de novo guideline production in GP professional organizations worldwide. There is substantial overlap in the areas of methods of production and publication and some variation in clinical topics and dissemination methods. Overall it indicates there is potential for collaboration between GP organizations worldwide to reduce duplication of effort, facilitate reproducibility, and identify areas of standardization internationally. Furthermore, there is an opportunity for leading guideline producing GP organizations to collaborate with countries where the focus remains on secondary/specialist care, helping to strengthen primary care in these countries.

Funding

This work was supported by the Irish College of General Practitioners and Health Research Board of Ireland Applied Partnership Award (APA-2019-001). BC is funded by the HRB Emerging Investigators Award (EIA-2019-09).

Acknowledgements

We thank Paul Murphy, Information specialist, RCSI University of Medicine and Health Sciences, Dublin, Ireland, Ciaran Brown, Medical student, research summer school, RCSI, University of Medicine and Health Sciences, Dublin, Ireland and Afrina Ariff, Medical student, research summer school, RCSI, University of Medicine and Health Sciences, Dublin, Ireland, for their contributions to this study.

Authors' contributions

Each author agrees to be personally responsible for their own contribution. EOB made substantial contribution to the design, acquisition, analysis, interpretation, and drafting of the manuscript, as well as revising and updating the final version for submission. SD made substantial contribution to the acquisition of data and revision of the manuscript and approval of the final version for submission. VH made substantial contribution to the conception and design of the work and approval of the final submitted version of the manuscript. SS made substantial contribution to the design and revising of the manuscript and approval of the final submitted version. NOH made substantial contribution to the design of the work and approval of the final submitted version. AW made substantial contribution to revising the manuscript and approval of the final submitted version. BC made substantial contribution to the design, interpretation, revision, and approval of the final version of the manuscript. EW made substantial contribution to the conception, design, acquisition, analysis, revisions of drafted work, and approval of the final version for submission.

Ethical approval and consent to participate

Ethical approval was not required for this scoping review of GP professional organizations.

Consent for publication

Not applicable.

Conflict of interest

None declared.

Data availability

The datasets generated and/or analysed during the current study are available in the Open Science Framework repository; https://osf.io/cedup/?view_only=12e4fe042a2443dea7345e193a4f5c4d.

Appendix 1. Search strategy

Bibliographic database search

Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations, Daily and Versions(R) 1946 to 12 April 2021		
1	general practice.mp. or exp Family Practice/ or exp General Practice/	97421
2	(general ADJ1 practitioner) OR (general ADJ1 practitioners) OR (family ADJ1 practi?e) OR (family ADJ1 physician*)	127219
3	primary health care.mp. or exp Primary Health Care/	182036
4	1 OR 2 OR 3	311989
5	exp Practice Guideline/OR exp Practice Guidelines as Topic/ OR (practice ADJ2 guideline\$)	166940
6	((quick adj2 reference adj2 guide*) or (quick adj2 reference) or (evidence adj1 reference) or (evidence adj1 guide*) or ("evidence based" adj1 reference) or ("evidence based" adj1 guide)).mp.	1535
7	5 OR 6	168082
8	4 AND 7	12225
9	LIMIT 8 TO 2010–2021	6214
EMBASE		
1	"general practice"/exp OR (general NEXT/1 practice) OR (family NEXT/1 practice) OR (general NEXT/1 practioner\$)	165374
2	"primary medical care"/exp OR (primary NEXT/2 care)	334254
3	1 OR 2	453204
4	"practice guideline"/mj OR (practice NEXT/1 guideline\$)	465331
5	(quick NEXT/1 reference) OR (evidence NEXT/2 reference) OR (evidence NEXT/2 guide*)	18087
6	4 OR 5	475402
7	3 AND 6	28513
8	LIMIT 7 TO 2010–2021	17207
9	LIMIT 8 TO EMBASE ONLY RECORDS, EXCLUDING MEDLINE	5285
COCHRANE LIBRARY		
1	(general NEAR/1 practice) OR (primary NEAR/2 care)	32360
2	practice NEAR/1 guidelin*	13412
3	1 AND 2	1664
4	LIMIT 3 TO REVIEWS OR PROTOCOLS, 2010–2021	260
WEB OF SCIENCE, Science and Social Citation Indexes LIMITED 2010–2021		
1	TS = (general NEAR/1 practice OR primary NEAR/2 care)	110331
2	TS = (practice NEAR/1 guideline*)	31599
3	1 AND 2	1826

Grey literature search

Evidence synthesis

Limited 2010–2021

1396 results for (“general practice” or “family practice” or “general practitioner” or “family practitioner” or “primary healthcare”) and (“practice guidelines” or “evidence based practice”)

17 duplicates removed

Total 1379

Guideline central

Search by Speciality, Family Medicine = 851

Filter by Organisation = American Academy of Family Medicine = 4

Appendix 2. GP organizations questions for supplementary information

- 1 In what country do you work?
 - 2 What general practice/family physician professional organization do you represent? (e.g. Royal College of General Practitioners, Irish College of General Practitioners, The Royal Australian College of General Practitioners, The Dutch College of General Practitioners)
 - 3 Is this general practice/family physician professional organisation a national organisation? Y/N
 - 4 For this scoping review doctors who work in general practice are defined as: “physicians who provide the first point of contact to patients in the community.” They are available for patients of all ages, with all conditions and provide continuity of care. They are primarily responsible for the “provision of comprehensive and continuing care to every individual seeking medical care irrespective of age, sex and illness” (WONCA 2015). Does this definition represent your general practice/family physician professional organisation’s member’s practice of medicine? Y/N
 - 5 *If you answered “No” to Q4, please provide a brief description of how it differs, otherwise write the word “not applicable” in the box to move to the next question.*
 - 6 Gatekeeping is a core function of general practice in some health systems defined as “matching patient’s needs and preferences with the judicious use of medical services.”³ As part of this role, general practitioners authorize access to speciality care, hospital care, and some diagnostic tests. Do the general practice/family physician members of your prof organisation have a gate-keeping role in their practice of medicine?
 - 7 *If you answered “No” to Q6, please provide a brief description of how it differs, otherwise write the word “not applicable” in the box to move to the next question.*
 - 8 Does your general practice/family physician professional organisation produce denovo (new) clinical practice guidelines/guidance documents? Y/N
 - 9 *If you answered “Yes” to Q8, please provide brief details of the process (level of evidence, development team, etc.) involved in the production of these guidelines/guidance documents, otherwise write the word “not applicable” in the box to move to the next question.*
 - 10 How often does your general practice/family physician professional organisation update existing guidelines/guidance documents?
 - 11 Please provide brief details of the process involved in updating existing guidelines, otherwise write the word “not applicable” in the box to move to the next question.
 - 12 Does your general practice/family physician professional organisation adapt existing international guidelines for use within the context of general practice in your healthcare system?
 - 13 Does the general practice/family physician professional organisation endorse national/international guidelines by providing links on the website to these guidelines?
 - 14 *If you answered “Yes” to Q13, what guidelines does your general practice/family physician professional organisation endorse? Please choose as many answers as applicable, or “not applicable” to move to the next question.*
 - 15 Does your general practice/family physician professional organisation allow public/open access to guidelines/guidance documents on their website?
 - 16 If a guideline/guidance document is published in the journal of the GP/FP professional organisation, is this considered a publication of the professional organisation? Y/N
-

- 17 We are seeking a list of clinical guideline/guidance categories (regardless of the production process) available to general practitioners/family physicians in your organisation. Please select from the list below or include other categories if not provided in the list
- Addiction Management
 - Cancer Care
 - Care of the older person
 - Chronic Diseases
 - Domestic Violence
 - Laboratory Investigations
 - Medication review
 - Mental Health
 - Musculoskeletal Medicine
 - Neurology
 - Pandemic and Immunisations
 - Pain Management
 - Paediatric Health
 - Preventive Health
 - Sexual Health
 - Women's Health
 - Other
- 18 For the majority of clinical guidelines/guidance documents produced what additional material is available? Please tick as many answers as applicable.
- Summary document
 - Patient information leaflet
 - Education/Teaching material
 - Audit ideas
 - Infographics
 - Other
- 19 What are the formats and modes of dissemination of the clinical practice guidelines/guidance documents? Please tick as many answers as applicable.
- Portable document format (Pdf)
 - Podcast
 - Peer reviewed publications
 - eLearning modules
 - Periodicals
 - Continued medical education meetings
 - Faculty study days
 - Webinars
 - Workshops
 - Other
- 20 What methods are used for implementation of the guidelines/guidance documents by your general practice/family physician professional organisation? Please tick as many answers as applicable.
- Clinical decision support system
 - Financial incentives
 - Other incentives
 - Other

Appendix 3. Covidence database and grey literature search charted guidelines

https://osf.io/cedup/?view_only=12e4fe042a2443dea7345e193a4f5c4d

Appendix 4. Key Informant Contact responses

https://osf.io/cedup/?view_only=12e4fe042a2443dea7345e193a4f5c4d

References

1. Vanselow NA, Donaldson MS, Yordy KD. A new definition of primary care. *JAMA*. 1995;273(3):192.
2. Wonca Europe. The European Definition of General Practice/Family Medicine. 2005. [accessed 2023 Feb 7]. <https://www.woncaeurope.org/page/definition-of-general-practice-family-medicine>
3. Franks P, Clancy CM, Nutting PA. Gatekeeping revisited—protecting patients from overtreatment. *N Engl J Med*. 1992;327(6):424–429.
4. Stiggelbout AM, Weijden T, Van derWit MPT, DeFrosch D, Légaré F, Montori VM, Trevena L, Elwyn G. et al. Shared decision making: really putting patients at the centre of healthcare. *BMJ*. 2012;344(7842):e256.
5. Légaré F, Adekpedjou R, Stacey D, Turcotte S, Kryworuchko J, Graham ID, Lyddiatt A, Politi MC, Thomson R, Elwyn G, et al. Interventions for increasing the use of shared decision making by healthcare professionals. *Cochrane Database Syst Rev*. 2018;19(7):CD006732. doi: 10.1002/14651858.CD006732.pub4.
6. Austad B, Hetlevik I, Mjølstad BP, Helvik A. et al. Applying clinical guidelines in general practice: a qualitative study of potential complications. *BMC Fam Pract*. 2016;17:92.
7. Biezen R, Roberts C, Buising K, Thursky K, Boyle D, Lau P, Clark M, Manski-Nankervis J-A. How do general practitioners access guidelines and utilise electronic medical records to make clinical decisions on antibiotic use? Results from an Australian qualitative study. *BMJ Open*. 2019;9(8):e028329.
8. Grol R, Dalhuijsen J, Thomas S, Veld C, Rutten G, Mokkink Het al. Attributes of clinical guidelines that influence use of guidelines in general practice: observational study. *BMJ*. 1998;317(7162):858–861.

9. Burgers JS, Grol RP, Zaat JO, Spies TH, van der Bij AK, Mokkink HGet al. Characteristics of effective clinical guidelines for general practice. *Br J Gen Pract.* 2003;53(486):15–19.
10. Lugtenberg M, Burgers JS, Westert GP. Effects of evidence-based clinical practice guidelines on quality of care: a systematic review. *Qual Saf Health Care.* 2009;18(5):385–392.
11. Department of Health, National Clinical Effectiveness Committee. *How to develop a National Clinical Guideline: a manual for guideline developers.* Dublin, Ireland. Department of Health (DoH); 2019.
12. Steel N, Abdelhamid A, Stokes T, Edwards H, Fleetcroft R, Howe A, Qureshi N. A review of clinical practice guidelines found that they were often based on evidence of uncertain relevance to primary care patients. *J Clin Epidemiol.* 2014;67(11):1251–1257.
13. Salisbury C, Johnson L, Purdy S, Valderas JM, Montgomery AA. Epidemiology and impact of multimorbidity in primary care: a retrospective cohort study. *Br J Gen Pract.* 2011;61(582):e12–e21.
14. Wallace E, Salisbury C, Guthrie B, Lewis C, Fahey T, Smith SM et al.. Managing patients with multimorbidity in primary care. *BMJ.* 2015;350:h176.
15. Luijckx H, Lucassen P, van Weel C, Loeffen M, Lagro-Janssen A, Schermer T. How GPs value guidelines applied to patients with multimorbidity: a qualitative study. *BMJ Open.* 2015;5(10):e007905.
16. Muth C, Blom JW, Smith SM, Johnell K, Gonzalez-Gonzalez AI, Nguyen TS, Brueckle MS, Cesari M, Tinetti ME, Valderas JMet al. Evidence supporting the best clinical management of patients with multimorbidity and polypharmacy: a systematic guideline review and expert consensus. *J Intern Med.* 2019;285(3):272–288.
17. Lugtenberg M, Zegers-van Schaick JM, Westert GP, Burgers JS et al. Why don't physicians adhere to guideline recommendations in practice? An analysis of barriers among Dutch general practitioners. *Implement Sci.* 2009;4(1):54.
18. Abdelhamid A, Howe A, Stokes T, Qureshi N, Steel N. Primary care evidence in clinical guidelines: a mixed methods study of practitioners' views. *Br J Gen Pract.* 2014;64(628):e719–e727.
19. Legido-Quigley H, Panteli D, Brusamento S, Knai C, Saliba V, Turk E, Solé M, Augustin U, Car J, McKee M, et al.. Clinical guidelines in the European Union: mapping the regulatory basis, development, quality control, implementation and evaluation across member states. *Health Policy.* 2012;107(2):146–156.
20. Allan GM, Aubrey-Bassler K, Cauchon M, Ivers NM, Katz A, Kirkwood J, Kuling PJ, Mang EJ, Moore S, Safarov A, et al. Endorsement of clinical practice guidelines: criteria from the College of Family Physicians of Canada. *Can Fam Physician.* 2021;67(7):499–502.
21. Graham ID, Harrison MB. Evaluation and adaptation of clinical practice guidelines. *Evid Based Nurs.* 2005;8(3):68–72.
22. Roberge P, Fournier L, Brouillet H, Delorme A, Beaucage C, Côté R, Demers P, Gervais M, Laflamme F, Latulippe L, et al. A provincial adaptation of clinical practice guidelines for depression in primary care: a case illustration of the ADAPTE method: depression guideline adaptation with ADAPTE method. *J Eval Clin Pract.* 2015;21(6):1190–1198.
23. Fervers B, Burgers JS, Haugh MC, Latreille J, Mlika-Cabanne N, Paquet L, Coulombe M, Poirier M, Burnand B. Adaptation of clinical guidelines: literature review and proposition for a framework and procedure. *Int J Qual Health Care.* 2006;18(3):167–176.
24. Grol R, Jones R. Twenty years of implementation research. *Fam Pract.* 2000;17(Suppl 1):S32–S35.
25. O'Brien E, Clyne B, Smith SM, O'Herlihy N, Harkins V, Wallace E et al. A scoping review protocol of evidence-based guidance and guidelines published by general practitioner professional organisations. *HRB Open Res.* 2021;4:53.
26. Peters MDJ, Marnie C, Tricco AC, Pollock D, Munn Z, Alexander L, McInerney P, Godfrey CM, Khalil H et al. Updated methodological guidance for the conduct of scoping reviews. *JBI Evid Synth.* 2020;18(10):2119–2126.
27. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol.* 2005;8(1):19–32.
28. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci.* 2010;5(1):69.
29. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MD, Horsley T, Weeks L, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med.* 2018;169(7):467–473.
30. Peters MD, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB et al. Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc.* 2015;13(3):141–146.
31. Micah DJ, Peters CG, Patricia M, Zachary M, Andrea CT, Hanan K. Chapter 11: Scoping Reviews (2020 version). In: Aromataris E, Munn Z (Editors). *JBI Manual for Evidence Synthesis*, JBI, 2020. Available from <https://synthesismanual.jbi.global>. <https://doi.org/10.46658/JBIMES-20-12>.
32. van Avendonk MJP, Hassink-Franke LJA, Terluin B, van Marwijk HWJ, Wiersma T, Burgers JS et al. [Summarisation of the NHG practice guideline 'Anxiety']. Samenvatting van de NHG-standaard 'Angst'. *Ned Tijdschr Geneesk.* 2012;156(34):A4509.
33. van Avendonk, M.J.P, van Weel-Baumgarten EM, van der Weel GM, Wiersma T, Burgers JSet al. [Summary of the Dutch College of General Practitioners' practice guideline 'Depression']. Samenvatting van de NHG-Standaard 'Depressie'. *Ned Tijdschr Geneesk.* 2012;156(38):A5101.
34. Damen-van Beek Z, Lucassen PLBJ, Gorgels W, Smelt AFH, Knuistingh Neven A, Bouma Met al. [The NHG guideline 'Sleep problems and sleeping pills']. De NHG-standaard 'Slaapproblemen en slaappmiddelen'. *Ned Tijdschr Geneesk.* 2015;159:A8679.
35. Sijborn M, Luijckx H, Boomsma L, Larsen IM, Burgers J, van der Weel G, et al. [The Dutch College of General Practitioners' practice guideline 'Problem drinking']. De NHG-standaard 'Problematisch alcoholgebruik'. *Ned Tijdschr Geneesk.* 2015;159:A8646.
36. Korownyk C, Perry D, Ton J, Kolber MR, Garrison S, Thomas B, Allan GM, Bateman C, de Queiroz R, Kennedy D, et al.. Managing opioid use disorder in primary care: PEER simplified guideline. *Can Fam Physician.* 2019;65(5):321–330.
37. van der Feltz-Cornelis CM, Swinkels JA, Blankenstein AHN, Hoedeman R, Keuter EJW, Werkgroep Somatoforme klachten en stoornissen et al. [The Dutch multidisciplinary guideline entitled 'Medically unexplained physical symptoms and somatoform disorder']. De Nederlandse multidisciplinaire richtlijn 'Somatisch onvoldoende verklaarde lichamelijke klachten en somatoforme stoornissen'. *Ned Tijdschr Geneesk.* 2011;155(18):A1244.
38. Frost JL, Rich RL, Jr, Robbins CW, Stevermer JJ, Chow RT, Leon KK, Bird MD. et al. Depression following acute coronary syndrome events: screening and treatment guidelines from the AAFP. *Am Fam Physician.* 2019;99(12).
39. Haasenritter J, Bösner S, Klug J, Ledig T, Donner-Banzhoff N et al. DEGAM-Guideline Number 15: chest pain in primary care. *Z Allgemeinmed.* 2011;87(4):182–191.
40. van den Donk M, Opstelten W; P. Dutch College of General. [Important changes in the Dutch College of General Practitioners (NHG) practice guideline 'Atrial fibrillation': revised practice guideline issued]. Belangrijke wijzigingen in de NHG-standaard 'Atriumfibrilleren': herziene versie verschenen. *Ned Tijdschr Geneesk.* 2013;157(38):A6697.
41. Hauk L. Newly detected atrial fibrillation: AAFP updates guideline on pharmacologic management. *Am Fam Physician.* 2017;96(5):332–333.
42. van Rijn-van Kortenof NMM, Wiersma T, van Hof N, Balak FSR, GoudswaardLANet et al. [Summary of the practice guideline 'Venous leg ulcer' from the Dutch College of General Practitioners]. Samenvatting van de standaard 'Ulcus cruris venosum' van het Nederlands Huisartsen Genootschap. *Ned Tijdschr Geneesk.* 2011;155:A3158.
43. de Jong J, Lucassen WAM, Geersing G-J, Burgers JS, and Wiersma T et al. [NHG guideline 'Deep-vein thrombosis and pulmonary embolus']. NHG-standaard 'Diepe veneuze trombose en longembolie'. *Ned Tijdschr Geneesk.* 2015;159:A8657.
44. Opstelten W, Boode, BSP, Heeringa JR, Frans H, Goudswaard ANLet et al. [Summary of the practice guideline 'Atrial fibrillation'

- (first revision) from the Dutch College of General Practitioners]. Samenvatting van de standaard 'Atriumfibrilleren' (eerste herziening) van het Nederlands Huisartsen Genootschap. *Ned Tijdschr Geneeskd.* 2010;154:A1570.
45. Klomp MAR, Verdaasdonk AL, Striekwold MP, Teunissen HE, Opstelten W, Goudswaard ANLet al. [Dutch College of General Practitioners' practice guideline 'Peripheral facial paralysis': a summary]. De NHG-standaard 'Perifere aangezichtsverlamming': samenvatting. *Ned Tijdschr Geneeskd.* 2010;154:A2439.
 46. Draijer LW, Kurver MJ, Opstelten W. [The NHG practice guideline 'Head injury']. De NHG-standaard 'Hoofdtrauma'. *Ned Tijdschr Geneeskd.* 2015;159:A8992.
 47. Barzel A, Hensler S, Konecny N, Muche-Borowski C, Scherer Met al. Updated DEGAM guideline stroke—what's new? *Z Allgemeinmed.* 2014;90(4):152–157.
 48. Verburg AFEA, Tjon-A-Tsien MRS, Verstappen WHJM, Beusmans GHMI, Wiersma T, and Burgers JSet al. [Summary of the 'Stroke' guideline of the Dutch College of General Practitioners' (NHG)]. Samenvatting van de NHG-standaard 'Beroerte'. *Ned Tijdschr Geneeskd.* 2014;158(1):A7022.
 49. Jendyk RM, Abholz HH. Acute dizziness in primary care offices—The German College of General Practitioners and family Physicians' new guideline. *Z Allgemeinmed.* 2017;93(1):12–16.
 50. van Asselt KM, de Jong-Potjer L, Beentjes M, Wiersma T, Goudswaard ANLet al. [Summary of the NHG Standard 'Preconception care']. Samenvatting van de NHG-standaard 'Preconceptiezorg'. *Ned Tijdschr Geneeskd.* 2012;155(35):A4680.
 51. Kurver MJ, van der Wijden CL, Burgers J. [Summary of the Dutch College of General Practitioners' practice guideline 'Contraception']. Samenvatting van de NHG-standaard 'Anticonceptie'. *Ned Tijdschr Geneeskd.* 2012;156(41):A5083.
 52. de Vries CJH, Eekhof H, Engel HF, Opstelten W, Burgers JSet al. [Summary of the Dutch College of General Practitioners' (NHG) practice guideline 'The menopausal transition']. Samenvatting van de NHG-standaard 'De overgang'. *Ned Tijdschr Geneeskd.* 2012;156(44):A5326.
 53. de Vries CJH, Meijer LJ, Janssen CAHI, Burgers JS, Opstelten Wet al. [Dutch College of General Practitioners' practice guideline on 'Vaginal bleeding']. NHG-standaard 'Vaginaal bloedverlies'. *Ned Tijdschr Geneeskd.* 2015;159:A8534.
 54. Hauk L. Planning for labor and vaginal birth after cesarean delivery: guidelines from the AAFP. *Am Fam Physician.* 2015;91(3):197–198.
 55. Egidi G, Ludt S, Angelow A, Baum E, Chenot JF, Feßler J, Haasenritter J, Donner-Banzhoff N, Popert Uet al. The new DEGAM guideline 'counseling on cardiovascular prevention in family medicine'. *Z Allgemeinmed.* 2017;93(7–8):312–318.
 56. Allan GM, Lindblad AJ, Comeau A, Coppola J, Hudson B, Mannarino M, McMinis C, Padwal R, Schelstraete C, Zarnke K, et al.. Simplified lipid guidelines: prevention and management of cardiovascular disease in primary care. *Can Fam Physician.* 2015;61(10):857–867.
 57. Harris M, Stocks N, Zwar N, Mazza D, Winzenberg T, Booth K, McPherson M. Smoking, nutrition, alcohol, physical activity (SNAP): a population health guide to behavioural risk factors in general practice. 2nd ed. Melbourne. Royal Australian College of General Practitioners; 2015.
 58. Zwar N, Richmond R, Borland R, Peters M, Litt J, Bell J, Caldwell B, Ferretter I. *Supporting smoking cessation: a guide for health professionals.* Melbourne. Royal Australian College of General Practitioners; 2014.
 59. *The Royal Australian College of General Practitioners. Guidelines for preventive activities in general practice.* 9th ed. East Melbourne, Vic. Royal Australian College of General Practitioners; 2016.
 60. Verlee EL, van Bergen JEAM, Dekker JH, Boeke AJPJ, Burgers JS, Bouma Met al. [Summary of the NHG guideline 'The STD consultation']. Samenvatting van de NHG-standaard 'Het soa-consult'. *Ned Tijdschr Geneeskd.* 2014;158:A7277.
 61. Damen-van Beek Z, Teunissen D, Dekker JH, Lagro-Janssen ALM, Berghmans LCM, Uijen JHJM, Mientjes GHC, Wiersma Tet al. [Practice guideline 'Urinary incontinence in women' from the Dutch College of General Practitioners]. NHG-standaard 'Incontinentie voor urine bij vrouwen'. *Ned Tijdschr Geneeskd.* 2016;160:D674.
 62. de Jong J, Leusink P, Wiersma T. [The Dutch College of General Practitioners practice guideline on 'Sexual problems']. NHG-standaard 'Seksuele klachten'. *Ned Tijdschr Geneeskd.* 2016;160:A9707.
 63. Blanker MH, de Reijke TM, van Moorselaar RJA, Opstelten Wet al. [Changes to Dutch College of General Practitioners guideline 'Micturition problems in men']. Beleid vroegdiagnostiek prostaatcarcinoom aangepast. NHG-standaard adviseert vaker verwijzen, minder invasief onderzoek. *Ned Tijdschr Geneeskd.* 2014;158:A8070.
 64. van Avendonk MJP, et al. [Summary of the Dutch College of General Practitioners' (NHG) practice guideline 'ADHD in children']. Samenvatting van de NHG-standaard 'ADHD bij kinderen'. *Ned Tijdschr Geneeskd.* 2015;159:A8395.
 65. van Avendonk MJP, Hassink-Franke LJA, Stijntjes F, Wiersma T, Burgers JSet al. [Summary of the Dutch College of General Practitioners' practice guidelines on obesity]. Samenvatting van de standaard 'Obesitas' van het Nederlands Huisartsen Genootschap. *Ned Tijdschr Geneeskd.* 2010;154:A2834.
 66. Lecomte S, Valentin S, Dromelet E, de Jonghe Met al. Prolonged cough in pediatric population first line care, Belgian guidelines. *Open Respir Med J.* 2017;11:54–66.
 67. Bindels PJE, van de Griendt E-J, Tuut MK, Steenkamer TA, Uijen JHJM, and Geijer RMMet al. [Dutch College of General Practitioners' practice guideline 'Asthma in children']. NHG-standaard 'Astma bij kinderen'. *Ned Tijdschr Geneeskd.* 2014;158:A7935.
 68. Loogman MCM, Bouma M, Burgers JS. [Practice guideline on 'Acute diarrhoea' from the Dutch College of General Practitioners]. NHG-standaard 'Acute diarree'. *Ned Tijdschr Geneeskd.* 2014;158:A8659.
 69. de Jongh E, Numans ME, de Wit NJ, Heemstra-Borst CG, Geijer RMM, Burgers JSet al. [Summary of the Dutch College of General Practitioners' (NHG) practice guideline 'Gastric symptoms']. Samenvatting van de NHG-standaard 'Maagklachten'. *Ned Tijdschr Geneeskd.* 2013;157(22):A6101.
 70. Berger M, de Wit N, Vogelenzang R, Wetzels R, Kortenhof NR, Opstelten W. The NHG guideline diverticulitis. *Huisarts Wet.* 2011;54(9):492–499.
 71. Weckmann G, Chenot J-F, Stracke S. The management of non-dialysis-dependent chronic kidney disease in primary care. *Dtsch Arztebl Int.* 2020;117(44):745–751.
 72. Verburg AFEA, Schaafstra A, Spinnewijn WEM, Kroes BW, Bouma M, Burgers JSet al. [Revised NHG practice guideline 'Lumbosacral radicular syndrome']. Herziene NHG-standaard 'Lumbosacraal radiculair syndroom'. *Ned Tijdschr Geneeskd.* 2015;159:A9122.
 73. Geraets JJXR, Geijer RMM, Goudswaard ANL. [Dutch College of General Practitioners' practice guideline on symptoms of the hand and wrist: a summary]. De NHG-standaard 'Hand-en polsklachten': samenvatting. *Ned Tijdschr Geneeskd.* 2010;154:A1795.
 74. The Royal Australian College of General Practitioners. *Guideline for the management of knee and hip osteoarthritis: communication, implementation and dissemination plan.* East Melbourne, Vic. 2018.
 75. van der Weele GM, Eizenga WH, Dautzenberg PLJ, Wiersma T, Burgers JSet al. [Practice guideline 'Delirium' from the Dutch College of General Practitioners]. NHG-standaard 'Delier'. *Ned Tijdschr Geneeskd.* 2014;158:A7766.
 76. Vollmar HC, Abholz H-H, Egidi G, Mand P, Butzlaff ME, Wilm Set al. [Development and implementation of the dementia guideline of the German Society of General Practice and Family Medicine]. Entwicklung und Implementierung der Leitlinie 'Demenz' der Deutschen Gesellschaft für Allgemeinmedizin und Familienmedizin. *Med Klin.* 2010;105:41–47.
 77. Luning-Koster M, Perry M, Moll van Charante EP, Vernooij-Dassen M, Wiersma T, Burgers JS, et al [Summary of Dutch College of General Practitioners' (NHG) practice guideline 'Dementia'].

- Samenvatting van de NHG-standaard 'Dementie'. *Ned Tijdschr Geneeskd.* 2012;156(49):A5323.
78. van den Aardweg JG. [The revised NHG guideline 'COPD': a new approach with old limitations]. De herziene NHG-standaard 'COPD'. Een nieuwe aanpak met oude beperkingen. *Ned Tijdschr Geneeskd.* 2015;159:A9056.
 79. Holzinger F, Beck S, Dini L, Stoter C, Heintze C et al. The diagnosis and treatment of acute cough in adults. *Dtsch Arztebl Int.* 2014;111(20):356–363.
 80. Geijer RMM, Tuut MK, in't Veen JCCM, Broekhuizen BDL, Chavannes NH, Smeele IJMet al. [The NHG guidelines 'Adult asthma' and 'COPD']. NHG-standaarden 'Astma bij volwassenen' en 'COPD'. *Ned Tijdschr Geneeskd.* 2015;159:A9076.
 81. Voigt K, Bergmann A, Bründel KH, Schübel J et al. Management of elevated TSH levels in adult patients in primary care: introduction of the New Guideline (S2k) of the German college of general practitioners and family physicians (DEGAM). *Z Allgemeinmed.* 2017;93(4):150–155.
 82. Sijbom M, van Lieshout J, Felix-Schollaart B, Burgers JS, Bouma Met al. [Summary of the 'Thyroid disorders' guideline of the Dutch College of General Practitioners (NHG)]. Samenvatting van de NHG-standaard 'Schildklierandoeningen'. *Ned Tijdschr Geneeskd.* 2013;157(35):A6667.
 83. Wachtler H, Chenot JF et al. [Guidelines for the management of sore throat from the German Society of General Practice and Family Medicine]. Die Halsschmerz-Leitlinie der Deutschen Gesellschaft für Allgemeinmedizin und Familienmedizin. *HNO.* 2011;59:480–484.
 84. de Jongh E, Opstelten W, Werkgroep NHGSA. [Revision of the Dutch College of General Practitioners practice guideline 'Acute sore throat']. Herziening van de NHG-standaard 'Acute keelpijn'. *Ned Tijdschr Geneeskd.* 2015;159:A9456.
 85. Luning-Koster MN, Lucassen PLBJ, Boukes FS, Goudswaard ANLet al. [Summary of the Dutch College of General Practitioners' practice guideline on food hypersensitivity]. Samenvatting van de NHG-Standaard 'Voedselovergevoeligheid'. *Ned Tijdschr Geneeskd.* 2011;155(18):A3063.
 86. Eizenga WH, Hakvoort L, Dubbeld P, Geijer RMM, Goudswaard ANLet al. [Dutch College of General Practitioner's practice guideline on polymyalgia rheumatica and temporal arteritis]. Samenvatting van de NHG-standaard 'Polymyalgia rheumatica en arteriitis temporalis'. *Ned Tijdschr Geneeskd.* 2010;154:A1919.
 87. de Vries CJH, de Witt-de Jong AWF, Dirven-Meijer PC, Burgers JS, Opstelten W et al. [The Dutch College of General Practitioners practice guideline 'Eczema']. NHG-standaard 'Eczem'. *Ned Tijdschr Geneeskd.* 2014;158:A8009.
 88. *The Royal Australian College of General Practitioners. Managing pandemic influenza in general practice: a guide for preparation, response and recovery.* East Melbourne, Vic. Royal Australian College of General Practitioners; 2014.
 89. Eizenga WE, Blom GH, Wiersma T. [The revised Dutch College of General Practitioners (NHG) practice guideline on 'Visual symptoms']. De NHG-standaard 'Visusklachten' herzien. *Ned Tijdschr Geneeskd.* 2016;160:D242.
 90. Allan GM, Ramji J, Perry D, Ton J, Beahm NP, Crisp N, Dockrill B, Dubin RE, Findlay T, Kirkwood J, et al. Simplified guideline for prescribing medical cannabinoids in primary care. *Can Fam Physician.* 2018;64(2):111–120.
 91. *The Royal Australian College of General Practitioners. Genomics in general practice.* East Melbourne, Vic. Royal Australian College of General Practitioners; 2018.
 92. Neumann I, Santesso N, Akl EA, Rind DM, Vandvik PO, Alonso-Coello P, Agoritsas T, Mustafa RA, Alexander PE, Schünemann H, et al. A guide for health professionals to interpret and use recommendations in guidelines developed with the GRADE approach. *J Clin Epidemiol.* 2016;72:45–55.
 93. Campos-Outcalt D. 8 USPSTF recommendations FPs need to know about. *J Fam Pract.* 2016;65:338–341.
 94. Geraets JJXR, Geijer RMM, Goudswaard ANLet al. [Dutch College of General Practitioners' practice guideline on symptoms of the hand and wrist: a summary]. De NHG-standaard 'Hand- en polsklachten': samenvatting. *Ned Tijdschr Geneeskd.* 2010;154:A1795.
 95. Allan GM, Ramji J, Perry D, Ton J, Beahm NP, Crisp N, Dockrill B, Dubin RE, Findlay T, Kirkwood J, et al. Simplified guideline for prescribing medical cannabinoids in primary care. *Can Fam Physician.* 2018;64(2):111–120.
 96. Cooke G, Valenti L, Glasziou P, Britt H. Common general practice presentations and publication frequency. *Aust Fam Physician.* 2013 Jan-Feb;42(1-2):65-8. .
 97. Xiong J, Lipsitz O, Nasri F, Lui LM, Gill H, Phan L, Chen-Li D, Iacobucci M, Ho R, Majeed A, et al. Impact of COVID-19 pandemic on mental health in the general population: a systematic review. *J Affect Disord.* 2020;277:55–64.
 98. Demyttenaere K, Bruffaerts R, Posada-Villa J, Gasquet I, Kovess V, Lepine JP, Angermeyer MC, Bernert S, de Girolamo G, Morosini P, et al. Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *JAMA.* 2004;291(21):2581–2590.
 99. Eccles MP, Grimshaw JM, Shekelle P, Schünemann HJ, Woolf S et al. Developing clinical practice guidelines: target audiences, identifying topics for guidelines, guideline group composition and functioning and conflicts of interest. *Implement Sci.* 2012;7(1):60.
 100. Fleury M-J, Imboua A, Aubé D, Farand L, Lambert Y et al. General practitioners' management of mental disorders: a rewarding practice with considerable obstacles. *BMC Fam Pract.* 2012;13(1):19.
 101. Damarell RA, Morgan DD, Tieman JJ. General practitioner strategies for managing patients with multimorbidity: a systematic review and thematic synthesis of qualitative research. *BMC Fam Pract.* 2020;21(1):131.
 102. Shekelle P, Woolf S, Grimshaw JM, Schünemann HJ, Eccles MP et al. Developing clinical practice guidelines: reviewing, reporting, and publishing guidelines; updating guidelines; and the emerging issues of enhancing guideline implementability and accounting for comorbid conditions in guideline development. *Implement Sci.* 2012;7:62.
 103. Woolf S, Schünemann HJ, Eccles MP, Grimshaw JM, Shekelle P et al. Developing clinical practice guidelines: types of evidence and outcomes; values and economics, synthesis, grading, and presentation and deriving recommendations. *Implement Sci.* 2012;7:61.
 104. Institute of Medicine (US) Committee to Advise the Public Health Service on Clinical Practice Guidelines. *Clinical Practice Guidelines: Directions for a New Program.* Field MJ, Lohr KN, editors. Washington (DC): National Academies Press (US); 1990.
 105. Qaseem A, Forland F, Macbeth F, Ollenschläger G, Phillips S, van der Wees P et al. Guidelines international network: toward international standards for clinical practice guidelines. *Ann Intern Med.* 2012;156(7):525–531.
 106. Schünemann HJ, Wiercioch W, Brozek J, Etzeandía-Ikobaltzeta I, Mustafa RA, Manja V, Brignardello-Petersen R, Neumann I, Falavigna M, Alhazzani W, et al. GRADE Evidence to Decision (EtD) frameworks for adoption, adaptation, and de novo development of trustworthy recommendations: GRADE-ADOLOPMENT. *J Clin Epidemiol.* 2017;81:101–110.
 107. Dizon JM, Machingaidze S, Grimmer K. To adopt, to adapt, or to contextualise? The big question in clinical practice guideline development. *BMC Res Notes.* 2016;9(1):442.
 108. de Fernelmont L, Laere SV, Devroey D. The quality of EBM sources perceived by Belgian family physicians. *Open Access Maced J Med Sci.* 2018;6(10):1918–1923.
 109. Gupta S, Rai N, Bhattacharaya O, Cheng AYY, Connelly KA, Boulet L-P, Kaplan A, Brouwers MC, Kastner Met al. Optimizing the language and format of guidelines to improve guideline uptake. *CMAJ.* 2016;188(14):E362–E368.
 110. Moran GM, Bakhai C, Song SH, Agwu JC et al. Type 2 diabetes: summary of updated NICE guidance. *BMJ.* 2022;377:o775.

111. Flodgren G, Hall AM, Goulding L, Eccles MP, Grimshaw JM, Leng GC, Shepperd S et al. Tools developed and disseminated by guideline producers to promote the uptake of their guidelines. *Cochrane Database Syst Rev*. 2016;(8):Cd010669.
112. Carlsen B, Bringedal B. Attitudes to clinical guidelines—do GPs differ from other medical doctors? *BMJ Qual Saf*. 2011;20(2):158–162.
113. Grimshaw J, Eccles M, Thomas R, MacLennan G, Ramsay C, Fraser C, Vale L. Toward evidence-based quality improvement. Evidence (and its limitations) of the effectiveness of guideline dissemination and implementation strategies 1966–1998. *J Gen Intern Med*. 2006;21(Suppl 2):S14–S20.
114. Brusamento S, Legido-Quigley H, Panteli D, Turk E, Knai C, Saliba V, Car J, McKee M, Busse R, et al. Assessing the effectiveness of strategies to implement clinical guidelines for the management of chronic diseases at primary care level in EU Member States: a systematic review. *Health Policy*. 2012;107(2):168–183.
115. Foy R, Willis T, Glidewell L, McEachan R, Lawton R, Meads D, Collinson M, Hunter C, Hulme C, West R, et al. Programme grants for applied research. In: *Developing and evaluating packages to support implementation of quality indicators in general practice: the ASPIRE research programme, including two cluster RCTs*. Southampton (UK): NIHR Journals Library, March 2020.
116. Unverzagt S, Oemler M, Braun K, Klement A. Strategies for guideline implementation in primary care focusing on patients with cardiovascular disease: a systematic review. *Fam Pract*. 2013;31(3):247–266.
117. Wang Z, Norris SL, Bero L. The advantages and limitations of guideline adaptation frameworks. *Implement Sci*. 2018;13(1):72.