

CORRECTIONS

Hyaluronan-binding region of aggrecan from pig laryngeal cartilage

Amino acid sequence, analysis of N-linked oligosaccharides and location of the keratan sulphate

F. P. BARRY, J. U. GAW, C. N. YOUNG and P. J. NEAME

Volume 286 (1992), pages 761–769

The following error has been found in the reported sequence.

page 762, Table 1: peptide T2; the sequence should read:

VLLGGSLTIPCYFIDPMHPVXTAPXTAPLAP

page 764, Fig. 2: residue 26 should be changed from threonine to glycine.

page 767, Fig. 9: the sequence in the top line should read:

...LLGGSLTIPC...

The corresponding regions in rat and human HABR have a threonine where a glycine occurs in the porcine sequence and would therefore read:

...LLGTSLTIPC...

page 768, column 2, line 23, should read: This cleavage occurred at the Asp³⁴¹-Phe³⁴² bond, which is 11 residues *N*-terminal to the site of KS attachment.

Molecular cloning and characterization of the promoter region of the calcineurin A α gene

C.-D. CHANG, T. TAKEDA, H. MUKAI, H. SHUNTOH, T. KUNO and C. TANAKA

Volume 288 (1992), pages 801–805

page 801: the accession number given in the final line of the footnotes is incorrect. The correct accession number is D10480.

Ca²⁺/calmodulin-dependent nitric oxide synthase activity in the human cervix carcinoma cell line ME-180

G. WERNER-FELMAYER, E. R. WERNER, D. FUCHS, A. HAUSEN, B. MAYER, G. REIBNEGGER, G. WEISS and H. WACHTER

Volume 289 (1993), pages 357–361

page 359, Table 2: the dilution of rabbit non-immune serum should be 1:10, not 1:1000.