

# Food-sourcing from on-farm trees mediates positive relationships between tree cover and dietary quality in Malawi

Received: 1 May 2023

Accepted: 11 July 2024

Published online: 13 August 2024

 Check for updates

Emilie Vansant<sup>1</sup>✉, Bowy den Braber<sup>1</sup>, Charlotte Hall<sup>1,2</sup>, Judith Kamoto<sup>3</sup>, Florian Reiner<sup>1</sup>, Johan Oldekop<sup>1,4</sup>✉ & Laura Vang Rasmussen<sup>1</sup>✉

Food security policies often overlook the potential of trees to provide micronutrient-rich foods. Here, through causal mediation analysis, we show the positive effect of tree cover on micronutrient adequacy, explained by people sourcing food from on-farm trees. Detailed survey data ( $n = 460$  households with repeated surveys) from Malawi were linked to high-resolution (3 m) tree-cover data to capture forest and non-forest trees. Our findings support integrating nutrition and landscape restoration policies.

Although global hunger rates have decreased in recent decades, one in three people experience some form of malnutrition, with the highest rates in low- and middle-income countries (LMICs)<sup>1</sup>. Trees are an important source of micronutrient-rich fruits, nuts, seeds and leafy vegetables, and there is increasing recognition of how forests can contribute to dietary quality through the provision of wild foods, income-generating products and ecosystem services that boost agricultural production<sup>2</sup>. However, less is known about the contribution of non-forest trees to improving diets in LMICs, despite the widespread prevalence of trees in agricultural landscapes. The few studies that successfully draw these links are restricted by the use of broad dietary quality indicators (for example, dietary diversity scores, food frequency questionnaires) and therefore do not indicate the degree to which trees, both in forests and outside of forests, support people's intake of key micronutrients.


In this study, we focus on Malawi, a densely populated country with high rates of poverty and malnutrition, where 51% of the population are classified as severely food insecure and 37% of children under 5 years are stunted<sup>3</sup>. Dependence on subsistence agriculture and wood-sourced fuel has contributed to high deforestation rates in recent decades, with a net loss of 224,000 ha of tree cover (15%) between 2000 and 2022 (ref. 4). These combined factors make Malawi an optimal site to study tree–diet linkages. Based on detailed surveys conducted with 460 women in the northern and southern parts of the country (Extended Data Fig. 1), we examine both the effect of overall tree cover and the

effect of sourcing food from on-farm trees on women's micronutrient adequacy (zinc, vitamin A, iron and folate). We conduct a causal mediation analysis to evaluate how the effect of tree cover on diets might be mediated by whether people obtain food from on-farm trees. Causal mediation analysis allows us to measure the extent to which our mediator (sourcing food from trees on farms) transmits the effect of our treatment variable (tree cover) on our dependent variables (zinc, vitamin A, iron and folate adequacy). This approach advances existing knowledge of the positive association between forests and dietary quality<sup>2</sup> by measuring how much of the tree cover–diet relationship can be explained by the sourcing of food from on-farm trees. Moreover, we advance previous studies by going beyond simple nutrition metrics and measuring women's micronutrient adequacy.

We base these analyses on an extensive, interdisciplinary dataset combining (1) repeated 24 h dietary recall surveys (on two non-consecutive days within a 7 day period) in both the dry and wet seasons, (2) survey data on households' decisions to use trees on and around their farmland for food provision, and (3) tree-cover estimates from 2019 PlanetScope imagery (3 m resolution)<sup>5</sup> within each household's surroundings.

Our causal mediation analysis (Fig. 1) reveals four key findings: (1) the amount of tree cover around households is positively associated with women's micronutrient adequacy levels (path C); (2) greater tree cover also increases the likelihood of whether households source food from their on-farm trees (path A); (3) sourcing food from on-farm trees

<sup>1</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark. <sup>2</sup>Biological and Environmental Sciences, University of Stirling, Stirling, UK. <sup>3</sup>Forestry Department, Lilongwe University of Agriculture and Natural Resources, Lilongwe, Malawi. <sup>4</sup>Global Development Institute, University of Manchester, Manchester, UK. ✉e-mail: [ecv@ign.ku.dk](mailto:ecv@ign.ku.dk); [Johan.Oldekop@manchester.ac.uk](mailto:Johan.Oldekop@manchester.ac.uk); [lr@ign.ku.dk](mailto:lr@ign.ku.dk)



**Fig. 1 | Hypothesized pathways linking tree cover, sourcing food from on-farm trees and people's dietary quality.** The causal mediation analysis<sup>21</sup> consists of three pathways (corresponding to arrows A, B and C), where the effect sizes for each pathway are used to measure the degree to which sourcing food

from on-farm trees (M) mediates the relationship between tree cover (X) and micronutrient adequacy (Y). <sup>a</sup>2019 PlanetScope data via Reiner et al.<sup>5</sup> <sup>b</sup>Household survey data.

improves women's micronutrient adequacy (path B); and (4) the effect of tree cover on women's adequacy levels for certain micronutrients is partly mediated by use of trees on farms as a food source (paths A + B). We control for the following variables in our models: household size, the Multidimensional Poverty Index (living standards dimension)<sup>6</sup>, farm size (area under cultivation), education level, livestock holdings (tropical livestock units)<sup>7</sup>, crop count and study region. This selection of covariates was informed by a synthesis of studies linking trees and dietary quality<sup>8</sup> as well as extensive fieldwork in Malawi.

We find positive statistically significant associations between the percentage of tree cover in households' surroundings and women's micronutrient adequacy (Fig. 1, path C). For example, we see that a 1% increase in tree cover is associated with a 0.25–0.37% average increase in iron adequacy rates, depending on the season. A jump from 0% to 62% tree cover (maximum observed) corresponds to an increase of 16–23% in overall iron adequacy (dry season,  $P = 0.0192$ ; wet season,  $P = 0.0019$ ) (Fig. 2a). Considering the low mean iron adequacy levels ranging from 64% to 74% (Supplementary Table 1), this indicates how tree cover can contribute to helping women meet WHO recommendations for iron intake (Supplementary Table 2).

When controlling for tree cover, women from households that source food from on-farm trees have higher levels of zinc, vitamin A, iron and folate adequacy in the dry season, and higher levels of zinc, vitamin A and folate adequacy in the wet season compared with women from households that do not source food from on-farm trees (Fig. 1, path B, and Fig. 2b). For example, women sourcing food from trees on their farms have on average 8–15% higher folate adequacy, depending on the season. This suggests that foods from on-farm trees are important sources of folate—especially in the wet season when, on average, folate adequacy is low (48%) (Supplementary Tables 1 and 3).

To better understand whether the effect of tree cover on micronutrient adequacy is mediated by the direct provision of food from on-farm trees, we also test the dependence of the treatment variable (tree cover) on the mediator variable (sourcing food from on-farm trees) (Fig. 1, path A). We find that if tree cover around the household increases by 10%, the probability of using trees on farms for food increases by 9–10% on average. That is, in areas with higher tree cover, households are more likely to source food from on-farm trees.


The causal mediation analysis reveals how the direct provision of food from on-farm trees helps to explain the effect of tree cover on micronutrient adequacy. We observe statistically significant average causal mediation effects (ACMEs) for zinc, vitamin A and folate adequacy in both seasons (Fig. 2c and Supplementary Table 4; see

Supplementary Table 5 for additional robustness checks). The fact that we do not observe a mediation effect for iron in the wet season—but a positive association between tree cover and iron adequacy—is most likely because areas with high tree cover host wild animals and green leafy vegetables rich in iron, whereas few of the commonly consumed on-farm tree fruits in our study areas have high iron content (for example, bananas, mangoes, oranges, papayas) (Extended Data Fig. 2 and Supplementary Table 3). Yet, high consumption of mangoes, avocado and guava across dry and wet seasons can contribute to improved vitamin A, zinc and folate adequacy (Supplementary Tables 3 and 6).

Ninety-seven percent of Malawians do not have sufficient income to afford a healthy diet on a daily basis<sup>3</sup>. Our results show the nutritional benefits of using trees in agricultural landscapes for direct food provision, thereby identifying an avenue by which households can improve their supply of nutrient-rich foods regardless of income level. However, we note that while trees may produce fruits or nuts annually, monetary and land tenure considerations can influence investment decisions due to the time lag between planting and harvest (Supplementary Information). Access to silvicultural extension services can help farmers reduce this initial investment by enhancing and accelerating tree productivity—which can help ensure the availability of wild and cultivated nutrient-rich foods, especially in areas with limited market access and infrastructure<sup>9</sup>.

Rural Malawians are vulnerable to seasonal fluctuations in food availability<sup>10</sup>. Food shortages in the pre-harvest period, coupled with staple crop price fluctuations, can increase household reliance on self-cultivated or wild foods at certain times of the year. Our results showing how women benefited from different fruits from their on-farm trees in dry and wet seasons (Extended Data Fig. 2 and Supplementary Table 6) can inform reforestation programmes and agroecological interventions<sup>11</sup>. For example, portfolios of socially and ecologically suitable food tree species with staggered harvest periods can be integrated into landscape restoration initiatives (such as AFRIOO) to leverage the co-benefits of trees for the environment and people's diets<sup>12</sup>.

Our causal mediation analysis shows how farmers' decisions to manage trees on their farm for food are influenced by tree cover in the wider landscape—and how trees from forest to farm affect people's nutrition. These findings lay the groundwork for conservation and restoration policies that address forest and agroforestry systems in tandem. Strengthening the evidence base for tree–diet linkages across more countries and contexts is a key step to aligning food and nutrition initiatives with forest and tree conservation policies and programmes.



**Fig. 2 | Effect estimates for each component of the causal mediation analysis.** a–c, Results are presented as coefficient effect estimates for linear regression models measuring the effect of tree cover in a 1 km radius around each household on women’s micronutrient adequacy (a), logistic regression models measuring the effect of sourcing food from on-farm trees on micronutrient adequacy (b) and mediation analysis outputs given as ACMEs, indicating the degree to

which sourcing food from on-farm trees mediates the effect of tree cover on micronutrient adequacy (c). In all plots, data points are presented as coefficient values within a 95% confidence interval error bar. *P* values are based on two-sided tests. No adjustments were made for multiple comparisons. *n* = 460 women. \*\*\**P* < 0.001, \*\**P* < 0.01, \**P* < 0.05.

## Methods

Malawi is one of the most densely populated countries in Africa, approaching 20 million people with a high annual population growth rate of 3.06%<sup>10</sup>. The majority of the population (83%) lives in rural areas and at least partially depend on smallholder agriculture for their livelihoods. Sixty-two percent of the population was classified as living in multidimensional poverty in 2021<sup>13</sup>, and only 3.7% of people in rural areas have access to electricity<sup>10</sup>. Increasing dependence on natural resources for food, fuelwood and livelihoods has driven widespread deforestation. Reliance on rain-fed food production exacerbates seasonal production variability, leaving farmers highly vulnerable to climatic events (such as droughts and floods) and persistent food insecurity. Consequently, malnutrition rates are high, with widespread deficiencies in zinc, vitamin A, iron and folate, especially among women and children. For example, anaemia prevalence is 31% for women of reproductive age<sup>3</sup>. Taken together, the high levels of natural resource reliance, poverty and malnutrition justify Malawi as a case study country to examine the links between tree cover and dietary quality.

Malawi's climate is characterized by a unimodal rainy season occurring from December to April. To account for dietary changes based on seasonal food availability, we conducted two rounds of fieldwork in the dry season (October 2021) and wet season (March 2022). Study areas were selected in both a northern region (Nkhata Bay district) and a southern region (Mulanje district) to expand our inferential potential to the country level (Extended Data Fig. 1). The northern study area has a greater percentage of tree cover on customary land, whereas the southern region has a higher population density and less tree cover. In light of recent evidence indicating a relationship between market access and dietary quality in Malawi<sup>14,15</sup>, we selected study sites with relatively equal distance to trading centres of at least 5,000 people.

To measure the extent to which sourcing food from on-farm trees affects people's dietary quality, we conducted a household survey with 515 initial respondents. Household surveys were administered using Qualtrics XM with questions pertaining to household characteristics, household assets, farming systems, forest use and respondents' food consumption. We selected women with at least one child between the ages of 2 and 5 years as the primary survey respondents in each household. This was to target women of reproductive age and focus on women feeding their children solid foods. In Malawi, women play an integral role in household food security and nutrition. They are traditionally responsible for food selection, preparation and feeding of dependents (elders/children). Women's dietary diversity has been shown to align with household dietary diversity<sup>16</sup>, which indicates that women are reliable representatives of dietary quality at the household level. Eligible respondents were selected using systematic sampling (taking every *n*th household on a list, depending on village size). In areas with lower population density, a greater percentage of eligible women were selected to comply with all selection criteria. In especially remote villages, systematic random sampling was not possible due to the limited number of eligible households. In two village areas, all available eligible households were sampled.

We used a quantitative, 24 h dietary recall survey to collect detailed information on the type, quantity and source of the foods people consumed the previous day. For both rounds of fieldwork, each household was visited on two non-consecutive days within a 7 day period. Multiple 24 h recalls at different times of the year have been shown to be useful in accounting for seasonal variation in food intake, and multiple 24 h recalls with the same individual are integral to capturing variability in food intake<sup>17</sup>. To reduce systematic error and bias in the dietary data collection, an interview protocol with culturally sensitive tools and methods was developed in close collaboration with local enumerators. The first visit consisted of conducting a combined household/dietary recall survey, and the second visit consisted of a follow-up dietary recall. For the follow-up dietary recall survey, we obtained attrition rates of 97% (*n* = 499 of 515) and 98% (*n* = 451 of 460) in the dry and wet

seasons, respectively. As such, our statistical modelling was based on the 460 respondents for whom we had complete data in both seasons<sup>18</sup>.

Photograph aids and local serving size aids (plates, bowls, cups) were used to help respondents estimate the quantities of food and drink items consumed. Using these same aids, we later converted the local portion sizes into standard units (grams). From the collected food consumption data, we estimated the dietary supply of four key micronutrients: zinc, vitamin A, iron and folate. These micronutrients are of critical nutritional importance and are all targeted by Malawian government policies and interventions due to persistent, widespread deficiencies in the population. Food composition tables for Malawi<sup>19</sup> were used to estimate the micronutrient content of all food and drink items reported by respondents, and the daily supply of each nutrient was estimated using the consumed quantities in grams. Note that some data were 'borrowed' from other food composition tables where data were missing in the Malawian tables (Supplementary Information).

Estimated usual intake values were generated using the multiple source method<sup>20</sup>. This method synthesizes the multiple dietary surveys per respondent to adjust for interpersonal variation in consumption patterns. To estimate the adequacy of each micronutrient to meet minimum requirements, we used recommended nutrient intake (RNI) values<sup>21</sup>. RNI values were estimated based on each respondent's age and pregnancy and breastfeeding status in both the dry and wet season surveys (Supplementary Table 2). We then compared the RNI against each respondent's estimated usual intake (intake/RNI, capped at 1) to calculate the nutritional adequacy ratio for each micronutrient. To assess the mean adequacy level for all micronutrients of interest, we also calculated the mean adequacy ratio for each respondent and for our study population overall in each season (Supplementary Table 1).

We use tree cover in the landscape (including trees inside and outside forests) as a treatment variable. Tree cover was calculated as a percentage within a 1 km radius around each household location, aggregated from a very high resolution map of African tree cover in 2019<sup>4</sup>. The continental tree-cover map was created using a deep learning model to segment tree cover at the individual tree level, based on 3 m resolution PlanetScope satellite imagery. The use of very high resolution imagery notably allows the mapping of individual non-forest tree crowns, such that both forest trees and trees outside forests are included in the tree cover. Note that while tree cover was used as a treatment variable for testing paths C and A, we also controlled for tree cover (as a covariate) for testing path B (Fig. 1).

We focused on the use of trees on farms for food as a binary mediator. In the dry season household survey, respondents were asked if they had trees on or around their farmland ('around' was defined as within 15 m of the field boundary). If they responded 'yes', they were asked if their household sources food from these trees for household consumption (not to be conflated with using food from trees on farms for commercial purposes or owning food tree seedlings from which they cannot yet harvest food). This variable therefore centres on the households' decision to use trees on and around their farmland for household food provision. As the question on tree use did not specify use during a specific time period, data on how households use on-farm trees were only collected in the dry season, with the assumption that presence/use of on-farm trees is not dependent on seasonality.

It is also acknowledged that women's access to resources is influenced by land tenure systems. In Malawi, land inheritance is patrilineal in the northern region and matrilineal in the southern region (Supplementary Information). Although different land tenure systems must be considered in any intervention, our analysis operates on the assumption that the participants, by saying they are using the trees on their farm for food, had access to those trees.

We used the 'mediation' package in R to evaluate the average causal mediation (that is, indirect) effect of our food tree variable<sup>22</sup>. We tested the significance of the indirect effects using bootstrapping procedures, where unstandardized indirect effects were computed for 1,000



bootstrapped samples with 95% confidence intervals. We then conducted sensitivity analyses to test for omitted variable bias (Extended Data Fig. 3), the robustness of the ACME estimates (strong confounding effects between the mediator and outcome) (Supplementary Table 7), and the consistency of model results using a 500 m tree-cover buffer radius around each household (Supplementary Table 4). As an additional robustness measure, we conducted the causal mediation analysis using the mean adequacy ratio as an outcome (Supplementary Table 5) to check for consistency in model trends.

In the dry season household survey, respondents also reported the different food tree species they currently cultivated. To explore if the diversity of on-farm food tree species could explain the relationship between tree cover and micronutrient adequacy, we conducted a supplementary analysis equivalent to the second step of a hurdle model. Here we only selected households that source food from their on-farm trees ( $n = 360$ ) to avoid conflation with food trees used only for income purposes, and we excluded households that did not source food from on-farm trees ( $n = 100$ ). We then conducted a causal mediation analysis with this subsample, using food tree species count as a mediator. The results (Extended Data Fig. 4) show that higher food tree diversity has a statistically significant positive effect on iron adequacy in the wet season ( $P = 0.0090$ ), but not for any of the other micronutrients (Supplementary Table 8). This indicates that the decision to source any food from on-farm trees has a larger effect on micronutrient adequacy than the number of food tree species.

### Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

### Data availability

The dataset generated by the survey research for the replication of this study is available in a Harvard Dataverse repository and can be accessed at <https://doi.org/10.7910/DVN/WBUTCK> (ref. 18). We also accessed very high-resolution (3 m) tree-cover data from 2019 PlanetScope nanosatellite constellation imagery<sup>4</sup>. Estimated nutrient intakes were calculated using the following publicly available food composition tables: Malawi, <http://hdl.handle.net/10427/D217R336D>; Tanzania, <https://nutritionsource.hsph.harvard.edu/food-tables/>; Zambia, <https://nfnz.org.zm/download/zambia-food-composition-tables-4th-edition/>; Mozambique, <http://hdl.handle.net/10138/337295>; Kenya, <https://nutritionhealth.or.ke/programmes/healthy-diets-physical-food-composition-tables/>; West Africa, <https://openknowledge.fao.org/server/api/core/bitstreams/c5b37ac2-7082-48ab-a4a5-68d27deb4849/content>; United States (USDA), <https://fdc.nal.usda.gov/>. Source data are provided with this paper.

### Code availability

The code written to complete the analyses and ensure the replication of this study is available in a Harvard Dataverse repository and can be accessed at <https://doi.org/10.7910/DVN/WBUTCK> (ref. 18).

### References

- High Level Panel of Experts *Food Security and Nutrition: Building a Global Narrative Towards 2030* (FAO, 2020).
- Gergel, S. et al. Conceptual links between landscape diversity and diet diversity: a roadmap for transdisciplinary research. *BioScience* **70**, 563–575 (2020).
- FAO, IFAD, UNICEF, WFP & WHO *The State of Food Security and Nutrition in the World 2022* (FAO, 2022); <https://doi.org/10.4060/cc0639en>
- Interactive World Forest Map & Tree Cover Change Data* (Global Forest Watch, 2023); <https://www.globalforestwatch.org/map/?menu=eyJkYXRhc2VOQ2F0ZWdvcnk0iJmb3Jlc3RDaGFuZ2UjLjZtZU51U2VjdGlvbil6ImRhdGFzZXZXRzInO%3D>
- Reiner, F. et al. More than one quarter of Africa's tree cover is found outside areas previously classified as forest. *Nat. Commun.* **14**, 2258 (2023).
- Alkire, S. & Santos, M. E. Measuring acute poverty in the developing world: robustness and scope of the multidimensional poverty index. *World Dev.* **59**, 251–274 (2014).
- Guidelines on Methods for Estimating Livestock Production and Productivity* (FAO, 2018); <https://www.fao.org/3/ca6400en/ca6400en.pdf>
- Vansant, E. et al. What are the links between tree-based farming and dietary quality for rural households? A review of emerging evidence in low- and middle-income countries. *People Nat.* **4**, 296–311 (2022).
- Bennett, A. et al. Spatial analysis of aquatic food access can inform nutrition-sensitive policy. *Nat. Food* **3**, 1010–1013 (2022).
- Climate-Smart Agriculture in Malawi* (CIAT and World Bank, 2018); [https://climateknowledgeportal.worldbank.org/sites/default/files/2019-06/CSA%20\\_Profile\\_Malawi.pdf](https://climateknowledgeportal.worldbank.org/sites/default/files/2019-06/CSA%20_Profile_Malawi.pdf)
- Bezner Kerr, R. et al. Participatory agroecological research on climate change adaptation improves smallholder farmer household food security and dietary diversity in Malawi. *Agric. Ecosyst. Environ.* **279**, 109–121 (2019).
- McMullin, S. et al. Developing fruit tree portfolios that link agriculture more effectively with nutrition and health: a new approach for providing year-round micronutrients to smallholder farmers. *Food Secur.* **11**, 1355–1372 (2019).
- Malawi Multidimensional Poverty Index Report* (National Statistical Office, 2021); <https://ophi.org.uk/malawi-mpi-report-2021/>
- Matita, M. et al. Does household participation in food markets increase dietary diversity? Evidence from rural Malawi. *Glob. Food Secur.* **28**, 100486 (2021).
- Koppmair, S., Kassie, M. & Qaim, M. Farm production, market access and dietary diversity in Malawi. *Public Health Nutr.* **20**, 325–335 (2017).
- Minimum Dietary Diversity for Women* (FAO, 2021).
- Dietary Assessment: A Resource Guide to Method Selection and Application in Low Resource Settings* (FAO, 2018).
- Vansant, E. Replication data for: food-sourcing from on-farm trees mediates positive relationships between tree cover and dietary quality in Malawi. *Harvard Dataverse* <https://doi.org/10.7910/DVN/WBUTCK> (2024).
- Malawian Food Composition Table 2019* (MAFOODS, 2019).
- Harttig, U. et al. The MSM program: web-based statistics package for estimating usual dietary intake using the Multiple Source Method. *Eur. J. Clin. Nutr.* **65**, S87–S91 (2011).
- WHO & FAO *Vitamin and Mineral Requirements in Human Nutrition* (WHO, 2004).
- Tingley, D., Yamamoto, T., Hirose, K., Keele, L. & Imai, K. mediation: R package for causal mediation analysis. *J. Stat. Softw.* **59**, 1–38 (2014).

### Acknowledgements

We gratefully acknowledge the research assistants from the Lilongwe University of Agriculture and Natural Resources who conducted the household and dietary surveys: R. Liguluwe, A. Posiano, T. Luhanga, F. Khoropa, G. Ntholo and J. Chautsi. This research was funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement number 853222 FORESTDIE) (E.V., B.d.B., C.H. and L.V.R.) and UKRI Frontier Research Grant (EP/X023222/1), which was selected by the European Research Council (J.O.).

### Author contributions

E.V. conceived the study within the guidelines of the ERC FORESTDIE project. E.V. designed the study in collaboration with L.V.R. and led the data collection in Malawi. F.R. prepared and contributed the tree-cover

data. E.V. and B.d.B. carried out the analysis, with help from L.V.R. and J.O. E.V. received inputs from co-authors L.V.R., C.H., J.O. and J.K. on the interpretation of the results. All co-authors (B.d.B., C.H., J.O., F.R., J.K., L.V.R.) contributed to the writing of the paper and approved its publication.

### Competing interests

The authors declare no competing interests.

### Additional information

**Extended data** is available for this paper at <https://doi.org/10.1038/s43016-024-01028-4>.

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s43016-024-01028-4>.

**Correspondence and requests for materials** should be addressed to Emilie Vansant, Johan Oldekop or Laura Vang Rasmussen.


**Peer review information** *Nature Food* thanks Ida Djenontin, Debbie Pierce and Bronwen Powell for their contribution to the peer review of this work.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.


**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024




Data Source: 2019 PlanetScope via Reiner et al., 2023

**Extended Data Fig. 1 | Map of Malawi indicating study sites with surveyed households.** Map features surveyed household points overlaid on 2019 PlanetScope data with tree cover predictions at 3 m resolution<sup>4</sup>. Each household is randomly displaced up to 500 m for confidentiality purposes.




**Extended Data Fig. 2 | The consumption frequency of different tree-based foods in the dry and wet seasons.** The alluvial flow diagram shows the number of women who reported eating different tree-based foods sourced from their own farms in the dry season (n = 441) and wet season (n = 168).





**Extended Data Fig. 3 | Sensitivity contour plots of t-values to test for omitted variable bias.** Sensitivity tests using tree cover (A) and sourcing food from on-farm trees (B) as treatments to model effects on micronutrient adequacy levels (n = 460 women).



**Extended Data Fig. 4 | Comparison of Average Causal Mediation Effects (ACMEs) when using ‘sourcing food from on-farm trees’ and ‘number of food tree species’ as mediator variables.** Coefficient plots comparing ACME estimates for how a) the degree to which sourcing food trees on farms (binary mediator) and b) the unique food tree species count per household (count mediator) mediates the relationship between tree cover and micronutrient

adequacy. Note: the analysis using the unique food tree species count mediator was conducted only with women from households that source food from on-farm trees (n= 360 women). In all plots, data points are presented as coefficient values within a 95% confidence interval error bar. P values are based on two-sided tests. No adjustments were made for multiple comparisons. See Supplementary Table 8 for exact P values. \*\*\* $P < 0.001$ ; \*\* $P < 0.01$ .

## Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

### Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

- | n/a                                 | Confirmed  |
|-------------------------------------|--|
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> The exact sample size ( $n$ ) for each experimental group/condition, given as a discrete number and unit of measurement  |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly  |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> The statistical test(s) used AND whether they are one- or two-sided<br><i>Only common tests should be described solely by name; describe more complex techniques in the Methods section.</i>   |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> A description of all covariates tested   |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons  |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals) |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> For null hypothesis testing, the test statistic (e.g. $F$ , $t$ , $r$ ) with confidence intervals, effect sizes, degrees of freedom and $P$ value noted<br><i>Give <math>P</math> values as exact values whenever suitable.</i>                            |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings  |
| <input type="checkbox"/>            | <input checked="" type="checkbox"/> For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes   |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Estimates of effect sizes (e.g. Cohen's $d$ , Pearson's $r$ ), indicating how they were calculated  |

*Our web collection on [statistics for biologists](#) contains articles on many of the points above.*

### Software and code

Policy information about [availability of computer code](#)

**Data collection** The household surveys were designed using online using Qualtrics XM (accessed 2021-2022, no specific version available). The Qualtrics XM license agreement package included access to an app through which to conduct offline surveys. We used this to administer our survey on tablets using the Qualtrics Offline Surveys App (Version 17).

**Data analysis** Survey data was downloaded to CSV format and accessed through Excel 2016 software. The CSV file was uploaded to ArcGIS Pro (Version 3.0.3) to displace and map household GPS points.

For each respondent's food consumption data, we also generated Estimated Usual Intake values using the Multiple Source Method (MSM). This method synthesizes the multiple dietary surveys per respondent to adjust for interpersonal variation in consumption patterns. We used a web-based statistics package developed by the German Institute of Human Nutrition (DIFE) to implement the MSM with our nutrition data: <https://nugo.dife.de/msm>.

To examine the relationship between tree cover, use of on-farm food trees, and dietary quality, we cleaned and analyzed our data in R (version 4.2.2). We ran linear regressions, controlling for a selection of covariates, to examine these relationships. Specifically, we used the 'mediation' package in R to evaluate the causal mediation (i.e. indirect) effect of our food tree variable. This package also allowed us to evaluate the robustness of our average causal mediation effect estimates by testing for the possible confounding effect of unmeasured pretreatment variables. We also tested for the impact of omitted variables using the 'sensmakr' package in R.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

## Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

The dataset generated by the survey research and the code to facilitate the replication of this study is publicly available at: <https://doi.org/10.7910/DVN/WBUTCK>  
We also accessed very high resolution (3m) tree cover data from 2019 PlanetScope nanosatellite constellation imagery: <https://www.nature.com/articles/s41467-023-37880-4>

Estimated nutrient intakes were calculated using the following publicly available Food Composition Tables:

Malawi: <http://hdl.handle.net/10427/D217R336D>

Tanzania: <https://nutritionsource.hsph.harvard.edu/food-tables/>

Zambia: <https://nfdc.org.zm/download/zambia-food-composition-tables-4th-edition/>

Mozambique: <http://hdl.handle.net/10138/337295>

Kenya: <https://nutritionhealth.or.ke/programmes/healthy-diets-physical/food-composition-tables/>

West Africa: <https://openknowledge.fao.org/server/api/core/bitstreams/c5b37ac2-7082-48ab-a4a5-68d27deb4849/content>

United States (USDA): <https://fdc.nal.usda.gov/>

## Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

### Reporting on sex and gender

For our household and dietary surveys, we targeted women in each household with children between the ages of 2-5 years. The focus was on gender, rather than biological sex, due to the cultural and societal role women play in Malawian households in determining patterns of food consumption. Consent was obtained for sharing individual-level data. As all of our individual data was collected from people who self-identified as women, no gender-based analysis was performed.

### Reporting on race, ethnicity, or other socially relevant groupings

In the survey, we asked respondents to self-report their ethnicity (the question was phrased as "What is your ethnic group?" and translated accordingly). No data was collected on race or any other social grouping. Reporting on ethnicity was not included in the data analysis or manuscript. This is because the village areas selected in each study region were ethnically homogenous, with the village area populations predominantly (>90%) identifying as either Tonga/Tumbuka (Nkhata Bay) or Lomwe (Mulanje). It is noted that in terms of food culture, Tonga and Tumbuka cultures are regarded as similar based on mutual linguistic intelligibility and shared cultural history relative to other ethnic groups in Malawi (Kamwendo, 2005). Therefore, given the strong correlation between ethnicity and study region, we decided to only use the latter variable in our analysis.

Kamwendo, G. (2005). Language, Identity and the Politics of Recognition in Post-Banda Northern Malawi. *Ufahamu: A Journal of African Studies*, 31(1–2). <https://doi.org/10.5070/F7311-2016524>

### Population characteristics

The human research participants were women, aged 18-47 (average: 28). At the time of the dry season survey, 38 women were pregnant, 143 reported to be breastfeeding. At the time of the wet season survey, 30 women were pregnant, 119 reported to be breastfeeding. We selected women with at least one child between the ages of 2 – 5 years as the primary survey respondents. This was to target women of reproductive age and focus on women feeding their children solid foods. In Malawi, women play an integral role in household food security and nutrition. They are traditionally responsible for food selection, preparation, and feeding of dependents (elders/children). Women's dietary diversity has been shown to align with household dietary diversity, which indicates that women are reliable representatives of dietary quality at the household level.

No other data was collected regarding respondent health status or other physical conditions.

### Recruitment

Prior to recruitment, local authorities in each village area were visited to obtain permission to conduct research activities in each community. Individual survey participants were then recruited by local Health Surveillance Assistants (HSAs), who had records on women in each village area corresponding to our recruitment criteria (women with children between the ages of 2-5 years). After obtaining a full list of eligible women in each village area, we conducted systematic sampling (choosing a random starting point and then selecting every nth woman on the list relative to list length/village size). The list of women was not in any particular order, so we can assume there was no hidden pattern that would skew the sample. In the event that a recruited respondent was not available for the survey, the following household on the HSA's list was selected. Consent forms in the local language were read and signed by recruited survey participants to ensure voluntary participation and optional withdrawal at any time in the study prior to data collection.

### Ethics oversight

The project has received approval from the European Research Council (funding the parent project under the European Union's Horizon 2020 research and innovation programme) and the Research Ethics Committee for Science and Health at the University of Copenhagen (hosting institution).

Note that full information on the approval of the study protocol must also be provided in the manuscript.



# Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences  Behavioural & social sciences  Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

## Behavioural & social sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	We analyzed the relationship between tree cover, household decisions to source food from on-farm trees and women's micronutrient adequacy in rural Malawi. As part of a mixed-methods case study, we collected quantitative and qualitative data in the dry season (October 2021) and wet season (March 2022). Our analysis includes data from 460 women with whom we conducted surveys in both seasons, including 24-hr dietary recall data on food consumption.
Research sample	Our survey was designed to collect data at the individual and household level. At the individual level, our research sample consisted of women (aged 18 - 48; average 28). Given the age range and average, we find this sample to be adequately representative of women of reproductive age in Malawi - our target demographic for the survey. Our sample consisted of women with at least one child between the ages of 2 - 5 years as the primary survey respondents. This was to target women of reproductive age and focus on women feeding their children solid foods. In Malawi, women play an integral role in household food security and nutrition. They are traditionally responsible for food selection, preparation, and feeding of dependents (elders/children). Women's dietary diversity has been shown to align with household dietary diversity, which indicates that women are reliable representatives of dietary quality at the household level.
Sampling strategy	Across the two districts, we initially surveyed 515 women (dry season) and followed-up with 460 women (wet season) (89% attrition rate). No statistical methods were used to predetermine sample size. The sample size for the dry season was set at 500, with 15 extra households added due to available time and resources. Upon consulting experts in this field of nutrition, this sample size was determined as sufficient for a study using a 24-hour dietary recall (a time-intensive survey tool). The sample size for the wet season was determined by the availability of respondents - as we had to track down all respondents we had previously visited for the first round of the survey.
Data collection	<p>The household and 24-hr dietary recall surveys were tested and validated by a team of six MSc Nutrition graduates from the Lilongwe University of Agriculture and Natural Resources (LUANAR), who were hired as enumerators to conduct the surveys in the local language, Chichewa. All enumerators also contributed to the translation and revision of the household and dietary data survey tools. Therefore no researchers were blind to the study design nor research hypotheses during data collection. Household surveys were administered using tablets (Qualtrics XM app) with questions pertaining to household characteristics, household assets, farming systems, forest use, and respondents' food consumption. We used a quantitative, 24-hr dietary recall survey to collect detailed information on the type, quantity, and source of the foods people consumed the previous day. Photo aids and local serving size aids (plates, bowls, cups) were used to help respondents estimate the quantities of food and drink items consumed.</p> <p>Enumerators were trained to interview participants in isolation to avoid biases, with the exception of minors under the care of the respondent. Multiple 24-hr recalls at different times of the year have shown to be useful in accounting for seasonal variation in food intake, and multiple day 24-hr recalls with the same individual are integral to capturing variability in food intake. To reduce systematic error and bias in the dietary data collection, an interview protocol with culturally sensitive tools and methods was developed in close collaboration with local enumerators. For both rounds of fieldwork, each household was visited on two non-consecutive days within a 7-day period. The first visit consisted of conducting a combined household/dietary recall survey, and the second visit consisted of a follow-up dietary recall. Conducting two dietary recalls per respondent allowed us to obtain a more accurate 'snapshot' of habitual food consumption for each respondent, thereby improving our ability to calculate their estimated usual intakes for each target nutrient in each season (see Data Analysis section).</p>
Timing	The first round of data collection was conducted from 20/10/2021 - 24/11/2021, and the second round was conducted from 01/03/2022 - 28/03/2022.
Data exclusions	In our analysis, we excluded the 55 households for which we only had household and dietary data from the dry season survey.
Non-participation	No participants declined participation in the survey. In the follow-up (wet season) round of data collection, we were unable to re-visit 55 households. This is because the woman who we had originally surveyed was unavailable for reasons including: seasonal migration, family illness, funeral attendance. When possible, all efforts were made to follow-up with individual respondents within the survey time-frame.
Randomization	Study sites (ie. village clusters) were intentionally chosen in each district based on proximity to forests and markets. Eligible respondents corresponding to our pre-determined study criteria were randomly selected from lists provided by local health authorities (taking every nth household on the list, depending on village size). Post data collection, we compared women in areas with different degrees of tree cover and controlled for potential confounding variables that could explain differences in food consumption, including: household size, the Multidimensional Poverty Index (MPI) living standard, farm size (acres under cultivation), education level, livestock holdings (TLU), crop count, and study region. This selection of covariates was informed by a synthesis of studies linking trees and dietary quality as well as extensive fieldwork in Malawi.

# Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

## Materials & experimental systems

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern
<input checked="" type="checkbox"/>	<input type="checkbox"/> Plants

## Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

## Plants

### Seed stocks

Report on the source of all seed stocks or other plant material used. If applicable, state the seed stock centre and catalogue number. If plant specimens were collected from the field, describe the collection location, date and sampling procedures.

### Novel plant genotypes

Describe the methods by which all novel plant genotypes were produced. This includes those generated by transgenic approaches, gene editing, chemical/radiation-based mutagenesis and hybridization. For transgenic lines, describe the transformation method, the number of independent lines analyzed and the generation upon which experiments were performed. For gene-edited lines, describe the editor used, the endogenous sequence targeted for editing, the targeting guide RNA sequence (if applicable) and how the editor was applied.

### Authentication

Describe any authentication procedures for each seed stock used or novel genotype generated. Describe any experiments used to assess the effect of a mutation and, where applicable, how potential secondary effects (e.g. second site T-DNA insertions, mosaicism, off-target gene editing) were examined.