

Biomarker Predictors of Clinical Efficacy of the Anti-IgE Biologic Omalizumab in Severe Asthma in Adults

Results of the SoMOSA Study

✉ Ratko Djukanović¹, Paul Brinkman², Johan Kolmert³, Cristina Gomez³, James Schofield^{1,4}, Joost Brandsma¹, Andy Shapanis⁴, Paul J. S. Skipp^{1,4}, Anthony Postle¹, Craig Wheelock³, Sven-Erik Dahlen³, Peter J. Sterk², Thomas Brown⁵, David J. Jackson⁶, Adel Mansur⁷, Ian Pavord⁸, Mitesh Patel⁹, Christopher Brightling¹⁰, Salman Siddiqui¹⁰, Peter Bradding¹⁰, Ian Sabroe¹¹, Dinesh Saralaya¹², Livingstone Chishimba¹³, Joanna Porter¹⁴, Douglas Robinson¹⁴, Stephen Fowler^{15,16}, Peter H. Howarth¹, Louisa Little¹⁷, Thomas Oliver¹⁷, Kayleigh Hill¹⁷, Louise Stanton¹⁷, Alexander Allen¹⁷, Deborah Ellis¹⁷, Gareth Griffiths¹⁷, Tim Harrison¹⁸, Ayobami Akenroye^{19,20}, Jessica Lasky-Su²⁰, Liam Heaney²¹, Rekha Chaudhuri²², and Ramesh Kurukulaaratchy¹; on behalf of the SoMOSA study team and the U-BIOPRED study team

Abstract

Background: The anti-IgE monoclonal antibody omalizumab is widely used for severe asthma. This study aimed to identify biomarkers that predict clinical improvement during 1 year of omalizumab treatment.

Methods: One-year open-label Study of Mechanisms of action of Omalizumab in Severe Asthma (SoMOSA) involving 216 patients with severe (Global Initiative for Asthma step 4/5) uncontrolled atopic asthma (at least two severe exacerbations in the previous year) taking high-dose inhaled corticosteroids and long-acting β -agonists with or without maintenance oral corticosteroids. It had two phases: 0–16 weeks, to assess early clinical improvement by Global Evaluation of Therapeutic Effectiveness (GETE); and 16–52 weeks, to assess late responses based on $\geq 50\%$ reduction in exacerbations or mOCS dose. All participants provided samples (exhaled breath, blood, sputum, urine) before and after 16 weeks of omalizumab treatment.

Measurements and Main Results: A total of 191 patients completed phase 1; 63% had early improvement. Of 173 who completed phase 2, 69% had reduced exacerbations by $\geq 50\%$ and 57% (37 of 65) taking mOCSs had reduced their dose by $\geq 50\%$. The primary outcomes 2,3-dinor-11- β -PGF₂ α , GETE score, and standard clinical biomarkers (blood and sputum eosinophils, exhaled nitric oxide, serum IgE) did not predict either clinical response. Five volatile organic compounds and five plasma lipid biomarkers strongly predicted the $\geq 50\%$ reduction in exacerbations (receiver operating characteristic areas under the curve of 0.780 and 0.922, respectively) and early responses (areas under the curve of 0.835 and 0.949, respectively). In an independent cohort, gas chromatography/mass spectrometry biomarkers differentiated between severe and mild asthma.

Conclusions: This is the first discovery of omics biomarkers that predict improvement in asthma with biologic agent treatment. Prospective validation and development for clinical use is justified.

(Received in original form October 3, 2023; accepted in final form April 18, 2024)

✉ This article is open access and distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>). For reprints, please e-mail Diane Gern (dgern@thoracic.org).

Supported by a Novartis investigator-led project.

Author Contributions: Conception and Study Design: R.D., A.P., C.W., P.J.S.S., P.J.S., P.B., L.L., G.G., L.H., and R.C. Study Management: R.D., T.O. (project manager), T.H., L.H., and R.C. Clinical Trial Unit Team: L.L., T.O. and G.G. Acquisition of Clinical Data/Samples: T.B., D.J.J., A.M., I.P., M.P., C.B., S.S., I.S., D.S., L.C., J.P., D.R. S.F., P.H.H., T.H., L.H., R.C., and R.K. Data Management: A. Allen, D.E., and G.G. Massachusetts General Brigham Biobank Study: A. Akenroye and J. L.-S. Statistical Analysis: R.D., P.B., J.K., J.S., J.B., A.S., L.L., K.H., and G.G. Review and Editing: All authors.

Correspondence and requests for reprints should be addressed to Ratko Djukanović, M.D., D.M., Mailpoint 810, Level F, Sir Henry Wellcome Laboratories, South Block, Southampton University Hospital, Tremona Road, Southampton SO16 6YD, United Kingdom. E-mail: r.djukanovic@soton.ac.uk.

This article has a related editorial.

A data supplement for this article is available via the Supplements tab at the top of the online article.

Am J Respir Crit Care Med Vol 210, Iss 3, pp 288–297, Aug 1, 2024

Copyright © 2024 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.202310-1730OC on April 18, 2024

Internet address: www.atsjournals.org

¹School of Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton and National Institute for Health and Care Research Southampton Biomedical Research Centre, Southampton, United Kingdom; ²Department of Respiratory Medicine, Amsterdam University Medical Center, University of Amsterdam, the Netherlands; ³Institute of Environmental Medicine, Karolinska Institutet, and the Department of Respiratory Medicine and Allergy, Karolinska University Hospital, Stockholm, Sweden; ⁴Biological Sciences, University of Southampton, Southampton, United Kingdom; ⁵Portsmouth Hospitals University National Health Service Trust, Queen Alexandra Hospital, Portsmouth, United Kingdom; ⁶Guy's Severe Asthma Centre, School of Immunology & Microbial Sciences, King's College London, London, United Kingdom; ⁷University of Birmingham and Heartlands Hospital, University Hospitals Birmingham National Health Service Foundation Trust, Birmingham, United Kingdom; ⁸Oxford Respiratory National Institute for Health and Care Research Biomedical Research Centre, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom; ⁹Respiratory Medicine and R&D, University Hospitals Plymouth National Health Service Trust, Plymouth, United Kingdom; ¹⁰Institute for Lung Health and Leicester National Institute for Health and Care Research Biomedical Research Centre, University of Leicester, Leicester, United Kingdom; ¹¹Clinical Research Facility, Sheffield Teaching Hospitals National Health Service Foundation Trust, Sheffield, United Kingdom; ¹²Bradford Institute for Health Research and the National Patient Recruitment Centre, Bradford, United Kingdom; ¹³Clinical Sciences, Liverpool University Hospitals National Health Service Foundation Trust, Liverpool, United Kingdom; ¹⁴University College London Respiratory and National Institute for Health and Care Research University College London Hospitals Biomedical Research Centre, London, United Kingdom; ¹⁵Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The University of Manchester, Manchester, United Kingdom; ¹⁶Manchester Academic Health Science Centre and National Institute for Health and Care Research Manchester Biomedical Research Centre, Manchester University Hospitals National Health Service Foundation Trust, Manchester, United Kingdom; ¹⁷Southampton Clinical Trials Unit, University of Southampton, and University Hospital Southampton National Health Service Foundation Trust, Southampton, United Kingdom; ¹⁸Nottingham Respiratory National Institute for Health and Care Research Biomedical Research Centre, University of Nottingham, Nottingham, United Kingdom; ¹⁹Division of Allergy and Clinical Immunology and ²⁰Channing Division of Network Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts; ²¹Wellcome-Wolfson Institute for Experimental Medicine, Belfast, Northern Ireland; and ²²Gartnavel General Hospital and School of Infection & Immunity, University of Glasgow, Glasgow, United Kingdom

ORCID IDs: 0000-0001-6039-5612 (R.D.); 0000-0002-8113-0653 (C.W.); 0000-0002-8615-8778 (A.M.); 0000-0002-7307-169X (J.P.); 0000-0002-4524-1663 (S.F.); 0000-0002-1909-1470 (A. Akenroye).

The anti-IgE monoclonal antibody omalizumab (Xolair) is widely used to reduce asthma exacerbations and the need for oral corticosteroids (OCSs) in severe allergic asthma (1–3), but there is no reliable way to predict its benefit. In current practice, patients with at least two severe exacerbations in the previous year requiring OCSs are given a 16-week therapeutic trial, and the response is assessed using the Global Evaluation of Treatment Effectiveness (GETE) (4), a clinical tool based solely on the physician's assessment. GETE responders are then advised to continue treatment and undergo review after 1 year of treatment for a reduction in severe acute asthma exacerbations or dose of maintenance OCSs (mOCSs). Although using GETE enriches the responder population (4), a significant proportion of selected patients do not benefit in the long term, and there may be GETE nonresponders who show a response later. Thus, there is an unmet need for predictive biomarkers to optimize the use of omalizumab.

Studies evaluating standard, simple-to-measure clinical biomarkers as predictors of clinical response to omalizumab have had inconsistent results (5); none have assessed biomarker combinations. To improve our understanding of the mechanisms of action of omalizumab and identify predictive biomarkers for clinical practice, we designed a real-world Study of Mechanisms of action of Omalizumab in Severe Asthma (SoMOSA). In this article, the focus is on identifying biomarkers that predict which patients show

improvement with treatment. We hypothesized that omics biomarkers ("breathomics," proteomics, lipidomics) and urine eicosanoids in readily obtained samples (exhaled breath, blood, sputum, urine) can predict early responses (using GETE at 16 wk) and late responses ($\geq 50\%$ reduction in acute exacerbations or mOCS dose during the first year of treatment), outcomes that are the rationale for prescribing biologic agents. We measured more than 1,400 omics variables developed by the Unbiased Biomarkers Predictive of Respiratory Disease outcomes (U-BIOPRED) program (6, 7), including the prostaglandin D₂ metabolite 2,3-dinor-11 β -PGF₂ α and LTE₄ (leukotriene E₄), whose concentrations we have previously found to be lower in patients with severe asthma treated with omalizumab than in patients receiving standard treatment (8). The predictive value of omics biomarkers was compared with GETE score and standard clinical practice biomarkers (fractional exhaled nitric oxide [F_ENO], blood and sputum eosinophil counts, serum IgE). Evidence of the clinical relevance of the identified predictive biomarkers was then sought in datasets from two independent cohorts: U-BIOPRED (6, 7) and the Massachusetts General Brigham (MGB) Biobank (9).

Methods

Study Design and Clinical Assessment in the Core SoMOSA Study

This was an open-label, real-world study; all participants received standard-of-care

omalizumab and met current inclusion criteria. After 16 weeks of treatment (study phase 1), patients were evaluated by GETE score for early responses. At study end (52 wk), late responses were defined as a $\geq 50\%$ decrease in asthma exacerbations or dose of mOCSs between 16 and 52 weeks of treatment (phase 2). Asthma severity and control were assessed using the Asthma Control Questionnaire 7, Asthma Control Test, and Standardized Asthma Quality of Life Questionnaire. In contrast to standard practice, patients considered nonresponders based on the GETE assessment were also invited to continue treatment in phase 2. The study protocol was approved by the Wales Research Ethics Committee 5, Bangor (15-WA-0302), and patients provided written informed consent.

Two independent cohorts, U-BIOPRED and the MGB Biobank, provided data used to seek additional clinical value of any identified predictive biomarkers in SoMOSA.

Participants

For the core SoMOSA study in patients from 17 tertiary severe asthma clinics, the inclusion criteria were severe asthma (Global Initiative for Asthma step 4/5) that was uncontrolled (Asthma Control Questionnaire score ≥ 1.5 , atopic, at least two severe exacerbations in the past year) despite high-dose inhaled corticosteroids and long-acting β -agonists with or without mOCSs, serum total IgE concentration 30–1,500 IU/ml,

At a Glance Commentary

Scientific Knowledge on the

Subject: The mechanisms of action of the anti-IgE biologic agent omalizumab in asthma are poorly understood, and commonly measured biomarkers (exhaled nitric oxide, serum IgE, eosinophils) cannot reliably predict the clinical response to treatment. In the age of stratified medicine, the search for reliable ways to predict clinical responses to biologic agents must be extended to the spectrum of omics biomarkers that have transformed our understanding of the mechanisms of asthma.

What This Study Adds to the

Field: This is the first study to provide proof of concept that omics methods can prospectively identify biomarkers that predict to a high degree whether patients respond to omalizumab based on a $\geq 50\%$ reduction in acute exacerbations. This study offers a set of volatile organic compounds as the most promising biomarkers for the prediction of clinical response and a set of plasma biomarkers for which laboratory methods to measure individual biomarkers would be needed. Prospective studies comparing clinical responses in patients selected by these biomarkers with those selected according to criteria used in current practice are needed to validate the candidate biomarkers identified in our study for use in clinical practice.

and age 18–70 years (*see* online supplement for complete criteria).

Biomarker datasets from two independent cohorts were identified as suitable for additional analysis of the biomarkers shown in the core SoMOSA cohort as predictive of clinical responses to omalizumab: the U-BIOPRED study (10) and the MGB Biobank (*see* online supplement for details of cohorts and methods).

Standard and Omics Biomarkers

In the SoMOSA study, patients provided exhaled breath, blood, induced sputum, and

morning urine samples before and after 16 weeks of treatment. Four analytical omics methods that are able to quantify large numbers of biomarkers (6, 7) were applied and compared for predictive efficacy with biomarkers often used in clinics (blood and sputum eosinophil counts and FE_{NO}) and with the GETE-based early clinical response tool. Ultra-high-performance liquid chromatography (LC)–tandem mass spectrometry (MS) measured urine concentrations of 14 arachidonic acid–derived eicosanoids (11). Exhaled breath was analyzed by two methods: 1) gas chromatography (GC)–MS for individual volatile organic compounds (VOCs) and 2) a combination of electronic nose (eNose) cross-reactive sensors (12) that produced signatures without VOC identities. Intact lipids in sputum and plasma were measured by ultra-high-performance supercritical fluid chromatography–ion mobility–tandem MS (13). Quantitative data-independent LC/HDMS^E (liquid chromatography/high-definition mass spectrometry) was used to measure proteins in sputum and morning urine (7).

The omics methods applied in the U-BIOPRED study were broadly the same as those used in SoMOSA, with some technical advances in the latter. Plasma samples from the MGB Biobank underwent global metabolomic profiling (Metabolon) using untargeted LC-MS platforms, which includes amines, amino acids, and polar and nonpolar lipids (14). *See* online supplement for more details of U-BIOPRED and MGB Biobank analytical methods.

Power Calculation and Statistical Analysis

The change in the urine prostaglandin D_2 metabolite 2,3-dinor-11 β -PGF_{2 α} from baseline to 16 weeks after the initiation of omalizumab was the selected as the primary outcome and for power calculation using data from a U-BIOPRED study comparing asthmatic subjects taking and not taking omalizumab (8). Omics biomarkers were prespecified as coprimary outcomes because power calculations are not possible for unbiased omics biomarkers. Assuming 66% of participants would show a response (2:1 ratio of responders to nonresponders), 194 completed participants were required, with sample size adjustment allowed depending on the final ratio of responders to nonresponders. The same calculation was used to compare exacerbation responders and nonresponders. The same

participant number was assumed to be required to test the hypothesis that 2,3-dinor-11 β -PGF_{2 α} in urine would be reduced in participants with a $\geq 50\%$ reduction in exacerbations (*see* online supplement for more details).

Initial analysis of treatment effects on patient-reported outcomes, FEV₁% predicted, FE_{NO} , and blood and sputum eosinophil counts was performed with analysis of covariance or quantile regression models depending on the distribution of the data. For the omics analysis, missing values were addressed as previously described (7, 13), excluding from analysis molecules with detection rates across samples below $< 40\%$ for proteins and 60% for lipids. Because of differences in methodology between lipidomics and proteomics, missingness was dealt with differently: lipidomics data were imputed using 50% of the lowest limit of detection, whereas, for proteomics, we used median levels to minimize identification of false-positive markers. Data were batch-corrected for location, defining GETE score and exacerbations as outcomes of interest to preserve variation. Features that detected contaminants due to sample collection and/or processing were removed. Data were then split 50/50 into training and test cohorts; the latter was analyzed after a final model was produced on the training cohort. Feature selection was performed on the training data. The equal Gini estimator sought to identify the top five predictive features for each omics platform data set, which were then used to train the final machine learning prediction model using a random forest algorithm, with fivefold cross-validation repeated three times. After training, the prediction model was tested on the test cohort, and the results were plotted as receiver operating characteristic (ROC) curves. Comparisons of identified predictive biomarkers from the core SoMOSA study were made using the U-BIOPRED and MGB Biobank datasets using two-sample Wilcoxon tests applied to participants with severe and mild to moderate asthma in the former and omalizumab responders and nonresponders in the latter. Sparse partial least squares discriminant analysis was applied to the U-BIOPRED data set to assess whether those groups of biomarkers identified by random forest analysis to predict clinical responses could differentiate between those with severe and mild/moderate asthma and between patients taking omalizumab and not taking omalizumab.

SoMOSA CONSORT Diagram




Figure 1. Consolidated Standards of Reporting Trials diagram. OCS = oral corticosteroid.

Results

Analysis of the SoMOSA Study Data

Of 811 initially assessed patients, 217 were enrolled; 191 successfully completed phase 1,

173 completed phase 2, and 43 withdrew (Figure 1 and Tables E1 and E2 in the online supplement). In keeping with the prespecified allowance to adjust the required number of patients completing the study,

recruitment stopped after 191 patients completed phase 1.

Clinical responses. Based on GETE score at 16 weeks, 121 of 191 patients (63%) were classified as early responders (Table 1).

Table 1. Demographic and Main Clinical Outcomes

	<i>n</i>	ACQ7	ACT	AQLQ	AcEx	Taking mOCSs, <i>n</i>	IgE, IU	F _{ENO} , ppb	Blood eos, x10 ⁹ /L	Sputum eos, %
Early (16 wk)										
Responders	121 (63.3%)									
Baseline		2.9 (2.0–3.6)	12.0 (9.0–17.0)	3.8 (2.9–5.0)	4.0 (3.0–6.0)	41/41 (33.1%)	231.0 (114.0–377.0)	33.5 (17.3–59.0)	0.26 (0.11–0.48)	6.0 (0.9–23.0)
16 wk	121	1.6 (1.0–2.1)	18.0 (14.0–21.0)	5.5 (4.5–6.2)	NA	41/41 (32.2%)	NA	23.5 (12.8–39.0)	0.18 (0.10–0.38)	1.6 (0.5–8.1)
52 wk	113	1.6 (0.9–2.3)	18.0 (15.0–22.0)	5.7 (4.8–6.3)	1.0 (0.0–2.0)	31/39 (30.6%)	NA	26.0 (16.0–48.5)	0.20 (0.11–0.37)	2.3 (1.0–12.0)
Nonresponders	70 (36.7%)									
Baseline		2.9 (2.4–3.7)	10.0 (9.0–13.0)	3.7 (2.9–4.7)	4.0 (2.0–6.0)	34/34 (48.6%)	157.0 (87.0–302.0)	34.3 (16.0–63.0)	0.23 (0.11–0.41)	2.3 (0.9–13.9)
16 wk	70	2.6 (2.0–3.3)	13.0 (10.0–16.0)	4.2 (3.4–5.2)	NA	34/34 (47.1%)	NA	29.8 (19.0–52.0)	0.15 (0.08–0.30)	2.3 (0.6–5.6)
52 wk	60	2.4 (1.1–3.0)	14.5 (10.5–19.0)	4.6 (3.7–5.8)	1.0 (0.0–3.0)	25/29 (40.0%)	NA	23.0 (15.0–41.0)	0.17 (0.07–0.38)	2.5 (0.5–10.0)
AcEx (52 wk)										
Responders	120 (71.0%)									
Baseline		2.7 (2.0–3.6)	12.0 (9.0–16.0)	3.8 (3.0–5.0)	4.0 (3.0–6.0)	38/38 (31.7%)	209.8 (112.5–326.4)	33.0 (16.0–56.0)	0.25 (0.11–0.44)	3.0 (1.0–17.8)
16 wk	120	1.9 (1.1–2.6)	17.5 (14.0–21.0)	5.2 (4.1–6.1)	NA	38/38 (30.0%)	NA	24.0 (14.0–37.0)	0.19 (0.10–0.35)	1.8 (0.5–5.8)
52 wk	120	1.6 (0.9–2.4)	18.5 (13.5–22.0)	5.6 (4.6–6.3)	1.0 (0.0–2.0)	29/38 (30.0%)	NA	22.5 (15.5–42.0)	0.20 (0.10–0.39)	2.3 (0.8–12.3)
Nonresponders	49 (29.0%)									
Baseline		3.0 (2.3–3.7)	10.0 (8.0–13.0)	3.8 (2.9–4.8)	4.0 (2.0–5.0)	26/26 (53.1%)	191.7 (98.0–360.0)	31.3 (18.0–53.0)	0.22 (0.11–0.50)	7.8 (1.3–30.5)
16 wk	49	2.4 (1.6–3.0)	13.0 (11.0–18.0)	4.8 (3.7–5.7)	NA	26/26 (51.0%)	NA	28.0 (17.5–51.0)	0.12 (0.09–0.30)	2.3 (0.8–8.5)
52 wk	49	2.4 (1.4–3.0)	14.0 (12.0–18.0)	4.6 (3.9–5.6)	3.0 (2.0–4.0)	24/26 (51.0%)	NA	31.0 (16.5–56.5)	0.18 (0.10–0.38)	2.9 (1.0–6.1)
mOCS (52 wk)										
Responders	37 (56.9%)									
Baseline		2.4 (2.0–3.6)	13.0 (10.0–19.0)	4.0 (3.2–5.5)	3.0 (2.0–4.0)	37/37 (100%)	177.0 (87.0–492.0)	36.0 (19.0–55.5)	0.24 (0.09–0.37)	3.9 (1.5–12.5)
16 wk	37	2.0 (1.3–2.6)	18.0 (13.0–21.0)	5.1 (4.4–6.1)	NA	37/37 (100%)	NA	30.0 (17.5–44.0)	0.16 (0.07–0.30)	3.5 (0.5–13.5)
52 wk	37	2.0 (0.9–2.9)	17.0 (13.0–19.0)	5.2 (4.6–6.1)	1.0 (0.0–2.0)	23/37 (62.2%)	NA	31.0 (19.5–63.0)	0.25 (0.16–0.42)	7.5 (1.0–21.0)
Nonresponder	28 (43.1%)									
Baseline		3.0 (2.4–3.9)	11.0 (9.0–17.0)	4.0 (3.2–4.6)	4.0 (2.0–6.0)	28/28 (100%)	152.0 (101.0–254.0)	30.0 (15.8–72.0)	0.13 (0.06–0.26)	11.0 (0.8–13.0)
16 wk	28	2.4 (1.1–3.5)	15.5 (10.5–21.5)	4.9 (3.5–6.0)	NA	28/28 (100%)	NA	27.5 (20.8–61.8)	0.10 (0.04–0.26)	2.6 (1.5–8.5)
52 wk	28	2.3 (1.4–3.9)	16.0 (10.5–22.0)	4.6 (3.5–6.1)	2.0 (1.0–3.5)	28/28 (100%)	NA	27.0 (15.5–63.5)	0.12 (0.06–0.32)	4.9 (1.3–24.8)
<i>P</i> values*										
16 wk	<0.001	0.005	<0.001	<0.001	NA	NA	NA	0.014	0.180	0.227
52-wk AcEx	0.479	0.002	0.001	0.001	NA	NA	NA	0.077	0.099	0.344
52-wk mOCS	0.479	0.853	0.442	0.442	NA	NA	NA	0.504	0.166	0.251

Definition of abbreviations: AcEx = acute exacerbation; ACQ = asthma control questionnaire; ACT = asthma control questionnaire; AQLQ-S = asthma quality of life questionnaire; F_{ENO} = fractional exhaled nitric oxide; eo = eosinophil; mOCS = maintenance oral corticosteroid.

Data in parentheses are IQRs. Model: comparison of changes in variable values from baseline to 16 or 52 weeks, with variable after 16 of the 52 weeks of treatment = intercept + response group + variable at baseline. The numbers of participants who enrolled and remained in the study at 16 and 52 weeks are shown in the second column. The mOCS use data are shown as the numbers of participants taking a mOCS at the time of assessment as a proportion of the total numbers of patients assessed (i.e., still in the study) at that time point.

**P* values determined by analysis of covariance/quantile regression depending on the distribution of the data.

The majority ($n = 173$; 91%) completed phase 2; of those, 120 (71%) were late responders based on a $\geq 50\%$ reduction in acute exacerbation (Table 1) unrelated to age, sex, smoking history, or body mass index (see Table E2). Of 65 patients taking mOCSs, 37 (57%) reduced the dose by $\geq 50\%$ without losing asthma control (Table 1). Among early responders not taking mOCSs, 71.6% also met late-responder criteria; similarly, 70.7% of late responders not initially taking mOCSs were also early responders. Among patients taking mOCSs at enrolment, 80% of early responders met the criteria for late responders by acute exacerbations or mOCS use. Taking these two late response criteria together, 62% of late responders were also early responders, whereas 63% of early non responders (44 of 70), who would normally be asked to stop treatment, were shown in phase 2 to be late responders based on reduced exacerbations, reduction in mOCS use, or both. Thus, of 36 GETE nonresponders not taking mOCSs, 24 (67%) had a positive response in phase 2. Of the 34 GETE nonresponders using mOCSs

before treatment, 20 (59%) had a positive response in phase 2.

The numbers of responders and nonresponders based on exacerbation reduction (120 and 49, respectively) or GETE score (121 and 70, respectively) were deemed sufficient to split the cohort into training and test sets. In contrast, the numbers of responders by mOCS reduction (37 and 28, respectively) were too small for analysis.

Biomarker measurements. A total of 1,408 variables passed quality control. Because individual biomarker molecules can result in multiple MS variables that require deconvolution to produce single variables, the 1,408 variables were reduced to 14 eicosanoids, 70 breath VOCs, 112 sputum proteins, and 147 urine proteins. A further 158 eNose variables provided signatures without molecular identities. Of the 589 lipid variables in plasma and 305 in sputum, identities were determined only if concentrations were different between responders and nonresponders (86 in plasma and 25 in sputum).

Baseline differences in biomarkers between responders and nonresponders.

Baseline concentrations of 2,3-dinor-11 β -PGF_{2 α} (primary outcome) did not differentiate early or late responders and nonresponders (Figure 2 and Figure E1). Even though baseline LTE₄ was significantly ($P = 0.018$) higher in early responders, LTE₄ and other eicosanoid levels did not differentiate late responders and nonresponders (Figure 2). The same was true for FeNO, blood and sputum eosinophil counts, and IgE (Table 1). In contrast, a total of 368 omics variables were different between responders and nonresponders across the four omics platforms (Figure 3): 103, 143, and 122 when comparing responses by GETE, exacerbation reduction, and mOCS use reduction, respectively, 67 being different for more than one outcome.

Prediction of clinical responses to omalizumab by random forest analysis. The 2,3-dinor-11 β -PGF_{2 α} did not predict the early GETE-based response (ROC AUC, 0.556) or $\geq 50\%$ exacerbation reduction during




Figure 2. Urine eicosanoids, 2,3-dinor-11 β -PGF_{2 α} , and LTE₄ (leukotriene E₄). (A) Baseline concentrations of 2,3-dinor-11 β -PGF_{2 α} (primary outcome) and LTE₄ (pg/ml) in patients defined as responders or nonresponders based on Global Evaluation of Therapeutic Effectiveness score. (B) Changes in concentrations of 2,3-dinor-11 β -PGF_{2 α} and LTE₄ in the entire cohort (responders and nonresponders) from baseline to 16 weeks analyzed by Mann-Whitney U test. (C) Receiver operating characteristic area under the curve for 2,3-dinor-11 β -PGF_{2 α} in respect to the prediction of early (Global Evaluation of Therapeutic Effectiveness-based) and late (acute exacerbation-based) response to omalizumab. AUC = area under the curve; GETE = Global Evaluation of Therapeutic Effectiveness; OCS = oral corticosteroid.




Figure 3. Volcano plots of baseline concentrations of all biomarker variables in responders and nonresponders. Responses shown include early response judged by GETE response (A) and late response defined by $\geq 50\%$ reduction in exacerbations (B). The red and blue biomarkers (all $P < 0.05$) are labeled by numbers (see Table E4 for identities). Green and red dots represent greater than onefold different biomarkers. The data are shown as the means of concentrations in the responders from which the means of the concentrations in the nonresponders have been subtracted (i.e., responder minus nonresponder). They are shown as log₂-transformed data. The P values were obtained by Mann-Whitney U test. GETE = Global Evaluation of Therapeutic Effectiveness.

phase 2 (ROC AUC, 0.542) (Figure 2), nor did the other urine eicosanoids (data not shown). Similarly, GETE, F_{ENO} , blood or sputum eosinophil counts, and serum IgE (Figure E2) did not predict exacerbation reductions.

Analysis of all the omics platforms showed that breathomics and plasma lipidomics predicted early and late responses (Figure 4), whereas the other omics platforms had weak predictive value (Table E3). One set of five exhaled breath VOCs (benzothiazole, acetophenone, 2-pentyl-furan, methylene chloride, 2-methylbutane) predicted early improvement (ROC AUC, 0.835); another set of VOCs (2-ethyl-1-hexanol, toluene, 2-pentene, nonanal, and a VOC of unknown identity detected as X79.175 by GC-MS) predicted a $\geq 50\%$ exacerbation reduction (ROC AUC, 0.780). Two sets of five plasma lipids were highly predictive of early and late clinical responses (ROC AUCs, 0.949 and 0.922, respectively). The plasma lipids that predicted early responses consisted of four triglycerides (TG[54:6], TG[56:7], TG[55:2], and

TG[52:3]) and a currently unidentified lipid. A further set predicted exacerbation reductions; of these, only one could be identified in lipid databases or the wider literature, namely the sphingomyelin peak for SM(d40:2), likely comprising a combination of SM(d18:2/22:0), SM(d16:1/24:1), and SM(d18:1/22:1) molecular species (15). Two further peaks were putatively identified as TG52:3 and ceramide.

Effect of treatment on eicosanoids and standard biomarkers. Urinary 2,3-dinor-11- β -PGF_{2 α} decreased significantly ($P = 0.029$) after 16 weeks of treatment, with no difference between responders and nonresponders (Figure 2). LTE₄ was also reduced ($P < 0.001$), but to a similar extent in responders and nonresponders (Figure 2). The other urine eicosanoids did not change (data not shown).

In the entire cohort, omalizumab reduced blood and sputum eosinophil numbers during phase 1 ($P < 0.001$ and $P < 0.023$, respectively) and F_{ENO} and blood eosinophils during phase 2 ($P = 0.022$, $P < 0.001$, respectively), but these changes

were not related to treatment responses except for F_{ENO} , which was reduced more in early responders ($P = 0.014$); however, neither F_{ENO} nor any of the other standard biomarkers discriminated late responders and nonresponders by ROC analysis in isolation or when combined (see Figure E2). We also stratified patients according to F_{ENO} and blood eosinophil count cutoff values used by Hanania and colleagues (2) as biomarker-high or -low when assessing their clinical response to omalizumab. We found that such stratification did not predict which stratum of patients would respond to omalizumab (Figure E3). Similarly, time to the first protocol-defined asthma exacerbation, as demonstrated by Kaplan-Meier curves, was no different (Figure E4) between these strata of patients.

Analysis of the identified predictive biomarkers in the U-BIOPRED and MGB Biobank. A search of the U-BIOPRED data undertaken for matching VOCs and plasma lipids showed that several of the candidate biomarkers we found to be predictive of responses to omalizumab were able to differentiate between individuals with severe atopic asthma and mild/moderate asthma (see online supplement for full details) in the U-BIOPRED cohort. In the MGB Biobank, the concentrations of plasma sphingomyelin (sphingomyelin d18:1/22:1, d18:2/22:0, d16:1/24:1) were significantly ($P = 0.03$) lower in responders to omalizumab than in nonresponders.

Discussion

To our knowledge, this is the first study to use a multi-omics approach to identify predictive biomarkers for severe asthma, providing proof of concept that breathomics and plasma lipidomics biomarkers can predict who benefits from omalizumab during the first 16 weeks of treatment and who shows a $\geq 50\%$ reduction in exacerbations during the first year of treatment. In an independent cohort, the biomarkers identified in SoMOSA were shown to differentiate between mild/moderate and severe asthma, including those with more frequent exacerbations who would be candidates for treatment with omalizumab. Development of these biomarkers has significant potential to give patients, their medical teams, and payers more certainty of achieving reduced exacerbations with omalizumab, a key objective of asthma treatment.




Figure 4. Breath volatile organic compounds (VOCs) and plasma lipids that predict early or late clinical responses. (A) The biomarker identities of the VOCs were derived from the variables detected by gas chromatography–mass spectrometry, whereas the identities of the plasma lipids were derived from the variables detected by ultra–high-performance supercritical fluid chromatography–ion mobility–tandem mass spectrometry. Receiver operating characteristic area under the curve figures show predictions by VOCs (B) and lipids (C) of early clinical responses judged by GETE score and late responses by reduction in asthma exacerbations. See Table E3 for the receiver operating characteristic area under the curve values for the other omics platforms (sputum lipids, sputum proteins, urine proteins, and eicosanoids). AUC = area under the curve; GETE = Global Evaluation of Therapeutic Effectiveness

Consistent with previously reported efficacy, 63% of patients showed improvement within 16 weeks of starting treatment, which suggests that the enrolled cohort is representative of the typical patient considered for omalizumab. In our study, GETE, the clinical tool widely used to assess clinical response to omalizumab, did not predict late improvement (see Figure E2); indeed, many patients classified by GETE as non–early responders had a late response (i.e., reduced exacerbations or mOCS use). Although 2,3-dinor-11 β -PGF₂ α , the coprimary outcome used for power calculation, was reduced significantly with treatment, the changes were similar in responders and nonresponders, and baseline concentrations did not predict early or late

improvement (Figure 2). Similarly, none of the standard biomarkers currently used in asthma management (FeNO, sputum and blood eosinophil counts, and serum IgE) had predictive value (see Figure E2).

Breathomics is a growing field in medicine (16). There are several types of eNoses that provide signatures, but not identities of VOCs, and MS methods like GC-MS effectively predict clinical and therapeutic outcomes. Whereas the combination of eNose cross-reactive sensors could not predict clinical improvement, five VOCs (2-ethyl-1-hexanol, toluene, 2-pentene, and one unknown VOC) derived by GC-MS confidently predicted the reductions in exacerbations, whereas a separate set of five GC-MS–derived VOCs

(benzothiazole, acetophenone, 2-pentyl-furan, methylene chloride, and 2-methyl-butane) predicted good early responses. Together, these VOCs differentiated between individuals with mild/moderate asthma and atopic severe asthma (ROC AUC, 0.931) and between mild/moderate asthma and severe asthma prone to exacerbations (at least two exacerbations per year), a cutoff for initiating treatment with a biologic agent. Many of these VOCs have been reported in respiratory studies. Nonanal is associated with neutrophilic asthma and smoking; it has been able to predict exacerbations and discriminate between allergic and nonallergic asthma in children (17–19). Toluene, a common organic solvent, is increased in smokers (20), is related to environmental

exposure (21), and has also been associated with asthma (22). We have previously found nonanal within a group of exhaled breath biomarkers in patients with cystic fibrosis with sputum positive for *Pseudomonas aeruginosa* (23). The predictive set in our study also included 2-ethyl-1-hexanol, for which there is prior evidence of a role in asthma and in lung cancer (reviewed by Sola-Martinez and coworkers [24]). It is a known indoor pollutant and the main metabolite of di(2-ethylhexyl)phthalate, a solvent and frequent plasticizer of polyvinylchloride.

Concentrations of 2-ethyl-1-hexanol sampled in ambient air are negligible compared with those in exhaled breath (24), suggesting that, if it is in part inhaled, it is concentrated in the lungs. 2-ethyl-1-hexanol is produced in greater quantities by cancer cells (25). Within the lungs, it acts as an endocrine-disrupting chemical and is associated with oxidative stress and modulation of immune responses (26). The hydrocarbon 2-pentene, also a solvent and known byproduct of thermal cracking of petroleum, is found in ambient air. It is also a volatile compound derived from lipid peroxidation, with increased concentrations found by GC-MS in the headspace of bacterial cultures (27). Among the GC-MS variables that predicted early improvement, three have been reported in respiratory conditions: acetophenone in patients with cystic fibrosis with *Pseudomonas aeruginosa* (23) and 2-pentylfuran in patients with *Aspergillus fumigatus* (28). Analysis of VOCs in exhaled breath that diagnose ventilator-associated pneumonia has proposed a set of 12 predictive VOCs, among them 2-methyl-butane (29). We could not find any similar reports for benzothiazole.

Lipidomic analysis of plasma also identified two sets of predictive biomarkers. Early improvement was predicted by four triglycerides and one unknown lipid species. In comparison to our understanding of the roles of leukotrienes, knowledge of other lipids in asthma is limited, although obesity is strongly associated with asthma. Serum triglyceride levels are higher in obese people with asthma, even when adjusted for body mass index, blood eosinophils, and statin treatment (30). A recent lipidomics study that identified more than 1,300 plasma lipid species showed that triglyceride levels, albeit different from the ones in our analysis, differentiated asthma from health and were related to asthma severity (31), with ceramides being related to asthma severity, in keeping with the findings in our study.

Ceramide exacerbates inflammation, mucus production, and endoplasmic reticulum stress, and increased levels are associated with airway hyperresponsiveness, a key feature of asthma (32). However, these lipids were not good at differentiating between severe and mild/moderate asthma and asthma with frequent exacerbations in the U-BIOPRED study; even though concentrations of plasma triglyceride 52:3 and one unidentified lipid were significantly higher in those with severe atopic asthma and in those with at least two exacerbations per year, the ROC AUC indicated weak differentiation (see online supplement). Of note, however, comparison of responders and nonresponders to omalizumab (defined by $\geq 50\%$ reduction in exacerbations) in the MGB cohort showed significantly lower concentrations of sphingomyelin (d18:1/22:1, d18:2/22:0, d16:1/24:1) in responders.

This study has limitations. It could be argued that we should have used a classical randomized controlled trial design, despite ample precedent of similar study design in oncology. Our discussions with the patient advisory group strongly favored a real-world study design, arguing that a placebo arm would be unethical because it would deny patients a drug known to improve a severe condition and that recruitment into a placebo-controlled trial would be difficult because omalizumab is readily available and patients expect to be treated. The fact that study recruitment took 26 months and required engagement of 17 severe asthma centers with exclusive rights to prescribe biologic agents justified this decision. The other limitation of the study is that there were too few patients in whom mOCS treatment was reduced by $\geq 50\%$, a measure that is very relevant to patients because of OCS side effects.

The identified biomarkers should be viewed as candidate biomarkers that require confirmation in a prospective study in which treatment efficacy in patients selected based on these biomarkers would be compared with efficacy in patients selected based on standard clinical criteria. Further studies are also needed to elucidate how these biomarkers are involved in asthma pathogenesis. Prospective validation of the candidate biomarkers should focus on breathomics, an easy-to-apply platform, possibly in combination with plasma lipid measurements. In view of the cost of developing routine analytical methods, the development of single-platform assays is likely to be easier, more acceptable to patients, and less expensive. Although lipids

had greater predictive power (AUC >0.9) than the VOC biomarkers (AUCs, 0.835 for early and 0.780 for late responses), breathomics is, in our view, a superior omics platform because of easier sample collection, more certainty about the VOC identities and, most importantly, easier development of point-of-care instruments for clinical use. Further elucidation of the detected lipids would likely be more complex and costly, and, with uncertain outcomes, riskier. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Acknowledgment: The authors thank the following for their excellent help to coordinate and deliver the study: University Hospital Southampton National Health Service Foundation Trust R&D and all R&D departments from individual study sites, National Institute for Health and Care Research Respiratory Translational Research Collaboration for facilitating contracts and for general support of the study's research objectives. The authors thank the Medical Research Council and the RASP-UK (Refractory Asthma Stratification Programme) Trial Steering Committee for oversight of the program during annual reviews and the patient advisory group for their decisive input into the study design. The authors wish to pay special tribute to Dr. Borislav Dimitrov, a statistician at the University of Southampton, who undertook the power calculations and helped to plan the statistical analysis but died before the end of the study.

Contributors to the SoMOSA Study are as follows.

Study design: Ratko Djukanović, Anthony Postle, Craig Wheelock, Peter J. Sterk, Paul Skipp, Gareth Griffiths, Louisa Little, Borislav Dimitrov (deceased), Liam Heaney, Rekha Chaudhuri, and Peter Bradding.

Study management group: Ratko Djukanović, Liam Heaney, Rekha Chaudhuri, Tim Harrison, Robert Niven, Tom Stokes (patient member of the Study Management Group), and Thomas Oliver.

Principal investigators at study sites: Rekha Chaudhuri, Robin Gore, Liam Heaney, Adel Mansur, Dinesh Saralaya, Ian Pavord, Mitesh Patel (initially Matthew Masoli), Christopher Brightling, David Jackson, Tim Harrison, Thomas Brown, Ian Sabroe, Livingstone Chishimba, Ramesh Kurukulaaratchy (initially Peter H. Howarth), Joanna Porter (initially Douglas Robinson), and Stephen Fowler (initially Robert Niven).

Study site clinical teams: Vanessa Brown, Alison Clinton, Christine McAnally (Belfast), Rahul Shrimanker, Gareth Hynes, Mary Green (Oxford), Salman Siddiqui, Peter Bradding, Christobelle White, Richard Russell, Michelle Bourne, Beverley Hargadon, Vijay Mistry, Tracey Thornton, Sarah Diver (Leicester), Joanne Kavanagh, Diana Roque (GKT, London), Ben Durham, Linda Kay, Matthew Austin, Ian Smith, Mercy Korley, Sam Anderson, Liam Haslam (Sheffield), Katherine Smith, Helen Smith (Nottingham), Christopher

Brereton, Kamran Tariq, Laura Presland, Clair Barber, Jon Ward (Southampton), Jagdeep Sahota, Joel Solis, Akif Khawaja (University College London), Valarie Magaya, Sarah Davies, Mary Bellamy, Lesley Horton, Tamika Thompson (Birmingham), Jacqui Galloway, Tom Dymond (Addenbrookes, Cambridge), Matthew Masoli, Julie Anderton, Natasha Wilmhurst, Jennifer Kingdon (Plymouth), Thomas Jones, Jonathon Winter, Scott Elliott (Portsmouth), Daniela Ferreira, Jesús Reiné, Esther German, John Blakey, Elena

Mitsi, Carla Solorzano, Elissavet Nikolaou, Catherine Lowe (Liverpool), Freda Yang, Steven Smith, Diane Murray, Tracyanne Grandison, and Marc Jones (Glasgow).

Trial management team: Gareth Griffiths, Thomas Oliver, Louisa Little, Jessica Rajaram, Nicholas Das, Zina Eminton, Jacqui Nuttall, Susi Renz, and Claire Forbes.

Data managements: Alexander Allen, Deborah Ellis, Emma Tilt, Emma Wrixon, and Nicola Scott.

Statistical analysis: Kayleigh Hill, Kerensa Thorne, Angeliki Galanopolous, Louise Stanton, Paul Brinkman, James Schofield, Johan Kolmert, Joost Brandsma, Gareth Griffith, Ratko Djukanović, and Borislav Dimitrov (deceased).

Massachusetts General Brigham Biobank study: Ayobami Akenroye and Jessica Lasky-Su.

Quality control: Kathryn Clark, Keira Fines, and Benjamin Sale.

References

- Hanania NA, Alpan O, Hamilos DL, Condeemi JJ, Reyes-Rivera I, Zhu J, *et al.* Omalizumab in severe allergic asthma inadequately controlled with standard therapy: a randomized trial. *Ann Intern Med* 2011;154:573–582.
- Hanania NA, Wenzel S, Rosén K, Hsieh HJ, Mosesova S, Choy DF, *et al.* Exploring the effects of omalizumab in allergic asthma: an analysis of biomarkers in the EXTRA study. *Am J Respir Crit Care Med* 2013;187:804–811.
- Bousquet J, Cabrera P, Berkman N, Buhl R, Holgate S, Wenzel S, *et al.* The effect of treatment with omalizumab, an anti-IgE antibody, on asthma exacerbations and emergency medical visits in patients with severe persistent asthma. *Allergy* 2005;60:302–308.
- Bousquet J, Rabe K, Humbert M, Chung KF, Berger W, Fox H, *et al.* Predicting and evaluating response to omalizumab in patients with severe allergic asthma. *Respir Med* 2007;101:1483–1492.
- Kroes JA, Zielhuis SW, van Roon EN, Ten Brinke A. Prediction of response to biological treatment with monoclonal antibodies in severe asthma. *Biochem Pharmacol* 2020;179:113978.
- Wheelock CE, Goss VM, Balgoma D, Nicholas B, Brandsma J, Skipp PJ, *et al.*; U-BIOPRED Study Group. Application of 'omics technologies to biomarker discovery in inflammatory lung diseases. *Eur Respir J* 2013;42:802–825.
- Schofield JPR, Burg D, Nicholas B, Strazzeri F, Brandsma J, Staykova D, *et al.*; U-BIOPRED Study Group. Stratification of asthma phenotypes by airway proteomic signatures. *J Allergy Clin Immunol* 2019;144:70–82.
- Kolmert J, Gómez C, Balgoma D, Sjödin M, Bood J, Konraden JR, *et al.* Urinary leukotriene E4 and prostaglandin D2 metabolites increase in adult and childhood severe asthma characterized by type-2 inflammation. *Am J Respir Crit Care Med* 2020;203:37–53.
- Boutin NT, Schechter SB, Perez EF, Tchamitchian NS, Cerretani XR, Gainer VS, *et al.* The evolution of a large biobank at Mass General Brigham. *J Pers Med* 2022;12:1323.
- Shaw DE, Sousa AR, Fowler SJ, Fleming LJ, Roberts G, Corfield J, *et al.*; U-BIOPRED Study Group. Clinical and inflammatory characteristics of the European U-BIOPRED adult severe asthma cohort. *Eur Respir J* 2015;46:1308–1321.
- Gómez C, Gonzalez-Riano C, Barbas C, Kolmert J, Hyung Ryu M, Carlsten C, *et al.* Quantitative metabolic profiling of urinary eicosanoids for clinical phenotyping. *J Lipid Res* 2019;60:1164–1173.
- Brinkman P, Wagener AH, Hekking PP, Bansal AT, Maitland-van der Zee AH, Wang Y, *et al.* Identification and prospective stability of electronic nose (eNose)-derived inflammatory phenotypes in patients with severe asthma. *J Allergy Clin Immunol* 2019;143:1811–1820.e1817.
- Brandsma J, Goss VM, Yang X, Bakke PS, Caruso M, Chanez P, *et al.*; U-BIOPRED Study Group. Lipid phenotyping of lung epithelial lining fluid in healthy human volunteers. *Metabolomics* 2018;14:123.
- Ryals J, Lawton K, Stevens D, Milburn M. Metabolon, Inc. *Pharmacogenomics* 2007;8:863–866.
- Quehenberger O, Armando AM, Brown AH, Milne SB, Myers DS, Merrill AH, *et al.* Lipidomics reveals a remarkable diversity of lipids in human plasma. *J Lipid Res* 2010;51:3299–3305.
- Ibrahim W, Natarajan S, Wilde M, Cordell R, Monks PS, Greening N, *et al.* A systematic review of the diagnostic accuracy of volatile organic compounds in airway diseases and their relation to markers of type-2 inflammation. *ERJ Open Res* 2021;7:00030-02021.
- Schleich FN, Zanella D, Stefanuto PH, Bessonov K, Smolinska A, Dallinga JW, *et al.* Exhaled volatile organic compounds are able to discriminate between neutrophilic and eosinophilic asthma. *Am J Respir Crit Care Med* 2019;200:444–453.
- Caldeira M, Perestrelo R, Barros AS, Bilelo MJ, Morête A, Câmara JS, *et al.* Allergic asthma exhaled breath metabolome: a challenge for comprehensive two-dimensional gas chromatography. *J Chromatogr A* 2012;1254:87–97.
- Jareño-Esteban JJ, Muñoz-Lucas MÁ, Carrillo-Aranda B, Maldonado-Sanz JA, de Granda-Orive I, Aguilar-Ros A, *et al.*; Study Group. Volatile organic compounds in exhaled breath in a healthy population: effect of tobacco smoking. *Arch Bronconeumol* 2013;49:457–461.
- Capone S, Tufariello M, Forleo A, Longo V, Giampetruzzi L, Radogna AV, *et al.* Chromatographic analysis of VOC patterns in exhaled breath from smokers and nonsmokers. *Biomed Chromatogr* 2018;32:e4132.
- Maniscalco M, De Laurentiis G, Pentella C, Mormile M, Sanduzzi A, Carratù P, *et al.* Exhaled breath condensate as matrix for toluene detection: a preliminary study. *Biomarkers* 2006;11:233–240.
- Dragonieri S, Schot R, Mertens BJ, Le Cessie S, Gauw SA, Spanevello A, *et al.* An electronic nose in the discrimination of patients with asthma and controls. *J Allergy Clin Immunol* 2007;120:856–862.
- Kos R, Brinkman P, Neerincx AH, Paff T, Gerritsen MG, Lammers A, *et al.*; Amsterdam Mucociliary Clearance Disease (AMCD) Research Group and the Amsterdam UMC Breath Research Group. Targeted exhaled breath analysis for detection of *Pseudomonas aeruginosa* in cystic fibrosis patients. *J Cyst Fibros* 2022;21:e28–e34.
- Sola-Martínez RA, Lozano-Terol G, Gallego-Jara J, Morales E, Cantero-Cano E, Sanchez-Solis M, *et al.*; NELA study group. Exhaled volatile analysis as a useful tool to discriminate asthma with other coexisting atopic diseases in women of childbearing age. *Sci Rep* 2021;11:13823.
- Sponring A, Filipiak W, Mikoviny T, Ager C, Schubert J, Miekisch W, *et al.* Release of volatile organic compounds from the lung cancer cell line NCI-H2087 *in vitro*. *Anticancer Res* 2009;29:419–426.
- Wakayama T, Ito Y, Sakai K, Miyake M, Shibata E, Ohno H, *et al.* Comprehensive review of 2-ethyl-1-hexanol as an indoor air pollutant. *J Occup Health* 2019;61:19–35.
- Boots AW, Smolinska A, van Berkel JJBN, Fijten RRR, Stobbering EE, Boumans MLL, *et al.* Identification of microorganisms based on headspace analysis of volatile organic compounds by gas chromatography-mass spectrometry. *J Breath Res* 2014;8:027106.
- Chambers ST, Syhre M, Murdoch DR, McCartin F, Epton MJ. Detection of 2-pentylfuran in the breath of patients with *Aspergillus fumigatus*. *Med Mycol* 2009;47:468–476.
- Schnabel R, Fijten R, Smolinska A, Dallinga J, Boumans M-L, Stobbering E, *et al.* Analysis of volatile organic compounds in exhaled breath to diagnose ventilator-associated pneumonia. *Sci Rep* 2015;5:17179.
- van Zelst CM, de Boer GM, Türk Y, van Huisstede A, In't Veen JCCM, Birnie E, *et al.* Association between elevated serum triglycerides and asthma in patients with obesity: an explorative study. *Allergy Asthma Proc* 2021;42:e71–e76.
- Jiang T, Dai L, Li P, Zhao J, Wang X, An L, *et al.* Lipid metabolism and identification of biomarkers in asthma by lipidomic analysis. *Biochim Biophys Acta Mol Cell Biol Lipids* 2021;1866:158853.
- Siddique MM, Li Y, Chaurasia B, Kaddai VA, Summers SA. Dihydroceramides: from bit players to lead actors. *J Biol Chem* 2015;290:15371–15379.