

<https://doi.org/10.1038/s44271-024-00124-2>

# A systematic review and Bayesian meta-analysis provide evidence for an effect of acute physical activity on cognition in young adults

Check for updates

Jordan Garrett <sup>1,2</sup>✉, Carly Chak <sup>1,2</sup>, Tom Bullock <sup>1,2</sup> & Barry Giesbrecht <sup>1,2</sup>✉

Physical exercise is a potential intervention for enhancing cognitive function across the lifespan. However, while studies employing long-term exercise interventions consistently show positive effects on cognition, studies using single acute bouts have produced mixed results. Here, a systematic review and meta-analysis was conducted to determine the impact of acute exercise on cognitive task performance in healthy young adults. A Bayesian hierarchical model quantified probabilistic evidence for a modulatory relationship by synthesizing 651 effect sizes from 113 studies from PsychInfo and Google Scholar representing 4,390 participants. Publication bias was mitigated using the trim-and-fill method. Acute exercise was found to have a small beneficial effect on cognition ( $g = 0.13 \pm 0.04$ ;  $BF = 3.67$ ) and decrease reaction time. A meta-analysis restricted to executive function tasks revealed improvements in working memory and inhibition. Meta-analytic estimates were consistent across multiple priors and likelihood functions. Physical activities were categorized based on exercise type (e.g., cycling) because many activities have aerobic and anaerobic components, but this approach may limit comparison to studies that categorize activities based on metabolic demands. The current study provides an updated synthesis of the existing literature and insights into the robustness of acute exercise-induced effects on cognition. Funding provided by the United States Army Research Office.

A single bout of exercise induces a cascade of neuromodulatory changes that influence multiple brain systems<sup>1,2</sup>. This includes an increase in the synthesis of neurotransmitters (e.g., acetylcholine, dopamine, GABA, glutamate) and neurotrophic factors (e.g., BDNF), which can occur in a brain-region-specific manner (see ref. 1 for review). Given these impacts on the brain, it would be reasonable to hypothesize that single brief bouts of exercise are associated with changes in performance across a range of cognitive domains. Consistent with this hypothesis, there is abundant evidence that attention<sup>3-6</sup>, working memory<sup>7-11</sup>, decision making<sup>12,13</sup>, and cognitive control<sup>14,15</sup> are facilitated by brief bouts of physical exercise. However, there is also evidence suggesting that exercise has little or no effect on cognitive task performance. For instance, Komiyama et al.<sup>16</sup> observed no difference in accuracy on a spatial delayed response task between exercise and rest conditions. Further, working memory performance has been shown to remain unchanged either during or after a single bout of exercise<sup>6</sup>. The discrepant pattern of results in the literature investigating the link between exercise and performance on

cognitive tasks is surprising given the consistent and robust physiological effects of even brief bouts of physical activity. However, it is unclear whether this limited impact of exercise on performance reflects the true state of affairs or whether the apparent lack of robust influence is due to vast empirical discrepancies across studies in the literature. Studying the impact of single exercise sessions on cognition can provide insight into how changes in our body's physiological state impact behavior. This understanding can then guide the creation of more effective longer-term exercise interventions, which essentially involve regularly repeating brief exercise sessions over an extended period.

Meta-analytic techniques are a set of powerful tools that can expose dominant trends within a methodologically heterogeneous literature. There is a consensus amongst narrative reviews and previous meta-analyses that an acute bout of exercise has a small positive influence on behavioral performance<sup>1,17-23</sup>. The nature of this effect is moderated by exercise protocol, cognitive tasks, and participant characteristics. For instance,

<sup>1</sup>Department of Psychological & Brain Sciences, University of California, Santa Barbara, CA, USA. <sup>2</sup>Institute for Collaborative Biotechnologies, University of California, Santa Barbara, CA, USA. ✉e-mail: [jordangarrett@ucsb.edu](mailto:jordangarrett@ucsb.edu); [giesbrecht@ucsb.edu](mailto:giesbrecht@ucsb.edu)

Lambourne & Tomporowski<sup>20</sup> observed that task performance during exercise was dependent on exercise modality, the type of cognitive task, and when it was completed relative to exercise onset. Similarly, post-exercise performance was moderated by exercise modality and the type of cognitive task. Chang et al.<sup>18</sup> reported that post-exercise cognitive performance was influenced by exercise intensity, duration, and the time of cognitive test relative to exercise cessation. Interestingly, the authors found that study sample age was a significant moderator, where larger positive effects were found for high school (14–17 years), adult (31–60 years), and older adult (>60 years) samples compared to elementary (6–13 years) and young adult (18–30 years) samples. Multiple meta-analyses have observed that the effect of exercise is dependent on cognitive domain, with measures of executive function, attention, crystallized intelligence, and information processing speed showing the largest gains<sup>18,19,24–26</sup>. Further, there is evidence that exercise has a differential influence on the speed and accuracy of cognitive processes. McMorris et al.<sup>21</sup> observed that acute, intermediate exercise facilitated response times on working memory tasks, while accuracy was compromised. In contrast, exercise has been shown to boost both the accuracy and speed of cognitive control<sup>23</sup>. Altogether, it is important to consider cognitive task, participant, and physical activity characteristics to develop a holistic model of the relationship between exercise and cognition.

While these earlier meta-analyses have provided unique insights into understanding the relationship between acute exercise and cognition, they have two major limitations. First, the most recent holistic quantitative synthesis of the extant literature was published over a decade ago<sup>18</sup>. Meanwhile, the exercise and cognition literature has grown drastically. According to the electronic database Web of Science, almost 6,000 articles associated with the search term “exercise and cognition” have been published since this last holistic meta-analysis. In addition, more recent meta-analyses have primarily focused on executive processes<sup>19,22,26,27</sup>. Thus, previous models may provide an outdated and limited account of exercise-induced influences on other aspects of cognition, such as perception, long-term memory, and learning. Second, previous meta-analytic approaches employed frequentist statistical methods, which are based on a decision threshold rather than a characterization of the relevant evidence. As a result, it is possible that acute exercise and moderator variables are deemed to have a significant influence on task performance despite the fact that there may only be a small degree of probabilistic evidence in favor of this notion. In addition, relying on a decision threshold prevents these models from conveying the likelihood that an exercise protocol elicits a change in cognitive task performance. Past frequentist meta-analytic models also treated heterogeneity parameters as a fixed quantity and utilize only a point estimate, which can lead to an underestimation of the variability either between or within studies<sup>28–31</sup>. This is especially true when the number of modeled studies is low<sup>32–34</sup>. When considered together, there is a clear need for an updated meta-analysis using an approach that addresses these limitations.

The current study addressed these limitations in two ways. First, a comprehensive literature search was conducted spanning the years 1995–2023. To quantify the influence of exercise on cognition in young healthy adults, the search was limited to non-clinical studies whose subjects were between 18–45 years old. The analysis focused on subjects within this age range since exercise research has predominantly been dedicated toward studying the effects in children and older adults<sup>35,36</sup>. Studies were required to be experimental in nature, and consist of both an acute exercise manipulation and cognitive task measurements. A broad range of cognitive domains encompassing tasks probing perception to executive function were included in the meta-analysis. Similarly, a wide range of exercise types and testing contexts were included. For example, traditional laboratory exposures to exercise (e.g., cycling, running) and sport activities in real-world settings were viable candidates for analysis. By casting a wide net, the current study provides a large scope and updated summary of the current state of the exercise and cognition literature.

Second, the current study uses a Bayesian meta-analytic approach to synthesize studies across the acute exercise and cognition literature. The Bayesian approach affords a flexible modeling framework that uses reported

effect sizes to characterize the relative evidence in favor of a modulatory account. Inherently, a random effects meta-analytic model is hierarchical in nature, making it well suited for Bayesian methods. When utilized within this statistical framework, priors are placed on parameters at the highest level of the model such as the estimated pooled effect size and measures of heterogeneity. This approach has several advantages compared to its frequentist counterpart. First, the use of priors on heterogeneity parameters can attenuate the underestimation of variation both between and within studies<sup>37,38</sup>, leading to a clearer understanding of sources of heterogeneity and an increased precision when estimating the pooled effect size<sup>39</sup>. Furthermore, priors provide additional constraints on low-level parameter estimates and a greater degree of “shrinkage” of outliers towards the overall pooled effect size or mode(s) of grouping variables<sup>39,40</sup>. Therefore, a Bayesian meta-analysis is more robust to outliers and can be more conservative when proper priors are employed. Second, the method yields a posterior distribution for all parameter estimates. This grants the capability of directly modeling the degree of uncertainty in heterogeneity estimates<sup>37</sup>. Posterior distributions can be used to compute the probability that an exercise protocol elicits a change in task performance of a given magnitude (e.g., large effect size). Compared to the approximation of *p*-values and confidence intervals, which require additional assumptions for hierarchical models, calculating the high-density interval (HDI), which indicates the most credible outcomes in the posterior distribution, for complex hierarchical models is seamless<sup>39</sup>. Third, it is possible to incorporate knowledge from previous meta-analyses when constructing prior distributions. This affords the ability to quantitatively compare the observed data to the predictions of previous models.

Considering the results of past meta-analyses, exercise was expected to have a small positive influence on cognition. Cognitive task and exercise characteristics were anticipated to moderate this relationship, as evidenced by nonzero parameter estimates, reflecting the selective nature of exercise-induced effects. Model comparisons were conducted to evaluate how moderator inclusion improved predictive performance, and robustness of parameter estimates were determined by employing multiple priors and likelihood functions.

## Methods

### Literature search

Studies investigating the impact of an acute bout of exercise on cognition were obtained through searches of the electronic databases PsychInfo and Google Scholar according to the PRISMA guidelines<sup>41</sup>. On 09 September 2023, databases were queried using a search string that combined the following physical activity and cognitive domain keywords: [“exercise” OR “physical activity” OR “physical exertion” OR “physical fatigue”] AND [“perception” OR “attention” OR “working memory” OR “executive function” OR “memory” OR “decision making” OR “motor skill” OR “skill acquisition” OR “language” OR “reasoning”]. For the PsychInfo search, the filters “journal article”, “English”, “empirical study”, “human”, and “peer reviewed” were applied. Search results were limited to studies published between 1995 and 2023 and whose subjects were between 18 and 45 years of age. Note, this literature search and analysis were not preregistered, nor was a review protocol prepared prior to the literature search.

### Eligibility criteria

Studies were deemed eligible for inclusion in the meta-analysis if they met all of the following criteria: assessed the influence of an acute bout of exercise on cognition, compared the effects of exercise with an active and/or passive control group(s), utilized cognitive tasks that measured reaction time (RT) and/or accuracy, tested cognition either during, pre-, or post-exercise and consisted of cognitively normal subjects. Note, an acute bout was defined as an instance of physical activity that occurred within a single 24-hour period<sup>18</sup>. Two researchers independently screened records based on their title, abstract, and full text. In the case of discrepancies, a third researcher resolved them by reading the full-text.

## Data extraction and coding

Information concerning experimental design and procedures, exercise details (i.e., type, intensity, duration), and sample characteristics were extracted from the final list of studies by a single researcher. Means and standard deviations of accuracy and/or RT measures on all cognitive tasks were inserted into an electronic spreadsheet for the calculation of effect sizes. The primary outcome measures for each domain were inserted separately if a task assessed multiple cognitive domains. Regarding studies that probed cognition at multiple time points during or post-exercise, measures for each time point were also recorded separately. If the statistics necessary for calculating effect sizes were not reported in the full-text of the article, the authors were contacted and asked to provide them.

All effect sizes were categorized into one of seven cognitive domains that were generally based on the DSM-5<sup>42</sup>: executive function, information processing, perception, attention, learning, motor skills, and memory. The classification criteria used for categorizing a cognitive task into a domain is provided in the Supplementary Table 1. To account for variability in the metric used to measure exercise intensity across studies (e.g., ventilatory threshold, heart rate), each intensity was labeled as either light, moderate, or vigorous according to the American College of Sports Medicine guidelines<sup>43</sup>. Exercise durations were grouped into one of five time bins:  $\leq 16$  minutes, 20–27 minutes, 30–35 minutes, 40–45 minutes,  $\geq 60$  minutes. In the event that a study did not provide the exercise duration, its time bin was labeled as “not provided”. Exercise types were based on the modality reported in each study, yielding the following categorizations: cycling, high intensity interval training (HIIT), running, walking, circuit training, resistance exercise, and sports activity. The latter category encompassed studies that used sports-related exercises that did not fit into the other labels, such as rock climbing or soccer. The time at which cognitive task performance was evaluated relative to exercise was categorized as either during exercise or 0, 15, 20–75, and  $\geq 180$  minutes after cessation. Lastly, effect sizes were also coded according to task performance dependent measures (i.e., RT vs accuracy). Note, the levels of each categorical moderator were chosen with the intention of achieving a balance between specificity and statistical power to yield reliable estimates that can inform the design of future exercise studies.

## Calculating effect sizes

Cohen’s  $d$  effect sizes were calculated for studies that tested cognition pre-/post-exercise without a control condition by dividing the mean change in performance by the standard deviation of the pre-test. If the study included a control group (e.g., rest), the mean change of the control condition was subtracted from the mean change of the exercise condition and divided by the pooled standard deviation of pretest scores<sup>20,44</sup>. For studies that tested cognition during, or only after exercise, the mean of the control condition was subtracted from the mean of the exercise condition and divided by the standard deviation of the control condition<sup>21</sup>. All effect sizes were converted into the bias-corrected standardized mean difference, Hedge’s  $g$ , by multiplying them by the correction factor  $J = 1 - \frac{3}{4df-1}$  where  $df$  is the degrees of freedom<sup>45</sup>. The sign of effect sizes for RT and error were reversed to reflect a positive influence of exercise on cognitive task performance. Once effect sizes were extracted from each study, inspection of a funnel plot and Egger’s regression test were conducted to assess the risk of publication bias.

## Bayesian hierarchical modeling

The overall effect of exercise on cognition was assessed using a Bayesian hierarchical model<sup>46,47</sup>, which was implemented through the R package *brms*<sup>48</sup>. In the first level of the model, a study’s observed effect size(s)  $\hat{\theta}_{ik}$  was assumed to be an estimate of the true effect size  $\theta_k$ . The observed effect(s)  $\hat{\theta}_{ik}$  were modeled as being sampled from a normally distributed population underlying study  $k$  with a mean equivalent to the true effect and a variance of  $\sigma_k^2$ . In the second level of the model, the true effect size  $\theta_k$  was assumed to have been drawn from an overarching distribution whose mean represented the overall pooled effect  $\mu$ , and whose variance depicted the degree of between-study heterogeneity  $\tau^2$ . The final level of the model contained weakly informative priors. A standard normal prior was used for the pooled

effect, while the prior for  $\tau^2$  was a Half-Cauchy distribution with location and scale parameters set to 0 and 0.5, respectively.

Following the main meta-analysis, subgroup analyses were conducted to determine potential moderators of the relationship between exercise and cognitive task performance. More specifically, we analyzed the influence of the following primary moderators: cognitive domain, time of cognitive test relative to exercise, task outcome measure, exercise intensity, duration, and type. The following secondary moderators were also analyzed to determine the influence of study and participant characteristics on the overall pooled effect size: average sample age, body mass index (BMI kg/m<sup>2</sup>), height (cm), weight (kg), VO<sub>2</sub> max (ml/kg/min), percentage of female participants, within- vs between-study design, and publication year. With the exception of publication year and the percentage of female participants, all secondary moderators were mean centered for interpretability. A standard normal distribution was used as a weakly informative prior for the difference in effect sizes between subgroups. When reporting model parameter estimates, we use the [mode  $\pm$  standard deviation] and the 89% HDI of posterior distribution.

## Statistical inference


For all estimated effect sizes, Bayes Factors (BFs) were used to determine the degree of evidence in favor of a difference from zero. BFs were approximated using the reciprocal of the Savage-Dickey density ratio, which was implemented using the function *bayesfactor\_parameters* from the *bayestestR* package<sup>49</sup>. This method involves dividing the height of the prior distribution for the null value by the height of the posterior distribution at the same value, and represents the credibility of the null value for a parameter once the data has been taken into consideration. BFs were also used to ascertain the predictive performance of subgroup models. After each model was compared to a null counterpart (i.e., moderator excluded) using the function *bayesfactor\_models*, an inclusion BF (*bayesfactor\_inclusion*) was estimated to determine if including a moderator improved predictive power<sup>50</sup>. To estimate stable BFs, a large number of sampling iterations (10,000) and warmup samples (2000) were used for each of four chains when estimating model parameters<sup>51</sup>. BFs were interpreted following the guidelines proposed by Jeffreys<sup>52</sup>. A BF between 1 and 3 indicates “anecdotal” evidence for the alternative hypothesis, between 3 and 10 indicates “moderate” evidence, between 10 and 30 indicates “strong” evidence, and greater than 30 indicates “very strong” evidence<sup>39,53–55</sup>. The reciprocal of these ranges signifies evidence in favor of the null hypothesis (e.g., 0.33–1 = anecdotal evidence). When conducting subgroup analyses with more than two factors, orthonormal coding was employed to ensure that an identical prior was used for each factor level and that estimated BFs were accurate<sup>56</sup>. Parameter estimates were extracted from all models using the R package *emmeans*.

## Sensitivity analysis

A popular criticism of the Bayesian approach is that priors are chosen subjectively, which in turn can bias parameter estimates and their corresponding BFs<sup>40,57</sup>. Although utilizing weakly informative priors mitigates bias, a sensitivity analysis that evaluates the contribution of both priors and the likelihood function must be conducted to determine if the model results are robust<sup>38,58–60</sup>. Thus, we replicated the previously described modeling approach with the exception of using two different priors for the overall pooled effect size. The first was a normal distribution with a mean of zero and standard deviation of  $\frac{1}{2}$ . Since this prior adds greater weight to the probability that exercise has no influence on task performance, we denoted it as the no effect (NE) prior. The second prior was constructed by synthesizing estimates from previous meta-analyses on acute exercise and cognition<sup>18,20–23</sup>, resulting in a normal distribution with a mean of 0.24 and standard deviation of 0.57. This prior was denoted as the positive effect (PE) prior.

The influence of the likelihood function was assessed by modeling study effect sizes as being sampled from a  $t$ -distribution. An advantage of using this likelihood function, compared to a normal distribution, is that

**Fig. 1 | PRISMA flow diagram of literature search results.** A total of 113 studies were deemed eligible for meta-analytic modeling.



model parameter estimates are influenced less by outliers<sup>40</sup>. The Half-Cauchy prior was used for the scale of the distribution, while a standard normal prior was used for its mean. For its shape (i.e., degree of freedom) an exponential distribution with a rate equal to 1/29 served as a prior. To determine if meta-analytic estimates were robust across the alternative priors and likelihood function, we visually compared the posterior distributions across models for large deviations<sup>58</sup>.

## Results

### Description of studies

The literature search yielded 15,900 peer reviewed journal articles, and after removing duplicates 8295 remained. Subsequent an initial screening based off the titles and abstracts, 805 studies were identified as potential candidates for modeling. 113 of these studies were deemed eligible for inclusion in the meta-analysis according to their full-text contents (Fig. 1). In total, 642 effect sizes were extracted from these studies, representing data from 4390 subjects. A majority of the effects measured the influence of exercise on executive function ( $k = 434$ ) and attention ( $k = 109$ ). Fewer effects were measured during exercise ( $k = 82$ ) relative to after the cessation of exercise ( $k = 560$ ). Visual inspection of a funnel plot suggested that the effect sizes were distributed symmetrically (Fig. 2a), however there was very strong evidence for asymmetry according to Egger's regression intercept ( $\beta = 1.18 \pm 0.25$ ; HDI = [0.78, 1.58]; BF = 253.24) suggesting the presence of publication bias. This was addressed by employing the trim and fill

approach, which imputes low-precision effect sizes until the funnel plot is symmetrical<sup>61</sup>.


### Overall effect

The meta-analysis indicated that there was moderate evidence for an acute bout of exercise to have a small positive influence on overall performance across cognitive domains ( $g = 0.13 \pm 0.04$ ; HDI = [0.06, 0.20]; BF = 3.67) (Fig. 2b, d). According to the posterior distribution, there was a low probability that the estimated pooled effect was less than or equal to zero ( $p = 0.01$ ) and an 80% chance that the effect size fell between the range of 0 to 0.2 (Fig. 2c). There was a large amount of heterogeneity within ( $\tau_{within} = 0.65 \pm 0.03$ ; HDI = [0.60, 0.70];  $I^2_{within} = 81.19\%$ ) and moderate amount between ( $\tau_{between} = 0.29 \pm 0.05$ ; HDI = [0.20, 0.38];  $I^2_{between} = 15.9\%$ ) studies (Fig. 2e). Effect size estimates for each individual study are presented in Table 1.

### Subgroup analyses

Primary subgroup analyses revealed that acute exercise reduced RT on cognitive tasks ( $g = 0.27$ ; HDI = [0.18, 0.36]; BF =  $6.71 \times 10^3$ ), but had no impact on accuracy ( $g = 0.04$ ; HDI = [-0.04, 0.12]; BF =  $6.15 \times 10^{-2}$ ) (Table 2) (Fig. 3a). Engaging in either cycling ( $g = 0.21$ ; HDI = [0.11, 0.32]; BF = 14.74) or HIIT ( $g = 0.73$ ; HDI = [0.40, 1.09]; BF = 26.05) was found to have an enhancing effect on performance in cognitive tasks (Fig. 3b). In regard to cognitive domain, there was evidence that acute exercise has a positive influence on executive processes ( $g = 0.18$ ; HDI = [0.10, 0.27];





**Fig. 2 | Meta-analysis of the effect of acute exercise on general cognitive task performance.** **a** Funnel plot of 642 study effect sizes (black circles). Imputed effect sizes after using the trim and fill method are represented by the unfilled circles ( $n = 9$ ). Vertical blue line indicates the estimated pooled effect sizes, while dashed black lines represent a pseudo 95% confidence limits. **b** Posterior distribution of estimated pooled effect. Horizontal black line indicates bounds of 89% HDI derived

using  $n = 651$  effect sizes. **c** Empirical cumulative density function of distribution in **b**, where the dashed black line indicates the pooled effect. **d** Representation of using the Savage-Dickey ratio to calculate BFs. The density of the null value in the prior distribution (red) is divided by its density in the posterior distribution (blue) to yield probabilistic evidence in favor of the alternative hypothesis. **e** Posterior distributions of between and within study heterogeneity.

BF = 36.97). Furthermore, behavioral performance was found to improve immediately after exercise cessation ( $g = 0.16$ ; HDI = [0.11, 0.30]; BF = 4.03) and in response to vigorous intensity exercises ( $g = 0.19$ ; HDI = [0.09, 0.28]; BF = 5.03). Lastly, at least moderate evidence in favor of non-zero parameter estimates were observed for the secondary moderators publication year, within-subjects design, age, percentage of female participants, and weight (Table 3).

To test for the possible contribution of a learning effect to the estimated overall pooled effect size, a separate meta-analysis was conducted on effects from studies employing a pre-/post-test design ( $N$  effect sizes = 298). Despite the estimated pooled effect size for this subset of data being nominally similar to the estimate for the entire dataset, there was anecdotal evidence in favor of the null hypothesis ( $g = 0.15 \pm 0.06$ ; HDI = [0.04, 0.24]; BF = 0.95). Moderator analyses indicated that there was no credible evidence for a difference in this estimated pooled effect size as a function of whether or not a control group was included in the study (BF<sub>Inclusion</sub> = 0.12; w/control :  $g = 0.18 \pm 0.10$ ; HDI = [0.03, 0.33]; BF = 0.51; w/o control :  $g = 0.11 \pm 0.13$ ; HDI = [-0.03, 0.26]; BF = 0.18), suggesting that the estimated influence of exercise on general cognitive performance is not driven by a learning effect.

### Model comparisons

Model comparisons were performed to determine if including a moderator improved predictive performance. Only a model that included task performance measure as a moderator was more likely when compared to a null counterpart (BF<sub>Inclusion</sub> = 357.10) (Table 4). This is likely due to a number of factors. First, acute exercise had a negligible impact on a majority of the levels in each subgroup. Second, there was a high degree of uncertainty in estimated model coefficients, as evidenced by their wide HDI intervals. Third, Bayesian inference automatically penalizes model complexity and favors more parsimonious models. If a model has many parameters, but a majority of them are nonzero, then a simpler counterpart will be favored.

### Interactions between moderators

An exploratory analysis was conducted to determine if the influence of moderator variables was contingent on one another. Due to the computationally intensive nature of Bayesian modeling, analyses were limited to the following pairs of moderators: (1) exercise intensity and type, (2) exercise intensity and duration, (3) exercise type and duration, (4) cognitive domain and exercise type, (5) cognitive domain and exercise intensity, (6) cognitive domain and task performance measure, (7) exercise type and task performance measure. Although none of the pairs of interaction models had more predictive power compared to a null counterpart (BF<sub>Inclusion</sub>: Model 1 =  $3.86 \times 10^{-6}$ ; Model 2 =  $1.66 \times 10^{-8}$ ; Model 3 =  $1.84 \times 10^{-3}$ ; Model 4 =  $1.31 \times 10^{-4}$ ; Model 5 =  $3.91 \times 10^{-5}$ ; Model 6 =  $7.1 \times 10^{-3}$ ; Model 7 =  $7.05 \times 10^{-4}$ ), there were two that had nonzero parameter estimates.

The first model included an interaction between cognitive domain and exercise type. There was evidence in favor of cycling improving performance on tasks that probed attention ( $g = 0.34$ ; HDI = [0.14, 0.56]; BF = 3.05) and executive function ( $g = 0.28$ ; HDI = [0.14, 0.40]; BF = 17.83). HIIT exercises were found to bolster executive function ( $g = 1.01$ ; HDI = [0.61, 1.43]; BF = 155.33), while resistance exercises had an aversive impact on attentional performance ( $g = -0.76$ ; HDI = [-1.20, -0.38]; BF = 18.07) (Fig. 4a). The second model included an interaction between cognitive domain and task performance measure and indicated that time-dependent measures of executive function are improved ( $g = 0.30$ ; HDI = [0.19, 0.39]; BF =  $1.10 \times 10^3$ ) (Fig. 4b).

### Sensitivity analyses

The estimated overall effect of acute exercise on cognition was consistent across the NE prior ( $g = 0.13 \pm 0.04$ ; HDI = [0.06, 0.20]; BF = 6.52), PE prior ( $g = 0.12 \pm 0.04$ ; HDI = [0.06, 0.19]; BF = 6.51), and  $t$  likelihood function ( $g = 0.12 \pm 0.04$ ; HDI = [0.06, 0.18]; BF = 8.77) (Fig. 5a). Interestingly, there was anecdotal-to-moderate evidence in favor of the synthesized estimate from previous meta-analyses (i.e.,  $g = 0.24$ ) across the PE (BF = 3.19), NE

**Table 1 | Study description and estimated effect sizes**

Study	N	Experimental design	Exercise type	Exercise intensity	Duration (min)	Cognitive domain	Test time	N effects	g	89% HDI
Hogervorst et al., <sup>124</sup>	15	Within	Cycling	Vigorous	60	Inhibition	Pre-Post	1	0.03	-0.39, 0.47
McMorris & Graydon, 1996	20	Within	Cycling	Moderate vigorous	2	Executive function	During	6	0.11	-0.19, 0.45
Brisswalter et al., 1997	20	Within	Cycling	Vigorous	40	Information processing	Pre-Post	4	0	-0.36, 0.35
Arcelin & Delignieres, 1998	22	Within	Cycling	Moderate	10	Information processing	During	2	0.03	-0.35, 0.44
McMorris et al., 1999	9	Within	Cycling	Moderate vigorous	Task completion	Executive function	During	4	0.01	-0.34, 0.38
Collardeau et al., 2001	11	Within	Running	Vigorous	90	Perception	Pre-Post	2	-0.05	-0.47, 0.34
Deuster et al., 2002	13	Within	Running	Vigorous	Volitional Exhaustion	Attention Learning Executive function	Pre-Post	24	0.16	-0.05, 0.39
Hillman et al., 2003	19	Within	Running	Vigorous	30	Executive function	Post	2	-0.08	-0.44, 0.35
Pesce et al., <sup>125</sup>	16	Within	Cycling	Moderate	Task completion	Attention	During	2	0.25	-0.15, 0.70
Davranche et al., 2005a	7	Within	Cycling	Maximal	Volitional exhaustion	Perception	Pre-Post	2	0.08	-0.32, 0.49
Davranche & Pichon, 2005b	12	Within	Cycling	Maximal	Volitional exhaustion	Perception	During	1	0.06	-0.37, 0.49
Pesce et al., <sup>126</sup>	48	Within	Cycling	Vigorous	Task duration	Attention	During	2	0.14	-0.24, 0.57
Vickers & Williams, <sup>127</sup>	10	Within	Cycling	Moderate Vigorous	2	Motor Skills	Pre-Post	4	-0.05	-0.43, 0.29
Lo Bue-Estes et al., <sup>128</sup>	26	Between	Running	Vigorous	20	Executive function Memory	Pre-Post	8	0.01	-0.28, 0.32
Coles & Tomporowski, <sup>129</sup>	18	Within	Cycling	Moderate	40	Executive function	Pre-Post	1	0	-0.41, 0.44
Fontana et al., <sup>130</sup>	32	Within	Running	Light Moderate Vigorous	4	Executive function	During	6	0.07	-0.23, 0.40
Luff et al., <sup>131</sup>	30	Within	Running	Vigorous	20	Attention information processing Memory	Pre-Post	10	-0.11	-0.39, 0.16
Pontifex et al., <sup>132</sup>	21	Within	Running Resistance	Vigorous	30	Executive function	Pre-Post	4	-0.31	-0.69, 0.05
Srygley et al., <sup>133</sup>	52	Within	Walking	Light	Task Completion	Executive function	During	2	0.03	-0.4, 0.39
Thomson et al., <sup>134</sup>	163	Within	Treadmill	Vigorous	27	Executive function	Pre-Post	2	0.1	-0.3, 0.49
Lambourne et al., <sup>6</sup>	19	Within	Cycling	Moderate	40	Executive function	Pre, During, Post	4	-0.36	-0.76, 0
Norling et al., <sup>135</sup>	121	Between	Running	Light Moderate Vigorous	30	Attention	Pre-Post	3	0.08	-0.28, 0.46
Young et al., <sup>136</sup>	27	Within	Running	Moderate	7	Motor skills	Pre-Post	1	-0.02	-0.47, 0.39
Chang et al., <sup>137</sup>	42	Between	Cycling	Vigorous	30	Executive function	Pre-Post	3	-0.02	-0.37, 0.35
Green & Helton, <sup>138</sup>	12	Within	Climbing	Moderate	3	Memory	During	1	-0.2	-0.71, 0.20
Ohlinger et al., <sup>139</sup>	50	Within	Walking	Light	Task completion	Attention Executive function	During	3	-0.01	-0.38, 0.35
Hope et al., <sup>140</sup>	52	Between	Punching	Vigorous	Volitional exhaustion	Memory	Post	3	-0.42	-0.83, -0.02
Lambourne, <sup>141</sup>	16	Within	Cycling	Vigorous	35	Executive function	Pre-Post	1	-0.01	-0.46, 0.39

**Table 1 (continued) | Study description and estimated effect sizes**

Study	N	Experimental design	Exercise type	Exercise intensity	Duration (min)	Cognitive domain	Test time	N effects	g	89% HDI
Moore et al., <sup>142</sup>	30	Between	Cycling	Moderate	6–54	Perception Executive function Attention	Pre-Post	5	-0.12	-0.43, 0.23
Queilhas Martins et al., <sup>9</sup>	24	Between	Cycling	Moderate	8	Executive function	During	1	0.1	-0.33, 0.53
Roberts & Cole, <sup>143</sup>	40	Within	Circuit	Light Moderate	1–5	Executive function	Pre-Post	8	-0.05	-0.34, 0.25
Bullock & Giesbrecht, <sup>144</sup>	26	Between	Cycling	Moderate	136	Perception	Pre, Post	8	-0.14	-0.44, 0.15
Byun et al., <sup>145</sup>	25	Within	Cycling	Light	10	Executive function	Pre-Post	2	0.17	-0.23, 0.58
Darling & Helton, <sup>146</sup>	12	Within	Climbing	Moderate	5	Memory	During	2	-0.1	-0.51, 0.29
Nibbeling et al., <sup>147</sup>	22	Between	Running	Vigorous	10	Motor Skills Executive function Memory Attention	Post	5	-0.08	-0.43, 0.24
Pontifex et al., <sup>148</sup>	34	Within	Running	Moderate	20	Attention	Pre-Post	2	0	-0.39, 0.4
Schmidt-Kassow et al., <sup>149</sup>	18; 31	Within	Walking	Light	30	Memory	During	2	0.05	-0.36, 0.43
Tsai et al., <sup>150</sup>	60	Between	Resistance	Moderate Vigorous	45	Executive function	Pre-Post	4	0.02	-0.33, 0.36
Bantoft et al., <sup>151</sup>	45	Within	Walking	Light	<60	Executive function Attention Information processing	During	4	-0.1	-0.39, 0.17
Larson et al., <sup>152</sup>	69	Between	Walking	Light	60	Executive function	During	2	-0.03	-0.43, 0.35
Osgood, <sup>153</sup>	86	Between	Sport Activity	Light	2	Attention	Post	1	-0.06	-0.47, 0.37
Perciavalle et al., <sup>154</sup>	30	Within	Cycling	Vigorous	Volitional Exhaustion	Executive function	Pre-Post	4	0.22	-0.11, 0.61
Shia et al., <sup>155</sup>	17	Within	Cycling	Vigorous	45	Attention	Pre-Post	1	-0.11	-0.55, 0.3
Stevens et al., <sup>156</sup>	35; 34	Between	Cycling	Light Moderate	15	Learning	During	4	0.04	-0.31, 0.4
Weng et al., <sup>157</sup>	26	Within	Cycling	Moderate	30	Executive function	Pre-Post	4	0.29	-0.07, 0.67
Alloway et al., <sup>158</sup>	72	Within	Running	Light	8	Executive function	Pre-Post	1	-0.01	-0.43, 0.42
Ando et al., <sup>159</sup>	14	Within	Cycling	Moderate Vigorous	6	Perception	During	2	-0.08	-0.48, 0.31
Brush et al., <sup>160</sup>	28	Within	Resistance	Light Vigorous	45	Executive function	Post	60	-0.1	-0.26, 0.04
Connell et al., <sup>161</sup>	24	Between	Cycling	Moderate	180	Attention	Pre-Post	4	-0.21	-0.6, 0.12
Hsieh et al., <sup>162</sup>	20	Within	Resistance	Moderate Vigorous	30	Executive function	Pre-Post	1	0.04	-0.39, 0.46
Hsieh et al., <sup>163</sup>	18	Within	Resistance	Moderate	30	Executive function	Post	1	0.03	-0.39, 0.47
Komiyama et al., <sup>16</sup>	10	Within	Cycling	Moderate	30	Executive function	During	3	0.04	-0.33, 0.42
Lowe et al., <sup>164</sup>	51	Within	Walking	Light Moderate	20	Executive function	Pre-Post	4	0.01	-0.34, 0.35
Torbeyns et al., <sup>165</sup>	23	Within	Cycling	Light	30	Memory Executive function Attention	During	6	-0.1	-0.42, 0.22

**Table 1 (continued) | Study description and estimated effect sizes**

Study	N	Experimental design	Exercise type	Exercise intensity	Duration (min)	Cognitive domain	Test time	N effects	g	89% HDI
Tsukamoto et al., <sup>166</sup>	12	Within	Cycling	Moderate Vigorous	40	Executive function	Pre-Post	16	-0.19	-0.45, 0.07
Zach & Shalom, <sup>167</sup>	20	Within	Running Volleyball Resistance	Vigorous	15; Game duration	Executive function	Pre-Post	6	0.37	0.02, 0.7
Chang et al., <sup>168</sup>	30	Within	Cycling	Vigorous	30	Executive function	Post	1	0.35	-0.07, 0.9
Crush & Loprinzi, <sup>169</sup>	352	Between	Running	Moderate	10; 20; 30; 45; 60	Executive function Attention	Post	75	-0.14	-0.28, 0.0
González Fernández et al., <sup>170</sup>	18	Within	Cycling	Light	45	Attention	During	1	0.02	-0.39, 0.45
Lindheimer et al., <sup>171</sup>	60	Between	Cycling	Light	25	Executive function	Pre-Post	16	-0.12	-0.35, 0.12
Lowe et al., <sup>172</sup>	28	Within	Walking	Light Moderate	20	Executive function	Pre-Post	4	-0.31	-0.68, 0.06
Luu & Hall, <sup>173</sup>	31	Within	Yoga	Light	25	Executive function	Pre-Post	2	-0.19	-0.62, 0.19
Randolph & O'Connor, <sup>174</sup>	18	Within	Walking	Light	10	Attention Executive function Information processing	Pre-Post	28	-0.06	-0.25, 0.14
Slutsky et al., <sup>175</sup>	24	Between	Cycling	Light	15	Attention Executive function	Pre-Post	7	-0.18	-0.47, 0.14
Sudo et al., <sup>176</sup>	32	Between	Cycling	Vigorous	Volitional exhaustion	Executive function	Pre-Post	3	0.28	-0.07, 0.7
Cuttler et al., <sup>177</sup>	120	Between	Resistance walking	Light moderate	30	Memory Attention	Pre-Post	6	-0.61	-0.98, -0.25
Daikoku et al., <sup>178</sup>	44	Between	Cycling	Light	3	Memory	During	1	-0.04	-0.5, 0.35
Elkana et al., <sup>179</sup>	69	Between	Cycling	Moderate	15	Executive function	Pre-Post	1	-0.03	-0.5, 0.35
Fenesi et al., <sup>180</sup>	77	Between	Calisthenics	Light	13	Learning Memory	Post	2	0.16	-0.24, 0.55
Kendall, <sup>181</sup>	60	Between	HIIT	Vigorous	20	Information processing Learning Cognitive Control	Post	8	0.82	0.42, 1.16
Legrand et al., <sup>182</sup>	101	Between	Running	Moderate	15	Attention Executive function	Pre-Post	2	0.05	-0.34, 0.44
Samani & Heath, <sup>183</sup>	14	Within	Cycling	Vigorous	10	Executive function	Pre-Post	1	0.08	-0.34, 0.52
Siddiqui & Loprinzi, <sup>184</sup>	20	Within	Walking	Moderate	20	Memory Executive function	Pre, During, Post	4	0.02	-0.31, 0.4
Sng et al., <sup>185</sup>	80	Between	Walking	Light	15	Memory	During, Post	4	-0.25	-0.62, 0.1
Wade & Loprinzi, <sup>186</sup>	34	Between	Walking	Moderate	15	Memory	Post	1	-0.13	-0.56, 0.3
Yamazaki et al., <sup>187</sup>	30	Within	Cycling	Light Moderate	10	Attention Executive function	Pre-Post	16	0.07	-0.16, 0.32
Baker et al., <sup>188</sup>	19	Within	Cycling	Light	120	Attention Executive function	During	3	-0.06	-0.44, 0.31
Du Pletz et al., <sup>189</sup>	26	Within	Cycling	Vigorous	20	Attention Executive function	Pre-Post	4	-0.06	-0.42, 0.28
Engeroff et al., <sup>190</sup>	26	Within	Resistance	Moderate Vigorous Maximal	60	Executive function	Pre-Post	4	0.1	-0.26, 0.45



**Table 1 (continued) | Study description and estimated effect sizes**

Study	N	Experimental design	Exercise type	Exercise intensity	Duration (min)	Cognitive domain	Test time	N effects	g	89% HDI
Haynes et al., <sup>191</sup>	24	Within	Walking	Light	15	Memory	During, Post	9	-0.04	-0.32, 0.24
Johnson & Loprinzi, <sup>192</sup>	40	Within	Walking	Moderate Vigorous	15	Memory	Post	2	-0.01	-0.04, 0.38
McGowan et al., <sup>193</sup>	58	Within	Running	Moderate	20	Executive function	Pre-Post	2	-0.02	-0.39, 0.39
Mehren et al., <sup>107</sup>	31; 32	Within	Cycling	Vigorous	30	Executive function	Post	8	0.15	-0.14, 0.45
Schmidt et al., <sup>194</sup>	15; 19	Within	Running	Vigorous	38	Attention	Pre-Post	6	-0.24	-0.57, 0.09
Sterling et al., <sup>195</sup>	32	Within	Walking	Moderate	3	Executive function Attention	Post	8	0	-0.29, 0.29
Wu et al., <sup>196</sup>	35	Within	Cycling Resistance	Light Vigorous	30	Executive function	Post	8	0.28	-0.03, 0.59
Zhou & Qin, <sup>197</sup>	72	Between	Cycling	Light	25	Executive function	Pre-Post	2	-0.05	-0.42, 0.37
Aly & Kojima, <sup>198</sup>	40	Between	Cycling	Light	20	Executive function	Pre-Post	2	0.07	-0.32, 0.47
Chacko et al., <sup>199</sup>	15	Within	Cycling	Vigorous	30	Executive function	Pre-Post	2	0.06	-0.32, 0.49
Kao et al., <sup>8</sup>	23	Within	Walking	Light	20	Executive function	Post	4	0.24	-0.10, 0.63
Morris et al., <sup>200</sup>	14	Within	Cycling	Moderate	30	Attention Executive function	Pre-Post	6	-0.01	-0.33, 0.3
Walsh et al., <sup>201</sup>	15; 13	Within	Walking	Moderate	20	Executive function Attention	Pre-Post	8	0.01	-0.31, 0.29
Kao et al., <sup>108</sup>	36	Within	HIIT Walking	Moderate	16	Executive function	Post	2	0.32	-0.06, 0.78
Kim et al., <sup>202</sup>	16	Within	Walking	Vigorous	10	Executive function	Pre-Post	2	-0.03	-0.38, 0.41
Klatt & Smeeton, <sup>203</sup>	27	Within	Cycling	Vigorous	Task Completion	Memory	During	4	-0.15	-0.5, 0.21
Kuhne et al., <sup>204</sup>	50	Between	Cycling	Moderate	40-55	Memory	Post	2	-0.09	-0.49, 0.29
Manocchio & Lowe, <sup>205</sup>	22	Within	Walking	Light Moderate	20	Attention Executive function	Pre-Post	8	-0.06	-0.34, 0.25
Miyashiro et al., <sup>206</sup>	17	Within	Push-Ups	Vigorous	20	Executive function	Post	1	0	-0.45, 0.41
Trammell & Aguilar, <sup>207</sup>	28	Within	Running	Vigorous	20	Memory Executive function	Pre-Post	5	-0.11	-0.47, 0.2
Zhu et al., <sup>208</sup>	16	Within	Cycling Running	Moderate Vigorous	40	Executive function	Pre-Post	16	0.1	-0.14, 0.34
Aguirre-Loaiza et al., <sup>209</sup>	19	Between	Cycling	Vigorous	20	Executive function	Pre-Post	4	-0.06	-0.41, 0.3
Drollette & Meadows, <sup>210</sup>	22	Within	HIIT	Vigorous	9	Executive function	Post	6	0.06	-0.24, 0.39
Engeroff et al., <sup>211</sup>	26	Within	Resistance	Moderate Vigorous	60	Attention	Pre-Post	16	-0.17	-0.39, 0.08
Frith et al., <sup>212</sup>	45	Within	Running	Moderate	15	Executive function	During	1	0.06	-0.4, 0.46
Kao et al., <sup>213</sup>	27	Within	Running	Moderate	24	Attention Executive function	Pre-Post	5	-0.04	-0.35, 0.31
LaCount et al., <sup>214</sup>	18	Within	HIIT	Vigorous	16	Attention Executive function Information processing	Post	3	-0.09	-0.46, 0.28
Loprinzi & Storm, <sup>215</sup>	180; 225; 158	Between	Treadmill	Moderate Vigorous	25	Memory	Post	6	-0.02	-0.33, 0.3

Table 1 (continued) | Study description and estimated effect sizes

Study	N	Experimental design	Exercise type	Exercise intensity	Duration (min)	Cognitive domain	Test time	N effects	g	89% HDI
Shirzad et al., <sup>216</sup>	28	Within	Cycling	Light	26	Executive function	Pre-Post	2	-0.23	-0.64, 0.17
Zheng et al., <sup>217</sup>	27	Within	Cycling	Moderate	15	Executive function	During	2	0.27	-0.11, 0.72
Chueh et al., <sup>218</sup>	30	Within	Step Exercise	Vigorous	20	Attention Executive function	Pre-Post	8	0.15	-0.15, 0.44
Yamada et al., <sup>219</sup>	85	Within	Recumbent Bike	Vigorous	20	Executive function	Pre-Post	4	0.02	-0.31, 0.38
B. Zhang et al., <sup>220</sup>	18	Within	Cycling	Moderate	25	Perception	Post	4	0.29	-0.09, 0.65
D. Zhang et al., <sup>221</sup>	76	Within	Cycling	Light	20	Attention	Pre-Post	4	0.03	-0.33, 0.36

(BF = 2.78), and standard normal (BF = 5.27) priors. Estimates of between-study heterogeneity were also robust across the NE prior ( $\tau_{between} = 0.29 \pm 0.05$ ; HDI = [0.20, 0.37]), the PE prior ( $\tau_{between} = 0.29 \pm 0.05$ ; HDI = [0.21, 0.37]), and *t* likelihood function ( $\tau_{between} = 0.31 \pm 0.03$ ; HDI = [0.26, 0.38]) (Fig. 5b). In contrast, within study heterogeneity was estimated to be lower when using the *t* likelihood function ( $\tau_{within} = 0.17 \pm 0.02$ ; HDI = [0.13, 0.19]) relative to the NE ( $\tau_{within} = 0.65 \pm 0.03$ ; HDI = [0.61, 0.70]) and PE ( $\tau_{within} = 0.65 \pm 0.02$ ; HDI = [0.60, 0.70]) priors (Fig. 5c). Note, this reduction reflects the diminished influence of outliers on variance estimates by the inclusion of the shape parameter for the *t* distribution ( $\nu = 1.52 \pm 0.14$ ; HDI = [1.30, 1.73]). In addition to testing the robustness of parameter estimates, a model comparison was conducted to determine if either the null or positive effect prior was more probable given the data. The *t*-likelihood function was not included in this comparison since it would only indicate if effect sizes were more likely to have been drawn from either a normal or *t*-distribution. When compared to a standard normal prior, there was anecdotal evidence in favor of both the PE (BF = 2.56) and NE (BF = 1.48) priors. Relative to the PE prior, there was anecdotal evidence against the NE prior (BF = 0.73). Altogether, parameter estimates were not biased by the prior or likelihood function.

### Executive function meta-analysis

Considering that the majority of the effect sizes were from tasks that probed executive function, and that this cognitive domain encompasses multiple sub-domains, a separate meta-analysis and set of meta-regressions were conducted on this subset of data. Categorization criteria from previous meta-analyses and systematic reviews<sup>17,24,26</sup> were used to classify effect sizes into the following sub-domains of executive function: working memory, cognitive control, decision making, planning, and inhibition. For completeness, the primary moderators used in the main meta-analysis were also tested.


The results were similar to the main meta-analysis. There was very strong evidence in favor of exercise having a small positive influence on overall task performance ( $g = 0.20 \pm 0.06$ ; HDI = [0.12, 0.30]; BF = 29.57), and a moderate degree of heterogeneity both within ( $\tau_{within} = 0.51 \pm 0.03$ ; HDI = [0.47, 0.57]) and between studies ( $\tau_{between} = 0.40 \pm 0.06$ ; HDI = [0.30, 0.48]). Subgroup analyses indicated that a model including the moderator task outcome measure had more predictive power relative to a null counterpart (BF<sub>Inclusion</sub> = 48.43). Paralleling the main meta-analysis, there was very strong evidence that acute exercise improved RT on executive function tasks ( $g = 0.32$ ; HDI = [0.21, 0.42]; BF = 748.18), but no credible evidence was observed for an effect on accuracy ( $g = 0.13$ ; HDI = [0.04, 0.23]; BF = 0.63) (Table 5). Furthermore, there was moderate evidence in favor of a positive impact of exercise on inhibition ( $g = 0.21$ ; HDI = [0.09, 0.33]; BF = 3.14) and working memory ( $g = 0.22$ ; HDI = [0.11, 0.34]; BF = 6.89) (Fig. 6). Yet, a model including executive function sub-domain as a moderator did not improve model performance (BF<sub>Inclusion</sub> =  $7.52 \times 10^{-4}$ ), nor did models including interactions between moderators.

### Discussion

A large corpus of empirical work has examined how a single bout of acute exercise modulates activity within multiple brain systems that underly cognition. Despite inconsistencies in results across empirical studies, there is consensus amongst previous reviews and meta-analyses that acute exercise impacts behavioral performance<sup>18,20,22</sup> and that this relationship is moderated by both exercise protocol and behavioral task characteristics. The goal of the present work was to address two key limitations of previous meta-analyses. First, recent meta-analyses have a narrower focus, often limited to a single cognitive domain or a specific subset of domains. In contrast, the current meta-analysis presents an updated synthesis of the literature spanning a much wider range of cognitive domains. Second, in contrast to previous frequentist approaches, a Bayesian framework was adopted allowing for the quantification of the degree of evidence in favor of the hypothesis that acute exercise influences cognition in young healthy adults.

**Table 2 | Primary moderator estimates**

Exercise moderator variable	N	g	89% HDI	BF <sub>10</sub>	Cognitive moderator variable	N	g	89% HDI	BF <sub>10</sub>
Intensity					Domain				
Light	167	0.10	-0.02, 0.22	0.13	Attention	109	0.06	-0.08, 0.17	0.08
Moderate	222	0.07	-0.03, 0.18	0.09	Executive function	434	0.18	0.10, 0.27	36.97
Vigorous	253	0.19	0.09, 0.28	5.03	Information processing	15	0.12	-0.17, 0.41	0.14
					Learning	12	0.24	-0.11, 0.59	0.25
Duration (minutes)					Memory	44	-0.06	-0.25, 0.13	0.08
≤16	161	0.14	0.02, 0.26	0.30	Motor skills	6	-0.03	-0.51, 0.49	0.20
20–27	152	0.15	0.02, 0.26	0.32	Perception	22	0.13	-0.17, 0.44	0.15
30–35	93	0.08	-0.08, 0.22	0.09					
40–45	113	0.04	-0.13, 0.21	0.09					
>60	48	-0.03	-0.21, 0.15	0.08	Task outcome				
Not provided	75	0.37	0.17, 0.57	6.21	Accuracy	377	0.04	-0.04, 0.12	6.15e-2
					Reaction time	265	0.27	0.18, 0.36	6.71e3
Type circuit	8	0.08	-0.33, 0.53	0.26					
Cycling	204	0.21	0.11, 0.32	14.74	Task completion time (relative to exercise)				
HIIT	18	0.73	0.40, 1.09	26.05	During	82	0.02	-0.18, 0.13	0.09
Resistance	97	-0.06	-0.29, 0.14	0.11	Immediately after	315	0.16	0.11, 0.30	4.03
Running	172	0.05	-0.10, 0.19	0.08	20–75 min post	94	0.22	0.13, 0.44	0.76
Sport activity	26	0.04	-0.23, 0.29	0.11	>180 min post	151	0.08	-0.03, 0.28	0.10
Walking	117	0.04	-0.10, 0.19	0.07					



**Fig. 3 | Subgroup analyses.** Posterior distributions of **a** cognitive and **b** exercise moderators. Horizontal black line indicates the 89% HDI interval, while the black dot represents the mode of the posterior distribution. Intervals derived using  $n = 651$  effect sizes.

The current meta-analysis observed that acute exercise has a small positive influence on overall cognitive task performance, and sensitivity analyses indicated that the alternative hypothesis was 6.51–8.77 times more likely than the null across multiple priors and likelihood functions. The magnitude and directionality of this effect were consistent with the results of previous

meta-analyses on acute exercise and cognition<sup>18,21,22,62</sup>. Subgroup analyses suggested that this relationship is moderated by task performance measure, cognitive domain, exercise type and intensity, and the time of task completion relative to exercise cessation. Model comparison results indicated that accounting for variations amongst moderator levels did not improve

**Table 3 | Secondary moderator estimates**

Moderator variable	N	g	89% HDI	BF <sub>10</sub>
Publication year	642	0.13	0.06, 0.20	3.25
Exp. design				
Between	193	0.03	-0.11, 0.16	0.07
Within	449	0.17	0.08, 0.24	12.18
Age (years) ( $\mu = 22.49$ )	599	0.14	0.06, 0.21	3.13
% Female	577	0.12	0.05, 0.20	4.23
BMI (kg/m <sup>2</sup> ) ( $\mu = 24.02$ )	378	0.20	0.08, 0.30	1.28
VO2 max (ml/kg/min) ( $\mu = 43.22$ )	299	0.19	0.05, 0.34	1.08
Height (cm) ( $\mu = 158.55$ )	275	0.21	0.09, 0.33	2.90
Weight (kg) ( $\mu = 65.02$ )	283	0.21	0.08, 0.33	4.28

**Table 4 | Subgroup model comparisons**

Model	BF <sub>Inclusion</sub>
No moderators	1
Exercise intensity	2.16e-4
Exercise duration	9.35e-4
Exercise type	2.00e-2
Cognitive domain	4.97e-3
Task outcome	357.10
Task completion time	5.52e-4
Publication date	9.05e-4
Experimental design	0.04
Age	0
% Female	4.53e-3
BMI	0.04
VO2	8.40e-3
Height	2.89e-3
Weight	0.01

predictive performance. Given our eligibility criteria, these results are limited to healthy individuals between the ages of 18–45 years old.



Similar to McMorris et al.<sup>63</sup>, acute exercise was found to improve RT but no credible evidence was observed for an influence on accuracy. A possible explanation for this differential impact on task outcome measures is that exercise modulates primary motor cortex (M1) excitability<sup>64</sup>. There is accumulating evidence that acute exercise increases M1 intracortical facilitation<sup>65–68</sup> and inhibition<sup>69,70</sup>. Yamazaki et al.<sup>68</sup> observed that the intracortical circuits of both exercised (i.e., legs) and non-exercised (i.e., hand) effectors are modulated by an acute bout of low intensity pedaling. Thus, alterations in the activity of excitatory or inhibitory circuits of non-exercised cortical representations may promote faster RT on cognitive tasks. However, the lack of concurrent changes in corticospinal excitability or motor-evoked potentials suggests that this explanation is not a viable account of a mechanism that engenders faster RTs. An alternative explanation is that exercise increases peripheral and central concentrations of catecholamines, such as norepinephrine, epinephrine, and dopamine, which in turn improves the speed of cognition<sup>1,71,72</sup>. Indeed, acute exercise has been found to improve response time on choice RT, decision-making, and interference tasks<sup>18,73,74</sup>. Yet, it is unclear as to why changes in neurochemical levels would facilitate RT but have no impact on accuracy. Considering that physical activity modulates population-level tuning in the sensory areas of nonhuman animals and invertebrates<sup>75–81</sup>, along with sensory responses in humans<sup>82–84</sup>, it stands to reason that the fidelity of stimulus representations would also be impacted, resulting in changes in

accuracy. Changes in the fidelity of feature selective stimulus representations can be determined by applying encoding models to recorded neural activity<sup>83,85–90</sup>. For instance, Garrett et al.<sup>91</sup> applied an inverted encoding model to topographical patterns of alpha band activity, recorded at the scalp, while subjects completed a spatial working memory task both at rest and during a bout of moderate intensity cycling. Notably, it was possible to reconstruct spatially selective responses during exercise, and the selectivity of these responses decreased during exercise relative to rest. Therefore, encoding models can be a powerful tool for future research to demystify how the precision of task-relevant representations is influenced by exercise. It is also important to keep in mind that many psychological tasks are relatively simple to do, which can lead to ceiling effects that may mask the influence of exercise on accuracy measures. Lastly, the differential impact of exercise on accuracy and RT may be due to the relative sensitivities of these dependent measures to modulations of different stages of information processing. For example, there is evidence that in near-threshold tasks accuracy is sensitive to perceptual manipulations, whereas in supra-threshold (i.e., perceptually easy tasks, including many of those used in the studies in this meta-analysis) RT is sensitive to modulations in both perceptual and post-perceptual processes<sup>92,93</sup>. Indeed, Davranche et al.<sup>73</sup> utilized a drift diffusion model to determine which aspects of decision-making are modulated by HIIT. Importantly, drift rate and decision response boundary size increased significantly after exercise relative to before, while non-decision time decreased. This suggests there was an improvement in perceptual discrimination, the efficiency of non-decisional processes (e.g., motor execution), and the adoption of a more conservative criterion. Future research employing computational models of response time and representational fidelity is needed to develop a comprehensive understanding of the selective influence exercise on information processing speed and accuracy.

Parameter estimates of a model including exercise modality as a moderator suggested that engaging in cycling or HIIT may beneficially impact cognition, especially on attentional and executive processes. Cycling is a commonly used modality in exercise and cognition research. Numerous empirical studies have found that a bout of cycling benefits inhibition, as measured using either the Stroop or Eriksen Flanker task<sup>15,94–99</sup>. Improvements in planning<sup>94,100</sup>, task-switching<sup>27,101,102</sup>, and the speed of decision making<sup>103</sup> have also been reported. In contrast to the ubiquity of cycling, the use of HIIT workouts in exercise and cognition research is a relatively recent practice, hence the small number of effect sizes from studies using this modality compared to other types of exercise. The number of effect sizes is important because low-level parameters in a hierarchical model are influenced both by the subset of data directly dependent on the low-level parameter, and by high-level parameter estimates that rely on all of the data. This makes low-level parameter estimates indirectly dependent on the entire dataset, and causes shrinkage in estimates at all levels of the model. In other words, the estimated relationship between HIIT and behavioral performance is derived directly from the few representative effect sizes and indirectly from the rest of the data. The observed positive effect of HIIT on cognition corroborates previous findings. For example, Alves et al.<sup>3</sup> observed that the time to complete a Stroop Task decreased after ten 1-minute bouts of exercising at 80% heart rate reserve relative to a control condition. Improvements in time-dependent measures on interference tasks (i.e., Stroop and flanker) have been correlated with an increase in left dorsolateral prefrontal cortex activity, as measured with functional near infrared spectroscopy (fNIRS), and a decrease in P3 latency measured with EEG<sup>104</sup>. Furthermore, enhancements have also been shown to coincide with an increase in peripheral levels of neural growth factors and lactate<sup>105</sup>. Lastly, a recent meta-analysis on elite athletes observed that HIIT team-based sports had a positive impact on cognitive task performance<sup>25</sup>. Interestingly, because of the small number of published studies in the literature, it is currently unclear if the type of exercise modality used for HIIT workouts (e.g., cycling, sprinting, resistance) differentially impacts cognition.

Behavioral task performance was found to be improved by engaging in vigorous intensity exercise. These results are surprising, considering that exercise intensity is believed to have an inverted-U relationship with

**Fig. 4 | Interactions between cognitive and exercise moderators.** Posterior mode estimates of models including interactions between cognitive domain and **a** exercise type and **b** task outcome measure. Width of line represents 89% HDI derived using  $n = 651$  effect sizes.



**Fig. 5 | Sensitivity analyses.** Estimates for the **a** overall pooled effect size, **b** between- and **c** within-study heterogeneity parameters across the t-likelihood function (TL), weakly informed, null effect (NE), and positive effect (PE) priors. Color dots

represent mode of posterior distributions, while color horizontal line depicts the 89% HDI derived using  $n = 651$  effect sizes.

performance; where moderate intensity exercise elicits the greatest enhancements while more intense, fatiguing exercise imposes decrements<sup>18,62,63,71,106,107</sup>. This effect could be driven by HIIT workouts, but may also depend on multiple cognitive task and exercise protocol characteristics. For instance, Chang et al.<sup>18</sup> observed that exercise intensity was only a significant moderator when cognition was tested post-exercise. Similarly, Oberste et al.<sup>23</sup> found that exercise intensity influenced time-dependent measures of interference control but not accuracy. When considering these results, one must also consider that both aforementioned meta-analyses included studies whose subjects were children, adolescents, and older adults. In contrast, the current study was limited to young adults, and there is evidence that the effect of exercise on cognition is comparatively smaller in this age group<sup>18,23</sup>. Thus, a model containing an interaction between cognitive domain, task outcome measure, and age groups across the lifespan may be required to observe evidence for an effect of intensity. In addition, there was evidence for the enhancing effects of exercise post-cessation, corroborating previous research<sup>1,18,94</sup>. Interestingly, in the current meta-analysis cognition was not found to be impacted during exercise. Prior meta-analytic findings on cognition during exercise are mixed, with some reporting that it is exacerbated<sup>20</sup>, while others that find evidence for an enhancement<sup>18</sup>.

Given that the majority of the effect sizes were from tasks that probed executive function, a separate meta-analysis was conducted on this subset of data. This analysis revealed that exercise has a small positive impact on RT


measures of executive processes. When looking at model parameters, there was evidence in favor of exercise-enhancing inhibition and working memory. Behavioral research has shown that both the accuracy<sup>9</sup> and speed of working memory<sup>108,109</sup> are facilitated by an instance of physical activity. What remains to be determined is the neural mechanisms that engender these behavioral effects. Kao et al.<sup>108</sup> observed that a reduction in RT on the Sternberg task post-HIIT corresponded to an increase in frontal alpha desynchronization during encoding, maintenance, and retrieval periods when working memory load is high. Neuroimaging studies have also found evidence for changes in the activation levels of frontal areas<sup>110</sup> and their connectivity with the intraparietal sulcus post-exercise<sup>111</sup>. These changes in neural activity were not accompanied by a change in behavior, suggesting that more research is needed to demystify the neuromodulatory effect of acute exercise on working memory.

Engaging in repeated bouts of acute exercise over a long period of time can have lasting changes on baseline neurochemical levels, cortical volume, and structural/functional connectivity, which can alter cognitive task performance<sup>1,112-115</sup>. Research investigating the influence of these long-term interventions on cognition has primarily focused on children or older adults. Systematic reviews and meta-analyses suggest that exercise has a small to moderate beneficial impact on general task performance for both of these age groups, with the largest effect sizes observed for measures of executive function, attention, and academic performance<sup>35</sup>. Despite the relative paucity of meta-analyses on how exercise interventions impact cognition in



**Table 5 | Executive function moderator estimates**

Exercise moderator variable	N	g	89% HDI	BF <sub>10</sub>	Cognitive moderator variable	N	g	89% HDI	BF <sub>10</sub>
Intensity					Sub-domain				
Light	107	0.23	0.08, 0.37	1.71	Cognitive control	55	0.23	0.06, 0.39	0.73
Moderate	151	0.15	0.01, 0.28	0.30	Decision making	28	0.10	-0.16, 0.39	0.14
Vigorous	175	0.24	0.12, 0.36	13.49	Inhibition	153	0.21	0.09, 0.33	3.14
					Planning	18	0.14	-0.12, 0.38	0.15
Duration (minutes)					Working memory	179	0.22	0.11, 0.34	6.89
≤16	93	0.25	0.09, 0.40	1.76					
20–27	99	0.13	-0.02, 0.27	0.19	Task outcome				
30–35	68	0.19	0.03, 0.36	0.44	Accuracy	253	0.13	0.04, 0.23	0.63
40–45	100	0.13	-0.06, 0.32	0.16	Reaction time	180	0.32	0.21, 0.42	749.18
>60	25	0.11	-0.12, 0.32	0.13					
Not provided	48	0.53	0.29, 0.83	24.40	Type				
					Circuit	8	0.15	-0.31, 0.65	0.31
Task completion time (relative to exercise)					Cycling	133	0.28	0.15, 0.42	19.58
During	39	0.23	0.004, 0.44	0.45	HIIT	12	0.96	0.56, 1.38	71.23
Immediately after	197	0.21	0.10, 0.31	4.17	Resistance	90	-0.07	-0.33, 0.17	0.12
20–75 min post	72	0.31	0.13, 0.47	4.11	Running	116	0.06	-0.12, 0.25	0.10
>180 min post	125	0.13	-0.03, 0.29	0.17	Sport activity	11	0.30	-0.07, 0.68	0.35
					Walking	63	0.14	-0.05, 0.34	0.18



**Fig. 6 | Subgroup analyses of effects from tasks testing executive function.** Posterior distributions for executive function sub-domain. Horizontal black line indicates the 89% HDI interval, while the black dot represents the mode of the posterior distribution, which was derived using  $n = 433$  effect sizes.

healthy young adults, recent work suggests that it may have a similar beneficial effect. Indeed, a recent meta-analysis, conducted by Ludyga et al.<sup>116</sup>, indicated that long-term exercise interventions have a small positive influence on general cognition regardless of age. The magnitude of this effect was dependent on the interaction between intervention length and exercise duration, with longer interventions and sessions producing greater benefits. Integrating these findings with the current meta-analysis, there is support for the notion that the beneficial impact of long-term interventions on cognition may be a product of repeated exposure to acute exercise induced effects.

There are a number of possible explanations as to why exercise induced effects are small. One possibility is that cognitive function is at its peak during young adulthood, leaving little room for improvements in task performance. Indeed, previous reviews and meta-analyses have observed that the effect of exercise is moderated by age<sup>35</sup>, with the greatest benefits observed for preadolescent children and older adults<sup>18,23,26</sup>. Contrary to this

account, though, the largest exercise induced effects were observed for executive processes, which are believed to be at peak efficiency during this period in the lifespan<sup>117,118</sup>. Furthermore, there was moderate evidence that the impact of exercise increased as the average age of sampled young adults also increased. Another explanation may be that cognition is resilient to slight or modest perturbations in overall global state. For example, Bullock et al.<sup>119</sup> demonstrated there was no change in accuracy or RT on a target detection task during experimentally induced hypoxia, hypercapnia, hypocapnia, and normoxia. Meta-analytic modeling of the influence of acute stress on executive function revealed that stress has a small negative impact on working memory and cognitive flexibility, but no impact on inhibition<sup>120</sup>. This suggests that cognition is able to selectively adapt to changes in physiological state caused by various types of stressors, including exercise. A final more intriguing and functional explanation for exercise not typically require the engagement of the body to execute the cognitive task, but rather have people engage in a cognitive task while exercising (or shortly thereafter). This experimental design contrasts real-world tasks that require engagement of the body in the service of the cognitive task. When components of the exercise are incorporated into task goals, then larger changes in performance may be observed. Empirical research investigating how exercise influences task performance in embodied settings versus classic laboratory settings (see<sup>121</sup> for review) is necessary to test the plausibility of this explanation. In addition, the notion that the integrated action of the body and the mind are required to produce the largest effects of exercise on cognition is consistent with a recent evolutionary account of the link between cognition and exercise<sup>122</sup>.

The discrepancy in moderator results between the current meta-analysis and previous meta-analyses could be due to differences in the statistical approach. Frequentist methods typically conduct an omnibus test to determine if levels of a moderator are significantly different from one another and as a measure of a model's goodness of fit. In contrast, the Bayesian approach determines how likely the observed effect sizes are under a model that includes a moderator and if predictive power is increased. There are a few key advantages to using the Bayesian approach compared to classical frequentist methods. First, it models the uncertainty involved in estimates of between- and within-study heterogeneity and returns a full

posterior distribution for both parameters<sup>123</sup>. With these posterior distributions, one can simulate possible pooled effect sizes across credible levels of heterogeneity and develop an informed hypothesis for a subsequent meta-analysis. Similarly, the posterior distributions of effect size estimates can be used as well-informed prior distributions for new data. Importantly, this facilitates the updating of meta-analyses as new research is published. It should be mentioned that the degree of between-study heterogeneity was numerically similar to previous meta-analyses<sup>18,22</sup>, implying that they did not suffer from an issue of underestimation by assuming heterogeneity to be a fixed quantity. Second, the Bayesian approach permits the inclusion of prior knowledge. Across all tested priors, there was evidence in favor of a pooled effect derived from averaging the reported estimates of previous meta-analyses. When comparing a prior distribution based on this knowledge to a null effect prior, the former was found to be more probable. Lastly, the posterior distribution of parameter estimates can be used to ascertain the likelihood that one will observe an effect size of a given magnitude for an exercise protocol and cognitive task combination. For example, a researcher could compute the probability that the influence of a bout of cycling on cognitive control will fall within the range of large effect sizes, even if that range does not encompass the maximum a posteriori probability estimate. In contrast, the frequentist approach only produces the maximum likelihood estimate and an interval around it based on fictitious repeats of the meta-analysis. Therefore, the Bayesian approach provides more information for designing future exercise and cognition studies.

### Limitations

A potential limitation in the current meta-analysis is the categorization of exercise type using the activity reported in each study. An alternative approach is to categorize exercise based on the theoretical and physiological distinctions between aerobic and anaerobic exercise. We did not adopt this approach here because many activities used in the literature typically include aerobic and anaerobic components, and basing their classification on what authors reported provides insights into the exercise modalities that have been predominantly used in the literature. Another limitation is the schema used to categorize exercise duration. In the event that a study did not report how long participants engaged in exercise, these effects were classified as “not provided”, rendering them as uninterpretable. Lastly, sensitivity analyses were not conducted for moderator parameter estimates due to the high degree of computational demands. However, considering that the pooled effect size estimate was robust across multiple priors and likelihood functions, it is likely that moderator parameter estimates are also consistent.

### Conclusions

In summary, the current meta-analytic examination has shown that there is moderate evidence for an acute bout of aerobic exercise inducing a small enhancement in overall performance on cognitive tasks, especially on those that probe executive function and measure response time. Incorporating computational models of decision-making processes, such as drift-diffusion or signal detection models, into exercise research may provide useful insights into the nature of speeded executive processes. Furthermore, testing performance in a real-world setting where individuals typically engage in physical activity may amplify exercise-induced effects.

### Data availability

Data have been made publicly available at <https://github.com/jggarrett23/PACMan>.

### Code availability

Code has been made publicly available at <https://github.com/jggarrett23/PACMan>.

Received: 12 June 2023; Accepted: 31 July 2024;

Published online: 28 August 2024

### References

1. Basso, J. C. & Suzuki, W. A. The effects of acute exercise on mood, cognition, neurophysiology, and neurochemical pathways: a review. *Brain Plast.* **2**, 127–152 (2017).
2. Hillman, C. H., Erickson, K. I. & Kramer, A. F. Be smart, exercise your heart: exercise effects on brain and cognition. *Nat. Rev. Neurosci.* **9**, 58–65 (2008).
3. Alves, C. R. R. et al. Influence of acute high-intensity aerobic interval exercise bout on selective attention and short-term memory tasks. *Percept. Mot. Skills* **118**, 63–72 (2014).
4. Chang, Y. K., Pesce, C., Chiang, Y. T., Kuo, C. Y. & Fong, D. Y. Antecedent acute cycling exercise affects attention control: an ERP study using attention network test. *Front. Hum. Neurosci.* **9**, 156 (2015).
5. Davranche, K. & Audiffren, M. Facilitating effects of exercise on information processing. *J. Sports Sci.* **22**, 419–428 (2004).
6. Lambourne, K., Audiffren, M. & Tomporowski, P. D. Effects of acute exercise on sensory and executive processing tasks. *Med. Sci. Sports Exerc* **42**, 0–00 (2010).
7. Chen, A. G., Zhu, L. N., Yan, J. & Yin, H. C. Neural basis of working memory enhancement after acute aerobic exercise: FMRI study of preadolescent children. *Front. Psychol.* **7**, 1804 (2016).
8. Kao, S. C., Wang, C. H. & Hillman, C. H. Acute effects of aerobic exercise on response variability and neuroelectric indices during a serial n-back task. *Brain Cogn.* **138**, 105508 (2020).
9. Quelhas Martins, A., Kavussanu, M., Willoughby, A. & Ring, C. Moderate intensity exercise facilitates working memory. *Psychol. Sport Exerc.* **14**, 323–328 (2013).
10. Roig, M., Nordbrandt, S., Geertsen, S. S. & Nielsen, J. B. The effects of cardiovascular exercise on human memory: a review with meta-analysis. *Neurosci. Biobehav. Rev.* **37**, 1645–1666 (2013).
11. Schaefer, S., Lövdén, M., Wieckhorst, B. & Lindenberger, U. Cognitive performance is improved while walking: differences in cognitive–sensorimotor couplings between children and young adults. *Eur. J. Dev. Psychol.* **7**, 371–389 (2010).
12. Kamijo, K. et al. Differential influences of exercise intensity on information processing in the central nervous system. *Eur. J. Appl. Physiol.* **92**, 305–311 (2004).
13. Kamijo, K. & Takeda, Y. General physical activity levels influence positive and negative priming effects in young adults. *Clin. Neurophysiol.* **120**, 511–519 (2009).
14. Hillman, C. H. et al. The effect of acute treadmill walking on cognitive control and academic achievement in preadolescent children. *Neuroscience* **159**, 1044–1054 (2009).
15. Kamijo, K., Nishihira, Y., Higashiura, T. & Kuroiwa, K. The interactive effect of exercise intensity and task difficulty on human cognitive processing. *Int. J. Psychophysiol.* **65**, 114–121 (2007).
16. Komiyama, T. et al. Cognitive function at rest and during exercise following breakfast omission. *Physiol. Behav.* **157**, 178–184 (2016).
17. Cantelon, J. A. & Giles, G. E. A review of cognitive changes during acute aerobic exercise. *Front. Psychol.* **12**, 653158 (2021).
18. Chang, Y. K., Labban, J. D., Gapin, J. I. & Etnier, J. L. The effects of acute exercise on cognitive performance: a meta-analysis. *Brain Res.* **1453**, 87–101 (2012).
19. Haverkamp, B. F. et al. Effects of physical activity interventions on cognitive outcomes and academic performance in adolescents and young adults: a meta-analysis. *J. Sports Sci.* **38**, 2637–2660 (2020).
20. Lambourne, K. & Tomporowski, P. The effect of exercise-induced arousal on cognitive task performance: a meta-regression analysis. *Brain Res.* **1341**, 12–24 (2010).
21. McMorris, T., Sproule, J., Turner, A. & Hale, B. J. Acute, intermediate intensity exercise, and speed and accuracy in working memory tasks: a meta-analytical comparison of effects. *Physiol. Behav.* **102**, 421–428 (2011).

22. Moreau, D. & Chou, E. The acute effect of high-intensity exercise on executive function: a meta-analysis. *Perspect. Psychol. Sci.* **14**, 734–764 (2019).
23. Oberste, M. et al. Effects and moderators of acute aerobic exercise on subsequent interference control: a systematic review and meta-analysis. *Front. Psychol.* **10**, 2616 (2019).
24. Ishihara, T., Drollette, E. S., Ludyga, S., Hillman, C. H. & Kamijo, K. The effects of acute aerobic exercise on executive function: a systematic review and meta-analysis of individual participant data. *Neurosci. Biobehav. Rev.* **128**, 258–269 (2021).
25. Logan, N. E., Henry, D. A., Hillman, C. H. & Kramer, A. F. Trained athletes and cognitive function: a systematic review and meta-analysis. *Int. J. Sport Exerc. Psychol.* **21**, 725–749 (2023).
26. Ludyga, S., Gerber, M., Brand, S., Holsboer-Trachsler, E. & Pühse, U. Acute effects of moderate aerobic exercise on specific aspects of executive function in different age and fitness groups: a meta-analysis. *Psychophysiology* **53**, 1611–1626 (2016).
27. Oberste, M., Sharma, S., Bloch, W. & Zimmer, P. Acute exercise-induced set shifting benefits in healthy adults and its moderators: a systematic review and meta-analysis. *Front. Psychol.* **12**, 2 (2021).
28. Borenstein, M., Hedges, L. V., Higgins, J. P. T. & Rothstein, H. R. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res. Synth. Methods* **1**, 97–111 (2010).
29. Hackenberger, B. K. Bayesian meta-analysis now – let’s do it. *Croat. Med. J.* **61**, 564 (2020).
30. Sutton, A. J. et al. Methods for meta-analysis in medical research methods for meta-analysis in medical research contents preface acknowledgements part a: meta-analysis methodology: the basics. <https://www.researchgate.net/publication/236905532> (2000).
31. Sutton, A. J. & Abrams, K. R. Bayesian methods in meta-analysis and evidence synthesis. *Stat. Methods Med. Res.* **10**, 277–303 (2001).
32. Chung, Y., Rabe-Hesketh, S. & Choi, I. H. Avoiding zero between-study variance estimates in random-effects meta-analysis. *Stat. Med.* **32**, 4071–4089 (2013).
33. Kontopantelis, E., Springate, D. A. & Reeves, D. A re-analysis of the Cochrane Library data: the dangers of unobserved heterogeneity in meta-analyses. *PLoS one* **8**, e69930 (2013).
34. Sidik, K. & Jonkman, J. N. A comparison of heterogeneity variance estimators in combining results of studies. *Stat. Med.* **26**, 1964–1981 (2007).
35. Erickson, K. I. et al. Physical activity, cognition, and brain outcomes: a review of the 2018 physical activity guidelines. *Med Sci. Sports Exerc.* **51**, 1242–1251 (2019).
36. Stillman, C. M., Esteban-Cornejo, I., Brown, B., Bender, C. M. & Erickson, K. I. Effects of exercise on brain and cognition across age groups and health states. *Trends Neurosci.* **43**, 533–543 (2020).
37. Thompson, S. G. & Sharp, S. J. Explaining heterogeneity in meta-analysis: a comparison of methods. *Stat. Med.* **18**, 2693–2708 (1999).
38. Williams, D. R., Rast, P. & Bürkner, P.-C. Bayesian meta-analysis with weakly informative prior distributions. <https://doi.org/10.31234/OSF.IO/7TBRM> (2018).
39. Kruschke, J. K. & Liddell, T. M. The Bayesian new statistics: hypothesis testing, estimation, meta-analysis, and power analysis from a bayesian perspective. *Psychon. Bull. Rev.* **25**, 178–206 (2018).
40. Kruschke, J. K. Doing Bayesian data analysis: a tutorial with R, JAGS, and Stan, 2nd Edn. <https://doi.org/10.1016/B978-0-12-405888-0.09999-2> (Elsevier Science, 2014).
41. Moher, D. et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst. Rev.* **20**, 148–160 (2015).
42. Sachdev, P. S. et al. Classifying neurocognitive disorders: the DSM-5 approach. *Nat. Rev. Neurol.* **10**, 634–642 (2014).
43. Garber, C. E. et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory. *Med. Sci. Sports Exerc.* **43**, 1334–1359 (2011).
44. Hedges, L. V. & Olkin, I. Statistical methods for meta-analysis. <https://doi.org/10.1016/c2009-0-03396-0> (Elsevier, 1985).
45. Borenstein, M., Hedges, L. V., Higgins, J. P. T. & Rothstein, H. Introduction to meta-analysis. (Wiley, 2021).
46. Higgins, J. P. T., Thompson, S. G. & Spiegelhalter, D. J. A re-evaluation of random-effects meta-analysis. *J. R. Stat. Soc. Ser. A, (Stat. Soc.)* **172**, 137 (2009).
47. Röver, C. Bayesian random-effects meta-analysis using the Bayesmeta R package. *J. Stat. Softw.* **93**, 1–51 (2020).
48. Bürkner, P. C. Advanced Bayesian multilevel modeling with the R package brms. *R. J.* **10**, 395–411 (2018).
49. Makowski, D., Ben-Shachar, M. & Lüdtke, D. bayestestR: describing effects and their uncertainty, existence and significance within the bayesian framework. *J. Open Source Softw.* **4**, 1541 (2019).
50. Hinne, M., Gronau, Q. F., van den Bergh, D. & Wagenmakers, E.-J. A conceptual introduction to bayesian model averaging. *Adv. Methods Pract. Psychol. Sci.* **3**, 200–215 (2020).
51. Schad, D. J., Betancourt, M. & Vasisht, S. Toward a principled Bayesian workflow in cognitive science. *Psychol. Methods* **26**, 103–126 (2021).
52. Jeffreys, H. Theory of probability. 3rd Ed., Clarendon Press, Oxford. (Oxford University Press, 1961).
53. Dienes, Z. How Bayes factors change scientific practice. *J. Math. Psychol.* <https://doi.org/10.1016/j.jmp.2015.10.003> (2016).
54. Kass, R. E. & Raftery, A. E. Bayes factors. *J. Am. Stat. Assoc.* <https://doi.org/10.1080/01621459.1995.10476572> (1995).
55. Wetzels, R. et al. Statistical evidence in experimental psychology: an empirical comparison using 855 t tests. *Perspect. Psychol. Sci.* **6**, 291–298 (2011).
56. Rouder, J. N., Morey, R. D., Speckman, P. L. & Province, J. M. Default Bayes factors for ANOVA designs. *J. Math. Psychol.* **56**, 356–374 (2012).
57. Goldstein, M. Subjective bayesian analysis: principles and practice. *Bayesian Anal.* **1**, 403–420 (2006).
58. Depaoli, S., Winter, S. D. & Visser, M. The importance of prior sensitivity analysis in bayesian statistics: demonstrations using an interactive shiny app. *Front. Psychol.* **11**, 3271 (2020).
59. Lambert, P. C., Sutton, A. J., Burton, P. R., Abrams, K. R. & Jones, D. R. How vague is vague? A simulation study of the impact of the use of vague prior distributions in MCMC using WinBUGS. *Stat. Med.* **24**, 2401–2428 (2005).
60. McElreath, R. Statistical rethinking: a Bayesian course with examples in R and Stan. <https://doi.org/10.1201/9781315372495> (2018).
61. Duval, S. & Tweedie, R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* **56**, 455–463 (2000).
62. Etnier, J. et al. The influence of physical fitness and exercise upon cognitive functioning: a meta-analysis. *J. Sport Exerc. Psychol.* **19**, 249–277 (1997).
63. McMorris, T. & Hale, B. J. Differential effects of differing intensities of acute exercise on speed and accuracy of cognition: a meta-analytical investigation. *Brain Cogn.* **80**, 338–351 (2012).
64. Neva, J. L. et al. Acute exercise modulates the excitability of specific interneurons in human motor cortex. *Neuroscience* **475**, 103–116 (2021).
65. Lulic, T., El-Sayes, J., Fassett, H. J. & Nelson, A. J. Physical activity levels determine exercise-induced changes in brain excitability. *PLoS One* **12**, e0173672 (2017).
66. Neva, J. L., Brown, K. E., Mang, C. S., Francisco, B. A. & Boyd, L. A. An acute bout of exercise modulates both intracortical and interhemispheric excitability. *Eur. J. Neurosci.* **45**, 1343–1355 (2017).
67. Singh, A. M., Duncan, R. E., Neva, J. L. & Staines, W. R. Aerobic exercise modulates intracortical inhibition and facilitation in a



- nonexercised upper limb muscle. *BMC Sports Sci. Med. Rehabil.* **6**, 23 (2014).
68. Yamazaki, Y. et al. Acute low-intensity aerobic exercise modulates intracortical inhibitory and excitatory circuits in an exercised and a non-exercised muscle in the primary motor cortex. *Front. Physiol.* **10**, 1361 (2019).
  69. Mooney, R. A. et al. Acute aerobic exercise modulates primary motor cortex inhibition. *Exp. Brain Res.* **234**, 3669–3676 (2016).
  70. Smith, A. E., Goldsworthy, M. R., Garside, T., Wood, F. M. & Ridding, M. C. The influence of a single bout of aerobic exercise on short-interval intracortical excitability. *Exp. Brain Res.* **232**, 1875–1882 (2014).
  71. McMorris, T., Tomporowski, P. D. & Audiffren, M. Exercise and cognitive function. *Wiley Blackwell*, 1–377 <https://doi.org/10.1002/9780470740668> (2009)
  72. McMorris, T. & Hale, B. J. Is there an acute exercise-induced physiological/biochemical threshold which triggers increased speed of cognitive functioning? A meta-analytic investigation. *J. Sport Health Sci.* **4**, 4–13 (2015).
  73. Davranche K., Giraud D., Hays A., Gajdos T. Exploring Decision-making Processes: The Impact of Acute High-Intensity Aerobic Exercise. bioRxiv; <https://doi.org/10.1101/2023.02.14.528466> (2023).
  74. Tomporowski, P. D. Effects of acute bouts of exercise on cognition. *Acta Psychol.* **112**, 297–324 (2003).
  75. Ayaz, A., Saleem, A. B., Schölvinck, M. L. & Carandini, M. Locomotion controls spatial integration in mouse visual cortex. *Curr. Biol.* **23**, 890–894 (2013).
  76. Fu, Y. et al. A cortical circuit for gain control by behavioral state. *Cell* **156**, 1139–1152 (2014).
  77. Kaneko, M., Fu, Y. & Stryker, M. P. Locomotion induces stimulus-specific response enhancement in adult visual cortex. *J. Neurosci.* **37**, 3532–3543 (2017).
  78. Keller, G. B., Bonhoeffer, T. & Hübener, M. Sensorimotor mismatch signals in primary visual cortex of the behaving mouse. *Neuron* <https://doi.org/10.1016/j.neuron.2012.03.040> (2012)
  79. Maimon, G., Straw, A. D. & Dickinson, M. H. Active flight increases the gain of visual motion processing in *Drosophila*. *Nat. Neurosci.* **13**, 393–399 (2010).
  80. Niell, C. M. & Stryker, M. P. Modulation of visual responses by behavioral state in mouse visual cortex. *Neuron* <https://doi.org/10.1016/j.neuron.2010.01.033> (2010)
  81. Polack, P.-O., Friedman, J. & Golshani, P. Cellular mechanisms of brain state-dependent gain modulation in visual cortex. *Nat. Neurosci.* **16**, 1331–1339 (2013).
  82. Bullock, T., Cecotti, H. & Giesbrecht, B. Multiple stages of information processing are modulated during acute bouts of exercise. *Neuroscience* **307**, 138–150 (2015).
  83. Bullock, T., Elliott, J. C., Serences, J. T. & Giesbrecht, B. Acute exercise modulates feature-selective responses in human cortex. *J. Cogn. Neurosci.* **29**, 605–618 (2017).
  84. Cao, L. & Händel, B. Walking enhances peripheral visual processing in humans. *PLOS Biol.* **17**, e3000511 (2019).
  85. Brouwer, G. J. & Heeger, D. J. Decoding and reconstructing color from responses in human visual cortex. *J. Neurosci.* **29**, 13992–14003 (2009).
  86. Bullock, T. et al. Eye movements disrupt EEG alpha-band coding of behaviorally relevant and irrelevant spatial locations held in working memory. *J. Neurophysiol.* **129**, 1191–1211 (2023).
  87. Foster, J. J., Sutterer, D. W., Serences, J. T., Vogel, E. K. & Awh, E. The topography of alpha-band activity tracks the content of spatial working memory. *J. Neurophysiol.* **115**, 168–177 (2016).
  88. MacLean, M. H., Bullock, T. & Giesbrecht, B. Dual process coding of recalled locations in human oscillatory brain activity. *J. Neurosci.* **39**, 6737–6750 (2019).
  89. Samaha, J., Sprague, T. C. & Postle, B. R. Decoding and reconstructing the focus of spatial attention from the topography of alpha-band oscillations. *J. Cogn. Neurosci.* **28**, 1090–1097 (2016).
  90. Sprague, T. C., Saproo, S. & Serences, J. T. Visual attention mitigates information loss in small- and large-scale neural codes. *Trends Cogn. Sci.* **19**, 215–226 (2015).
  91. Garrett, J., Bullock, T. & Giesbrecht, B. Tracking the contents of spatial working memory during an acute bout of aerobic exercise. *J. Cogn. Neurosci.* **33**, 1271–1286 (2021).
  92. Mordkoff, J. T. & Egeth, H. E. Response time and accuracy revisited: converging support for the interactive race model. *J. Exp. Psychol. Hum. Percept. Perform.* **19**, 981–991 (1993).
  93. Santee, J. L. & Egeth, H. E. Do reaction time and accuracy measure the same aspects of letter recognition? *J. Exp. Psychol. Hum. Percept. Perform.* **8**, 489–501 (1982).
  94. Basso, J. C., Shang, A., Elman, M., Karmouta, R. & Suzuki, W. A. Acute exercise improves prefrontal cortex but not hippocampal function in healthy adults. *J. Int. Neuropsychol. Soc.* **21**, 791–801 (2015).
  95. Chang, Y. K. et al. Effect of acute aerobic exercise on cognitive performance: role of cardiovascular fitness. *Psychol. Sport Exerc.* **15**, 464–470 (2014).
  96. Davranche, K., Hall, B. & McMorris, T. Effect of acute exercise on cognitive control required during an Eriksen Flanker task. *J. Sport Exerc. Psychol.* **31**, 628–639 (2009).
  97. Faulkner, J., Lambrick, D., Kaufmann, S. & Stoner, L. Effects of upright and recumbent cycling on executive function and prefrontal cortex oxygenation in young healthy men. *J. Phys. Act. health* **13**, 882–887 (2016).
  98. Kunzler, M. R. & Carpes, F. P. Intense cycling exercise improves acute cognitive responses. *Int. J. Sports Med.* **41**, 879–884 (2020).
  99. Yanagisawa, H. et al. Acute moderate exercise elicits increased dorsolateral prefrontal activation and improves cognitive performance with Stroop test. *NeuroImage* **50**, 1702–1710 (2010).
  100. Hung, T. M., Tsai, C. L., Chen, F. T., Wang, C. C. & Chang, Y. K. The immediate and sustained effects of acute exercise on planning aspect of executive function. *Psychol. Sport Exerc.* **14**, 728–736 (2013).
  101. Bae, S. & Masaki, H. Effects of acute aerobic exercise on cognitive flexibility required during task-switching paradigm. *Front. Hum. Neurosci.* **13**, 1–9 (2019).
  102. Pesce, C. & Audiffren, M. Does acute exercise switch off switch costs? A study with younger and older athletes. *J. Sport Exerc. Psychol.* **33**, 609–626 (2011).
  103. McMorris, T. Exercise and decision-making in team games. *Exercise and cognitive function* 179–192 <https://doi.org/10.1002/9780470740668.CH9> (2009)
  104. Kao, S. C., Drollette, E. S., Ritondale, J. P., Khan, N. & Hillman, C. H. The acute effects of high-intensity interval training and moderate-intensity continuous exercise on declarative memory and inhibitory control. *Psychol. Sport Exerc.* **38**, 90–99 (2018).
  105. Kujach, S. et al. Acute sprint interval exercise increases both cognitive functions and peripheral neurotrophic factors in humans: the possible involvement of lactate. *Front. Neurosci.* **13**, 1455 (2020).
  106. Dietrich, A. & Audiffren, M. The reticular-activating hypofrontality (RAH) model of acute exercise. *Neurosci. Biobehav. Rev.* **35**, 1305–1325 (2011).
  107. Mehren, A. et al. Intensity-dependent effects of acute exercise on executive function. *Neural Plast.* **2019**, 1–17 (2019).
  108. Kao, S.-C., Wang, C.-H., Kamijo, K., Khan, N. & Hillman, C. Acute effects of highly intense interval and moderate continuous exercise on the modulation of neural oscillation during working memory. *Int. J. Psychophysiol.* **160**, 10–17 (2021).

109. Rattray, B. & Smee, D. J. The effect of high and low exercise intensity periods on a simple memory recognition test. *J. Sport Health Sci.* **5**, 342 (2016).
110. Li, L. et al. Acute aerobic exercise increases cortical activity during working memory: a functional MRI study in female college students. *PLoS One* **9**, e99222 (2014).
111. Weng, T. B. et al. The acute effects of aerobic exercise on the functional connectivity of human brain networks. *Brain Plast.* **2**, 171–190 (2017).
112. Baniqued, P. L. et al. Brain network modularity predicts exercise-related executive function gains in older adults. *Front. Aging Neurosci.* **9**, 426 (2018).
113. Firth, J. et al. Effect of aerobic exercise on hippocampal volume in humans: a systematic review and meta-analysis. *NeuroImage* **166**, 230–238 (2018).
114. Voss, M. W., Nagamatsu, L. S., Liu-Ambrose, T. & Kramer, A. F. Exercise, brain, and cognition across the life span. *J. Appl. Physiol.* **111**, 1505–1513 (2011).
115. Voss, M. W. et al. Neurobiological markers of exercise-related brain plasticity in older adults. *Brain Behav. Immun.* **28**, 90–99 (2013).
116. Ludyga, S., Gerber, M., Pühse, U., Looser, V. N. & Kamijo, K. Systematic review and meta-analysis investigating moderators of long-term effects of exercise on cognition in healthy individuals. *Nat. Hum. Behav.* **4**, 603–612 (2020).
117. Ferguson, H. J., Brunson, V. E. A. & Bradford, E. E. F. The developmental trajectories of executive function from adolescence to old age. *Sci. Rep.* **11**, 1382 (2021).
118. Hartshorne, J. K. & Germine, L. T. When does cognitive functioning peak? The asynchronous rise and fall of different cognitive abilities across the lifespan. *Psychol. Sci.* **26**, 433–443 (2015).
119. Bullock, T., Giesbrecht, B., Beaudin, A. E., Goodyear, B. G. & Poulin, M. J. Effects of changes in end-tidal PO<sub>2</sub> and PCO<sub>2</sub> on neural responses during rest and sustained attention. *Physiol. Rep.* **9**, e15106 (2021).
120. Shields, G. S., Sazma, M. A. & Yonelinas, A. P. The effects of acute stress on core executive functions: a meta-analysis and comparison with cortisol. *Neurosci. Biobehav. Rev.* **68**, 651–668 (2016).
121. Gordon, J. et al. The road towards understanding embodied decisions. *Neurosci. Biobehav. Rev.* **131**, 722–736 (2021).
122. Raichlen, D. A. & Alexander, G. E. Adaptive capacity: an evolutionary-neuroscience model linking exercise, cognition, and brain health. *Trends Neurosci.* **40**, 408–421 (2017).
123. Raudenbush, S. W. & Bryk, A. S. Hierarchical linear models: applications and data analysis methods, 485 (2002).
124. Hogervorst, E., Riedel, W., Jeukendrup, A. & Jolles, J. Cognitive performance after strenuous physical exercise. *Percept. Mot. Skills* **83**, 479–488 (1996).
125. Pesce, C., Capranica, L., Tessitore, A. & Figura, F. Focusing of visual attention under submaximal physical load. *Int. J. Sport Exerc. Psychol.* **1**, 275–292 (2003).
126. Pesce, C., Tessitore, A., Casella, R., Pirritano, M. & Capranica, L. Focusing of visual attention at rest and during physical exercise in soccer players. *J. Sports Sci.* **25**, 1259–1270 (2007).
127. Vickers, J. N. & Williams, A. M. Performing under pressure: the effects of physiological arousal, cognitive anxiety, and gaze control in biathlon. *J. Mot. Behav.* **39**, 381–394 (2007).
128. Lo Bue-Estes, C. et al. Short-term exercise to exhaustion and its effects on cognitive function in young women. *Percept. Mot. Skills* **107**, 933–945 (2008).
129. Coles, K. & Tomporowski, P. D. Effects of acute exercise on executive processing, short-term and long-term memory. *J. Sports Sci.* **26**, 333–344 (2008).
130. Fontana, F. E., Mazzardo, O., Mokgothu, C., Furtado, O. & Gallagher, J. D. Influence of exercise intensity on the decision-making performance of experienced and inexperienced soccer players. *J. Sport Exerc. Psychol.* **31**, 135–151 (2009).
131. Luft, C. D. B., Takase, E. & Darby, D. Heart rate variability and cognitive function: effects of physical effort. *Biol. Psychol.* **82**, 186–191 (2009).
132. Pontifex, M. B., Hillman, C. H., Fernhall, B., Thompson, K. M. & Valentini, T. A. The effect of acute aerobic and resistance exercise on working memory. *Med. Sci. Sports Exerc.* **41**, 927–934 (2009).
133. Srygley, J. M., Mirelman, A., Herman, T., Giladi, N. & Hausdorff, J. M. When does walking alter thinking? Age and task associated findings. *Brain Res.* **1253**, 92–99 (2009).
134. Thomson, K., Watt, A. P. & Liukkonen, J. Differences in ball sports athletes speed discrimination skills before and after exercise induced fatigue. *J. Sports Sci. Med.* **8**, 259–264 (2009).
135. Norling, J. C., Sibthorp, J., Suchy, Y., Hannon, J. C. & Ruddell, E. The benefit of recreational physical activity to restore attentional fatigue: the effects of running intensity level on attention scores. *J. Leis. Res.* **42**, 135–152 (2010).
136. Young, W. et al. Acute effect of exercise on kicking accuracy in elite Australian football players. *J. Sci. Med. Sport* **13**, 85–89 (2010).
137. Chang, Y.-K. et al. Effects of acute exercise on executive function: a study with a tower of London task. *J. Sport Exerc. Psychol.* **33**, 847–865 (2011).
138. Green, A. L. & Helton, W. S. Dual-task performance during a climbing traverse. *Exp. Brain Res.* **215**, 307–313 (2011).
139. Ohlinger, C. M., Horn, T. S., Berg, W. P. & Cox, R. H. The effect of active workstation use on measures of cognition, attention, and motor skill. *J. Phys. Act. Health* **8**, 119–125 (2011).
140. Hope, L., Lewinski, W., Dixon, J., Blocksidge, D. & Gabbert, F. Witnesses in action: the effect of physical exertion on recall and recognition. *Psychol. Sci.* **23**, 386–390 (2012).
141. Lambourne, K. The effects of acute exercise on temporal generalization. *Q J. Exp. Psychol.* **65**, 526–540 (2012).
142. Moore, R. D., Romine, M. W., O’connor, P. J. & Tomporowski, P. D. The influence of exercise-induced fatigue on cognitive function. *J. Sports Sci.* **30**, 841–850 (2012).
143. Roberts, A. P. J. & Cole, J. C. The effects of exercise and body armor on cognitive function in healthy volunteers. *Mil. Med.* **178**, 479–486 (2013).
144. Bullock, T. & Giesbrecht, B. Acute exercise and aerobic fitness influence selective attention during visual search. *Front. Psychol.* **5**, 1290 (2014).
145. Byun, K. et al. Positive effect of acute mild exercise on executive function via arousal-related prefrontal activations: an fNIRS study. *NeuroImage* **98**, 336–345 (2014).
146. Darling, K. A. & Helton, W. S. Dual-task interference between climbing and a simulated communication task. *Exp. Brain Res.* **232**, 1367–1377 (2014).
147. Nibbeling, N., Oudejans, R. R. D., Ubink, E. M. & Daanen, H. A. M. The effects of anxiety and exercise-induced fatigue on shooting accuracy and cognitive performance in infantry soldiers. *Ergonomics* **57**, 1366–1379 (2014).
148. Pontifex, M. B., Parks, A. C., Henning, D. A. & Kamijo, K. Single bouts of exercise selectively sustain attentional processes. *Psychophysiology* **52**, 618–625 (2015).
149. Schmidt-Kassow, M. et al. Treadmill walking during vocabulary encoding improves verbal long-term memory. *Behav. Brain Funct.* **10**, 24 (2014).
150. Tsai, C. L. et al. Executive function and endocrinological responses to acute resistance exercise. *Front. Behav. Neurosci.* **8**, 1–12 (2014).
151. Bantoft, C. et al. Effect of standing or walking at a workstation on cognitive function: a randomized counterbalanced trial. *Hum. Factors* **58**, 140–149 (2016).
152. Larson, M. J. et al. Slow walking on a treadmill desk does not negatively affect executive abilities: an examination of cognitive



- control, conflict adaptation, response inhibition, and post-error slowing. *Front. Psychol.* **6**, 723 (2015).
153. Osgood, J. M. Acute cardiovascular exercise counteracts the effect of ego-depletion on attention: how ego-depletion increases boredom and compromises directed attention. *Int. J. Psychol. Stud.* **7**, 85–96 (2015).
  154. Perciavalle, V., Maci, T., Perciavalle, V., Massimino, S. & Coco, M. Working memory and blood lactate levels. *Neurol. Sci.* **36**, 2129–2136 (2015).
  155. Shia, R. M. et al. Individual differences in biophysiological toughness: sustaining working memory during physical exhaustion. *Mil. Med.* **180**, 230–236 (2015).
  156. Stevens, D. J., Arciuli, J. & Anderson, D. I. Concurrent movement impairs incidental but not intentional statistical learning. *Cogn. Sci.* **39**, 1081–1098 (2015).
  157. Weng, T. B., Pierce, G. L., Darling, W. G. & Voss, M. W. Differential effects of acute exercise on distinct aspects of executive function. *Med. Sci. Sports Exerc.* **47**, 1460–1469 (2015).
  158. Alloway, R. G., Alloway, T. P., Magyari, P. M. & Floyd, S. An exploratory study investigating the effects of barefoot running on working memory. *Percept. Mot. Skills* **122**, 432–443 (2016).
  159. Ando, S. et al. Effects of strenuous exercise on visual perception are independent of visual resolution. *Physiol. Behav.* **106**, 117–121 (2012).
  160. Brush, C. J. et al. Dose–response and time course effects of acute resistance exercise on executive function. *J. Sport Exerc. Psychol.* **38**, 396–408 (2016).
  161. Connell, C. J. W., Thompson, B., Kuhn, G. & Gant, N. Exercise-induced fatigue and caffeine supplementation affect psychomotor performance but not covert visuo-spatial attention. *PLoS One* **11**, e0165318 (2016).
  162. Hsieh, S.-S., Chang, Y.-K., Hung, T.-M. & Fang, C.-L. The effects of acute resistance exercise on young and older males' working memory. *Psychol. Sport Exerc.* **22**, 286–293 (2016).
  163. Hsieh, S.-S. Acute resistance exercise facilitates attention control in adult males without an age-moderating effect. *J. Sport Exerc. Psychol.* **38**, 247–254 (2016).
  164. Lowe, C. J., Kolev, D. & Hall, P. A. An exploration of exercise-induced cognitive enhancement and transfer effects to dietary self-control. *Brain Cogn.* **110**, 102–111 (2016).
  165. Torbeyns, T. et al. Cycling on a bike desk positively influences cognitive performance. *PLoS One* <https://doi.org/10.1371/journal.pone.0165510> (2016).
  166. Tsukamoto, H. et al. Greater impact of acute high-intensity interval exercise on post-exercise executive function compared to moderate-intensity continuous exercise. *Physiol. Behav.* **155**, 224–230 (2016).
  167. Zach, S. & Shalom, E. The influence of acute physical activity on working memory. *Percept. Mot. Skills* **122**, 365–374 (2016).
  168. Chang, Y.-K. et al. Acute exercise has a general facilitative effect on cognitive function: a combined ERP temporal dynamics and BDNF study. *Psychophysiology* **54**, 289–300 (2017).
  169. Crush, E. A. & Loprinzi, P. D. Dose-response effects of exercise duration and recovery on cognitive functioning. *Percept. Mot. Skills* **124**, 1164–1193 (2017).
  170. González Fernández, F. T., Etnier, J., Zabala, M. & Sanabria, D. Vigilance performance during acute exercise. *Int. J. Sport Psychol.* **48**, (2017).
  171. Lindheimer, J. B., O'Connor, P. J., McCully, K. K. & Dishman, R. K. The effect of light-intensity cycling on mood and working memory in response to a randomized, placebo-controlled design. *Psychosom. Med.* **79**, 243–253 (2017).
  172. Lowe, C. J., Staines, W. R. & Hall, P. A. Effects of moderate exercise on cortical resilience: a transcranial magnetic stimulation study targeting the dorsolateral prefrontal cortex. *Psychosom. Med.* **79**, 143–152 (2017).
  173. Luu, K. & Hall, P. A. Examining the acute effects of hatha yoga and mindfulness meditation on executive function and mood. *Mindfulness* **8**, 873–880 (2017).
  174. Randolph, D. D. & O'Connor, P. J. Stair walking is more energizing than low dose caffeine in sleep deprived young women. *Physiol. Behav.* **174**, 128–135 (2017).
  175. Slutsky, A. B. et al. The effects of low-intensity cycling on cognitive performance following sleep deprivation. *Physiol. Behav.* **180**, 25–30 (2017).
  176. Sudo, M. et al. Executive function after exhaustive exercise. *Eur. J. Appl Physiol.* **117**, 2029–2038 (2017).
  177. Cuttler, C., Connolly, C. P., LaFrance, E. M. & Lowry, T. M. Resist forgetting: effects of aerobic and resistance exercise on prospective and retrospective memory. *Sport Exerc. Perform. Psychol.* **7**, 205 (2017).
  178. Daikoku, T., Takahashi, Y., Tarumoto, N. & Yasuda, H. Auditory statistical learning during concurrent physical exercise and the tolerance for pitch, tempo, and rhythm changes. *Mot. Control* **22**, 233–244 (2018).
  179. Elkana, O. et al. Conscientiousness is associated with improvement in visuospatial working memory and mood following acute physical exercise: a randomized controlled trial. *Pers. Individ. Differ.* **132**, 126–132 (2018).
  180. Fenesi, B., Lucibello, K., Kim, J. A. & Heisz, J. J. Sweat so you don't forget: exercise breaks during a university lecture increase on-task attention and learning. *J. Appl. Res. Mem. Cogn.* **7**, 261–269 (2018).
  181. Kendall, B. The effects of acute exercise on postural control, information processing, motor skill acquisition, and executive function. *Wayne State University Dissertations* (2018).
  182. Legrand, F. D. et al. Brief aerobic exercise immediately enhances visual attentional control and perceptual speed. Testing the mediating role of feelings of energy. *Acta Psychol. (Amst.)* **191**, 25–31 (2018).
  183. Samani, A. & Heath, M. Executive-related oculomotor control is improved following a 10-min single-bout of aerobic exercise: evidence from the antisaccade task. *Neuropsychologia* **108**, 73–81 (2018).
  184. Siddiqui, A. & Loprinzi, P. D. Experimental investigation of the time course effects of acute exercise on false episodic memory. *J. Clin. Med.* **7**, 157 (2018).
  185. Sng, E., Frith, E. & Loprinzi, P. D. Temporal effects of acute walking exercise on learning and memory function. *Am. J. Health Promot.* **32**, 1518–1525 (2018).
  186. Wade, B. & Loprinzi, P. D. The experimental effects of acute exercise on long-term emotional memory. *J. Clin. Med.* **7**, 486 (2018).
  187. Yamazaki, Y. et al. Inter-individual differences in working memory improvement after acute mild and moderate aerobic exercise. *PLoS One* **13**, e0210053 (2018).
  188. Baker, R., Coenen, P., Howie, E., Williamson, A. & Straker, L. The musculoskeletal and cognitive effects of under-desk cycling compared to sitting for office workers. *Appl. Ergon.* **79**, 76–85 (2019).
  189. Du Rietz, E. et al. Beneficial effects of acute high-intensity exercise on electrophysiological indices of attention processes in young adult men. *Behav. Brain Res.* **359**, 474–484 (2019).
  190. Engeroff, T., Niederer, D., Vogt, L. & Banzer, W. Intensity and workload related dose–response effects of acute resistance exercise on domain-specific cognitive function and affective response – a four-armed randomized controlled crossover trial. *Psychol. Sport Exerc.* **43**, 55–63 (2019).

191. Haynes, J. T., Frith, E., Sng, E. & Loprinzi, P. D. Experimental effects of acute exercise on episodic memory function: considerations for the timing of exercise. *Psychol. Rep.* **122**, 1744–1754 (2019).
192. Johnson, L. & Loprinzi, P. D. The effects of acute exercise on episodic memory function among young university students: moderation considerations by biological sex. *Health Promot. Perspect.* **9**, 99–104 (2019).
193. McGowan, A. L., Chandler, M. C. & Brascamp, J. W. & Pontifex, M. B. Pupillometric indices of locus-coeruleus activation are not modulated following single bouts of exercise. *Int. J. Psychophysiol.* **140**, 41–52 (2019).
194. Schmidt, S. L. et al. Decrease in attentional performance after repeated bouts of high intensity exercise in association-football referees and assistant referees. *Front. Psychol.* **10**, 2014 (2019).
195. Stenling, A., Moylan, A., Fulton, E. & Machado, L. Effects of a brief stair-climbing intervention on cognitive performance and mood states in healthy young adults. *Front. Psychol.* **10**, 2300 (2019).
196. Wu, C.-H. et al. Effects of acute aerobic and resistance exercise on executive function: an ERP study. *J. Sci. Med. Sport* **22**, 1367–1372 (2019).
197. Zhou, F. & Qin, C. Acute moderate-intensity exercise generally enhances attentional resources related to perceptual processing. *Front. Psychol.* **10**, 2547 (2019).
198. Aly, M. & Kojima, H. Acute moderate-intensity exercise generally enhances neural resources related to perceptual and cognitive processes: a randomized controlled ERP study. *Ment. Health Phys. Act.* **19**, 100363 (2020).
199. Chacko, S. C. et al. A single bout of vigorous-intensity aerobic exercise affects reactive, but not proactive cognitive brain functions. *Int. J. Psychophysiol.* **147**, 233–243 (2020).
200. Morris, T. P. et al. Light aerobic exercise modulates executive function and cortical excitability. *Eur. J. Neurosci.* **51**, 1723–1734 (2020).
201. Walsh, J. J. et al. Acute aerobic exercise impairs aspects of cognitive function at high altitude. *Physiol. Behav.* **223**, 112979 (2020).
202. Kim, H. J., Bae, S., Huh, J. H., Lee, J. W. & Han, D. H. Hemodynamic changes in response to aerobic exercise: near-infrared spectroscopy study. *Int. J. Sports Med.* **42**, 377–385 (2021).
203. Klatt, S. & Smeeton, N. J. Attentional and perceptual capabilities are affected by high physical load in a simulated soccer decision-making task. *Sport Exerc. Perform. Psychol.* **10**, 205–216 (2021).
204. Kuhne, L. A. et al. The effects of acute cardiovascular exercise on memory and its associations with exercise-induced increases in neurotrophic factors. *Front. Aging Neurosci.* **13**, 750401 (2021).
205. Manocchio, F. & Lowe, C. J. Investigating cortical buffering effects of acute moderate intensity exercise: a cTBS study targeting the left dorsolateral prefrontal cortex. *Front. Hum. Neurosci.* **15**, 645326 (2021).
206. Miyashiro, S. et al. Activation of the orbitofrontal cortex by both meditation and exercise: a near-infrared spectroscopy study. *PLoS One* **16**, e0247685 (2021).
207. Trammell, J. P. & Aguilar, S. C. Natural is not always better: the varied effects of a natural environment and exercise on affect and cognition. *Front. Psychol.* **11**, 575245 (2020).
208. Zhu, Y., Sun, F., Chiu, M. M. & Siu, A. Y.-S. Effects of high-intensity interval exercise and moderate-intensity continuous exercise on executive function of healthy young males. *Physiol. Behav.* **239**, 113505 (2021).
209. Aguirre-Loaiza, H. et al. Effect of acute physical exercise on inhibitory control in young adults: high-intensity indoor cycling session. *Physiol. Behav.* **254**, 113902 (2022).
210. Drollette, E. S. & Meadows, C. C. The effects of acute high-intensity interval exercise on the temporal dynamics of working memory and contralateral delay activity. *Psychophysiology* **59**, e14112 (2022).
211. Engeroff, T., Banzer, W. & Niederer, D. The impact of regular activity and exercise intensity on the acute effects of resistance exercise on cognitive function. *Scand. J. Med. Sci. Sports* **32**, 94–105 (2022).
212. Frith, E., Miller, S. E. & Loprinzi, P. D. Effects of verbal priming with acute exercise on convergent creativity. *Psychol. Rep.* **125**, 375–397 (2022).
213. Kao, S.-C. et al. Acute effects of aerobic exercise on conflict suppression, response inhibition, and processing efficiency underlying inhibitory control processes: an ERP and SFT study. *Psychophysiology* **59**, e14032 (2022).
214. LaCount, P. A. et al. Acute effects of physical exercise on cognitive and psychological functioning in college students with attention-deficit/hyperactivity disorder. *Ment. Health Phys. Act.* **22**, 100443 (2022).
215. Loprinzi, P. D. & Storm, B. C. Acute exercise on memory: application of the retrieval-induced forgetting paradigm. *Psychol. Res.* **87**, 1417–1428 (2023).
216. Shirzad, M. et al. Passive exercise increases cerebral blood flow velocity and supports a postexercise executive function benefit. *Psychophysiology* **59**, e14132 (2022).
217. Zheng, K. et al. Changes in working memory performance and cortical activity during acute aerobic exercise in young adults. *Front. Behav. Neurosci.* **16**, 884490 (2022).
218. Chueh, T.-Y., Hung, C.-L., Chang, Y.-K., Huang, C.-J. & Hung, T.-M. Effects of cognitive demand during acute exercise on inhibitory control and its electrophysiological indices: a randomized crossover study. *Physiol. Behav.* **265**, 114148 (2023).
219. Yamada, Y. et al. Improved interference control after exercise with blood flow restriction and cooling is associated with but not mediated by increased lactate. *Physiol. Behav.* **270**, 114291 (2023).
220. Zhang, B., Meng, X., Yu, Y., Han, Y. & Liu, Y. Effects of moderate-to-vigorous acute exercise on conscious perception and visual awareness. *J. Mot. Behav.* **55**, 262–268 (2023).
221. Zhang, D., Jin, X., Wang, L. & Jin, Y. Form and color visual perception in green exercise: positive effects on attention, mood, and self-esteem. *J. Environ. Psychol.* **88**, 102028 (2023).

## Acknowledgements

This work was funded by the U.S. Army Combat Capabilities Development Command Soldier Center Measuring and Advancing Soldier Tactical Readiness and Effectiveness (MASTR-E) program through award W911NF-19F-0018 under US Army Research Office contract W911NF-19-D-0001 for the Institute for Collaborative Biotechnologies. The views expressed in this article are solely those of the authors and do not reflect the official policies or positions of the Department of Army, the Department of Defense, or any other department or agency of the U.S. government. The funders had no role in study design, data collection, and analysis, the decision to publish, or preparation of the manuscript. Thank you to Grace Giles, Ph.D., Julie Cantelon, Ph.D., and Neil Dundon, Ph.D., for providing guidance in conducting analyses and interpreting results. Lastly, thank you to Emily Machniak and Riddhima Chandra for assisting in data collection efforts.

## Author contributions

J.G., C.C., T.B., and B.G., designed the study, created the literature search criterion, and performed initial article screenings. J.G., C.C., and B.G. performed screenings of articles based on their full text. J.G., extracted effect sizes, conducted statistical analyses, and wrote the initial draft of the manuscript. B.G., T.B., and C.C. provided edits to the manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s44271-024-00124-2>.

**Correspondence** and requests for materials should be addressed to Jordan Garrett or Barry Giesbrecht.

**Peer review information** *Communications Psychology* thanks Sebastian Ludyga and Nicole Logan for their contribution to the peer review of this work. Primary Handling Editors: Daniel Quintana, Antonia Eisenkoeck and Jennifer Bellingier. A peer review file is available.

**Reprints and permissions information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024