



## Consideration on immunotherapy of liver metastases of malignant tumors

Chuang Jiang, Zhi-Hong Zhang, Jia-Xin Li

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's classification**

**Scientific Quality:** Grade C

**Novelty:** Grade C

**Creativity or Innovation:** Grade C

**Scientific Significance:** Grade C

**P-Reviewer:** Gupta D, India

**Received:** March 10, 2024

**Revised:** May 5, 2024

**Accepted:** June 3, 2024

**Published online:** August 27, 2024

**Processing time:** 159 Days and 0.5 Hours



**Chuang Jiang, Zhi-Hong Zhang, Jia-Xin Li**, Division of Liver Surgery, Department of General Surgery and Laboratory of Liver Surgery, and State Key Laboratory of Biotherapy and Collaborative Innovation Center of Biotherapy, West China Hospital, Sichuan University, Chengdu 610041, Sichuan Province, China

**Jia-Xin Li**, Department of General Surgery, Dafang County People's Hospital, Bijie 551600, Guizhou Province, China

**Corresponding author:** Jia-Xin Li, MD, Assistant Professor, Chief Doctor, Division of Liver Surgery, Department of General Surgery and Laboratory of Liver Surgery, and State Key Laboratory of Biotherapy and Collaborative Innovation Center of Biotherapy, West China Hospital, Sichuan University, No. 37 Guo Xue Alley, Chengdu 610041, Sichuan Province, China. [lijiaxin@scu.edu.cn](mailto:lijiaxin@scu.edu.cn)

### Abstract

In this editorial, we comment on the article "Analysis of the impact of immunotherapy efficacy and safety in patients with gastric cancer and liver metastasis" by Liu *et al* that was published in the recent issue of the *World Journal of Gastrointestinal Surgery*. It has prompted us to think and summarize some thoughts on immunotherapy for malignant tumor liver metastasis. Immunotherapy plays a crucial role in the treatment of malignant tumors; however, the presence of liver metastases in advanced tumors may impact its efficacy. Although patients with liver metastases can still benefit from immunotherapy, multiple clinical studies have indicated that, compared to other sites of metastasis, liver metastases may diminish the effectiveness of immunotherapy. The efficacy of immune checkpoint inhibitors in patients with liver metastases often fails to reach the ideal level, primarily due to the liver metastases exploiting the host's peripheral immune tolerance mechanisms to promote systemic CD8(+) T cell exhaustion, resulting in a systemic immune-tolerant environment. This article aims to summarize the reasons for the decreased efficacy of immunotherapy following liver metastasis in various malignant tumors and propose potential clinical strategies for management.

**Key Words:** Liver metastasis; Immunotherapy; Immune tolerance; Cancer; Treatment

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Immunotherapy is an important treatment method for solid tumors nowadays. However, liver metastasis leads to a decrease in the efficacy of immunotherapy for malignant tumors. The main reason is that liver metastasis can promote the depletion of systemic CD8(+) T cells by utilizing the host's peripheral immune tolerance mechanism, thereby leading to a systemic immune tolerance environment. In the dilemma of limited efficacy of immunotherapy alone, the combination with chemotherapy, radiotherapy, targeted therapy, and other anti-tumor methods can significantly enhance the efficacy of immunotherapy, holding important clinical prospects.

**Citation:** Jiang C, Zhang ZH, Li JX. Consideration on immunotherapy of liver metastases of malignant tumors. *World J Gastrointest Surg* 2024; 16(8): 2374-2381

**URL:** <https://www.wjgnet.com/1948-9366/full/v16/i8/2374.htm>

**DOI:** <https://dx.doi.org/10.4240/wjgs.v16.i8.2374>

## INTRODUCTION

### Liver metastasis

Metastases often occur in the advanced stage of malignant tumors. Although different tumors often have different tendencies towards the metastatic organs, the liver is the organ most susceptible to metastasis due to its unique anatomical position, cell metabolism, and immune microenvironment[1]. Liver metastasis substantially impacts patient survival. Studies have shown that patients with liver metastasis have a poorer prognosis compared to those with other organ metastases[2]. There are different pathological types of liver metastasis, with the most common pathologic type reported being not otherwise specified (NOS) adenocarcinoma (74.9%). Most NOS adenocarcinomas originate from the digestive tract (68.2%), especially from the colorectum (46.1%)[3]. There are three different histopathological growth patterns (HGPs) of metastatic cancer cells, including: (1) Desmoplastic type; (2) Replacement type; and (3) Pushing type [4]. Different HGP subtypes have different cytokine profiles and varying levels of lymphocyte infiltration[5]. The HGP of liver metastasis in different tumors is also different. For example, the most common manifestation of liver metastasis in uveal melanoma and breast cancer is replacement type[6,7]; in liver metastases of digestive tract tumors such as gastrointestinal and pancreatic malignancies, the occurrence frequency of the three HGP subtypes is approximately equal [8]. Multiple studies have shown that the prognosis of different HGP subtypes of liver metastasis varies, with patients with desmoplastic HGP having the best prognosis[7,9]. However, some studies have found that patients with non-desmoplastic HGP are more likely to benefit from adjuvant chemotherapy[10]. If HGP can be included in the treatment decisions for liver metastasis patients, it may benefit patients and thus improve their prognosis.

### Liver metastasis and immune tolerance

Despite distant metastasis indicating the advanced stage of malignant tumors, in recent years, immunotherapy represented by immune checkpoint inhibitors (ICIs) has become a major breakthrough in the field of malignant tumor treatment, changing the treatment decisions for various solid tumors[11]. For the overall population, immunotherapy provides survival benefits for patients with liver metastasis, but it has been observed in various malignant tumors that the benefits of immunotherapy are significantly reduced compared to patients without liver metastasis. For example, Botticelli *et al*[12] found that the metastasis of urothelial carcinoma to lymph nodes was completely relieved by ICIs treatment, while the metastasis to the liver showed resistance to such treatment. Takeyasu *et al*'s[13] study on non-small cell lung cancer (NSCLC) found that liver metastasis was associated with poorer median progression-free survival (PFS) after treatment with pembrolizumab (3.4 mo *vs* 9.4 mo,  $P = 0.0018$ ). Similar findings were reported by Tumei *et al*[14], who found that pembrolizumab immunotherapy for melanoma also showed poorer immunotherapy outcomes in patients with liver metastases. A multicenter clinical study on stage IV melanoma showed that the liver metastasis group had poorer response to anti PD-1 monoclonal antibody therapy compared to the non-liver metastasis group, with a significantly lower objective response rate (ORR; 4.3% *vs* 20.7%,  $P < 0.05$ )[15]. In the recent issue of the *World Journal of Gastroenterology Surgery*, Liu *et al*[16] published an interesting paper revealing that in patients with advanced gastric cancer who received immunotherapy, PFS was significantly worse in the liver metastasis group compared to the non-liver metastasis group (5.0 mo *vs* 11.2 mo,  $P < 0.05$ ). The reason for this clinical phenomenon is currently believed to be liver metastasis leading to immune tolerance.

Liver immune tolerance is a recognized concept[17,18], and it is currently believed that the liver is a key frontline immune organ, with its default immune state mainly being anti-inflammatory or immune tolerance. Under appropriate conditions, the liver is capable of mounting rapid and robust immune responses as well. The balance between immunity and tolerance is crucial for liver function[19]. Peripheral T cells are key participants in adaptive immune responses, and multiple mechanisms lead to depletion or inhibition of peripheral T cells in the liver, resulting in immune tolerance. Preclinical studies have found that metastatic liver cancer can induce systemic tumor-specific CD8(+) T cell loss through the Fas-FasL pathway, leading to reduced efficacy of immunotherapy[20]. Comprehensive sequencing of primary tumors (including melanoma, NSCLC, *etc.*) from patients with metastatic cancer showed a decrease in T cell clonality and diversity, and a decrease in T cell effector capacity in patients with liver metastasis[21]. Tumei *et al*'s[14] study on biopsy specimens of melanoma liver metastasis found that compared to the non-liver metastasis group, the CD8(+) T cell count at the infiltrating edge of the liver metastasis group was significantly reduced (547 *vs* 1441,  $P < 0.05$ ). Analysis of the distri-

bution of primary tumor and liver metastatic tumor cells in patients with liver metastasis of colorectal cancer by He *et al* [22], revealed that there were more immunosuppressive cells in liver metastases, among which the distribution difference of macrophages was the most obvious. This may be related to intrahepatic immunosuppression and poor immunotherapy efficacy. The lymph nodes, skin, and lungs are areas where anti-tumor immunity is easily beneficial, which may be attributed to the presence of a large number of immune cells in these specific organs[23].

## COMBINATION THERAPY OVERCOMES IMMUNE TOLERANCE

Reversing the immunosuppressive state and improving the efficacy of immunotherapy in patients with liver metastasis are the key to treating liver metastasis. At present, the main approach is to combine another treatment method to improve the efficacy of immunotherapy. However, clinical research on the immunotherapy-based combination therapy for liver metastasis of malignant tumors is limited, with more focus on preclinical studies or case reports (Table 1).

### **Immunotherapy combined with chemotherapy**

The effectiveness of immunotherapy combined with chemotherapy has been demonstrated by multiple studies targeting various malignant tumors. A subgroup analysis of 115 NSCLC liver metastasis patients in the KEYNOTE-189 study found that the combination of pembrolizumab and chemotherapy significantly prolonged the median overall survival (OS) and PFS of patients compared to the chemotherapy combined with placebo group[24]. A meta-analysis that included eight randomized controlled clinical studies showed that PD-1/PD-L1 inhibitors + chemotherapy can reduce the risk of tumor progression (hazard ratio [HR] = 0.60, 95% confidence interval [CI]: 0.55-0.65) and mortality (HR = 0.71, 95% CI: 0.58-0.90) [25]. A study by Inoue *et al*[26] on liver metastasis in colorectal cancer found that the infiltration of CD8(+), CD3(+), and CD56(+) cells in the cetuximab + chemotherapy group was higher than that in the chemotherapy group and no chemotherapy group ( $P < 0.05$  for all). The efficacy of combination immunotherapy regimen for liver metastasis of many malignant tumors is worth further investigation. Some phase III clinical studies[27,28] have shown that chemotherapy combined with immunotherapy can improve treatment efficacy and prognosis in advanced gastric cancer patients, but clinical studies on immunotherapy for liver metastasis of gastric cancer are extremely limited. Preclinical studies have also yielded some clinically instructive results. Ho *et al*[29] observed that gemcitabine + PD-1 antibody significantly prolonged the median survival time (66 d *vs* 56 d) of mice with liver metastasis of pancreatic cancer compared with gemcitabine alone. They also found that the combination therapy can enhance the anti-cancer effect and Th1 Lymphocytes response of M1 macrophages.

### **Immunotherapy combined with radiotherapy**

Studies have shown that liver targeted radiation therapy can reshape the liver immune microenvironment, reduce T cell consumption, and restore the effectiveness of immunotherapy[20]. Similar studies on NSCLC have shown that local liver radiation therapy can lead to a higher proportion of CD8(+) T cells and CD4(+) T cells in tumor tissue, and the surface of tumor cells will also express higher levels of PD-1/PD-L1. Therefore, radiotherapy combined with immunization may also be an effective treatment for NSCLC patients with liver metastases[30]. In recent years, the effectiveness of stereotactic body radiation therapy (SBRT) has been confirmed for lung cancer liver metastasis. The addition of SBRT in NSCLC liver metastasis patients can promote the anti-angiogenic effect, enhance the immunogenicity of tumors in chemotherapy and immunotherapy, and thus achieve a synergistic effect with systemic therapy[31]. Xu *et al*[32] reported a case of gastric cancer with liver metastasis, which progressed after chemotherapy and targeted therapy, but, after changing the treatment strategy to PD-1 antibody combined with SBRT, partial response was achieved.

### **Immunotherapy combined with targeted therapy**

In a mouse model of colon cancer liver metastasis, it was found that the anti-tumor effect of blocking PD-1 checkpoint alone was not significant, while immunotherapy combined with VEGF inhibitors significantly improved the therapeutic effect. This indicates that VEGF inhibitors may sensitize the antitumor effects of immunotherapy[33]. At present, the combination of angiogenesis inhibitors and anti-PD-1/PD-L1 has become the first-line treatment for patients with metastatic renal cell carcinoma. In the IMpower150 study[34] for metastatic NSCLC, it was found that for patients with liver metastasis, the atezolizumab + bevacizumab + carboplatin + paclitaxel group (ABCP group) had longer OS and PFS compared to the bevacizumab + carboplatin + paclitaxel group (BCP group). Immunotherapy combined with antivascular therapy is of great significance for patients with liver metastasis.

### **Dual immunotherapy**

Considering the immune tolerance of the liver, the combination therapy of dual drug inhibition at immune checkpoints is expected to have a synergistic effect on liver metastasis patients and achieve better therapeutic effects. The CheckMate 277 study[35] evaluated the efficacy in the nivolumab combined with ipilimumab group and chemotherapy group of advanced or metastatic NSCLC patients. The results showed that the dual immunotherapy significantly improved patient efficacy compared to the chemotherapy group (2-year ORR: 40.0% *vs* 32.8%; median OS: 17.1 mo *vs* 13.9 mo,  $P = 0.007$ ). Although there have been no reports of dual immunotherapy for melanoma liver metastases, CheckMate067 study[36] found that the combined treatment of nivolumab and ipilimumab compared with the monotherapy could improve the OS of melanoma patients, which also suggests that the potential value of dual immunotherapy is worth further exploration.

Table 1 Immunotherapy-based combination therapy for malignant liver metastasis

Cancer	Ref.	Type	Combination	Treatment strategy
Lung cancer	Ma <i>et al</i> [45]	Clinical study	Immunotherapy + targeted therapy	Anti-PD-1 + anlotinib
	Jiang <i>et al</i> [46]	Preclinical study	Immunotherapy + targeted therapy + radiotherapy	Anti-PD-1 + anlotinib + radiotherapy
	Gadgeel <i>et al</i> [24]	Clinical study	Immunotherapy + chemotherapy	Pembrolizumab + pemetrexed
	Socinski <i>et al</i> [34]	Clinical study	Immunotherapy + targeted therapy + chemotherapy	Atezolizumab + bevacizumab + carboplatin + paclitaxel
	Hellmann <i>et al</i> [35]	Clinical study	Dual immunotherapy	Nivolumab + ipilimumab
Colorectal cancer	Ragusa <i>et al</i> [33]	Preclinical study	Immunotherapy + targeted therapy	Anti-PD-1 + VEGF inhibitors
	Song <i>et al</i> [39]	Preclinical study	Immunotherapy + targeted therapy	Anti-PD-L1 + lipopolysaccharide block
	Wang <i>et al</i> [47]	Clinical study	Immunotherapy + targeted therapy	Toripalimab + regorafenib
	Eng <i>et al</i> [48]	Clinical study	Immunotherapy + targeted therapy	Atezolizumab + cobimetinib
	Hu <i>et al</i> [40]	Preclinical study	Immunotherapy + gene therapy	Anti-PD-1 + relaxin gene therapy
	Kadota <i>et al</i> [49]	Preclinical study	Immunotherapy + targeted therapy	Anti-PD-1 + dasatinib
	Inoue <i>et al</i> [26]	Clinical study	Immunotherapy + chemotherapy	Cetuximab + chemotherapy
Pancreatic cancer	Ho <i>et al</i> [29]	Preclinical study	Immunotherapy + chemotherapy	Anti-PD-1 + gemcitabine
	Qiu <i>et al</i> [50]	Clinical study	Immunotherapy + targeted therapy	Sintilimab + anlotinib + S-1
	Blair <i>et al</i> [51]	Preclinical study	Immunotherapy + targeted therapy	Anti-PD-1 + focal adhesion kinase inhibitor (FAKi) + Anti-CXCR4 antibody
	Matsumoto <i>et al</i> [38]	Preclinical study	Immunotherapy + targeted therapy	Alpha-galactosylceramide (KRN7000) + angiogenesis inhibitor AGM-1470 (TNP470)
	Hu <i>et al</i> [40]	Preclinical study	Immunotherapy + gene therapy	Anti-PD-1 + relaxin gene therapy
	Zhang <i>et al</i> [52]	Case report	Immunotherapy + chemotherapy	Penpulimab + modified FOLFIRINOX
Breast cancer	Schmid <i>et al</i> [53]	Clinical study	Immunotherapy + chemotherapy	Atezolizumab + nab-paclitaxel
	Yu <i>et al</i> [20]	Preclinical study	Immunotherapy + radiotherapy	Anti-PD-L1 + radiotherapy
	Lee <i>et al</i> [21]	Preclinical study	Immunotherapy + targeted therapy	Anti-PD-1 + regulatory T cells targeting therapy
	Hu <i>et al</i> [40]	Preclinical study	Immunotherapy + gene therapy	Anti-PD-1 + relaxin gene therapy
	Ozaki <i>et al</i> [54]	Clinical study	Immunotherapy + targeted therapy + chemotherapy	Nivolumab + bevacizumab + paclitaxel
Melanoma	Tang <i>et al</i> [55]	Clinical study	Immunotherapy + targeted therapy + chemotherapy	Anti-PD-1 + axitinib + TACE
	Hong <i>et al</i> [56]	Case report	Immunotherapy + cryoablation	Anti-PD-1 + cryoablation
	Blomen <i>et al</i> [57]	Clinical study	Dual immunotherapy + liver-specific therapies	Nivolumab + ipilimumab + liver-specific therapies
Gastric cancer	Xu <i>et al</i> [32]	Case report	Immunotherapy + radiotherapy	Anti-PD-L1 + radiotherapy
	Wang <i>et al</i> [58]	Case report	Immunotherapy + targeted therapy	Camrelizumab + lenvatinib

	Peng <i>et al</i> [59]	Case report	Immunotherapy + chemotherapy	AK104 (a PD-1/CTLA-4 bispecific antibody) + mXELOX
Gastric hepatoid adenocarcinoma	Liu <i>et al</i> [60]	Case report	Immunotherapy + targeted therapy + chemotherapy	Pembrolizumab + bevacizumab + epirubicin + albumin binding paclitaxel
Nasopharyngeal cancer	Zhang <i>et al</i> [61]	Case report	Immunotherapy + chemotherapy	Anti-PD-1 + TP regimen (nab-paclitaxel + cisplatin)
Cervical cancer	Nance <i>et al</i> [62]	Case report	Immunotherapy + radiotherapy	Pembrolizumab + Yttrium-90 (Y90)

### Other combination therapies

In clinical practice, in order to improve efficacy, there are often not only dual combination therapies, but also multiple combination therapy methods. For example, the Impower150 study used a combination of four treatment methods to significantly improve the survival of patients[37]. A meta-analysis showed that despite the presence of liver metastasis, the efficacy of bevacizumab + chemotherapy + immunotherapy did not significantly decrease, indicating that the use of targeted therapy and chemotherapy can reverse the systemic immune tolerance caused by liver metastasis in lung cancer[25].

There are still many combination therapy methods that are still being explored in clinical research or continuously attempted in preclinical studies. Matsumoto *et al*[38] found that, the combination of an immune stimulator [alpha-galactosylceramide (KRN7000)] and an angiogenesis inhibitor [AGM-1470 (TNP470)] can significantly inhibit the growth of liver metastases from pancreatic cancer, produce synergistic effects, and improve the therapeutic effect; in a mouse model of colon cancer liver metastasis, the use of lipopolysaccharide trapping system can improve the efficacy of PD-L1 monoclonal antibody[39]. Hu *et al*[40] confirmed that up-regulating the expression of the relaxin (*RLN*) gene can further produce synergistic anti-metastasis effect with PD-L1 in the mouse models of liver metastasis of colorectal cancer, pancreatic cancer, and breast cancer. Although these treatment methods have not yet been translated into clinical applications, basic research is more conducive to developing more scientific combination therapy strategies from the mechanism level, which is conducive to further improving the treatment prognosis of liver metastasis patients through clinical translation.

## SIDE EFFECTS OF COMBINATION THERAPY

The side effects of combination therapy in patients are also an important evaluation indicator for the selection of treatment strategy. Although different combination therapy modes can enhance anti-tumor effects, they inevitably bring different side effects. Compared to using ICIs alone, the incidence of toxicity in combination therapy is higher[41]. The incidence of adverse reactions in immunotherapy combined with chemotherapy is higher than that in combination with targeted therapy or radiotherapy. The combination of ICIs and platinum containing chemotherapy is mainly related to the cytotoxicity of chemotherapy drugs; the combination of VEGF and VEGFR inhibitors can cause hypertension and proteinuria; the combination of liver metastasis radiotherapy is mainly manifested as adverse reactions of the digestive tract[42-44]; and in dual immunotherapy, the combined treatment strategy can also cause relatively more adverse reactions compared to monotherapy[35].

## CONCLUSION

Patients with malignant tumor liver metastasis have a poor prognosis. The liver, as an immune tolerant organ, presents a challenge in clinical treatment due to the decreased efficacy of immunotherapy after liver metastasis. It is of great significance to conduct in-depth research on the characteristics of the liver immune microenvironment and identify potential combination therapy targets for the development of new combination therapy strategies for liver metastasis patients. The combination of immunotherapy with chemotherapy, anti-vascular therapy, or radiotherapy, and dual immunotherapy may all improve the anti-tumor effect of monotherapy. However, current clinical research is still relatively limited, and the safety and effectiveness of combination therapy still need to be further evaluated. In the future, more clinical and translational studies should be carried out to optimize immunotherapy strategies through systematic evaluation schemes, so that liver metastasis patients can gain greater benefits from immunotherapy.

## FOOTNOTES

**Author contributions:** Jiang C, Zhang ZH, and Li JX contributed to this paper; Jiang C designed the overall concept and outline of the manuscript; Zhang ZH and Li JX contributed to the discussion and design of the manuscript; Jiang C and Li JX contributed to the writing and editing of the manuscript, and review of the literature.

**Supported by** the Project of Guizhou Provincial Department of Science and Technology, No. Qian Ke He Cheng Guo LC[2024]109.

**Conflict-of-interest statement:** All the authors declare no conflict of interest related to this article.



**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country of origin:** China

**ORCID number:** Jia-Xin Li [0009-0005-8538-5900](https://orcid.org/0009-0005-8538-5900).

**S-Editor:** Lin C

**L-Editor:** Wang TQ

**P-Editor:** Zhang L

## REFERENCES

- Li X, Ramadori P, Pfister D, Seehawer M, Zender L, Heikenwalder M. The immunological and metabolic landscape in primary and metastatic liver cancer. *Nat Rev Cancer* 2021; **21**: 541-557 [PMID: [34326518](https://pubmed.ncbi.nlm.nih.gov/34326518/) DOI: [10.1038/s41568-021-00383-9](https://doi.org/10.1038/s41568-021-00383-9)]
- Horn SR, Stoltzfus KC, Lehrer EJ, Dawson LA, Tehelebi L, Gusani NJ, Sharma NK, Chen H, Trifiletti DM, Zaorsky NG. Epidemiology of liver metastases. *Cancer Epidemiol* 2020; **67**: 101760 [PMID: [32562887](https://pubmed.ncbi.nlm.nih.gov/32562887/) DOI: [10.1016/j.canep.2020.101760](https://doi.org/10.1016/j.canep.2020.101760)]
- de Ridder J, de Wilt JH, Simmer F, Overbeek L, Lemmens V, Nagtegaal I. Incidence and origin of histologically confirmed liver metastases: an explorative case-study of 23,154 patients. *Oncotarget* 2016; **7**: 55368-55376 [PMID: [27421135](https://pubmed.ncbi.nlm.nih.gov/27421135/) DOI: [10.18632/oncotarget.10552](https://doi.org/10.18632/oncotarget.10552)]
- Oliveira RC, Alexandrino H, Cipriano MA, Alves FC, Tralhão JG. Predicting liver metastases growth patterns: Current status and future possibilities. *Semin Cancer Biol* 2021; **71**: 42-51 [PMID: [32679190](https://pubmed.ncbi.nlm.nih.gov/32679190/) DOI: [10.1016/j.semcancer.2020.07.007](https://doi.org/10.1016/j.semcancer.2020.07.007)]
- Garcia-Vicién G, Mezheyeuski A, Bañuls M, Ruiz-Roig N, Molleví DG. The Tumor Microenvironment in Liver Metastases from Colorectal Carcinoma in the Context of the Histologic Growth Patterns. *Int J Mol Sci* 2021; **22** [PMID: [33546502](https://pubmed.ncbi.nlm.nih.gov/33546502/) DOI: [10.3390/ijms22041544](https://doi.org/10.3390/ijms22041544)]
- Liang Y, Zhang H, Song X, Yang Q. Metastatic heterogeneity of breast cancer: Molecular mechanism and potential therapeutic targets. *Semin Cancer Biol* 2020; **60**: 14-27 [PMID: [31421262](https://pubmed.ncbi.nlm.nih.gov/31421262/) DOI: [10.1016/j.semcancer.2019.08.012](https://doi.org/10.1016/j.semcancer.2019.08.012)]
- Barnhill R, van Dam PJ, Vermeulen P, Champenois G, Nicolas A, Rawson RV, Wilmott JS, Thompson JF, Long GV, Cassoux N, Roman-Roman S, Busam KJ, Scolyer RA, Lazar AJ, Lugassy C. Replacement and desmoplastic histopathological growth patterns in cutaneous melanoma liver metastases: frequency, characteristics, and robust prognostic value. *J Pathol Clin Res* 2020; **6**: 195-206 [PMID: [32304183](https://pubmed.ncbi.nlm.nih.gov/32304183/) DOI: [10.1002/cjp2.161](https://doi.org/10.1002/cjp2.161)]
- Latacz E, Höppener D, Bohlak A, Leduc S, Tabariès S, Fernández Moro C, Lugassy C, Nyström H, Bozóky B, Floris G, Geyer N, Brodt P, Llado L, Van Mileghem L, De Schepper M, Majeed AW, Lazaris A, Dirix P, Zhang Q, Petrillo SK, Vankerckhove S, Joye I, Meyer Y, Gregorieff A, Roig NR, Vidal-Vanaclocha F, Denis L, Oliveira RC, Metrakos P, Grünhagen DJ, Nagtegaal ID, Mollevi DG, Jarnagin WR, D'Angelica MI, Reynolds AR, Doukas M, Desmedt C, Dirix L, Donckier V, Siegel PM, Barnhill R, Gerling M, Verhoef C, Vermeulen PB. Histopathological growth patterns of liver metastasis: updated consensus guidelines for pattern scoring, perspectives and recent mechanistic insights. *Br J Cancer* 2022; **127**: 988-1013 [PMID: [35650276](https://pubmed.ncbi.nlm.nih.gov/35650276/) DOI: [10.1038/s41416-022-01859-7](https://doi.org/10.1038/s41416-022-01859-7)]
- Galjart B, Nierop PMH, van der Stok EP, van den Braak RRJC, Höppener DJ, Daelemans S, Dirix LY, Verhoef C, Vermeulen PB, Grünhagen DJ. Angiogenic desmoplastic histopathological growth pattern as a prognostic marker of good outcome in patients with colorectal liver metastases. *Angiogenesis* 2019; **22**: 355-368 [PMID: [30637550](https://pubmed.ncbi.nlm.nih.gov/30637550/) DOI: [10.1007/s10456-019-09661-5](https://doi.org/10.1007/s10456-019-09661-5)]
- Buisman FE, van der Stok EP, Galjart B, Vermeulen PB, Balachandran VP, Coebergh van den Braak RRJ, Creasy JM, Höppener DJ, Jarnagin WR, Kingham TP, Nierop PMH, Sadot E, Shia J, Groot Koerkamp B, Grünhagen DJ, D'Angelica M, Verhoef C. Histopathological growth patterns as biomarker for adjuvant systemic chemotherapy in patients with resected colorectal liver metastases. *Clin Exp Metastasis* 2020; **37**: 593-605 [PMID: [32691187](https://pubmed.ncbi.nlm.nih.gov/32691187/) DOI: [10.1007/s10585-020-10048-w](https://doi.org/10.1007/s10585-020-10048-w)]
- Pao W, Ooi CH, Birzele F, Ruefli-Brasse A, Cannarile MA, Reis B, Scharf SH, Schubert DA, Hatje K, Pelletier N, Spleiss O, Reed JC. Tissue-Specific Immunoregulation: A Call for Better Understanding of the "Immunostat" in the Context of Cancer. *Cancer Discov* 2018; **8**: 395-402 [PMID: [29545369](https://pubmed.ncbi.nlm.nih.gov/29545369/) DOI: [10.1158/2159-8290.CD-17-1320](https://doi.org/10.1158/2159-8290.CD-17-1320)]
- Botticelli A, Cirillo A, Scagnoli S, Cerbelli B, Strigari L, Cortellini A, Pizzuti L, Vici P, De Galitiis F, Di Pietro FR, Cerbelli E, Ghidini M, D'Amati G, Della Rocca C, Mezi S, Gelibter A, Giusti R, Cortesi E, Ascierto PA, Nuti M, Marchetti P. The Agnostic Role of Site of Metastasis in Predicting Outcomes in Cancer Patients Treated with Immunotherapy. *Vaccines (Basel)* 2020; **8** [PMID: [32353934](https://pubmed.ncbi.nlm.nih.gov/32353934/) DOI: [10.3390/vaccines8020203](https://doi.org/10.3390/vaccines8020203)]
- Takeyasu Y, Yoshida T, Shibaki R, Matsumoto Y, Goto Y, Kanda S, Horinouchi H, Yamamoto N, Motoi N, Ohe Y. Differential Efficacy of Pembrolizumab According to Metastatic Sites in Patients With PD-L1 Strongly Positive (TPS  $\geq$  50%) NSCLC. *Clin Lung Cancer* 2021; **22**: 127-133.e3 [PMID: [33183972](https://pubmed.ncbi.nlm.nih.gov/33183972/) DOI: [10.1016/j.clcc.2020.10.002](https://doi.org/10.1016/j.clcc.2020.10.002)]
- Tumeh PC, Hellmann MD, Hamid O, Tsai KK, Loo KL, Gubens MA, Rosenblum M, Harview CL, Taube JM, Handley N, Khurana N, Nosrati A, Krummel MF, Tucker A, Sosa EV, Sanchez PJ, Banayan N, Osorio JC, Nguyen-Kim DL, Chang J, Shintaku IP, Boasberg PD, Taylor EJ, Munster PN, Algazi AP, Chmielowski B, Dummer R, Grogan TR, Elashoff D, Hwang J, Goldinger SM, Garon EB, Pierce RH, Daud A. Liver Metastasis and Treatment Outcome with Anti-PD-1 Monoclonal Antibody in Patients with Melanoma and NSCLC. *Cancer Immunol Res* 2017; **5**: 417-424 [PMID: [28411193](https://pubmed.ncbi.nlm.nih.gov/28411193/) DOI: [10.1158/2326-6066.CIR-16-0325](https://doi.org/10.1158/2326-6066.CIR-16-0325)]
- Wang X, Ji Q, Yan X, Lian B, Si L, Chi Z, Sheng X, Kong Y, Mao L, Bai X, Tang B, Li S, Zhou L, Cui C, Guo J. The Impact of Liver Metastasis on Anti-PD-1 Monoclonal Antibody Monotherapy in Advanced Melanoma: Analysis of Five Clinical Studies. *Front Oncol* 2020; **10**: 546604 [PMID: [33117684](https://pubmed.ncbi.nlm.nih.gov/33117684/) DOI: [10.3389/fonc.2020.546604](https://doi.org/10.3389/fonc.2020.546604)]
- Liu K, Wu CX, Liang H, Wang T, Zhang JY, Wang XT. Analysis of the impact of immunotherapy efficacy and safety in patients with gastric cancer and liver metastasis. *World J Gastrointest Surg* 2024; **16**: 700-709 [PMID: [38577087](https://pubmed.ncbi.nlm.nih.gov/38577087/) DOI: [10.4240/wjgs.v16.i3.700](https://doi.org/10.4240/wjgs.v16.i3.700)]
- Thomson AW, Knolle PA. Antigen-presenting cell function in the tolerogenic liver environment. *Nat Rev Immunol* 2010; **10**: 753-766 [PMID: [20444444](https://pubmed.ncbi.nlm.nih.gov/20444444/) DOI: [10.1038/nri2879](https://doi.org/10.1038/nri2879)]

- 20972472 DOI: [10.1038/nri2858](https://doi.org/10.1038/nri2858)]
- 18 **Krenkel O**, Tacke F. Liver macrophages in tissue homeostasis and disease. *Nat Rev Immunol* 2017; **17**: 306-321 [PMID: [28317925](https://pubmed.ncbi.nlm.nih.gov/28317925/) DOI: [10.1038/nri.2017.11](https://doi.org/10.1038/nri.2017.11)]
  - 19 **Kubes P**, Jenne C. Immune Responses in the Liver. *Annu Rev Immunol* 2018; **36**: 247-277 [PMID: [29328785](https://pubmed.ncbi.nlm.nih.gov/29328785/) DOI: [10.1146/annurev-immunol-051116-052415](https://doi.org/10.1146/annurev-immunol-051116-052415)]
  - 20 **Yu J**, Green MD, Li S, Sun Y, Journey SN, Choi JE, Rizvi SM, Qin A, Waninger JJ, Lang X, Chopra Z, El Naqa I, Zhou J, Bian Y, Jiang L, Tezel A, Skvarce J, Achar RK, Sitto M, Rosen BS, Su F, Narayanan SP, Cao X, Wei S, Szeliga W, Vatan L, Mayo C, Morgan MA, Schonewolf CA, Cuneo K, Kryczek I, Ma VT, Lao CD, Lawrence TS, Ramnath N, Wen F, Chinnaiyan AM, Cieslik M, Alva A, Zou W. Liver metastasis restrains immunotherapy efficacy *via* macrophage-mediated T cell elimination. *Nat Med* 2021; **27**: 152-164 [PMID: [33398162](https://pubmed.ncbi.nlm.nih.gov/33398162/) DOI: [10.1038/s41591-020-1131-x](https://doi.org/10.1038/s41591-020-1131-x)]
  - 21 **Lee JC**, Mehdizadeh S, Smith J, Young A, Mufazalov IA, Mowery CT, Daud A, Bluestone JA. Regulatory T cell control of systemic immunity and immunotherapy response in liver metastasis. *Sci Immunol* 2020; **5** [PMID: [33008914](https://pubmed.ncbi.nlm.nih.gov/33008914/) DOI: [10.1126/sciimmunol.aba0759](https://doi.org/10.1126/sciimmunol.aba0759)]
  - 22 **He Y**, Han Y, Fan AH, Li D, Wang B, Ji K, Wang X, Zhao X, Lu Y. Multi-perspective comparison of the immune microenvironment of primary colorectal cancer and liver metastases. *J Transl Med* 2022; **20**: 454 [PMID: [36195882](https://pubmed.ncbi.nlm.nih.gov/36195882/) DOI: [10.1186/s12967-022-03667-2](https://doi.org/10.1186/s12967-022-03667-2)]
  - 23 **Fares J**, Fares MY, Khachfe HH, Salhab HA, Fares Y. Molecular principles of metastasis: a hallmark of cancer revisited. *Signal Transduct Target Ther* 2020; **5**: 28 [PMID: [32296047](https://pubmed.ncbi.nlm.nih.gov/32296047/) DOI: [10.1038/s41392-020-0134-x](https://doi.org/10.1038/s41392-020-0134-x)]
  - 24 **Gadgeel S**, Rodríguez-Abreu D, Speranza G, Esteban E, Felip E, Dómine M, Hui R, Hochmair MJ, Clingan P, Powell SF, Cheng SY, Bischoff HG, Peled N, Grossi F, Jennens RR, Reck M, Garon EB, Novello S, Rubio-Viqueira B, Boyer M, Kurata T, Gray JE, Yang J, Bas T, Pietanza MC, Garassino MC. Updated Analysis From KEYNOTE-189: Pembrolizumab or Placebo Plus Pemetrexed and Platinum for Previously Untreated Metastatic Nonsquamous Non-Small-Cell Lung Cancer. *J Clin Oncol* 2020; **38**: 1505-1517 [PMID: [32150489](https://pubmed.ncbi.nlm.nih.gov/32150489/) DOI: [10.1200/JCO.19.03136](https://doi.org/10.1200/JCO.19.03136)]
  - 25 **Qin BD**, Jiao XD, Liu J, Liu K, He X, Wu Y, Ling Y, Duan XP, Qin WX, Wang Z, Zang YS. The effect of liver metastasis on efficacy of immunotherapy plus chemotherapy in advanced lung cancer. *Crit Rev Oncol Hematol* 2020; **147**: 102893 [PMID: [32065969](https://pubmed.ncbi.nlm.nih.gov/32065969/) DOI: [10.1016/j.critrevonc.2020.102893](https://doi.org/10.1016/j.critrevonc.2020.102893)]
  - 26 **Inoue Y**, Hazama S, Suzuki N, Tokumitsu Y, Kanekiyo S, Tomochika S, Tsunedomi R, Tokuhisa Y, Iida M, Sakamoto K, Takeda S, Ueno T, Yoshino S, Nagano H. Cetuximab strongly enhances immune cell infiltration into liver metastatic sites in colorectal cancer. *Cancer Sci* 2017; **108**: 455-460 [PMID: [28075526](https://pubmed.ncbi.nlm.nih.gov/28075526/) DOI: [10.1111/cas.13162](https://doi.org/10.1111/cas.13162)]
  - 27 **Janjigian YY**, Shitara K, Moehler M, Garrido M, Salman P, Shen L, Wyrwicz L, Yamaguchi K, Skoczylas T, Campos Bragagnoli A, Liu T, Schenker M, Yanez P, Tehfe M, Kowalyszyn R, Karamouzis MV, Bruges R, Zander T, Pazo-Cid R, Hitre E, Feeney K, Cleary JM, Poulart V, Cullen D, Lei M, Xiao H, Kondo K, Li M, Ajani JA. First-line nivolumab plus chemotherapy versus chemotherapy alone for advanced gastric, gastro-oesophageal junction, and oesophageal adenocarcinoma (CheckMate 649): a randomised, open-label, phase 3 trial. *Lancet* 2021; **398**: 27-40 [PMID: [34102137](https://pubmed.ncbi.nlm.nih.gov/34102137/) DOI: [10.1016/S0140-6736\(21\)00797-2](https://doi.org/10.1016/S0140-6736(21)00797-2)]
  - 28 **Kang YK**, Chen LT, Ryu MH, Oh DY, Oh SC, Chung HC, Lee KW, Omori T, Shitara K, Sakuramoto S, Chung IJ, Yamaguchi K, Kato K, Sym SJ, Kadowaki S, Tsuji K, Chen JS, Bai LY, Oh SY, Choda Y, Yasui H, Takeuchi K, Hirashima Y, Hagihara S, Boku N. Nivolumab plus chemotherapy versus placebo plus chemotherapy in patients with HER2-negative, untreated, unresectable advanced or recurrent gastric or gastro-oesophageal junction cancer (ATTRACTION-4): a randomised, multicentre, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2022; **23**: 234-247 [PMID: [35030335](https://pubmed.ncbi.nlm.nih.gov/35030335/) DOI: [10.1016/S1470-2045\(21\)00692-6](https://doi.org/10.1016/S1470-2045(21)00692-6)]
  - 29 **Ho TTB**, Nasti A, Seki A, Komura T, Inui H, Kozaka T, Kitamura Y, Shiba K, Yamashita T, Mizukoshi E, Kawaguchi K, Wada T, Honda M, Kaneko S, Sakai Y. Combination of gemcitabine and anti-PD-1 antibody enhances the anticancer effect of M1 macrophages and the Th1 response in a murine model of pancreatic cancer liver metastasis. *J Immunother Cancer* 2020; **8** [PMID: [33188035](https://pubmed.ncbi.nlm.nih.gov/33188035/) DOI: [10.1136/jitc-2020-001367](https://doi.org/10.1136/jitc-2020-001367)]
  - 30 **Su TS**, Liang P, Liang J, Lu HZ, Jiang HY, Cheng T, Huang Y, Tang Y, Deng X. Long-Term Survival Analysis of Stereotactic Ablative Radiotherapy Versus Liver Resection for Small Hepatocellular Carcinoma. *Int J Radiat Oncol Biol Phys* 2017; **98**: 639-646 [PMID: [28581406](https://pubmed.ncbi.nlm.nih.gov/28581406/) DOI: [10.1016/j.ijrobp.2017.02.095](https://doi.org/10.1016/j.ijrobp.2017.02.095)]
  - 31 **Klement RJ**, Guckenberger M, Alheid H, Allgäuer M, Becker G, Blanck O, Boda-Heggemann J, Brunner T, Duma M, Gerum S, Habermehl D, Hildebrandt G, Lewitzki V, Ostheimer C, Papachristofilou A, Petersen C, Schneider T, Semrau R, Wachter S, Andratschke N. Stereotactic body radiotherapy for oligo-metastatic liver disease - Influence of pre-treatment chemotherapy and histology on local tumor control. *Radiother Oncol* 2017; **123**: 227-233 [PMID: [28274491](https://pubmed.ncbi.nlm.nih.gov/28274491/) DOI: [10.1016/j.radonc.2017.01.013](https://doi.org/10.1016/j.radonc.2017.01.013)]
  - 32 **Xu J**, Liu H, Ni G, Huang Y, Liang H, Ni Y, Huang Q, Yang Z. Clinical efficacy of PD-1 inhibitor combined with radiotherapy in a multi-drug resistant patient with liver metastasis from gastric cancer. *Front Surg* 2023; **10**: 1101294 [PMID: [37151866](https://pubmed.ncbi.nlm.nih.gov/37151866/) DOI: [10.3389/fsurg.2023.1101294](https://doi.org/10.3389/fsurg.2023.1101294)]
  - 33 **Ragusa S**, Prat-Luri B, González-Loyola A, Nassiri S, Squadrito ML, Guichard A, Cavin S, Gjorevski N, Barras D, Marra G, Lutolf MP, Perentes J, Corse E, Bianchi R, Wetterwald L, Kim J, Oliver G, Delorenzi M, De Palma M, Petrova TV. Antiangiogenic immunotherapy suppresses desmoplastic and chemoresistant intestinal tumors in mice. *J Clin Invest* 2020; **130**: 1199-1216 [PMID: [32015230](https://pubmed.ncbi.nlm.nih.gov/32015230/) DOI: [10.1172/JCI129558](https://doi.org/10.1172/JCI129558)]
  - 34 **Socinski MA**, Jotte RM, Cappuzzo F, Orlandi F, Stroyakovskiy D, Nogami N, Rodríguez-Abreu D, Moro-Sibilot D, Thomas CA, Barlesi F, Finley G, Kelsch C, Lee A, Coleman S, Deng Y, Shen Y, Kowanetz M, Lopez-Chavez A, Sandler A, Reck M; IMpower150 Study Group. Atezolizumab for First-Line Treatment of Metastatic Nonsquamous NSCLC. *N Engl J Med* 2018; **378**: 2288-2301 [PMID: [29863955](https://pubmed.ncbi.nlm.nih.gov/29863955/) DOI: [10.1056/NEJMoa1716948](https://doi.org/10.1056/NEJMoa1716948)]
  - 35 **Hellmann MD**, Paz-Ares L, Bernabe Caro R, Zurawski B, Kim SW, Carcereny Costa E, Park K, Alexandru A, Lupinacci L, de la Mora Jimenez E, Sakai H, Albert I, Vergnenegre A, Peters S, Syrigos K, Barlesi F, Reck M, Borghaei H, Brahmer JR, O'Byrne KJ, Geese WJ, Bhagavatheswaran P, Rabindran SK, Kasinathan RS, Nathan FE, Ramalingam SS. Nivolumab plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer. *N Engl J Med* 2019; **381**: 2020-2031 [PMID: [31562796](https://pubmed.ncbi.nlm.nih.gov/31562796/) DOI: [10.1056/NEJMoa1910231](https://doi.org/10.1056/NEJMoa1910231)]
  - 36 **Hodi FS**, Chiarion-Sileni V, Gonzalez R, Grob JJ, Rutkowski P, Cowey CL, Lao CD, Schadendorf D, Wagstaff J, Dummer R, Ferrucci PF, Smylie M, Hill A, Hogg D, Marquez-Rodas I, Jiang J, Rizzo J, Larkin J, Wolchok JD. Nivolumab plus ipilimumab or nivolumab alone versus ipilimumab alone in advanced melanoma (CheckMate 067): 4-year outcomes of a multicentre, randomised, phase 3 trial. *Lancet Oncol* 2018; **19**: 1480-1492 [PMID: [30361170](https://pubmed.ncbi.nlm.nih.gov/30361170/) DOI: [10.1016/S1470-2045\(18\)30700-9](https://doi.org/10.1016/S1470-2045(18)30700-9)]
  - 37 **Reck M**, Mok TSK, Nishio M, Jotte RM, Cappuzzo F, Orlandi F, Stroyakovskiy D, Nogami N, Rodríguez-Abreu D, Moro-Sibilot D, Thomas CA, Barlesi F, Finley G, Lee A, Coleman S, Deng Y, Kowanetz M, Shankar G, Lin W, Socinski MA; IMpower150 Study Group. Atezolizumab plus bevacizumab and chemotherapy in non-small-cell lung cancer (IMpower150): key subgroup analyses of patients with EGFR

- mutations or baseline liver metastases in a randomised, open-label phase 3 trial. *Lancet Respir Med* 2019; **7**: 387-401 [PMID: 30922878 DOI: 10.1016/S2213-2600(19)30084-0]
- 38 **Matsumoto G**, Nagai S, Muta M, Tsuruta K, Okamoto A, Toi M. Survival benefit of KRN7000 immune therapy in combination with TNP470 in hamster liver metastasis model of pancreatic cancer. *Oncol Rep* 2003; **10**: 1201-1206 [PMID: 12883681]
- 39 **Song W**, Tiruthani K, Wang Y, Shen L, Hu M, Dorosheva O, Qiu K, Kinghorn KA, Liu R, Huang L. Trapping of Lipopolysaccharide to Promote Immunotherapy against Colorectal Cancer and Attenuate Liver Metastasis. *Adv Mater* 2018; **30**: e1805007 [PMID: 30387230 DOI: 10.1002/adma.201805007]
- 40 **Hu M**, Wang Y, Xu L, An S, Tang Y, Zhou X, Li J, Liu R, Huang L. Relaxin gene delivery mitigates liver metastasis and synergizes with check point therapy. *Nat Commun* 2019; **10**: 2993 [PMID: 31278269 DOI: 10.1038/s41467-019-10893-8]
- 41 **Zhou X**, Yao Z, Bai H, Duan J, Wang Z, Wang X, Zhang X, Xu J, Fei K, Zhang Z, Tan F, Xue Q, Gao S, Gao Y, Wang J, He J. Treatment-related adverse events of PD-1 and PD-L1 inhibitor-based combination therapies in clinical trials: a systematic review and meta-analysis. *Lancet Oncol* 2021; **22**: 1265-1274 [PMID: 34391508 DOI: 10.1016/S1470-2045(21)00333-8]
- 42 **Vogelhuber T**, Eitz KA, Oechsner M, Vogel MME, Combs SE. Analysis of using high-precision radiotherapy in the treatment of liver metastases regarding toxicity and survival. *BMC Cancer* 2021; **21**: 780 [PMID: 34229642 DOI: 10.1186/s12885-021-08488-y]
- 43 **Wu S**, Kim C, Baer L, Zhu X. Bevacizumab increases risk for severe proteinuria in cancer patients. *J Am Soc Nephrol* 2010; **21**: 1381-1389 [PMID: 20538785 DOI: 10.1681/ASN.2010020167]
- 44 **An MM**, Zou Z, Shen H, Liu P, Chen ML, Cao YB, Jiang YY. Incidence and risk of significantly raised blood pressure in cancer patients treated with bevacizumab: an updated meta-analysis. *Eur J Clin Pharmacol* 2010; **66**: 813-821 [PMID: 20401474 DOI: 10.1007/s00228-010-0815-4]
- 45 **Ma K**, Guo Q, Li X. Efficacy and safety of combined immunotherapy and antiangiogenic therapy for advanced non-small cell lung cancer: a real-world observation study. *BMC Pulm Med* 2023; **23**: 175 [PMID: 37208639 DOI: 10.1186/s12890-023-02470-z]
- 46 **Jiang Y**, Qiao S, Li L, Zhu X. Combination of radiotherapy and Anlotinib enhances benefit from immunotherapy to liver metastasis and abscopal tumor from lung cancer. *Int Immunopharmacol* 2024; **128**: 111441 [PMID: 38171056 DOI: 10.1016/j.intimp.2023.111441]
- 47 **Wang F**, He MM, Yao YC, Zhao X, Wang ZQ, Jin Y, Luo HY, Li JB, Wang FH, Qiu MZ, Lv ZD, Wang DS, Li YH, Zhang DS, Xu RH. Regorafenib plus toripalimab in patients with metastatic colorectal cancer: a phase Ib/II clinical trial and gut microbiome analysis. *Cell Rep Med* 2021; **2**: 100383 [PMID: 34622226 DOI: 10.1016/j.xcrm.2021.100383]
- 48 **Eng C**, Kim TW, Bendell J, Argilés G, Tebbutt NC, Di Bartolomeo M, Falcone A, Fakih M, Kozloff M, Segal NH, Sobrero A, Yan Y, Chang I, Uyei A, Roberts L, Ciardiello F; IMblaze370 Investigators. Atezolizumab with or without cobimetinib versus regorafenib in previously treated metastatic colorectal cancer (IMblaze370): a multicentre, open-label, phase 3, randomised, controlled trial. *Lancet Oncol* 2019; **20**: 849-861 [PMID: 31003911 DOI: 10.1016/S1470-2045(19)30027-0]
- 49 **Kadota H**, Yuge R, Shimizu D, Miyamoto R, Otani R, Hiyama Y, Takigawa H, Hayashi R, Urabe Y, Kitada Y, Oka S, Tanaka S. Anti-Programmed Cell Death-1 Antibody and Dasatinib Combination Therapy Exhibits Efficacy in Metastatic Colorectal Cancer Mouse Models. *Cancers (Basel)* 2022; **14** [PMID: 36551634 DOI: 10.3390/cancers14246146]
- 50 **Qiu X**, Lu C, Sha H, Zhu Y, Kong W, Tong F, Wang Q, Meng F, Liu B, Du J. Efficacy and safety of second-line therapy by S-1 combined with sintilimab and anlotinib in pancreatic cancer patients with liver metastasis: a single-arm, phase II clinical trial. *Front Immunol* 2024; **15**: 1210859 [PMID: 38361920 DOI: 10.3389/fimmu.2024.1210859]
- 51 **Blair AB**, Wang J, Davelaar J, Baker A, Li K, Niu N, Shao Y, Funes V, Li P, Pachter JA, Maneval DC, Dezem F, Plummer J, Chan KS, Gong J, Hendifar AE, Pandol SJ, Burkhart R, Zhang Y, Zheng L, Osipov A. Dual Stromal Targeting Sensitizes Pancreatic Adenocarcinoma for Anti-Programmed Cell Death Protein 1 Therapy. *Gastroenterology* 2022; **163**: 1267-1280.e7 [PMID: 35718227 DOI: 10.1053/j.gastro.2022.06.027]
- 52 **Zhang B**, Tang K, Dong X. A case of pancreatic cancer treated with chemotherapy combined with immunotherapy and targeted therapy. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2023; **52**: 578-582 [PMID: 37899397 DOI: 10.3724/zdxbyxb-2023-0258]
- 53 **Schmid P**, Rugo HS, Adams S, Schneeweiss A, Barrios CH, Iwata H, Diéras V, Henschel V, Molinero L, Chui SY, Maiya V, Husain A, Winer EP, Loi S, Emens LA; IMpassion130 Investigators. Atezolizumab plus nab-paclitaxel as first-line treatment for unresectable, locally advanced or metastatic triple-negative breast cancer (IMpassion130): updated efficacy results from a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2020; **21**: 44-59 [PMID: 31786121 DOI: 10.1016/S1470-2045(19)30689-8]
- 54 **Ozaki Y**, Tsurutani J, Mukohara T, Iwasa T, Takahashi M, Tanabe Y, Kawabata H, Masuda N, Futamura M, Minami H, Matsumoto K, Yoshimura K, Kitano S, Takano T. Data of programmed death-ligand 1 expression and VEGF: Nivolumab, bevacizumab and paclitaxel For HER2-negative metastatic breast cancer. *Data Brief* 2022; **45**: 108558 [PMID: 36118297 DOI: 10.1016/j.dib.2022.108558]
- 55 **Tang B**, Mo J, Yan X, Duan R, Chi Z, Cui C, Si L, Kong Y, Mao L, Li S, Zhou L, Lian B, Wang X, Bai X, Xu H, Li C, Dai J, Guo J, Sheng X. Real-world efficacy and safety of axitinib in combination with anti-programmed cell death-1 antibody for advanced mucosal melanoma. *Eur J Cancer* 2021; **156**: 83-92 [PMID: 34425407 DOI: 10.1016/j.ejca.2021.07.018]
- 56 **Hong H**, Shen L, Tan H, Wu Y, Liu Y, Fan W. Local treatment of liver metastasis in a patient with advanced malignant melanoma: A case report. *J Cancer Res Ther* 2023; **19**: 1675-1679 [PMID: 38156937 DOI: 10.4103/jert.jert\_394\_22]
- 57 **Blomen CL**, Kött J, Hartung TI, Torster LK, Gebhardt C. Combination of Immune Checkpoint Inhibitors and Liver-Specific Therapies in Liver-Metastatic Uveal Melanoma: Can We Thus Overcome Its High Resistance? *Cancers (Basel)* 2021; **13** [PMID: 34945010 DOI: 10.3390/cancers13246390]
- 58 **Wang K**, Wang H, Lv Y, Liu H, Liu J, Zhang Y. Camrelizumab combined with lenvatinib in the treatment of gastric cancer with liver metastasis: a case report. *Ann Palliat Med* 2021; **10**: 803-809 [PMID: 33545802 DOI: 10.21037/apm-20-2572]
- 59 **Peng J**, Zhu Q, Peng Z, Chen Z, Liu Y, Liu B. Patients with positive HER-2 amplification advanced gastroesophageal junction cancer achieved complete response with combined chemotherapy of AK104/cadonilimab (PD-1/CTLA-4 bispecific): A case report. *Front Immunol* 2022; **13**: 1049518 [PMID: 36569905 DOI: 10.3389/fimmu.2022.1049518]
- 60 **Liu M**, Luo C, Xie ZZ, Li X. Treatment of gastric hepatoid adenocarcinoma with pembrolizumab and bevacizumab combination chemotherapy: A case report. *World J Clin Cases* 2022; **10**: 5420-5427 [PMID: 35812689 DOI: 10.12998/wjcc.v10.i16.5420]
- 61 **Zhang J**, Tang H, Li H, Jiang D, Luo H. Case report of chemoradiotherapy combined with immunotherapy for liver metastasis and lymph node metastases in the head of the pancreas of nasopharyngeal carcinoma. *Arch Med Sci* 2022; **18**: 1413-1419 [PMID: 36160356 DOI: 10.5114/aoms/152106]
- 62 **Nance ME**, Biedermann GB, Bhat AP, Davis RM. Chemorefractory liver metastasis from cervical cancer successfully treated with a combination of yttrium-90 and immunotherapy. *Radiol Case Rep* 2020; **15**: 1359-1365 [PMID: 32636973 DOI: 10.1016/j.radcr.2020.06.010]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

