


CORRECTIONS

Platelet-activating factor: the biosynthetic and catabolic enzymes


F. SNYDER

Volume 305 (1995), pages 689–705

In Scheme 13 on p. 697 CDP should be replaced by CMP to give the following reaction:



In Scheme 14 on p. 697 CMP should be replaced by PP_i to give the following reaction:



Inhibition of specific binding of okadaic acid to protein phosphatase 2A by microcystin-LR, calyculin-A and tautomycin: method of analysis of interactions of tight-binding ligands with target proteins

A. TAKAI, K. SASAKI, H. NAGAI, G. MIESKES, M. ISOBE, K. ISONO and T. YASUMOTO

Volume 306 (1995), pages 657–665

p. 661, first column, line 8:

$$v = \frac{p}{3} + \cos\left(\frac{\arccos u}{3}\right) \text{ should read } v = \frac{p}{3} + 2 \left| \frac{s}{3} \right|^{\frac{1}{2}} \cos\left(\frac{\arccos u}{3}\right)$$

p. 663, Figure 3: on the ordinate of the inset (tautomycin) [EL₁]/E_i should read φ/E_i

p. 664, second column, line 8: (1.0 ± 10⁶ – 2.5 × 10⁸) should read (1.0 × 10⁶ – 2.5 × 10⁸)

p. 665, first column, line 1: markedly higher affinity for PP1 and for PP2 should read markedly higher affinity for PP1 than for PP2