

RESEARCH ARTICLE

Evaluating dyadic factors associated with self-care in patients with heart failure and their family caregivers: Using an Actor-Partner Interdependence Model

JinShil Kim¹, Kye Hun Kim², Mi-Seung Shin³, Seongkum Heo⁴, Jung-Ah Lee⁵, KyungAh Cho¹, Minjeong An^{6*}

1 College of Nursing, Gachon University, Incheon, South Korea, **2** Department of Cardiovascular Medicine, Medical School, Chonnam National University, Gwangju, South Korea, **3** Gil Medical Center, Division of Cardiology, Department of Internal Medicine, College of Medicine, Gachon University, Incheon, South Korea, **4** Georgia Baptist College of Nursing, Mercer University, Atlanta, GA, United States of America, **5** Sue & Bill Gross School of Nursing, University of California, Irvine, Irvine, CA, United States of America, **6** College of Nursing, Chonnam National University, Gwangju, South Korea

* anminjeong@jnu.ac.kr



OPEN ACCESS

Citation: Kim J, Kim KH, Shin M-S, Heo S, Lee J-A, Cho K, et al. (2024) Evaluating dyadic factors associated with self-care in patients with heart failure and their family caregivers: Using an Actor-Partner Interdependence Model. *PLoS ONE* 19(9): e0308515. <https://doi.org/10.1371/journal.pone.0308515>

Editor: Shigao Huang, Xijing Hospital, Air Force Medical University, CHINA

Received: September 14, 2023

Accepted: July 24, 2024

Published: September 19, 2024

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0308515>

Copyright: © 2024 Kim et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and [Supporting Information](#) files.

Abstract

Dyadic conditions of patients with heart failure and their caregivers may affect both patient self-care and caregiver contribution to patient self-care (CCPS). The purpose of this study was to examine the relationships of patient-caregiver physical function and depressive symptoms to the patient self-care (maintenance and management) and CCPS. Data from 55 were analyzed using an Actor-Partner Interdependence Model to address the aim through AMOS. Patient self-care was very poor. Better patient physical function was related to better patient self-care management (actor effect) and poorer CCPS maintenance (partner effect). Better caregiver physical function was related to CCPS management (actor effect). Severer patient depressive symptoms were related to poorer patient self-care maintenance (actor effect) and poorer CCPS management (partner effect). Physical function and depressive symptoms in patient-caregiver dyads were related to patient self-care and CCPS. To improve patient self-care and CCPS, dyadic support for physical function and depressive symptoms is needed.

Introduction

Heart failure (HF) is a chronic debilitating condition, which commonly results in poor quality of life [1–3] and high rates of hospitalization and mortality [4, 5]. Because self-care affects the status of these health outcomes [1–4], patients with HF need life-long efforts to adhere to the therapeutic guidelines, which is self-care. However, self-care in patients with HF is considerably poor [6–9]. Thus, self-care in patients with HF needs to be improved.

One important factor affecting patient self-care is social support from significant others, including family caregivers, because patients with HF need to adhere to a wide spectrum of

Funding: The National Research Foundation of Korea for Jane Kim supported part of this work for JinShil Kim (Grant #: NRF-2021R1A2C2009491). URL: <http://nrf.re.kr> The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

treatment regimens [10, 11]. In patients with HF, a caregiver can support and assist the patient to understand and follow complex therapeutic recommendations to maintain the body stability (self-care maintenance), recognize signs and symptoms (symptom perception), and to manage symptoms when they occur (self-care management) [12–16]. A caregiver can contribute to the patient's self-care maintenance by reminding and encouraging the patient to take medication as prescribed, encouraging the patient to perform regular physical activities and doing them with the patient, and preparing for and providing low sodium foods to the patient [13]. A caregiver can contribute to the patient's symptom recognition by monitoring and recognizing signs and symptoms of patients [13, 16]. A caregiver can contribute to the patient's self-care management by assisting the patient to determine and manage the causes of signs and symptoms [13, 16]. Thus, in the literature, caregivers' active involvement in patient self-care improved patient self-care and reduced patient hospitalization rates [12, 17–20]. Therefore, patient-caregiver dyad needs to be considered to improve patient self-care and health outcomes.

Both in patients with HF and in their caregivers, depressive symptoms are prevalent and a factor affecting both patient self-care and caregiver contribution to patient self-care (CCPS). In patients with HF, the prevalence of depressive symptoms ranged from 32% to 44% [21, 22], in caregivers of patients with HF, the prevalence ranged from 6% to 64% [23–25]. In another study [26], approximately 21% of both patients with HF and their caregivers had depressive symptoms. High prevalence of depressive symptoms in patients may be due to the debilitating nature of HF symptoms [27]. The prevalence of depressive symptoms in caregivers may be due to increase in caregiving burden [25, 28], which is increased with becoming more severe HF symptoms in patients [29]. In patients with HF, depressive symptoms were consistently associated with poor self-care in a systematic literature review [30] and in caregivers of HF patients, the absence of depressive symptoms or anxiety was associated with an improved quality of life for the patients [31]. In addition, depressive symptoms in patients and in caregivers were associated with increased caregiver burden [28, 32], which may impact patient self-care [33, 34]. Therefore, depressive symptoms in patients with HF and in their caregivers can negatively affect patient self-care and CCPS. However, these relationships in patient-caregiver dyad have not been frequently examined.

Another factor possibly affecting both patient self-care and CCPS is physical functional status of both patients and their caregivers. Physical function in patients with HF is considerably impaired due to HF and HF symptoms [35–37]. Impaired physical function in patients with HF can positively or negatively affect patient self-care [38–40]. For example, patients with impaired physical function due to development or worsening of HF symptoms may feel urgent needs to be involved in self-care to reduce HF symptoms. However, in some cases, patients with impaired physical function may not be involved in self-care actively because of limited physical capacities in preparing and following a low sodium diet and performing regular exercise. Patients with impaired physical function can also affect CCPS because of increased demands and time for caregiving [41].

On the other hand, physical function in caregivers of patients with HF could be impaired because of lack of time and effort for their own self-management due to caregiving [42, 43]. Impaired physical function of caregivers can adversely affect CCPS because of their limited capacity for shopping and preparing low sodium foods for patients, and of recognizing of patients' signs and symptoms and supporting symptom management, which will also adversely affect patient self-care. The relationships of physical function of caregivers to CCPS or patient self-care have been rarely examined in HF. In one study, physical aspects of quality of life assessing the effects of physical function on daily lives were significantly associated with patient self-care management [44]. In addition, the relationships of physical function of

patients and caregivers to patient self-care also have not been frequently examined in HF. Therefore, further research is needed to examine those relationships in patients with HF and in their caregivers.

The purpose of this study was to examine the relationships of depressive symptoms and physical function to CCPS (i.e., contribution to patient self-care maintenance and contribution to patient self-care management) and patient self-care (self-care maintenance and self-care management) in patient-caregiver dyads using an Actor-Partner-Interdependence Model (APIM). An APIM is a useful method to promote understanding of the dyadic experiences and their interrelated characteristics [31, 45]. Specifically, we aimed to (1) compare the levels of dyadic physical function, depressive symptoms, and self-care in HF, and (2) examine whether physical function and depressive symptoms in patient-caregiver dyads were associated with patient self-care and CCPS. We hypothesized that physical function and depressive symptoms in patients with HF and in their caregivers as dyads would affect patient self-care and CCPS, respectively (i.e., actor effects), and their partner's outcome (i.e., partner effects), respectively.

Methods

Study design and setting

A cross-sectional, correlational study design was used to examine the physical function and depressive symptom status of patients with HF and their caregivers as dyads, and their actor and partner associations with patient self-care and CCPS using the data of a parent study. The purpose of the parent study was to examine patient and caregiver factors related to different types of self-care and self-care confidence in patients with HF using non-dyadic analysis methods [46]. The dyadic data of patients and their caregivers for this secondary analysis study were obtained from four university-affiliated hospitals in South Korea. Physicians and nurses at each hospital screened and enrolled patients with HF and their caregivers when they visited hospitals from June 12, 2019 to October 29, 2021.

Sample

Patients were eligible to participate in this study if the following criteria were met: adult patients (≥ 19 years) with a diagnosis of HF and prescription of any HF medication(s) based on the professional guidelines (e.g., beta-blockers, angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, and diuretics) [47]. Caregivers of patients were eligible if the following criteria were met: adults (≥ 19 years), an informal primary caregiver of a patient based on self-report, and no diagnosis of HF. Communal exclusion criteria for both patients and caregivers were end-stage of HF, terminal comorbid conditions (e.g., cancer or organ failure with life expectancy < 6 months), or no documented or self-reported neurocognitive disorders (e.g., stroke, Alzheimer, dementia, or memory disorder, or mental disorders), which could possibly accompany cognitive impairment and preclude understanding of and following the recommended guidelines to manage the care of HF. The judgement of these exclusion criteria was done by the physicians and nurses at each hospital who screened and recruited patients through reviews of the electronic medical records.

For this dyadic analysis, APIM with distinguishable dyads was used [48]. A minimum sample size was estimated to detect the actor and partner effects, given a power of .80 and an alpha of .05. In a prior study [49], 41 patient-caregiver dyads were estimated to examine the effects of patients' and caregivers' emotional distress (independent variables) on their own quality of life as well as their partner's (dependent variable) using APIM regression for distinguishable dyads [49]. Considering the prior study [49], 49 patient-caregiver dyads were needed to have

adequate power (80%) to detect the average effect sizes of the Person 1 and Person 2 actor effects (.275, .250). In this study, data from 55 patient-caregiver dyad were analyzed to address the purpose of this study.

Measurements

The patient and caregiver dyads provided data on patient self-care (patients only), CCPS (caregivers only), physical function (patients and caregivers), and depressive symptoms (patients and caregivers).

Patient self-care. Patient self-care was assessed by the Revised Self-Care of Heart Failure Index, v7.2 (the revised SCHFI) [1]. The revised SCHFI, v7.2 consists of 29 items in the three scales, including the Self-care Maintenance (10 items), the Symptom Perception (11 items), and the Self-care Management (8 items) [1]. For this dyadic analysis, the Self-care Maintenance and Self-care Management scales were used. Each scale of the revised SCHFI basically adopts a five-point Likert scale. Following the scoring algorithm, 0 to 100 standardized scores for each of the scales are computed [50], with higher scores indicating higher levels of self-care maintenance and self-care management, respectively. A cut-point of 70 or greater in each scale indicates as adequate self-care [1]. The reliability and validity of the original version [1, 3] and of the Korean version [51] were supported.

Caregiver contribution to patient self-care. Caregiver contribution to patient self-care was assessed by the Caregiver Contribution to Self-care of Heart Failure Index, v1.0 [CC-SCHFI] [52]. The CC-SCHFI consists of 22 items in the three scales, including the Caregiver Contribution to Self-care Maintenance (10 items), the Caregiver Contribution to Self-care Management (6 items), and the Caregiver Contribution to Self-Care Self-Efficacy (6 items). For this dyadic study, the Caregiver Contribution to Self-care Maintenance and the Caregiver Contribution to Self-care Management scales were used. One item in the CC-SCHFI Self-care Management assessing recognition of patient symptoms when they present was excluded with the approval from the developer in order to include all caregivers of patients with and without HF symptoms. Each scale of the CC-SCHFI basically adopts a four-point Likert scale. Following the scoring algorithm, 0 to 100 standardized scores for each scale was computed [50, 52], with higher scores indicating higher levels of caregiver contribution to self-care maintenance and caregiver contribution to self-care management, respectively. A cut-point of 70 or greater in each scale is assumed as adequate CCPS [1]. The reliability and validity of the original version were supported [53]. The Korean version of the CC-SCHFI was approved by the developer and posted on the website [54], while its psychometric testing is undertaking by authors of this study.

Physical function. Physical function was assessed by the Korean Activity Status Index (KASI) in patients with HF and their caregivers [55]. The KASI consists of 15 daily activities, which has a weighted value for the energy expenditure of each activity [55]. Using a dichotomous option (yes or no), participants are asked whether they were able to perform these activities. A total score is a sum of these weighted values. The possible total scores range from 0 to 79, with higher scores indicating higher levels of physical function. The validity of the KASI was supported by a moderate correlation coefficient between the KASI score and treadmill exercise time ($r_s = 0.62$) and 77% agreement between the KASI-estimated functional class and functional class estimated by exercise [55].

Depressive symptoms. Depressive symptoms were assessed by the Patient Health Questionnaire-9 (PHQ-9) in patients with HF and in their caregivers [56, 57]. Each item adopts a four-point Likert scale (from 0 = *never* to 3 = *always*). The possible total scores range from 0 to 27, with higher scores indicating more severe depressive symptoms. The reliability and validity

of the English version in patients with HF and the Korean version in a geriatric population have been supported [57, 58].

Demographic characteristics. Demographic characteristics of both groups were collected using a standard demographic questionnaire. The demographic characteristics included age, sex, marital status, education level, and the patient-caregiver relationship.

Clinical characteristics. Clinical characteristics of patients were collected by the trained research nurses at each of the four sites using a standardized data abstraction form through medical record reviews. Clinical characteristics included etiology of HF, left ventricular ejection fraction, and prescribed medications. The New York Heart Association (NYHA) functional class of patients was determined by the trained nurses. Caregivers also self-reported if they had any diseases.

Data analysis

Data were analyzed using IBM SPSS Statistics for Windows [59] and AMOS 27.0 software [60]. Patient and caregiver data were paired before data analyses and a significance level of $< .05$ was used in all tests. Descriptive statistics, such as mean with standard deviation or frequency with percent, were performed to describe the sample characteristics of patient and caregiver dyads and research variables. Independent t tests and χ^2 tests were conducted to compare the differences in continuous and categorical sample characteristics between patients and caregivers, respectively. Independent t tests or Mann-Whitney U tests were used to compare the differences in physical function, depressive symptoms, and self-care maintenance and management between patients and caregivers depending on the normality in the distribution of each variable.

A structural equation modeling (SEM) approach was employed to estimate the APIM with distinguishable dyads. In this method, actor effects (associations between an individual's physical function or depressive symptoms and his or her own self-care maintenance and management or contributions) and partner effects (associations between an individual's physical function or depressive symptoms and his or her partner's self-care maintenance and management or contributions). Both actor and partner effects of physical function and depressive symptoms were examined as predictors of patient self-care or CCPS. To report standardized coefficients as well as unstandardized coefficients, second SEM was conducted after standardizing variables with mean and pooled standardization of combined patients and caregivers [61].

Ethical consideration

This study was approved by four Institutional Review Boards of the University affiliated hospitals (GAIRB 2019–187, HC19QEDI0070, HYUH 2019-12-017-008 and CNUH 2021–055). Patients with HF and their caregivers were screened and recruited during their hospital visits. After a patient and his/her caregiver signed a written informed consent form, trained research nurses at each hospital conducted face-to-face interviews to collect data on variables of interest in this study adhering to the research protocol to assure of data quality and avoid any biases. All the research processes were performed according to the principles of the Declaration of Helsinki [62].

Results

Sample characteristics of the patient-caregiver dyads

Fifty-five dyads (patients: 40.0% female, mean age = 65.4 ± 14.6 years; caregivers: 96.4% female, mean age = 52.3 ± 12.6 years) participated in this study (Table 1). Regarding patient clinical

Table 1. Demographic and disease-related characteristics of patients and caregivers (N = 55 dyads).

	Patients	Caregivers
	n (%) or Mean \pm SD	n (%) or Mean \pm SD
Age	65.4 \pm 14.6	52.3 \pm 12.6
Sex (female)	22 (40.0)	53 (96.4)
Marital (married)	41 (74.5)	45 (81.8)
Education		
< High school	28 (50.9)	8 (14.5)
High school	13 (23.6)	30 (54.5)
College and higher	14 (25.5)	17 (30.9)
NYHA (I/II)	37 (67.3)	
Etiology		
Hypertension	12 (21.8)	
Ischemic	24 (43.6)	
Cardiomyopathy	5 (9.1)	
Valvular heart disease	2 (3.6)	
Alcoholic	2 (3.6)	
Arrhythmia	2 (3.6)	
Idiopathic	8 (14.5)	
LVEF	40.7 \pm 13.3	
<40	28 (50.9)	
\geq 40	27 (49.1)	
Relationship		
Spouse		30 (54.5)
Children		17 (30.9)
Siblings		2 (3.6)
Parent		1 (1.8)
Grandchild		3 (5.5)
Missing		2 (3.6)
Disease (yes)		28 (50.9)

Note. LVEF = left ventricular ejection fraction assessed by Echo. NYHA = New York Heart Association. SD = standard deviation.

<https://doi.org/10.1371/journal.pone.0308515.t001>

characteristics, approximately two-thirds of patients (67.3%) were asymptomatic (NYHA functional class I) or mildly symptomatic (NYHA class II). The mean left ventricular ejection fraction was 40.7% (\pm 13.3). Slightly less than half of the patients (43.6%) had an ischemic origin of HF. The patient-caregiver relationships were spouses (54.5%), adult children (30.9%), and other informal family caregivers (10.9%).

Comparisons of physical function, depressive symptoms, patient self-care, and caregiver contribution to patient self-care

In comparisons of physical function and depressive symptoms in patient-caregiver dyads (Table 2), physical function of patients with HF was significantly poorer than that of caregivers (33.5 vs. 67.2, $t = -9.70$, $p < .001$). Depressive symptoms of patients were also significantly severer than those of caregivers (8.9 vs. 2.9, $t = -6.25$, $p < .001$).

Patient self-care (maintenance $M \pm SD = 52.7 \pm 20.2$; management = 46.5 ± 18.0) and CCPS (maintenance = 45.8 ± 19.6 ; management = 33.4 ± 23.7) were poor (Table 2). The scores of self-care management in patients and CCPS management in caregivers were significantly

Table 2. Differences in physical function, depressive symptoms, self-care maintenance, and self-care management between patients with heart failure and their caregivers (N = 55 dyads).

	Patients		Caregivers		t or U or χ^2	p
	n (%) or Mean \pm SD	Range	n (%) or Mean \pm SD	Range		
Physical function	33.5 \pm 21.8	0–76.8	67.2 \pm 13.7	29.8–76.8	-9.70	< .001
Depressive symptoms	8.9 \pm 6.3	0.0–24.0	2.9 \pm 3.3	0.0–17.0	-6.25	< .001
Self-care maintenance [‡]	52.7 \pm 20.2	7.5–92.5	45.8 \pm 19.6	3.3–90.0	-1.83 [†]	.067
<70 (inadequate level)	44 (80.0)		48 (87.3)		1.06	.440
\geq 70 (adequate level)	11 (20.0)		7 (12.7)			
Self-care management [§]	46.5 \pm 18.0	15.6–100.0	33.4 \pm 23.7	0.0–100.0	-3.47 [†]	< .001
<70 (inadequate level)	49 (89.1)		52 (94.5)		1.09	.489
\geq 70 (adequate level)	6 (10.9)		3 (5.5)			

Note. [†]Mann-Whitney U

[‡]Self-care maintenance in patients and contribution to self-care maintenance in caregivers

[§]Self-care management in patients and contribution to self-care management in caregivers

SD = standard deviation.

<https://doi.org/10.1371/journal.pone.0308515.t002>

different in patient-caregiver dyads (46.5 vs. 33.4, $p < .001$), but not the scores of patient self-care maintenance and caregiver contribution to self-care maintenance (52.7 vs. 45.8, $p = .067$). Using a cut-point of 70 or greater, small percentages of dyads demonstrated self-care adequacy. In patients, 20.0% and 10.9% showed self-care adequacy in the maintenance and management, respectively. In caregivers, 12.7% and 5.5% showed CCPS adequacy in the maintenance and management, respectively.

Actor and partner effects of physical function and depressive symptoms for patient self-care and caregiver contribution to patient self-care

Fig 1 illustrates significant actor-partner relationships of physical function and depressive symptoms to patient self-care and CCPS. There were significant actor effects of patient and caregiver physical function on patient self-care management and CCPS management, respectively (Table 3). Better patient physical function was related to better patient self-care management ($b = 0.30$, $p = .005$, actor effect), and better caregiver physical function was related to better CCPS management ($b = 0.54$, $p = .015$, actor effect) (Table 3). In addition, there was a significant partner effect of patient physical function on CCPS maintenance. Better patient




Fig 1. Relationships of physical function and depressive symptoms to self-care. Note. * $p < .05$, ** $p < .01$, *** $p < .001$. PT, patients; CG, caregivers; e1, error term of patients' self-care maintenance; e2, error term of caregivers' contribution to self-care maintenance; e3, error term of patients' self-care management; e4, error term of caregivers' contribution to self-care management. Continuous lines indicate significant relationships, and dotted lines indicate non-significant relationships.

<https://doi.org/10.1371/journal.pone.0308515.g001>

Table 3. Actor-Partner Interdependence Model parameter estimates for self-care maintenance and management.

	Self-Care Maintenance [†]				Self-Care Management [‡]			
	b	beta	p	SE	b	beta	p	SE
Physical function								
Actor effects								
Patients	-0.07	-0.05	.548	.11	0.30	0.30	.005	.11
Caregivers	-0.24	-0.17	.190	.18	0.54	0.48	.015	.22
Partner effects								
Patients on caregivers	-0.37	-0.38	.001	.11	-0.22	-0.19	.121	.14
Caregivers on patients	0.27	0.21	.136	.18	-0.02	-0.01	.926	.17
Depressive symptoms								
Actor effects								
Patients	-1.37	-1.34	< .001	.39	-0.22	-0.22	.536	.36
Caregivers	0.22	0.26	.770	.74	0.01	-0.05	.989	.90
Partner effects								
Patients on caregivers	-0.52	-0.53	.186	.39	-1.11	-1.05	.019	.47
Caregivers on patients	-0.19	-0.24	.799	.74	-0.11	-0.11	.869	.68

Note.

[†]Self-care maintenance in patients and contribution to self-care maintenance in caregivers

[‡]Self-care management in patients and contribution to self-care management in caregivers

SE = standard error. b = unstandardized coefficient. beta = standardized coefficient.

<https://doi.org/10.1371/journal.pone.0308515.t003>

physical function was related to poorer CCPS maintenance ($b = -0.37, p = .001$) (partner effect). There was a significant actor effect of patient depressive symptoms on patient self-care maintenance and a partner effect on CCPS management. Severe depressive symptoms were related to poorer patient self-care maintenance ($b = -1.37, p < .001$; actor effect) and poorer CCPS management ($b = -1.11, p = .019$; partner effect).

Discussion

To the best of our knowledge, we initially examined whether physical function and depressive symptoms of patient-caregiver dyads were related to patient self-care maintenance and management and CCPS maintenance and management in HF. The findings of this study partially supported the hypotheses suggested. In this study, actor-partner effects of physical function and depressive symptoms of dyads for self-care differed by self-care types. Actor effects in the relationships of better physical function to better patient self-care management and CCPS management were found, while a partner effect in the relationship of better patient physical function to poorer CCPS maintenance was found. An actor effect in the relationship of severe depressive symptoms to poorer patient self-care maintenance and a partner effect in the relationship of severe depressive symptoms to poorer CCPS management were found. However, caregiver physical function and depressive symptoms were not related to patient self-care maintenance or management (i.e., partner effects). The overall findings of this study suggest that clinicians and researchers need to consider patient-caregiver dyads and their physical and psychological conditions to improve patient self-care and CCPS.

According to a recent meta-analysis mostly from the western countries [63], self-care in patients with HF was poor in all three self-care dimensions (maintenance, management and confidence). Adequate self-care maintenance (≥ 70 out of 100) was found only in 4 out of 36 studies, and adequate self-care management (≥ 70) was found in none of 35 studies. Patients

from Eastern Asian countries, including Korea, China, and Taiwan, also demonstrated poor self-care with mean scores of patient self-care maintenance ranging from 39.7 (China) to 66.0 (Taiwan) and patient self-care management ranging from 29.7 (Taiwan) to 63.1 (Taiwan) [63]. In this study, patient self-care was also very poor. The mean scores of patient self-care maintenance and management were 52.7 and 46.5, respectively. Using the cut-point of 70, only 20.0% and 10.9% of patients showed adequate patient self-care maintenance and management, respectively. Caregivers can contribute to patient self-care by helping their patients maintain recommended behavioral recommendations and perform management strategies when symptoms occur [13, 16]. However, CCPS maintenance (55.9–70.0 out of 100) and management (55.3–67.6 out of 100) were also poor in Italian caregivers of patients with HF [52, 53]. Caregiver contribution to patient self-care has not been frequently examined in East Asian countries. In China [64], the mean scores of CCPS maintenance and management were lower than those in Italy. In this study, the mean scores of CCPS maintenance (45.8) and management (33.4) were even slightly lower than those in China. Using the cut-point of 70, only 12.7% and 5.5% of caregivers showed adequate CCPS maintenance and management, respectively. The overall findings in the prior studies and in this study strongly demonstrate the needs for improvement in patient self-care and CCPS.

Poor self-care can be attributed to patients' and/or their caregivers' depressive symptoms or impaired physical function [30, 34, 38, 65]. Depressive symptoms are very common in patients with HF in East Asians and Western countries, ranged from 32% to 46% [21, 22, 66]. In this study, the prevalence of depressive symptoms in patients with HF (45.5%) was similar to that in the prior studies. Depressive symptoms are also common in caregivers of patients with HF in Eastern and Western countries, ranging from 6% to 64% [23–25]. The prevalence of depressive symptoms in both patients and their caregivers in patient-caregiver dyads was 21% [26]. In this study, the prevalence of depressive symptoms in caregivers (3.6%) was similar to the lowest prevalence in other studies. On the other hand, impaired physical function in patients with HF and their caregivers has been reported in prior studies [35–37, 42, 43]. In this study, physical function score in patients was 33.5 out of 79 (≥ 46 : no functional limitation), indicating considerable impairment of physical function, while physical function score in caregivers (67.2) was considerably better than that in patients and indicated no functional limitation in caregivers. Thus, overall, our findings in conjunction with prior studies demonstrate high prevalence of depressive symptoms and impaired physical function in patients, while relatively varied prevalence in caregivers depending on studies. However, in prior studies, the prevalence of both depressive symptoms and physical function in patient-caregiver dyads has not been frequently examined.

In this study, each of physical function and depressive symptoms in patients had both actor and partner effects on one type of patient self-care and one type of CCPS. These findings indicate that the impaired physical and psychological conditions of patients adversely affect not only patient self-care but also CCPS. Contrastingly, caregiver physical function had only the actor effect on CCPS, while caregiver depressive symptoms showed no actor or partner effects on either patient self-care or CCPS. These findings provide valuable insights regarding the dyadic relationships of physical function and depressive symptoms to patient self-care or CCPS that have not been tested fully in HF, especially using APIM. Previous studies [30, 38] examined and supported the relationships of physical function and/or depressive symptoms in patients with patient self-care, consistent with our study findings. However, the relationships of physical function and depressive symptoms in patients to CCPS have been rarely examined in HF. Nevertheless, the study findings suggest their effects on CCPS maintenance and management. Considering the significant relationships between CCPS and patient self-care [52], these findings emphasize the importance of improvements in physical function and depressive

symptoms in patients with HF to improve patient self-care. On the other hand, the relationships of physical symptoms and depressive symptoms in caregivers to patient self-care and CCPS have been rarely examined among patient-caregiver dyad in HF. The finding of this study demonstrated the importance of caregiver physical function in CCPS maintenance. One possible reason for no other actor and partner effects of caregiver physical function and depressive symptoms on patient self-care and CCPS may be relatively good physical function and a very low prevalence of depressive symptoms in this caregiver sample. In other studies, impaired physical function and high prevalence of depressive symptoms in caregivers of patients with HF have been reported [23–25, 42, 43]. Therefore, further studies are needed to examine the comprehensive relationships of physical function and depressive symptoms to patients' self-care and CCPS in patient-caregiver dyads in larger sample studies with appropriate prevalence of impaired physical function and depressive symptoms.

Dyadic perspectives of HF self-care and influential factors for self-care have received increased attention over last decade, while evidence remains lacking. Our results indicate several implications for clinical practice and research. The findings of this study imply that assessment of physical and psychological conditions of both patients and caregivers are critically important to improve patient self-care and caregiver contribution in clinical practice. Poor self-care and CCPS in Korean patients and their caregivers also warrant educational support to promote their understanding of the self-care concept and the importance of caregiver assistance in the self-care process, especially in caregivers who showed substantially poor CCPS in clinical practice.

The findings of this study suggest the directions for practice and future studies. In this study, the prevalence of inadequate patient self-care, inadequate CCPS, impaired physical function in patients, and depressive symptoms in patients was high. In addition, the findings also suggest the actor and partner effects of physical function and depressive symptoms in patients on patient self-care and CCPS. Furthermore, physical function in caregivers also showed the actor effect on CCPS management. All these findings suggest the needs for improvements in physical function in patients and in caregivers and depressive symptoms in patients. Thus, clinicians need to assess these variables and develop and deliver appropriate interventions to patient-caregiver dyads in order to improve these variables, and, in turn, to improve patient self-care. The intervention effects on both physical function and depressive symptoms in patient-caregiver dyad have not been frequently tested in HF. A mobile health intervention study was provided to both patients and caregivers, and improved patients' self-care and emotional status, but the effects on caregivers have not been examined [67]. An education and psychosocial support intervention was also provided to both patients with HF and their caregivers, but did not affect depressive symptoms in patients and caregivers, self-care in patients, and caregiver burden in caregivers [68]. Thus, researchers and clinicians need to develop more effective interventions to improve physical function and depressive symptoms, and, in turn, CCPS and patient self-care in patient-caregiver dyads. In addition, further research is also needed to test the hypothesized actor and partner relationships in patient-caregiver dyads using more rigorous study designs, such as longitudinal study designs and larger samples with higher prevalence of impaired physical function and depressive symptoms, especially in caregivers.

This study had several limitations. Although our sample for the dyadic analysis was justified with the adequate power, the validation of the results is warranted in a larger sample study with appropriate prevalence of impaired physical function and depressive symptoms in caregivers. Patients and their caregivers were recruited using a convenience sampling method, limiting the generalization of the study findings. Further, the nature of the cross-sectional design in this study remains unknown whether improvements in physical and psychological

conditions of patient-caregiver dyads will lead to improvements in the dyadic patient self-care and CCPS. Moreover, the study findings may be influenced by cultural factors impacting patient self-care and CCPS, limiting their direct applicability to other cultural or geographical contexts.

The study findings propose several approaches for future research. Firstly, exploring larger samples with random sampling can enhance the robustness of conclusions. Secondly, employing a longitudinal design to examine changes over time and establish causality is recommended. Thirdly, considering the incorporation of various characteristics, including socioeconomic status, education, and social support, can contribute to gaining a more comprehensive understanding of factors associated with self-care.

Conclusions

The findings of this study partially supported the hypothesized actor and partner effects of physical function and depressive symptoms on patient self-care and CCPS in patient-caregiver dyads. Impaired physical function and depressive symptoms in patients showed both the actor and partner effects on poor self-care in patients and also poor CCPS in caregivers. The partner effect of physical function in patients and the actor and partner effects of depressive symptoms in caregivers and in patients, respectively, were not supported. However, the relatively low prevalence of impaired physical function and depressive symptoms in this caregiver sample warrants further studies to examine the hypothesized actor and partner effects using caregiver samples with relatively high prevalence of impaired physical function and depressive symptoms.

Supporting information

S1 Dataset. Study dataset.
(ZIP)

Author Contributions

Conceptualization: JinShil Kim.

Formal analysis: Minjeong An.

Funding acquisition: JinShil Kim.

Investigation: JinShil Kim, Kye Hun Kim, Mi-Seung Shin, Minjeong An.

Project administration: JinShil Kim, Kye Hun Kim, Mi-Seung Shin, KyungAh Cho.

Resources: KyungAh Cho.

Supervision: JinShil Kim.

Validation: Seongkum Heo, Jung-Ah Lee, KyungAh Cho.

Writing – original draft: JinShil Kim, Seongkum Heo, Minjeong An.

Writing – review & editing: JinShil Kim, Kye Hun Kim, Mi-Seung Shin, Seongkum Heo, Jung-Ah Lee, KyungAh Cho.

References

1. Riegel B, Barbaranelli C, Carlson B, Sethares KA, Daus M, Moser DK, et al. Psychometric testing of the revised Self-Care of Heart Failure Index. *J Cardiovasc Nurs.* 2019; 34(2):183–192. <https://doi.org/10.1097/JCN.0000000000000543> PMID: 30303894

2. Seid MA. Health-related quality of life and extent of self-care practice among heart failure patients in Ethiopia. *Health and Quality of Life Outcomes*. 2020; 18(1):27. <https://doi.org/10.1186/s12955-020-01290-7> PMID: 32059670
3. Vellone E, De Maria M, Iovino P, Barbaranelli C, Zeffiro V, Pucciarelli G, et al. The Self-Care of Heart Failure Index version 7.2: further psychometric testing. *Res Nurs Health*. 2020; 43(6):640–650. <https://doi.org/10.1002/nur.22083> PMID: 33141495
4. Lee CS, Bidwell JT, Paturzo M, Alvaro R, Cocchieri A, Jaarsma T, et al. Patterns of self-care and clinical events in a cohort of adults with heart failure: 1 year follow-up. *Heart Lung*. 2018; 47(1):40–46. <https://doi.org/10.1016/j.hrtlng.2017.09.004> PMID: 29054487
5. Pobrotyn P, Mazur G, Kaluzna-Oleksy M, Uchmanowicz B, Lomper K. The level of self-care among patients with chronic heart failure. *Healthcare (Basel)*. 2021; 9(9):1179. <https://doi.org/10.3390/healthcare9091179> PMID: 34574953
6. Buck HG, Hupcey J, Wang H-L, Fradley M, Donovan KA, Watach A. Heart failure self-care within the context of patient and informal caregiver dyadic engagement: a mixed methods study. *Journal of Cardiovascular Nursing*. 2018; 33(4):384–391. <https://doi.org/10.1097/JCN.0000000000000465> PMID: 29601368
7. Lee KS, Choi JO, Jeon ES, Park JH, Jung MS, Lee Z, et al. Distinct factors associated with better self-care in heart failure patients with and without mild cognitive impairment. *J Cardiovasc Nurs*. 2019; 34(6):440–447. <https://doi.org/10.1097/JCN.0000000000000600> PMID: 31373955
8. Massouh A, Abu Saad Huijer H, Meek P, Skouri H. Determinants of self-care in patients with heart failure: observations from a developing country in the Middle East. *J Transcult Nurs*. 2020; 31(3):294–303. <https://doi.org/10.1177/1043659619865587> PMID: 31359818
9. Tawalbeh LI, Al Qadire M, Ahmad MM, Aloush S, Abu Sumaqa Y, Halabi M. Knowledge and self-care behaviors among patients with heart failure in Jordan. *Res Nurs Health*. 2017; 40(4):350–359.
10. Fivecoat HC, Sayers SL, Riegel B. Social support predicts self-care confidence in patients with heart failure. *Eur J Cardiovasc Nurs*. 2018; 17(7):598–604. <https://doi.org/10.1177/1474515118762800> PMID: 29533083
11. Jaarsma T, Hill L, Bayes-Genis A, La Rocca HB, Castiello T, Celutkiene J, et al. Self-care of heart failure patients: practical management recommendations from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail*. 2021; 23(1):157–174. <https://doi.org/10.1002/ehfj.2008> PMID: 32945600
12. Dunbar SB. Family caregivers for persons with heart failure: opportunities to improve outcomes for both. 2023 April 24 [cited 2024 March 22]. In: American Heart Association [Internet]. Available from: <https://professional.heart.org/en/science-news/family-caregiving-for-individuals-with-heart-failure/Commentary>.
13. Durante A, Paturzo M, Mottola A, Alvaro R, Vaughan Dickson V, Vellone E. Caregiver contribution to self-care in patients with heart failure: a qualitative descriptive study. *J Cardiovasc Nurs*. 2019; 34(2):E28–E35. <https://doi.org/10.1097/JCN.0000000000000560> PMID: 30589655
14. Kitko L, McIvannan CK, Bidwell JT, Dionne-Odom JN, Dunlay SM, Lewis LM, et al. Family caregiving for individuals with heart failure: a scientific statement from the American Heart Association. *Circulation*. 2020; 141(22):e864–e878. <https://doi.org/10.1161/CIR.0000000000000768> PMID: 32349542
15. Riegel B, Dunbar SB, Fitzsimons D, Freedland KE, Lee CS, Middleton S, et al. Self-care research: where are we now? where are we going? *Int J Nurs Stud*. 2021; 116:103402. <https://doi.org/10.1016/j.ijnurstu.2019.103402> PMID: 31630807
16. Vellone E, Riegel B, Alvaro R. A Situation-Specific Theory of caregiver contributions to heart failure self-care. *J Cardiovasc Nurs*. 2019; 34(2):166–173. <https://doi.org/10.1097/JCN.0000000000000549> PMID: 30363017
17. Agarwal KS, Bhimaraj A, Xu J, Bionat S, Pudlo M, Miranda D, et al. Decreasing heart failure readmissions among older patients with cognitive impairment by engaging caregivers. *J Cardiovasc Nurs*. 2020; 35(3):253–261. <https://doi.org/10.1097/JCN.0000000000000670> PMID: 32221145
18. Alnomasy N. Impact of caregiver support on patient self-care outcomes with heart failure: a systematic review. *J Nurs Pract* 2020; 3(1):151–161.
19. Clements L. Caregivers' influence on patients' heart failure self-care, hospital readmission and mortality Dissertations, University of Kentucky. 2019. Available from: https://uknowledge.uky.edu/nursing_etds/48/
20. Deek H, Chang S, Newton PJ, Nouredine S, Inglis SC, Arab GA, et al. An evaluation of involving family caregivers in the self-care of heart failure patients on hospital readmission: randomised controlled trial (the FAMILY study). *Int J Nurs Stud*. 2017; 75:101–111. <https://doi.org/10.1016/j.ijnurstu.2017.07.015> PMID: 28772186

21. Mills PJ, Wilson K, Iqbal N, Iqbal F, Alvarez M, Pung MA, et al. Depressive symptoms and spiritual well-being in asymptomatic heart failure patients. *J Behav Med.* 2015; 38(3):407–415. <https://doi.org/10.1007/s10865-014-9615-0> PMID: 25533643
22. Shen BJ, Maeda U. Psychosocial predictors of self-reported medical adherence in patients with heart failure over 6 months: an examination of the influences of depression, self-efficacy, social support, and their changes. *Ann Behav Med.* 2018; 52(7):613–619. <https://doi.org/10.1093/abm/kay003> PMID: 29635447
23. Clements L, Frazier SK, Moser DK, Lennie TA, Chung ML. The mediator effects of depressive symptoms on the relationship between family functioning and quality of life in caregivers of patients with heart failure. *Heart Lung.* 2020; 49(6):737–744. <https://doi.org/10.1016/j.hrtlng.2020.08.011> PMID: 32977035
24. Lacerda MS, Prado PRD, Barros A, Lopes JL. Depressive symptoms in the family caregivers of patients with heart failure: an integrative review. *Rev Gaucha Enferm.* 2019; 40:e20180057.
25. Hu X, Huang W, Su Y, Qu M, Peng X. Depressive symptoms in Chinese family caregivers of patients with heart failure: a cross-sectional study. *Medicine (Baltimore).* 2017; 96(13):e6480. <https://doi.org/10.1097/MD.0000000000006480> PMID: 28353589
26. Liljeroos MA, Miller JL, Lennie TA, Chung ML. Quality of life and family function are poorest when both patients with heart failure and their caregivers are depressed. *Eur J Cardiovasc Nurs.* 2022; 21(3):220–226. <https://doi.org/10.1093/eurjcn/zvab071> PMID: 34436564
27. Fan X, Meng Z. The mutual association between depressive symptoms and dyspnea in Chinese patients with chronic heart failure. *Eur J Cardiovasc Nurs.* 2015; 14(4):310–316. <https://doi.org/10.1177/1474515114528071> PMID: 24634388
28. Bozkurt Zincir S, Sunbul M, Zincir S, Aydin Sunbul E, Oguz M, Feriha Cengiz F, et al. Burden and depressive symptoms associated with adult-child caregiving for individuals with heart failure. *Scientific World Journal.* 2014; 2014(1):641817. <https://doi.org/10.1155/2014/641817> PMID: 25431793
29. Lahoz R, Proudfoot C, Fonseca AF, Loeffroth E, Corda S, Jackson J, et al. Caregivers of patients with heart failure: burden and the determinants of health-related quality of life. *Patient Preference and Adherence.* 2021; 15:1153–1164. <https://doi.org/10.2147/PPA.S297816> PMID: 34079236
30. Sedlar N, Lainscak M, Mårtensson J, Strömberg A, Jaarsma T, Farkas J. Factors related to self-care behaviours in heart failure: a systematic review of European Heart Failure Self-Care Behaviour Scale studies. *Eur J Cardiovasc Nurs.* 2017; 16(4):272–282. <https://doi.org/10.1177/1474515117691644> PMID: 28168895
31. Uchmanowicz I, Faulkner KM, Vellone E, Siennicka A, Szczepanowski R, Olchowska-Kotala A. Heart failure care: testing dyadic dynamics using the Actor-Partner Interdependence Model (APIM)—a scoping review. *International Journal of Environmental Research and Public Health.* 2022; 19(4):1919. <https://doi.org/10.3390/ijerph19041919> PMID: 35206131
32. Chung ML, Lennie TA, Mudd-Martin G, Dunbar SB, Pressler SJ, Moser DK. Depressive symptoms in patients with heart failure negatively affect family caregiver outcomes and quality of life. *Eur J Cardiovasc Nurs.* 2016; 15(1):30–38. <https://doi.org/10.1177/1474515114535329> PMID: 24829295
33. Watson R, Modeste NN, Catolico O, Crouch M. The relationship between caregiver burden and self-care deficits in former rehabilitation patients. *Rehabil Nurs.* 1998; 23(5):258–262. <https://doi.org/10.1002/j.2048-7940.1998.tb01796.x> PMID: 10067641
34. Buck HG, Mogle J, Riegel B, McMillan S, Bakitas M. Exploring the relationship of patient and informal caregiver characteristics with heart failure self-care using the Actor-Partner Interdependence Model: implications for outpatient palliative care. *J Palliat Med.* 2015; 18(12):1026–1032. <https://doi.org/10.1089/jpm.2015.0086> PMID: 26540092
35. Heo S, Lennie TA, Moser DK, Dunbar SB, Pressler SJ, Kim J. Testing of a health-related quality of life model in patients with heart failure: a cross-sectional, correlational study. *Geriatr Nurs.* 2022; 44:105–111. <https://doi.org/10.1016/j.gerinurse.2022.01.008> PMID: 35104725
36. Kitzman DW, Whellan DJ, Duncan P, Pastva AM, Mentz RJ, Reeves GR, et al. Physical rehabilitation for older patients hospitalized for heart failure. *N Engl J Med.* 2021; 385(3):203–216. <https://doi.org/10.1056/NEJMoa2026141> PMID: 33999544
37. McHorney CA, Mansukhani SG, Anatchkova M, Taylor N, Wirtz HS, Abbasi S, et al. The impact of heart failure on patients and caregivers: a qualitative study. *PLoS One.* 2021; 16(3):e0248240. <https://doi.org/10.1371/journal.pone.0248240> PMID: 33705486
38. Ryou C, Kang SM, Jang Y. Factors associated with self-care behaviours among Koreans with heart failure. *Eur J Cardiovasc Nurs.* 2021; 20(1):64–70. <https://doi.org/10.1177/1474515120934060> PMID: 33570596
39. Heo S, Moser DK, Lennie TA, Riegel B, Chung ML. Gender differences in and factors related to self-care behaviors: a cross-sectional, correlational study of patients with heart failure. *Int J Nurs Stud.* 2008; 45(12):1807–1815. <https://doi.org/10.1016/j.ijnurstu.2008.05.008> PMID: 18674762

40. Yoshinaga R, Tomita K, Wakayama K, Furuta S, Miyamoto K, Matsuda Y, et al. Factors related to self-care behaviors among hospitalized patients with heart failure in Japan, based on the European Heart Failure Self-Care Behaviour Scale. *J Phys Ther Sci*. 2022; 34(6):416–421. <https://doi.org/10.1589/jpts.34.416> PMID: 35698558
41. Hu X, Hu X, Su Y, Qu M. Quality of life among primary family caregivers of patients with heart failure in Southwest China. *Rehabil Nurs*. 2018; 43(1):26–34. <https://doi.org/10.1002/rmj.290> PMID: 27349203
42. Jackson JD, Cotton SE, Bruce Wirta S, Proenca CC, Zhang M, Lahoz R, et al. Burden of heart failure on caregivers in China: results from a cross-sectional survey. *Drug Des Devel Ther*. 2018; 12:1669–1678. <https://doi.org/10.2147/DDDT.S148970> PMID: 29922041
43. Ahmad FS, Barg FK, Bowles KH, Alexander M, Goldberg LR, French B, et al. Comparing perspectives of patients, caregivers, and clinicians on heart failure management. *J Card Fail*. 2016; 22(3):210–217. <https://doi.org/10.1016/j.cardfail.2015.10.011> PMID: 26505810
44. Bidwell JT, Vellone E, Lyons KS, D'Agostino F, Riegel B, Juárez-Vela R, et al. Determinants of heart failure self-care maintenance and management in patients and caregivers: a dyadic analysis. *Res Nurs Health*. 2015; 38(5):392–402. <https://doi.org/10.1002/nur.21675> PMID: 26355702
45. Lyons KS, Lee CS. Understanding the family care dyad: a comparison of two multilevel models. *Eur J Cardiovasc Nurs*. 2020; 19(2):178–184. <https://doi.org/10.1177/1474515120902368> PMID: 32041428
46. Heo S, Shin MS, Lee MO, Kim S, Kim SH, Randolph J, et al. Factors related to patients' self-care and self-care confidence in Korean patients with heart failure and their caregivers: a cross-sectional, correlational study. *J Cardiovasc Nurs*. 2023; 38(2):140–49. <https://doi.org/10.1097/JCN.0000000000000922> PMID: 35507026
47. Kim MS, Lee JH, Kim EJ, Park DG, Park SJ, Park JJ, et al. Korean guidelines for diagnosis and management of chronic heart failure. *Korean Circ J*. 2017; 47(5):555–643. <https://doi.org/10.4070/kcj.2017.0009> PMID: 28955381
48. Ackerman RA, Kenny DA. APIMPower: an interactive tool for Actor-Partner Interdependence Model power analysis [Computer software]. 2016 December [cited 2023 March 22]. Available from: <https://robert-a-ackerman.shinyapps.io/apimpower/>.
49. Thomson P, Howie K, Leslie SJ, Angus NJ, Andreis F, Thomson R, et al. Evaluating emotional distress and health-related quality of life in patients with heart failure and their family caregivers: testing dyadic dynamics using the Actor-Partner Interdependence Model. *PLoS One*. 2020; 15(1):e0227129. <https://doi.org/10.1371/journal.pone.0227129> PMID: 31914152
50. Riegel B. Self-care of heart failure index—patient version. 2016 [cited 2024 March 22]. In: Self-Care Measures [Internet]. Available from: <https://self-care-measures.com/available-self-care-measures/self-care-of-heart-failure-index-2/>.
51. Kim J, Kim KH, Lim YH, Heo S, Moon K, Oh MS, et al. Validity and reliability of the Korean version of the revised Self-Care of Heart Failure Index v7.2. *Clin Nurs Res*. 2022; 31(7):1296–1307. <https://doi.org/10.1177/10547738221106590> PMID: 35726492
52. Vellone E, Barbaranelli C, Pucciarelli G, Zeffiro V, Alvaro R, Riegel B. Validity and reliability of the caregiver contribution to self-care of heart failure index version 2. *J Cardiovasc Nurs*. 2020; 35(3):280–290. <https://doi.org/10.1097/JCN.0000000000000655> PMID: 32084084
53. Vellone E, Riegel B, Cocchieri A, Barbaranelli C, D'Agostino F, Glaser D, et al. Validity and reliability of the caregiver contribution to self-care of heart failure index. *J Cardiovasc Nurs*. 2013; 28(3):245–255. <https://doi.org/10.1097/JCN.0b013e318256385e> PMID: 22760172
54. Vellone E. Self care heart failure index—caregiver contribution. n.d. [cited 2024 March 22]. In: Self-Care Measures [Internet]. Available from: <https://self-care-measures.com/available-self-care-measures-caregiver-contributions/caregiver-contribution-self-care-heart-failure-index-2/>.
55. Sung J, On Y, Kim H, Chae I, Sohn D, Oh B, et al. Development of Korean Activity Scale/Index (KASI). *Korean Circulation J*. 2000; 30(8):1004–1009.
56. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001; 16(9):606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x> PMID: 11556941
57. Han C, Jo SA, Kwak J-H, Pae C-U, Steffens D, Jo I, et al. Validation of the Patient Health Questionnaire-9 Korean version in the elderly population: the Ansan Geriatric study. *Comprehensive Psychiatry*. 2008; 49(2):218–223. <https://doi.org/10.1016/j.comppsy.2007.08.006> PMID: 18243897
58. Hammash MH, Hall LA, Lennie TA, Heo S, Chung ML, Lee KS, et al. Psychometrics of the PHQ-9 as a measure of depressive symptoms in patients with heart failure. *Eur J Cardiovasc Nur*. 2013; 12(5):446–453. <https://doi.org/10.1177/1474515112468068> PMID: 23263270
59. IBM Corp. IBM SPSS Statistics for Windows, Version 27.0. Armonk, NY: IBM Corp; 2020.
60. Arbuckle JL. Amos (Version 27.0) [Computer Program]. Chicago: IBM SPSS; 2020.

61. Kenny DA, Kashy DA, Cook WL. Dyadic data analysis. 1st ed. New York: Guilford Publications; 2006.
62. World Medical Association. World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human subjects. *JAMA*. 2013; 310(20):2191–2194. <https://doi.org/10.1001/jama.2013.281053> PMID: 24141714
63. Aghajanloo A, Negarandeh R, Janani L, Tanha K, Hoseini-Esfidarjani S-S. Self-care status in patients with heart failure: systematic review and meta-analysis. *Nursing Open*. 2021; 8(5):2235–2248. <https://doi.org/10.1002/nop2.805> PMID: 33619877
64. Chen Y, Zou H, Zhang Y, Fang W, Fan X. Family caregiver contribution to self-care of heart failure: an application of the Information-Motivation-Behavioral Skills Model. *J Cardiovasc Nurs*. 2017; 32(6):576–583. <https://doi.org/10.1097/JCN.0000000000000398> PMID: 28181927
65. Gary R, Dunbar SB, Higgins M, Butts B, Corwin E, Hepburn K, et al. An intervention to improve physical function and caregiver perceptions in family caregivers of persons with heart failure. *J Appl Gerontol*. 2020; 39(2):181–191. <https://doi.org/10.1177/0733464817746757> PMID: 29347863
66. Lin X-x Gao B-B, Huang J-y. Prevalence of depressive symptoms in patients with heart failure in China: a meta-analysis of comparative studies and epidemiological surveys. *Journal of Affective Disorders*. 2020; 274:774–783. <https://doi.org/10.1016/j.jad.2020.05.099> PMID: 32664014
67. Piette JD, Striplin D, Marinec N, Chen J, Trivedi RB, Aron DC, et al. A mobile health intervention supporting heart failure patients and their informal caregivers: a randomized comparative effectiveness trial. *J Med Internet Res*. 2015; 17(6):e142. <https://doi.org/10.2196/jmir.4550> PMID: 26063161
68. Ågren S, Evangelista LS, Hjelm C, Strömberg A. Dyads affected by chronic heart failure: a randomized study evaluating effects of education and psychosocial support to patients with heart failure and their partners. *J Card Fail*. 2012; 18(5):359–366. <https://doi.org/10.1016/j.cardfail.2012.01.014> PMID: 22555264