



## Original articles

## Analysis of correlation between medial joint line change and lower limb coronal alignment after Oxford unicompartmental knee arthroplasty

Kung-Tseng Hung <sup>a,b</sup>, Kuo-Yao Hsu <sup>a,b</sup>, Chieh-Ming Cheng <sup>a,b</sup>, Yi-Jou Chen <sup>a,b</sup>,  
Chih-Hao Chiu <sup>a,b</sup>, Yi-Shen Chan <sup>b,c</sup>, Alvin Chao-Yu Chen <sup>a,b</sup>, Cheng Pang Yang <sup>a,b,\*</sup><sup>a</sup> Department of Orthopaedic Surgery, Chang Gung Memorial Hospital Linkou Branch, Taoyuan, Taiwan (ROC)<sup>b</sup> Bone and Joint Research Center, Chang Gung Memorial Hospital Linkou Branch, Taoyuan, Taiwan (ROC)<sup>c</sup> Department of Orthopedic Surgery, Keelung Chang Gung Memorial Hospital, Keelung City, Taiwan

## H I G H L I G H T S

- Postoperative HKAA in Oxford UKA had no correlation with the medial joint line change, the thickness of tibial resection and tibial implant.
- Preoperative HKAA had a strong correlation with postoperative HKAA in Oxford UKA.
- Preoperative smaller LDFA and larger MPTA had a moderate correlation with postoperative HKAA in Oxford UKA.

## A R T I C L E I N F O

## Keywords:

Unicompartmental knee arthroplasty  
Joint line  
Hip knee ankle angle  
Correlation

## A B S T R A C T

**Background:** Lower limb coronal alignment was thought to be a predictive factor for Unicompartmental Knee Arthroplasty (UKA) result. The tibial bony resection and implant position lead to joint line change postoperatively. Analysis was done to find out the correlation between these factors.

**Methods:** From 2019 to 2021, 90 medial Oxford UKA were implanted by a single surgeon. Hip Knee Ankle Angle (HKAA), Lateral Distal Femoral Angle (LDFA), Medial Proximal Tibial Angle (MPTA), and intraoperative bony resection thickness were measured. The medial joint line change was calculated. The correlation between joint line change and alignment change was evaluated.

**Results:** The mean tibial resection thickness was 4.3 mm. The mean tibial joint line was elevated by 2.3 mm, while the mean femoral joint line proximalized by 0.8 mm. HKAA changed from 8.4° varus preoperatively to 3.6° varus postoperatively. LDFA changed from 89.0° to 86.7°. MPTA changed from 85.6° to 86.6°. Preoperative HKAA showed a strong correlation with postoperative HKAA ( $p < 0.001$ ), and preoperative MPTA showed a positive correlation with postoperative HKAA ( $p < 0.001$ ). While preoperative LDFA had a negative correlation with postoperative HKAA ( $p < 0.001$ ). The femoral joint line change and LDFA change had a significant correlation with HKAA change ( $p < 0.05$ ).

**Conclusion:** The change of joint line had no correlation with postoperative HKAA in Oxford UKA. Preoperative HKAA strongly correlated with postoperative HKAA; while preoperative smaller LDFA and larger MPTA had a moderate correlation with postoperative HKAA. The femoral joint line change and LDFA change had a weak to moderate correlation with HKAA change.

## Introduction

Unicompartmental Knee Arthroplasty (UKA) serves as a viable solution for isolated compartmental knee osteoarthritis. UKA is characterized by the preservation of bone and ligaments, faster recovery,

improved function, and reduced invasiveness [1,2]. The current phase 3 Oxford medial Unicompartmental Knee Arthroplasty (OUKA) has demonstrated positive survivorship, with a 94% success rate at 10 years across various populations [1-3]. In the realm of UKA, the most common failure modes include aseptic loosening, osteoarthritis progression in the

**Abbreviations:** UKA, Unicompartmental Knee Arthroplasty; OUKA, Oxford medial Unicompartmental Knee Arthroplasty; AMOA, Anteromedial Osteoarthritis; MRI, Magnetic Resonance Imaging; MCL, Medial Collateral Ligament; HKAA, Hip Knee Ankle Angle; LDFA, Mechanical Lateral Distal Femoral Angle; MPTA, Mechanical medial Proximal Tibial Angle; MTJLC, Medial Tibial Joint Line Change; MFJLC, Medial Femoral Joint Line Change; sMTJLC, simulated Medial Tibial Joint Line Change; sMFJLC, simulated Medial Femoral Joint Line Change; MJLET, Medial Joint Line Elevation of the Tibia

\*Corresponding author.

E-mail address: [ronnie80097@gmail.com](mailto:ronnie80097@gmail.com) (C.P. Yang).<https://doi.org/10.1016/j.clinsp.2024.100478>

Received 23 April 2024; Accepted 23 July 2024

contralateral compartment, and bearing dislocation in mobile-bearing systems [1-4].

Valgus malalignment, which places excessive load on the lateral compartment, has been identified as a risk factor for lateral cartilage damage and the progression of knee osteoarthritis [5]. The coronal alignment of the lower limb, particularly the hip-knee-ankle angle, has been reported as a more critical determinant of UKA outcomes than tibial component alignment [6]. In fact, postoperative minor to moderate varus alignment has been associated with positive long-term outcomes, in contrast to severe undercorrection or overcorrection [7,8]. Furthermore, it has been reported that the degree of tibiofemoral angle correction positively correlates with the thickness of the tibial implant in fixed-bearing UKA systems [9]. Postoperative limb alignment in OUKA has also been found to correlate more with the thickness of the insert rather than the alignments of the femoral and tibial components [10].

On another note, variations in implant size and the thickness of intraoperative bony resection could lead to changes in the medial joint line, potentially affecting limb alignment [9,11,12]. To our knowledge, no study has yet evaluated the relationship between changes in the medial joint line and limb alignment in mobile-bearing UKA. This study hypothesized that joint line changes induced by OUKA correlate with limb coronal alignment and may be associated with the development of valgus deformity.

## Materials and methods

### Patient characteristics

This study was approved by the local medical foundation institutional review board with IRB Approval n° 202001568B0C101. Between February 2019 and January 2021, a consecutive series of 103 Oxford UKAs performed by a single senior surgeon was retrospectively reviewed.

UKA was indicated for patients with radiographic evidence of Anteromedial Osteoarthritis (AMOA), functionally intact cruciate and collateral ligaments, passively correctable varus deformity of less than 15°, flexion contracture of less than 15°, a range of motion exceeding 90°, and an asymptomatic patellofemoral joint. Patients with inflammatory arthritis were contraindicated. For comprehensive knee assessment and preoperative surgical planning, each patient underwent Magnetic Resonance Imaging (MRI) to evaluate cartilage loss, anterior cruciate ligament condition, and meniscal lesions. Functional outcomes were assessed postoperatively at two years using the Lysholm score, Oxford Knee Score, and Western Ontario and McMaster University Arthritis (WOMAC) index. The endpoint was set for July 2023.

### Surgical procedure

All surgeries were conducted under general anesthesia using the minimally invasive technique with the mobile-bearing medial Oxford UKA (Zimmer Biomet, Warsaw, IN, USA). Both femoral and tibial components were cemented in each case. The intraoperative tibial resection thickness was measured with a caliper during the operation and recorded in the operative record. Specifically, the anteromedial part of the resected tibia was measured, as this area typically exhibits full-thickness cartilage wear in AMOA. The actual tibial resection thickness was determined by adding 0.89 mm to the recorded thickness to account for the saw blade thickness.

### Radiographic evaluation

Plain films included lower extremity split scanography, knee anteroposterior, lateral, and Merchant radiographs. Preoperatively, a valgus stress view was routinely used to assess deformity correctability and Medial Collateral Ligament (MCL) tension [13,14]. For lower limb coronal alignment, the following indices were measured: 1) Hip Knee Ankle

Angle (HKAA), 2) Mechanical Lateral Distal Femoral Angle (LDFA), 3) Mechanical medial Proximal Tibial Angle (MPTA). HKAA was measured by the angle between the mechanical axes of the femur and tibia, with valgus deformity defined as HKAA > 180°. LDFA was determined by the lateral angle between the distal femoral joint line and the mechanical axis of the femur, while MPTA was defined as the medial angle between the tibial joint line and the mechanical axis of the tibia. Postoperatively, the tibial joint line was measured as a line parallel to the tibial tray (Figs. 1 and 2). Coronal alignment change was calculated by subtracting the preoperative value from the postoperative value [15-17].

### Joint line assessment

Medial Tibial Joint Line Change (MTJLC) was defined as the tibial implant thickness (comprising the tibial tray and insert) minus the actual tibial resection thickness. All tibial trays had a uniform thickness of 2.95 mm, and all inserts had an additional thickness of 0.5 mm, according to the manufacturer's specifications. A positive MTJLC value indicated joint line elevation or proximalization. Medial Femoral Joint Line Change (MFJLC) was defined as the distal femoral implant thickness minus the final spigot number. A positive MFJLC value indicated joint line distalization, while a negative value indicated joint line proximalization or reduction [12,18].

The lateral tibial and femoral cartilage thickness was measured on the preoperative MRI at the T2 coronal slice where intact full-thickness cartilage could be identified (Fig. 3). The authors also calculated the simulated joint line, taking cartilage thickness into account, to represent the healthy condition of the operated knee. The simulated Medial Tibial Joint Line Change (sMTJLC) was defined as MTJLC minus the simulated medial tibial cartilage thickness, which was calculated as the preoperative lateral tibial thickness divided by 1.37, following the study by Eckstein F et al. [19]. The simulated Medial Femoral Joint Line Change (sMFJLC) was defined as MFJLC minus the preoperative lateral femoral cartilage thickness since no significant difference between medial and lateral femoral cartilage thicknesses was found in previous studies [19,20].

### Statistical analysis

Data were reported as means ± Standard Deviations (SDs). The Shapiro-Wilk test was used to assess data normality [21]. Pearson's correlation coefficient ( $r$ ) was applied for normally distributed data, while Spearman's rank correlation was used for nonparametric data. A threshold of  $\alpha = 0.05$  was set for statistical significance. Data analysis was conducted using SPSS 20 (IBM Corp., Armonk, NY, USA).


## Results

### Patient characteristics

A total of 90 knees from 81 patients with complete data were eligible for analysis. The average follow-up duration was 3.3 years (range: 2.5 to 4.4 years). The average age of the patients was 68.8±7.56 years (range: 50 to 85 years). One patient experienced bearing dislocation at two years, which necessitated conversion to total knee arthroplasty. Demographic details are presented in Table 1.

### Radiographic evaluation

The HKAA changed from a preoperative average of 171.6°±3.96° to 176.4°±3.84° postoperatively. The average preoperative and postoperative MPTAs were 85.6°±2.80° and 86.6° ± 2.98°, respectively. The LDFA changed from 89.0° ± 2.50° preoperatively to 86.7° ± 3.03° postoperatively (Table 1). All patients showed significant changes in HKAA, LDFA, and MPTA ( $p < 0.05$ ) (Table 2). Twelve patients had a postoperative HKAA greater than 180°, but no early failures were recorded.



**Fig. 1.** Demonstration of pre-OP and post-OP HKAA measurement. HKAA was measured by the angle between the mechanical axes of the femur and tibia.

### Joint line assessment

The mean medial tibial joint line elevation was  $2.3 \pm 0.68$  mm, and the mean medial femoral joint line proximalization was  $0.8 \pm 1.38$  mm. The simulated tibial joint line elevation was  $0.9 \pm 0.72$  mm, and the simulated femoral joint line reduction was  $2.6 \pm 1.44$  mm (Table 1).

In terms of correlation with postoperative HKAA, preoperative HKAA showed a strong correlation ( $r = 0.777$ ,  $p < 0.001$ ). Preoperative LDFA had moderate negative correlations with postoperative HKAA ( $r = -0.549$ ,  $p < 0.001$ ). Preoperative MPTA demonstrated positive correlations with postoperative HKAA ( $r = 0.637$ ,  $p < 0.001$ ) (Table 3).

Regarding the correlation with HKAA change (Table 4), spigot size, medial femoral joint line change, postoperative LDFA, and LDFA change were statistically significant factors ( $r = -0.272$ ,  $p = 0.009$ ;  $r = 0.300$ ,  $p = 0.004$ ;  $r = -0.251$ ,  $p = 0.017$ ;  $r = -0.252$ ,  $p = 0.016$ , respectively).


### Discussion

In this study, changes in the medial tibial and femoral joint lines, implant size, and tibial resection thickness were not significant factors for the postoperative Hip Knee Ankle Angle (HKAA). However, both preoperative mechanical Lateral Distal Femoral Angle (LDFA) and Mechanical Medial Proximal Tibial Angle (MPTA) correlated with postoperative HKAA. Additionally, spigot size, medial femoral joint line change, and LDFA change were identified as factors correlated with HKAA change.


The joint line represents the physiological status of an individual knee. Malposition of the knee joint line during arthroplasty can result in instability, decreased range of motion, and altered stress in other knee compartments [18]. Finite element models on mobile-bearing UKA have shown that contact stress in the insert and lateral articular cartilage was sensitive to changes in the joint line [22]. Furthermore, Takayama K et al. reported that medial tibial joint line elevation over 5 mm could restrict knee extension and may cause postoperative flexion contracture [23].

The hypothesis that medial tibial joint line elevation correlated with postoperative coronal limb alignment, potentially leading to valgus alignment, was disproven by the present results. It was assumed that MCL tension was the most important factor, as the flexion-extension gap and joint stability after UKA were primarily affected by the MCL. If the tibial joint line was elevated by the implant, femoral milling must be adjusted to achieve optimal MCL tension, thereby keeping the postoperative HKAA within the patient's neutral knee condition. In some cases, a 0.5 mm increment of spigot was used, though not in an officially proven way, to attain optimal gap balance. The overall dislocation rate was 1.1% in the limited case series, which is satisfactory compared to the average 2.4% in the East Asian population who have a higher prevalence of dislocation [24].

Lower limb coronal alignment has been found to be a more determining factor for UKA outcomes than implant component alignments [6]. In terms of mobile-bearing UKA, Kim KT et al. reported that a



**Fig. 2.** Demonstration of pre-OP and post-OP LDFA and MPTA measurement. LDFA was determined by the lateral angle between the distal femoral joint line and the mechanical axis of the femur, while MPTA was defined as the medial angle between the tibial joint line and the mechanical axis of the tibia.



**Fig. 3.** Demonstration of MRI measurement. The lateral femoral and tibial cartilage thickness was measured on T2 coronal slice where cartilage-to-cartilage contact point was noticed. The measurement was made from the contact point to individual subchondral bone plate as red line. Note that medial compartment full cartilage wear with meniscus extrusion was highlighted by the yellow circle.

**Table 1**  
Descriptive statistics.

	All patients (n = 90)
<b>Demographics</b>	
Male / Female	27 / 63
Age, year, mean ± SD	68.8 ± 7.56 (50 ~ 85)
BMI, kg/m <sup>2</sup> , mean ± SD	27.7 ± 3.73 (21.2 ~ 39.0)
Lateral femoral cartilage thickness, mm, mean ± SD	1.8 ± 0.36 (1.0 ~ 3.1)
Lateral tibial cartilage thickness, mm, mean ± SD	2.0 ± 0.40 (1.0 ~ 2.9)
Simulated medial tibial cartilage thickness, mm, mean ± SD	1.4 ± 0.29 (0.8 ~ 2.1)
<b>Intraoperative findings</b>	
<i>Tibial implant size</i>	
AA	3
A	32
B	33
C	13
D	8
E	1
<i>Femoral implant size</i>	
XS	12
S	55
M	20
L	3
<i>Spigot size, mm, mean ± SD</i>	
1	4.1 ± 1.39
1.5	1
2	1
2.5	8
3	2
3.5	18
4	2
4.5	21
5	3
5.5	20
6	1
7	9
8	3
<i>Insert size, mm, mean ± SD</i>	
3	3.2 ± 0.40
4	76
5	13
<i>Tibial resection with saw blade thickness, mm, mean ± SD</i>	
	4.3 ± 0.64 (2.9 ~ 5.9)
<i>Tibial implant thickness, mm, mean ± SD</i>	
	6.6 ± 0.40 (6.5 ~ 8.5)
<i>Femoral implant distal thickness</i>	
	3.3 ± 0.18 (3.0 ~ 3.8)
<b>Joint line change</b>	
Medial Tibial Joint Line Change (MTJLC)	2.3 ± 0.68 (0.6 ~ 3.6)
Medial Femoral Joint Line Change (MFJLC)	-0.8 ± 1.38 (-4.7 ~ 2.3)
Simulated Medial Tibial Joint Line Change (sMTJLC)	0.9 ± 0.72 (-1.1 ~ 2.4)
Simulated Medial Femoral Joint Line Change (sMFJLC)	-2.6 ± 1.44 (-7.2 ~ 0.5)
<b>Preoperative radiographic measurement</b>	
HKAA, degrees, mean ± SD	171.6 ± 3.96 (162.0~180.1)
LDFA, degrees, mean ± SD	89.0 ± 2.50 (82.9~94.8)
MPTA, degrees, mean ± SD	85.6 ± 2.80 (77.3~93.7)
<b>Postoperative radiographic measurement</b>	
HKAA, degrees, mean ± SD	176.4 ± 3.84 (166.2~185.5)
LDFA, degrees, mean ± SD	86.7 ± 3.03 (80.3~95.4)
MPTA, degrees, mean ± SD	86.6 ± 2.98 (80.2~94.1)
<b>Postoperative functional score</b>	
Lysholm	88.7 ± 11.81 (24~100)
Oxford	41.5 ± 4.35 (24~47)
WOMAC	91.1 ± 6.60 (56~100)

SD, Standard Deviation; BMI, Body Mass Index; HKAA, Hip-Knee-Ankle Angle; LDFA, Lateral Distal Femoral Angle; MPTA, Medial Proximal Tibial Angle.

postoperative femorotibial angle of 4 to 6 degrees of valgus had the highest survival rate [25]. In previous studies, Kuwashima et al. concluded that femorotibial angle change and medial joint line elevation of the tibia (MJLET) had a significant correlation [11]. Kuroda et al. reported that a 5.0° change in HKAA was moderately correlated with an MJLET of 4.4 mm [12]. The results differed from the present study; however, the measurement methods for joint line elevation varied. In

**Table 2**  
Coronal alignment change.

	All patients (n = 63)	p-value
HKAA change	4.8 ± 2.61 (-1.1~11.8)	<0.001
LDFA change	-2.3 ± 2.98 (-8.8~5.6)	<0.001
MPTA change	1.0 ± 3.54 (-6.9~11.5)	0.01

HKAA, Hip-Knee-Ankle Angle; LDFA, Lateral Distal Femoral Angle; MPTA, Medial Proximal Tibial Angle.

**Table 3**  
Correlation analysis between different factors and post-op HKAA.

	Coefficient of correlation (r)	p-value
Tibial resection with saw blade thickness <sup>a</sup>	-0.072	0.502
Insert size <sup>a</sup>	-0.048	0.654
Tibial implant thickness <sup>a</sup>	-0.048	0.654
Spigot <sup>a</sup>	-0.012	0.909
Femoral implant distal thickness <sup>a</sup>	-0.096	0.367
Medial Tibial Joint Line Change (MTJLC) <sup>a</sup>	0.062	0.563
Medial Femoral Joint Line Change (MFJLC)	0.005	0.962
Simulated Medial Tibial Joint Line Change (sMTJLC)	0.162	0.127
Simulated Medial Femoral Joint Line Change (sMFJLC)	0.052	0.623
Pre-OP HKAA	0.777	<0.001
HKAA change	0.293	0.005
Pre-OP LDFA	-0.549	<0.001
Post-OP LDFA	-0.389	<0.001
LDFA change	0.063	0.555
Pre-OP MPTA	0.637	<0.001
Post-OP MPTA	0.297	0.005
MPTA change	-0.254	0.016

BMI, Body Mass Index; HKAA, Hip-Knee-Ankle Angle; LDFA, Lateral Distal Femoral Angle; MPTA, Medial Proximal Tibial Angle.

<sup>a</sup> Using Spearman's ρ for nonparametric data.

**Table 4**  
Correlation analysis between different factors and HKAA change.

	Coefficient of correlation	p-value
Tibial resection with saw blade thickness <sup>a</sup>	-0.143	0.178
Insert size <sup>a</sup>	-0.069	0.518
Tibial implant thickness <sup>a</sup>	-0.069	0.518
Spigot <sup>a</sup>	-0.272	0.009
Femoral implant distal thickness <sup>a</sup>	0.137	0.199
Medial tibial joint line change (MTJLC) <sup>a</sup>	0.098	0.360
Medial femoral joint line change (MFJLC)	0.300	0.004
Simulated medial tibial joint line change (sMTJLC)	0.020	0.852
Simulated medial femoral joint line change (sMFJLC)	0.300	0.004
Pre-OP HKAA	-0.375	<0.001
Post-OP HKAA	0.293	0.005
Pre-OP LDFA	-0.003	0.980
Post-OP LDFA	-0.251	0.017
LDFA change	-0.252	0.016
Pre-OP MPTA	-0.024	0.822
Post-OP MPTA	-0.075	0.483
MPTA change	-0.044	0.679

BMI, Body Mass Index; HKAA, Hip-Knee-Ankle Angle; LDFA, Lateral Distal Femoral Angle; MPTA, Medial Proximal Tibial Angle.

<sup>a</sup> Using Spearman's ρ.



Kuwashima's study,[11] the measurement was based solely on radiographic findings, which could be affected by image quality, angular projection, and measurement bias. In Kuroda's study,[12] although intraoperative bony resection thickness was measured as in this study, they did not specify which part of the resected bone was measured, leading to potential measurement bias. In these cases, the authors measured the anteromedial part of the tibia, where full-thickness cartilage wear was noted.

The differences in joint line change measurements, and the fact that the studies by Kuroda and Kuwashima were conducted on fixed-bearing UKAs,[11,12] were noteworthy. While both procedures resurface the damaged compartment, mobile-bearing UKA and fixed-bearing UKA have different traits. Inoue A et al. found that the femorotibial angle changed differently between the two UKA systems, with MCL tension being a determining factor [26]. In mobile-bearing UKA, adequate MCL tension and integrity are essential to provide knee stability and prevent insert dislocation. Using the Oxford knee instrument, incremental milling of the femur could be made to meet the optimal MCL tension and gap balance. However, the fixed-bearing design uses cutting jigs and saw blades for bony resection, which could make it difficult to make accurate adjustments for gap balance.

In this study, the medial tibial joint line elevated by approximately  $2.3 \pm 0.68$  mm, which was less than previously reported [11,12]. However, when accounting for healthy cartilage thickness, the simulated medial tibial joint line was only about  $0.9 \pm 0.72$  mm. Thus, the medial tibial joint line was reconstructed to within less than 1 mm deviation from the simulated healthy knee condition. Additionally, femoral resection in OUKA was performed by milling, allowing surgeons to adjust the depth of femoral milling in a more controlled way. It was found that spigot size, LDFA change, and femoral joint line change all correlated with HKAA change. These findings suggested that the major change in HKAA in OUKA originated from the femoral side rather than the tibial side in this limited case series, contrary to previous studies where tibial joint line change correlated with HKAA change [11,12].

In this series, the only factor that had a strong correlation with postoperative HKAA was preoperative HKAA ( $r = 0.777$ ,  $p < 0.001$ ), suggesting that inherent individual deformity largely determined postoperative alignment. Mullaji AB et al. reported that only preoperative HKAA was predictive of postoperative HKAA [27]. Since UKA mainly involves resurfacing, it was reasonable that more severe preoperative varus deformities tended to remain in varus alignment postoperatively. Additionally, LDFA and MPTA were found to correlate mildly to moderately with postoperative HKAA, with LDFA showing a negative correlation. LDFA and MPTA reflected individual bony morphology; smaller LDFA and larger MPTA biomechanically led to more valgus alignment. In this study, the preoperative LDFA and MPTA were both within normal ranges, but the LDFA of  $89.0^\circ$  was closer to the upper limit, while the MPTA of  $85.6^\circ$  was closer to the lower limit, indicating a preoperative imbalance [15]. Postoperatively, the MPTA and LDFA were  $86.6^\circ$  and  $86.7^\circ$ , respectively, suggesting that the imbalance and malalignment from cartilage wear were greatly reduced by bony resection and implant thickness. Zhang Q et al. also revealed that preoperative smaller LDFA, larger MPTA, and less medial tibial cut thickness were associated with postoperative valgus deformity with HKAA  $> 180^\circ$  in mobile-bearing UKA [28]. Although similar findings were noted in this study, a comparison of the patients into postoperative valgus versus non-valgus groups was not performed, as only twelve of the patients had postoperative HKAA greater than  $180^\circ$ , with an average of  $182.0^\circ$ , representing only a small population.

There were limitations to this study. It was a non-randomized retrospective study with a relatively small sample size. Preoperative functional outcomes of the patients were not presented due to incompleteness. However, all patients underwent comprehensive preoperative studies to confirm isolated compartmental damage with full-thickness cartilage wear. Although the short-term outcomes were favorable, long-term outcomes may differ, and further follow-up is needed.

Moreover, while a simulated healthy joint line was proposed through MRI measurements of cartilage thickness, there could still be significant variance. The most accurate method would involve using a three-dimensional analysis model, but this was too costly for routine use, and currently, no available model can simulate the entire healthy knee cartilage from a damaged medial compartment.

## Conclusion

The medial joint line change, the thickness of tibial resection, and the tibial implant had no correlation with postoperative hip knee ankle angle in Oxford UKA. The only factor of positive strong correlation with postoperative HKAA was preoperative HKAA; while preoperative smaller LDFA and larger MPTA also had moderate correlation with postoperative HKAA. With regard to the HKAA change, femoral joint line change and LDFA change had a weak to moderate correlation.

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgments

Not applicable

## References

- Xue H, Tu Y, Ma T, Wen T, Yang T, Cai M. Up to twelve-year follow-up of the Oxford phase three unicompartmental knee replacement in China: seven hundred and eight knees from an independent centre. *Int Orthop* 2017;41(8):1571–7.
- Pandit H, Hamilton TW, Jenkins C, Mellon SJ, Dodd CA, Murray DW. The clinical outcome of minimally invasive Phase 3 Oxford unicompartmental knee arthroplasty: a 15-year follow-up of 1000 UKAs. *Bone Joint J* 2015;97-b(11):1493–500.
- Mohammad HR, Strickland L, Hamilton TW, Murray DW. Long-term outcomes of over 8,000 medial Oxford Phase 3 Unicompartmental Knees—a systematic review. *Acta Orthop* 2018;89(1):101–7.
- van der List JP, Zuiderbaan HA, Pearle AD. Why do medial unicompartmental knee arthroplasties fail today? *J Arthroplasty* 2016;31(5):1016–21.
- Felson DT, Niu J, Gross KD, Englund M, Sharma L, Cooke TD, et al. Valgus malalignment is a risk factor for lateral knee osteoarthritis incidence and progression: findings from the Multicenter Osteoarthritis Study and the Osteoarthritis Initiative. *Arthritis Rheum* 2013;65(2):355–62.
- Slaven SE, Cody JP, Sershon RA, Ho H, Hopper Jr. RH, Fricka KB. Alignment in medial fixed-bearing unicompartmental knee arthroplasty: the limb has a leg up on the component. *J Arthroplasty* 2021;36(12):3883–7.
- Vasso M, Del Regno C, D'Amelio A, Viggiano D, Corona K, Schiavone Panni A. Minor varus alignment provides better results than neutral alignment in medial UKA. *Knee* 2015;22(2):117–21.
- Hernigou P, Deschamps G. Alignment influences wear in the knee after medial unicompartmental arthroplasty. *Clin Orthop Relat Res* 2004(423):161–5.
- Hopgood P, Martin CP, Rae PJ. The effect of tibial implant size on post-operative alignment following medial unicompartmental knee replacement. *Knee* 2004;11(5):385–8.
- Kim SJ, Bae JH, Lim HC. Factors affecting the postoperative limb alignment and clinical outcome after Oxford unicompartmental knee arthroplasty. *J Arthroplasty* 2012;27(6):1210–5.
- Kuwashima U, Okazaki K, Tashiro Y, Mizu-Uchi H, Hamai S, Okamoto S, et al. Correction of coronal alignment correlates with reconstruction of joint height in unicompartmental knee arthroplasty. *Bone Joint Res* 2015;4(8):128–33.
- Kuroda Y, Takayama K, Ishida K, Hayashi S, Hashimoto S, Tsubosaka M, et al. Medial joint line elevation of the tibia measured during surgery has a significant correlation with the limb alignment changes following medial unicompartmental knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2018;26(11):3468–73.
- Gibson P, Goodfellow J. Stress radiography in degenerative arthritis of the knee. *J Bone Joint Surg Brit* 1986;68-B(4):608–9.
- Tashiro Y, Matsuda S, Okazaki K, Mizu-Uchi H, Kuwashima U, Iwamoto Y. The coronal alignment after medial unicompartmental knee arthroplasty can be predicted: usefulness of full-length valgus stress radiography for evaluating correctability. *Knee Surg Sports Traumatol Arthrosc* 2014;22(12):3142–9.
- Paley D. Normal lower limb alignment and joint orientation. *Principles of Deformity Correction*. Springer; 2002. p. 1–18.
- Marx RG, Grimm P, Lillemo KA, Robertson CM, Ayeni OR, Lyman S, et al. Reliability of lower extremity alignment measurement using radiographs and PACS. *Knee Surg Sports Traumatol Arthrosc* 2011;19(10):1693–8.
- Asada S, Inoue S, Tsukamoto I, Mori S, Akagi M. Obliquity of tibial component after unicompartmental knee arthroplasty. *Knee* 2019;26(2):410–5.
- Weber P, Schroder C, Laubender RP, Baur-Melnyk A, von Schulze Pellengahr C, Jansson V, et al. Joint line reconstruction in medial unicompartmental knee arthroplasty:

- development and validation of a measurement method. *Knee Surg Sports Traumatol Arthrosc* 2013;**21**(11):2468–73.
19. Eckstein F, Winzheimer M, Hohe J, Englmeier KH, Reiser M. Interindividual variability and correlation among morphological parameters of knee joint cartilage plates: analysis with three-dimensional MR imaging. *Osteoarthritis Cartilage* 2001;**9**(2):101–11.
  20. Eckstein F, Winzheimer M, Westhoff J, Schnier M, Haubner M, Englmeier K-H, et al. Quantitative relationships of normal cartilage volumes of the human knee joint – assessment by magnetic resonance imaging. *Anatomy Embryol* 1998;**197**(5):383–90.
  21. Ghasemi A, Zahediasl S. Normality tests for statistical analysis: a guide for non-statisticians. *Int J Endocrinol Metab* 2012;**10**(2):486–9.
  22. Kang KT, Kwon OR, Son J, Suh DS, Kwon SK, Koh YG. Effect of joint line preservation on mobile-type bearing unicompartmental knee arthroplasty: finite element analysis. *Australas Phys Eng Sci Med* 2018;**41**(1):201–8.
  23. Takayama K, Ishida K, Muratsu H, Kuroda Y, Tsubosaka M, Hashimoto S, et al. The medial tibial joint line elevation over 5 mm restrained the improvement of knee extension angle in unicompartmental knee arthroplasty. *Knee Surg Sport Traumatol Arthrosc* 2018;**26**(6):1737–42.
  24. Sun X, Liu P, Lu F, Wang W, Guo W, Zhang Q. Bearing dislocation of mobile bearing unicompartmental knee arthroplasty in East Asian countries: a systematic review with meta-analysis. *J Orthop Surg Res* 2021;**16**(1):28.
  25. Kim KT, Lee S, Kim TW, Lee JS, Boo KH. The influence of postoperative tibiofemoral alignment on the clinical results of unicompartmental knee arthroplasty. *Knee Surg Relat Res* 2012;**24**(2):85–90.
  26. Inoue A, Arai Y, Nakagawa S, Inoue H, Yamazoe S, Kubo T. Comparison of alignment correction angles between fixed-bearing and mobile-bearing UKA. *J Arthroplasty* 2016;**31**(1):142–5.
  27. Mullaji AB, Shetty GM, Kanna R. Postoperative limb alignment and its determinants after minimally invasive Oxford medial unicompartmental knee arthroplasty. *J Arthroplasty* 2011;**26**(6):919–25.
  28. Zhang Q, Zhang Q, Guo W, Gao M, Ding R, Wang W. Risk factors of postoperative valgus malalignment in mobile-bearing medial unicompartmental knee arthroplasty. *Arch Orthop Trauma Surg* 2019;**139**(2):241–8.