

A mixed-methods study of VA video connect utilization among veterans with diabetes experiencing housing instability during the pandemic

DIGITAL HEALTH
Volume 10: 1–14
© The Author(s) 2024
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20552076241282629
journals.sagepub.com/home/dhj



Rebecca L Kinney^{1,2,3} , Laurel A Copeland² , Jack Tsai^{1,4,5},
Alice A Abbott², Kate Wallace², Lorrie A Walker², Jillian Weber⁶,
Shara Katsos¹ and Donald K McInnes^{7,8}

Abstract

Introduction: Prior to the coronavirus disease-2019 (COVID-19) pandemic the U.S. Department of Veterans Affairs (VA) had the largest telehealth program in the United States. The pandemic motivated providers within the VA to expand telehealth in effort to reduce disrupted care while mitigating risks. The pandemic provides a rare opportunity to examine how to better engage veterans experiencing housing instability (HI) in telehealth diabetes care.

Methods: Mixed methods design to examine VA video connect (VVC) diabetes care utilization among veterans experiencing HI from March 1, 2019, to March 1, 2022, combining multivariable regression analyses of VA administrative data with semi-structured interviews. Study aims included: (a) examine changes in diabetes care delivery mode over the peri-pandemic timeframe; (b) identify sociodemographic and clinical characteristics associated with VVC care among veterans with HI; and (c) understand the facilitators and barriers of VVC utilization.

Results: Totally, 5904 veterans were eligible for study analysis. Veterans who are female (OR: 1.63; 95% CI: 1.3, 2.0; $p < 0.0001$), self-identify as Hispanic (OR: 1.44; 95% CI: 1.1, 1.9; $p = 0.02$), are married (OR: 1.39; 95% CI: 1.2, 1.6; $p < 0.0001$), and are in VA priority group 1 (OR: 1.21; 95% CI: 1.1, 1.4; $p = 0.004$) were more likely to use VVC the pandemic. Veterans of older age (OR: 0.97; 95% CI: 0.97, 0.98; $p < .0001$) and rural dwelling (OR: 0.85; 95% CI: 0.7, 1.2; $p = 0.04$), were less likely to use VVC. Thirteen VA providers and 15 veterans were interviewed. Veterans reported that decisions about using VVC were driven by limitations in in-person care availability, safety, and convenience.

Discussion: Telehealth played an important role in providing veterans with HI access to diabetes care during the pandemic. Future interventions should seek to increase education and technology in effort to increase VVC uptake into routine diabetes care to ensure veterans' optimal and equitable access.

Keywords

Video telehealth, diabetes, homelessness, veterans, COVID-19

Submission date: 27 March 2024; Acceptance date: 20 August 2024

¹VA National Center on Homelessness Among Veterans, U.S. Department of Veterans Affairs, Washington, DC, USA

²VA Central Western Massachusetts Healthcare System, Leeds, MA, USA

³Department of Population and Quantitative Health Sciences, University of Massachusetts Medical School, Worcester, MA, USA

⁴Department of Management, Policy, and Community Health, School of Public Health, University of Texas Health Science Center at Houston, Houston, TX, USA

⁵Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA

⁶VHA Homeless Programs Office, U.S. Department of Veterans Affairs, Washington, DC, USA

⁷Center for Healthcare Organization and Implementation Research, VA Bedford Healthcare System, Bedford, MA, USA

⁸Department of Health Law, Policy and Management, Boston University School of Public Health, Boston, MA, USA

Corresponding author:

Rebecca Kinney, National Center on Homelessness among Veterans (NCHAV), VA Central Western Massachusetts Healthcare System, 421 N Main (151), Leeds, MA 01053, USA.
Email: Rebecca.Kinney@va.gov



Introduction

The U.S. Department of Veterans Affairs (VA) has the largest telehealth program in the United States, with over 900,000 veterans receiving 2 million remote care episodes.¹ Prior to 2019, telehealth was available in over 50 VA specialty care areas² (Appendix A), although the majority of remote utilization was for mental health and primary care.^{3,4} Veterans experiencing housing instability (HI) may benefit from telehealth encounters, particularly during times of crisis.⁵ Yet there is a paucity of research examining diabetes care telehealth utilization among veterans experiencing housing instability. The coronavirus disease-2019 (COVID-19) pandemic motivated VA providers (i.e. medical doctors, nurse practitioners, case managers, etc.) to expand telehealth rapidly to reduce disrupted care while mitigating COVID-19 risks.^{6,7} During March and April 2020, weekly VA Video Connect (VVC) encounters rose from 1102 to 13,068 in primary care; 1238 to 21,215 in specialty care.⁸ The pandemic presented a natural experiment in how veterans experiencing HI and their providers responded to a change in diabetes care modality, providing a rare opportunity to examine how the VA can better engage a population with social/economic challenges.

Among veterans experiencing HI, the pandemic exacerbated disparities and potentiated examining barriers to diabetes care.⁹ In the United States, the pandemic revealed wide disparities in infection and recovery rates by place of dwelling, race, ethnicity, socioeconomic status, and comorbidities.¹⁰ For veterans with diabetes experiencing HI, the impact of COVID-19 was two-fold: first, veterans were at increased risk for severe coronavirus infection and its complications;^{11,12} and second, diabetes management was disrupted by reduced access to health care, food security, shelter, and transportation.¹³ While confinement measures may reduce infection, isolation interferes with comprehensive diabetes care which requires regular patient-provider interactions.¹² To address this problem among veterans, clinicians turned to telehealth. Yet, technology barriers and internet unaffordability may have furthered a “digital divide”—disparate access to telehealth for veterans with HI.⁸

We examined the extent to which VVC can be leveraged to mitigate diabetes care disruptions among veterans with HI during the pandemic. Our aims were to: (a) examine changes in diabetes care delivery mode over the peri-pandemic timeframe; (b) identify sociodemographic and clinical characteristics associated with VVC care among veterans with HI; and (c) understand the facilitators and barriers of VVC utilization. Identify the characteristics associated with VVC diabetes care utilization among veterans with HI and VA providers to inform interventions which seek to increase uptake.

Methods

We used a mixed-methods design to examine VVC diabetes care utilization among veterans experiencing HI between

March 1, 2019, through March 1, 2022. Given the Quality Improvement nature of this National Center on Homelessness for Veterans (NCHAV) project, it was deemed exempt from research oversight by VA Central Western Massachusetts ethical review board. This determination included approval to verbally consent participants in the qualitative aim of this study.

Quantitative aim (aims 1 and 2)

To examine changes in diabetes care delivery mode over the peri-pandemic timeframe and to identify the veteran characteristics associated with VVC care. Veterans were identified by diagnostic and stop codes commonly used for housing instability in the VA’s electronic medical record.¹⁴ Stop codes, also referred to as decision support system (DSS) Identifiers, is a VHA term that characterizes VHA Outpatient Clinics by a six-character descriptor that is transmitted to the National Patient Care Database (NPCD) with each separate outpatient encounter. We examined access to telehealth (video and phone) and in-person care among veterans using the Corporate Data Warehouse (CDW), a national repository of clinical, administrative, and financial data from VA medical facilities including detailed information on each clinical visit.¹⁵

Participant eligibility

Eligible Veterans Integrated Services Network (VISN) 1 Veterans included those (a) experiencing homelessness (Stop codes housing instability: 37, 504, 507, 508, 511, 522, 528–530, 555, 556, etc.); (b) with a diabetes diagnosis (ICD-10 codes E11.XX, Z59.XX); (c) who accessed care from a VA between March 1, 2019, and March 1, 2022, in at least one of the healthcare fields of interest (endocrinology, pharmacy, primary care, optometry, other diabetes care specialty, podiatry, MOVE!) utilizing any of the three modalities; and (d) have an electronic health record (EHR) record in the CDW. Veterans were excluded if they did not access VA care between March 1, 2019, and March 1, 2022, or had missing data on the outcome measures described below. Given the data was analyzed retrospectively in aggregate, informed consent was not obtained.

Quantitative analysis

The outcome of outpatient care encounters was classified into type and modality using VA stop codes. Encounters were classified into mutually exclusive categories based on primary, endocrinology, and specialty care services for three years: pre-pandemic year (3/1/2019–2/29/2020), pandemic year 1, PY1 (3/1/2020–2/28/2021), and pandemic year 2, PY2 (3/1/2021–2/28/2022). Visit modality was categorized as in-person, VVC, and phone utilization to form exclusive categories. Bivariate comparisons using Pearson’s chi-square compared

veterans' characteristics by care modality. Of interest were veteran sociodemographics (e.g. age, sex, race/ethnicity, marital status, residential location) and clinical characteristics (e.g. previous outpatient encounters, condition severity, VA priority, and Elixhauser comorbidities).¹⁶ VA priority is a rating from 1 to 6 based on service-connected disability, income, military service.¹⁷ VA priority 1 represents the most economically disadvantaged Veterans versus those in priority groups 2–6. Multivariable logistic regressions were conducted to model associations between the veteran characteristics and the primary outcome, VVC utilization. Adjusted odds ratios or incident rate ratios and 95% confidence intervals (CIs) for each characteristic were reported. All analyses were conducted in SAS Enterprise Guide version 8.3 (SAS Institute, Cary, NC).

Qualitative aim (aim 3)

We conducted semi-structured interviews with both diabetes care providers and veterans to better understand their perceptions of VVC encounters.

Participant eligibility

Providers. For the purposes of this study, participants included VISN 1 VA providers in primary care, endocrinology, podiatry, optometry, pharmacy, and case management. Working with VA medical chiefs and the VA's Deputy Network Homeless Coordinator, we purposively selected providers and case managers throughout the VISN to ensure diverse care delivery settings. Providers and case managers were emailed an invitation to participate in a one-time phone interview, along with a study information sheet. Once individuals consented to participate, we scheduled the 30-minute interview in Microsoft Teams. At the beginning of each interview, a research coordinator reviewed the study with each participant and audio recorded the consent process before starting the interview. We conducted interviews with thirteen VA providers, the point which thematic saturation was reached.¹⁸

Veterans. Participants were VISN 1 veterans experiencing diabetes and homelessness who were offered telehealth encounters within the study timeframe. We purposively selected a diverse group of 15–20 Veterans from our Aim 1 cohort representing different telehealth experiences (users and non-users) and sociodemographics (i.e. age, race, gender). Eligible participants were mailed a study invitation and information sheet. Once a veteran expressed an interest in participating, the individual was contacted by the study coordinator who reviewed the study information sheet, consented the participant, and scheduled the Microsoft Teams interview. Like the provider participants, a research coordinator reviewed the study with each veteran and audio recorded the consent process before starting the interview. After interviewing 15 veterans, we had reached saturation and ceased recruiting more participants.

Qualitative analysis

Interviews queried providers and veterans on their perceptions of the barriers and facilitators to VVC utilization during the pandemic. Using past methods employed by our team, both interview guides were piloted among our expert panel, and revised as needed prior to implementation. Digitally recorded interviews were 90 minutes in length, professionally transcribed, and then analyzed in Atlas.ti.¹⁹ *A priori* codes were based on the interview guide questions and COVID-19 care disruption literature. Taking into consideration these themes, one PhD-level investigator (R.K.) employed open coding of the responses, creating code definitions and schemes as concepts emerged inductively from the data.^{20,21} This approach helped to identify relationships generated through the responses and from theories based on them. Codes with similar meanings were merged and organized into a broader category that covered the meanings. Applying content analysis, the investigator and lead research coordinator (R.K.; L.W.), through an iterative process, independently coded each of the responses and conferred to achieve consensus pertaining to the underlying themes.²² Additional team members (D.K.M. (PhD researcher); A.A. (endocrinologist)) were consulted when discrepancies arose.²³ Collectively, the team systematically and rigorously analyzed the data to reflect the views of all participants and validate salient findings.

Results

Quantitative analysis

Eligible for this analysis were 5904 VISN 1 veterans who were experiencing diabetes and HI during the timeframe of interest and had any relevant visit data, which specified type of care as well as modality, for all three years of the study. Given that all these 5904 had complete data, no one was excluded from the analysis. Approximately, 62,673 clinical encounters were made by 5904 veterans during the study period. Of these encounters, 6267 (13%) visits were exclusively telehealth (VVC or phone); 23,897 (38%) visits were both telehealth and in-person modality; and 30,482 (49%) were only in-person visits. Figure 1 shows that across specialties, the telehealth-only demonstrated increase in the raw count for PY1, followed by a decrease in PY2 as care returned to in-person visits.

There were 1503 (26%) veterans experiencing HI who had at least one VVC diabetes care encounter during PY1. Bivariate comparisons found veteran characteristics associated with VVC utilization included, younger age (60.5 years; SD: 11.2; $p < 0.001$), male (89% vs. 11%, female; $p < 0.0001$), self-identified non-Hispanic White (67% vs 6%, self-identified Hispanic, any race; $p < 0.0001$), and non-married (67% vs. 33%, married; $p < 0.0001$) (Table 1). Residing in an urban area was also associated with VVC utilization (78%




Figure 1. Clinical care modality by specialty among veterans with diabetes and experiencing housing instability across both pandemic years.

vs. 22%, rural; $p = 0.005$). Veterans assigned to VA priority groups 2–6 (51% vs. 49%, Priority 1; $p < 0.0001$) and who had higher comorbidities (88% ≥ 3 vs. 12% 2 or fewer; $p = 0.0006$) were more likely to utilize VVC.

Characteristics of VVC utilization remained constant in PY2, the only difference being veterans in priority group 1 (51%) utilized telehealth slightly more than those in priority groups 2–6 (49%; $p < 0.0001$).

Table 2 shows hierarchical multivariable analyses of the predisposing, enabling, and need characteristics associated with telehealth utilization in PY1. VVC utilization was more likely for those who were female (OR: 1.63; 95% CI: 1.3, 2.0; $p < 0.0001$), self-identified Hispanic (OR: 1.44; 95% CI: 1.1, 1.9; $p = 0.02$), or married (OR: 1.39; 95% CI: 1.2, 1.6; $p < 0.0001$). Veterans who were in VA priority group 1 (OR: 1.21; 95% CI 1.1, 1.4; $p = 0.004$) were more likely to use VVC, as were those seeing a provider in the year prior to the pandemic. Veterans who were older (OR: 0.97; 95% CI 0.97, 0.98; $p < 0.0001$), or residing in rural areas had reduced odds of VVC (OR: 0.85; 95% CI 0.7, 1.0; $p = 0.04$).

In PY2, the count indicates a slight decrease in telehealth utilization (Table 3). Predisposing characteristics of utilizing

VVC included being female (OR: 1.59; 95% CI 1.2, 2.1; $p = 0.0006$) and married (OR: 1.28; 95% CI 1.1, 1.5; $p = 0.003$). Veterans who were more likely to use VVC had needs characteristics which included 3 or more Elixhauser comorbidities (OR: 1.64; 95% CI 1.3, 2.1; $p = 0.07$), VA priority group 1 (OR: 1.28; 95% CI 1.1, 1.5; $p = 0.002$), and having a VVC encounter in PY1 (OR: 1.86; 95% CI 1.5, 2.3; $p < 0.0001$), specifically, a Pharmacy VVC encounter (OR: 6.24; 95% CI 4.1, 9.5; $p < 0.0001$). Veterans residing in a rural location had reduced odds of video telehealth in PY2. There were no significant differences in VVC utilization across race and ethnicity in PY2.

Qualitative analysis

We interviewed 13 VA providers and 15 veterans with diabetes with HI during the study timeframe. VA providers included five physicians (two primary care, two endocrinologists, one podiatrist), one diabetes educator, two dietitians, a VISN 1 Network Homeless Coordinator, two pharmacists, an optometrist, and a physical therapist. Providers were chosen to represent major care needs of veterans with diabetes.

Table 1. Demographics for overall sample, any VVC and/or phone telehealth use among diabetic veterans experiencing housing instability across both pandemic years.

Characteristic	Full sample N = 5904	VVC use, PY1 N = 1503	p-Value	Phone use, PY1 N = 5744	p-Value	VVC use, PY2 N = 1289	p-Value	Phone use, PY2 N = 5468	p-Value
Age (Mean [SD], range)	63.33 (11.18), 21.00-100.00	60.54 (11.18), 23.0-100.0	<0.0001	63.36 (11.11), 21.0-100.0	<0.0001	59.97 (11.38), 23.0-92.0	<0.0001	63.17 (11.04), 21.0-100.0	0.0007
Sex									
Male	5483 (92.9)	1334 (88.8)	<0.0001	5332 (92.8)	<0.0001	1136 (88.1)	<0.0001	5069 (92.7)	0.08
Female	421 (7.1)	169 (11.2)		412 (7.2)		153 (11.9)		399 (7.3)	
Race/ethnicity ^a									
Missing	194 (3.3)	52 (3.5)	<0.0001	179 (3.1)	<0.0001	31 (2.4)	0.04	170 (3.1)	0.03
White, non-Hispanic	4176 (70.7)	1010 (67.2)		4060 (70.7)		886 (68.7)		3849 (70.4)	
Black, non-Hispanic	1171 (19.8)	324 (21.6)		1149 (20.0)		285 (22.1)		1108 (20.3)	
Hispanic, any race	236 (4.0)	87 (5.8)		232 (4.0)		62 (4.8)		223 (4.1)	
Other, Non-Hispanic	127 (2.2)	30 (2.0)		124 (2.2)		25 (1.9)		118 (2.2)	
Marital status									
Missing	28 (0.5)	9 (0.6)	<0.0001	28 (0.5)	<0.0001	5 (0.4)	<0.0001	24 (0.4)	0.27
Married	1699 (28.8)	491 (32.7)		1645 (28.6)		428 (33.2)		1564 (28.6)	
Single/divorced/widow(ed)	4177 (70.8)	1003 (66.7)		4071 (70.9)		856 (66.4)		3880 (71.0)	
VA priority									
Group 1	2396 (40.6)	735 (48.9)	<0.0001	2342 (40.8)	<0.0001	659 (51.1)	<0.0001	2262 (41.4)	<0.0001
Groups 2, 3, 4, 5, 6	3508 (59.4)	768 (51.1)		3402 (59.2)		630 (48.9)		3206 (58.6)	
Last residence									

(continued)

Table 1. Continued.

Characteristic	Full sample N = 5904	VVC use, PY1 N = 1503	Phone use, PY1 N = 5744	VVC use, PY2 N = 1289	Phone use, PY2 N = 5468
Missing	27 (0.5)	5 (0.3)	25 (0.4)	4 (0.3)	27 (0.5)
Urban	4448 (75.3)	1174 (78.1)	4329 (75.4)	1005 (78.0)	4148 (75.9)
Rural	1429 (24.2)	324 (21.6)	1390 (24.2)	280 (21.7)	1293 (23.7)
Elixhauser group					
≤2	880 (14.9)	183 (12.2)	817 (14.2)	155 (12.0)	776 (14.2)
≥3	5024 (85.1)	1320 (87.8)	4927 (85.8)	1134 (88.0)	4692 (85.8)
Prior utilization, Pre-COVID					
Primary care					
Face to face	5560 (94.2)	1460 (97.1)	5436 (94.6)	1239 (96.1)	5176 (94.7)
Video	87 (1.5)	49 (3.3)	86 (1.5)	42 (3.3)	87 (1.6)
Phone	3964 (67.1)	1106 (73.6)	3911 (68.1)	944 (73.2)	3742 (68.4)
Endocrinology					
Face to face	940 (15.9)	342 (22.8)	927 (16.1)	283 (22.0)	906 (16.6)
Video	3 (0.1)	3 (0.2)	3 (0.1)	1 (0.1)	3 (0.1)
Phone	-	-	-	-	-
Pharmacy					
Face to face	2666 (45.2)	782 (52.0)	2634 (45.9)	687 (53.3)	2526 (46.2)
Video	26 (0.4)	16 (1.1)	26 (0.5)	15 (1.2)	26 (0.5)
Phone	-	-	-	-	-

aSelf-identified veteran race/ethnicity.

Table 2. Demographics and clinic variables associated with VVC and phone telehealth utilization among diabetic veterans experiencing housing instability in pandemic year 1.

Characteristic	VVC use, PY1 N= 1437 Odds ratio (95% CI)	p-Value	Phone use, PY1 N= 5517 Odds ratio (95% CI)	p-Value
Age	0.97 (0.97–0.98)	<0.0001	1.01 (0.99–1.02)	0.31
Race/ethnicity ^a				
Non-Hispanic, White	Ref.		Ref.	
Non-Hispanic Black	1.01 (0.86–1.19)	0.89	1.58 (0.97–2.59)	0.07
Non-Hispanic Other	0.92 (0.59–1.42)	0.69	1.59 (0.49–5.17)	0.45
Hispanic, any race	1.44 (1.07–1.93)	0.02	1.79 (0.64–5.05)	0.27
Sex				
Male	Ref.		Ref.	
Female	1.63 (1.31–2.04)	<0.0001	1.22 (0.58–2.57)	0.60
Marital status				
Single/Divorced/Widow(ed)	Ref.		Ref.	
Married	1.39 (1.21–1.59)	<0.0001	0.68 (0.47–0.97)	0.03
Last residence				
Urban	Ref.		Ref.	
Rural	0.85 (0.73–1.00)	0.04	1.07 (0.72–1.60)	0.74
Elixhauser group (Pre-COVID)				
Less than or equal to 2	Ref.		Ref.	
3 or more	0.97 (0.79–1.19)	0.78	1.67 (1.23–2.47)	0.01
VA priority				
Group 1	1.21 (1.06–1.38)	0.004	1.10 (0.76–1.60)	0.60
Groups 2, 3, 4, 5, 6	Ref.		Ref.	
Prior utilization, pre-COVID				
Primary care				
Face to face	2.10 (1.47–3.00)	<0.0001	2.03 (1.27–3.26)	0.003
Video	1.98 (1.23–3.19)	0.005	–	
Phone	1.21 (1.05–1.40)	0.01	2.35 (1.61–3.44)	<.0001

(continued)

Table 2. Continued.

Characteristic	VVC use, PY1 N= 1437	p-Value	Phone use, PY1 N= 5517	p-Value
	Odds ratio (95% CI)		Odds ratio (95% CI)	
Endocrinology face to face	1.56 (1.33–1.83)	<0.0001	–	
Pharmacy				
Face to face	–		1.89 (1.21–2.96)	0.005
Video	3.24 (1.39–7.55)	0.006	–	
Mental health visit	1.21 (1.03–1.43)	0.02	–	
Pulmonary clinic visit	1.27 (1.09–1.49)	0.003	–	
Cardiology clinic visit	1.24 (1.07–1.43)	0.004	3.12 (1.61–6.07)	0.0008
Pain clinic visit	1.35 (1.16–1.58)	0.0001	5.79 (1.82–18.43)	0.003
Telehealth visit	1.41 (1.21–1.65)	<0.0001	–	

^aSelf-identified veteran race/ethnicity.

The fifteen veteran participants were more likely to be ≥ 55 years in age (87%); male (80%); self-identified non-Hispanic, White in race (53%); divorced, separated (53%); 60% had a current “place to stay” (40% unstable housing; 13% homelessness); and were living in an urban location (60%; rural, 20%; suburban, 20%).

Qualitative analysis revealed four common themes reported by both VA providers and veterans, including: (a) overall VCC experience; (b) internet service and technology barriers; (c) the need for VVC education; and (d) variation by specialty and care type. Each of these themes will be discussed in more detail in the subsequent paragraphs.

Veteran and provider VVC experience

The majority of veterans and providers reported satisfaction with using VVC for diabetes care. Across VA clinics, when providers utilized VVC, they claimed that they were able to learn more about their patient’s environment.

“Many veterans are in their room or a small space ...the whole place is in such disarray that I just don’t know how they even get through on a daily basis.” (Endocrinologist)

Veterans voiced confidence that VVC is a convenient way to stay connected to care, particularly since transportation was also barrier to care during the pandemic. Most veterans reported residing in a location that was more than 30 minutes from a VA clinic. Therefore, veterans appreciated the opportunity to be seen by a provider without having to go into the clinic.

“One nice thing about doing VVC through your iPhone is you don’t need to worry about getting a ride to the clinic.” (Veteran, male)

A few veterans reported being very satisfied with the VVC encounters and preferred to continue with the visits post-pandemic.

“I’d prefer to keep that [VVC] going for anything that I don’t need to be physically present for... I’m just as good over the laptop.” (Veteran, male)

Respondents reported that VVC satisfaction was driven by care availability and convenience. Providers mentioned the importance of VVC in facilitating routine patient care during a time of isolation. Veterans reported using VVC because it was a safe care option. Providers reported that they appreciated that VVC provided an option to connect with patients when in-person visits were not an option.

“It doesn’t surprise me that patients, whether they’re homeless or not, were happy to connect...and were not so worried about the mode as they were lonely and needed care.” (Diabetes Nurse, female)

Internet service and technology barriers

Lack of technology and internet service were the predominant barriers to VVC for providers and veterans. Forty percent of the veterans interviewed did not have a

Table 3. Demographics and clinic variables associated with VVC and phone telehealth utilization among diabetic veterans experiencing housing instability in pandemic year 2.

Characteristic	VVC use, PY2 N= 1075		Phone use, PY2 N= 5246	
	Odds ratio (95% CI)	p-Value	Odds ratio (95% CI)	p-Value
Age	0.97 (0.96–0.98)	<0.0001	0.99 (0.98–0.99)	0.02
Race/ethnicity ^a				
Non-Hispanic White	Ref.		Ref.	
Non-Hispanic Black	1.01 (0.84–1.23)	0.35	1.32 (0.97–1.78)	0.07
Non-Hispanic Other	0.96 (0.58–1.60)	0.84	1.23 (0.58–2.60)	0.59
Hispanic, any race	0.75 (0.51–1.09)	0.17	1.37 (0.72–2.61)	0.33
Sex				
Male	Ref.		Ref.	
Female	1.59 (1.22–2.08)	0.0006	1.13 (0.71–1.79)	0.62
Marital status				
Single/divorced/widow(ed)	Ref.		Ref.	
Married	1.28 (1.09–1.52)	0.003	0.96 (0.76–1.22)	0.75
Last residence				
Urban	Ref.		Ref.	
Rural	0.89 (0.74–1.08)	0.23	0.76 (0.60–0.97)	0.03
Elixhauser group, Pre-COVID				
Less than or equal to 2	Ref.		Ref.	
3 or more	1.33 (0.98–1.80)	0.07	0.89 (0.67–1.19)	0.44
Elixhauser group, PY1				
Less than or equal to 2	Ref.		Ref.	
3 or more	1.64 (1.25–2.14)	0.0004	1.25 (0.95–1.64)	0.11
VA priority				
Group 1	1.28 (1.09–1.49)	0.002	1.28 (1.02–1.62)	0.03
Groups 2, 3, 4, 5, 6	Ref.		Ref.	
A1c levels pre-diabetic range, Pre-COVID	0.84 (0.71–0.99)	0.04	-	

(continued)

Table 3. Continued.

Characteristic	VVC use, PY2 N= 1075		Phone use, PY2 N= 5246	
	Odds ratio (95% CI)	p-Value	Odds ratio (95% CI)	p-Value
Prior utilization				
Primary care, pre-COVID phone	-		1.41 (1.12-1.76)	0.003
Primary care, PY1				
Face to face	1.33 (1.04-1.71)	0.02	-	
Video	2.49 (2.08-2.98)	<0.0001	-	
Phone	0.67 (0.49-0.92)	0.01	2.44 (1.81-3.28)	<0.0001
Endocrinology, PY1 video	3.04 (2.08-4.45)	<0.0001	-	
Pharmacy, PY1				
Face to face	-		1.51 (1.18-1.94)	0.001
Video	6.24 (4.11-9.48)	<0.0001	-	
Telehealth visit, pre-COVID	1.26 (1.06-1.50)	0.009	-	
Telehealth visit, PY1	1.86 (1.53-2.25)	<0.0001	1.62 (1.28-2.05)	<0.0001
Surgery visit, pre-COVID	1.20 (1.00-1.42)	0.05	-	
Mental health visit, PY1	0.81 (0.67-0.98)	0.03	-	
Cardiology clinic visit, PY1	1.22 (1.03-1.44)	0.02	1.65 (1.20-2.27)	0.002
Pulmonary clinic visit, PY1	-		1.49 (1.03-2.16)	0.03
Primary care visit, PY1	-		1.88 (1.41-2.49)	<0.0001
Diabetes visit, PY1	-		2.27 (1.44-3.58)	0.0004

^aSelf-identified veteran race/ethnicity.

smartphone or tablet to utilize for VVC care. Veterans also commented that internet services were often not available at the time of care resulting an access barrier.

“Video depends on where you are at the time...most of the time when I’m talking with Dr [name] it fades in and out, we lose each other... I have to go somewhere with good Wi-Fi.” (Veteran, female)

The majority of providers mentioned that having Information Technology (IT) support on site influenced staff willingness to use VVC with their patients. IT support was often needed in

the first month of the pandemic to ensure providers and/or veterans were able to connect to the encounter.

“The days when VVC wasn’t working, we could still do a phone call ... but someone to assist [with VVC] is helpful.” (Pharmacist, female)

VVC education

VVC training would increase utilization among providers and veterans. Providers reported variability in VVC

knowledge and training, resulting to varying levels of uptake. Onsite technicians and trainings to prepare providers for VVC encounters were essential to utilization.

“Prior [to COVID] I really hadn’t pursued [VVC] very much...but after the training was done, going through the good and bad of learning... when technology works, it’s great.” (Primary Care Clinician, male)

Providers mentioned that test sessions prior to the veteran’s visit aided in the implementation of VVC encounter(s). Providers discussed how important these trial sessions were to VVC efficiency.

[After the training] “I had an 80-year-old figure out how to get on the video call and it was a success.” (Diabetes Nurse, female)

Many veterans reported needing help to log onto a VVC appointment, and over a quarter of individuals reported “not knowing what to do” if the video dropped.

“I probably would need some help with telehealth; I’m not very tech savvy... when I was in college, the technology we had was an ink pen, a calculator, and a jacket.” (Veteran, male)

Veterans reported frustration with providers who were not familiar with VVC. One veteran underscored the bidirectional need for education to successfully implement VVC encounters.

“Not everybody at the other end knows how to use technology...especially with some of the older doctors, teach them to keep up with the technology.” (Veteran, male)

Utilization variation: care and specialty type

Providers reported using VVC for follow-up or screening appointments while urgent acute care remained in-person during the pandemic. One provider described how she was able to safely continue the majority of clinical services through VVC during the pandemic.

“I did things by VVC that I did not think could happen... sensor trainings, pump trainings, blood glucose testing... never, never in a million years would I have thought that those things could happen without having somebody right next to me.” (Diabetes Nurse Educator, female)

Other specialties, such as optometry, podiatry, and audiology, were very limited in the care they could provide through VVC. While VVC provided a convenient option for clinical encounters, the lack of diagnostic tools that could be used for virtual encounters limited some specialty care use.

“Other than doing basic follow-up it is hard to treat patients in optometry by telehealth...we just don’t have the right equipment, but it is possible [with the right equipment].” (Optometrist, female)

Discussion

Among veterans with diabetes experiencing HI during the pandemic, those who utilized VVC were more likely to be female, younger, self-identified Hispanic, married, and residing in an urban dwelling. Need characteristics included having prior VVC visits, more comorbidities, and VA priority group 1, compared to their counterparts. In the qualitative study, many participants reported that VVC is an effective and convenient mode to conduct clinical encounters, especially when in-person visits are not safe. Veterans using VVC were satisfied with their encounters but also underscored barriers to overcome, including access to stable internet service and devices such as smartphones, and VVC training.

The present study’s findings may reflect healthcare-seeking traits of the broader veteran population, in which females are more likely to engage in virtual visits.^{24,25} Studies have identified several factors driving VVC utilization among female veterans including higher technology comfort level, experience with telehealth, and convenience.^{26,27} VVC utilization was also more common among married veterans with diabetes experiencing HI. Significant others may provide social support and encourage veterans’ engagement in VVC, although this relationship in the context of HI needs further research.^{24,28,29}

We observed that VVC users experiencing HI were more likely to be self-identified Hispanic, while in-person visits were more common in their non-Hispanic, White counterparts. Racial and ethnic variations in VVC utilization shifted during the COVID pandemic; whether video utilization will persist among self-identified Hispanic veterans is unknown.^{23,30} In general, self-identified Hispanic patients tend to report lower rates of video care due to language barriers and lack of health insurance.³¹ Our findings corroborate recent studies showing that veterans who self-identify from racial/ethnic minority backgrounds used VVC more than self-identified non-Hispanic White veterans during the pandemic.^{8,32} This difference in VVC utilization may be reflective of self-identified Hispanic veterans’ access to VA healthcare compared to the general population in which a higher proportion of self-identified Hispanics are uninsured.^{31,33} Additionally, English proficiency is needed for U.S. military service, so language may be less of a VVC barrier for veterans compared to civilians.³² Future research should examine VVC uptake among all self-identified Hispanic veterans, not just those experiencing HI.

Age is inversely associated with VVC utilization among veterans with diabetes experiencing HI. Older patients are less likely to participate in VVC, and the reasons are

multifactorial including inexperience and lack of technology.^{34,35} Still, VA providers and veterans with diabetes reported willingness to use VVC when equipped with adequate resources and training, despite age.

Aligned with previous literature, our veterans reported that VVC reduces the travel distance, time, and costs to access their local VA.^{36,37}

The VA has made large efforts to mitigate the digital divide among racial/ethnic, geographical, and age diverse veteran populations,^{38,39} but we found some of our participants voiced not knowing how to access technology resources. Prior to the pandemic, the VA expanded its video-enabled tablet program to a cell phone option available to eligible veterans with clinical access barriers.^{36,40} Recent trends demonstrate that following the tablet or cell-phone receipt, veterans' engagement in VVC increased across VA clinical settings.⁴⁰ These trends coincide with previous work which reported that veterans experiencing HI welcome technology-based care to improve access to services.^{41,42} Our interviews report that many veterans with HI do not know how to access VA digital technology. While the VA is committed to increasing virtual care for all veterans, our data suggests that initiatives which aim to increase veterans' technology access and digital education are still needed.

While this study contributes to the current literature and our understanding of VVC utilization among veterans experiencing HI, there are limitations. First, our analysis was limited to veterans with diabetes in New England, so our findings may not be generalizable outside of this geographical region. Second, analyses were limited to veterans who had a care encounter between March 1, 2019, and March 1, 2022; therefore, it is possible that we may not have captured changes in use, clinical modality, and experiences among all veterans with diabetes experiencing HI during the pandemic. Third, interviews have the potential for recall and social desirability bias, although most interviews took place within the same month as the VVC visit. Furthermore, our research team has a strong track-record in maintaining the confidentiality and anonymity of qualitative responses, and querying with neutral questions in which the interviewee has been instructed that there are no right or wrong answers. Notwithstanding, this study uncovered some new characteristics of veterans with diabetes with HI who are willing to engage in VVC encounters and provided some insights into how to effectively engage this population in future utilization.

Conclusion

This study examined health care utilization and its modality among veterans with diabetes with housing instability and found that VVC played an important role in care access during the pandemic. Employing both quantitative and qualitative methods, we identified some facilitators and

barriers to VVC utilization among providers and veterans with diabetes. A comprehensive knowledge of VVC utilization patterns by veteran characteristics may inform tailored interventions that seek to close identified disparities gaps. While convenience and safety were facilitators of VVC care, enhanced technology and education are essential to future uptake and to mitigating the digital divide. Future work should assess how best to integrate VVC into routine diabetes care as in-person care is restored to ensure equitable access to care.

Acknowledgement: The authors would like to acknowledge the National Center on Homelessness Among Veterans (NCHAV) which made this project possible.

Contributorship: RK, AA, LC, and DM researched literature and conceived the study. All authors were involved in protocol development, gaining ethical approval, patient recruitment and data analysis. RK wrote the first draft of the manuscript. All authors reviewed and edited the manuscript and approved the final version of the manuscript.

Data availability: The data that support the findings of this study are available from the corresponding author, RLK, upon reasonable request.

Declaration of conflicting interests: The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval: This National Center on Homelessness for Veterans (NCHAV) Quality Improvement project was deemed exempt from oversight by VA Central Western Massachusetts ethical review board.

Funding: This work was supported with an intramural grant from the National Center on Homelessness among Veterans.

Guarantor: RK will take full responsibility for the article, including for the accuracy and appropriateness of the reference list.

ORCID iDs: Rebecca L Kinney  <https://orcid.org/0000-0001-6752-5715>

Laurel A Copeland  <https://orcid.org/0000-0002-9478-0209>

Supplemental material: Supplemental material for this article is available online.

References

1. Veteran Healthcare Administration. VA's telehealth program is already the largest in the nation. It's about to get bigger. Accessed September 1, 2023, <https://federalnewsnetwork.com>.

- com/veterans-affairs/2018/12/vas-telehealth-program-is-already-the-largest-in-the-nation-its-about-to-get-bigger/
2. Veterans Healthcare Administration. VA Telehealth Services. Accessed: September 15, 2023 <https://telehealth.va.gov/>
 3. Stroupe KT, Martinez R, Hogan TP, et al. Health care utilization and costs of veterans evaluated for traumatic brain injury through telehealth. *Telemed J E Health* 2019; 25: 1144–1153.
 4. Adams SV, Mader MJ, Bollinger MJ, et al. Utilization of interactive clinical video telemedicine by rural and urban veterans in the veterans health administration health care system. *J Rural Health* 2019; 35: 308–318.
 5. Garvin LA, Hu J, Slightam C, et al. Use of video telehealth tablets to increase access for veterans experiencing homelessness. *J Gen Intern Med* 2021; 36: 2274–2282.
 6. Hare N, Bansal P, Bajowala SS, et al. Work group report: COVID-19: unmasking telemedicine. *J Allergy Clin Immunol Pract* 2020; 8: 2461–2473.e3.
 7. Shaker MS, Oppenheimer J, Grayson M, et al. COVID-19: pandemic contingency planning for the allergy and immunology clinic. *J Allergy Clin Immunol Pract* 2020; 8: 1477–1488.e5.
 8. Heyworth L, Kirsh S, Zulman D, et al. Expanding access through virtual care: the VA's early experience with COVID-19. *NEJM Catalyst in Care Delivery* 2020.
 9. Abrams EM and Szeffler SJ. COVID-19 and the impact of social determinants of health. *Lancet Respir Med* 2020; 8: 659–661.
 10. Perry BL, Aronson B and Pescosolido BA. Pandemic precarity: COVID-19 is exposing and exacerbating inequalities in the American heartland. *Proc Natl Acad Sci U S A* 2021; 118: e2020685118.
 11. Tsai J and Rosenheck RA. Risk factors for homelessness among US veterans. *Epidemiol Rev* 2015; 37: 177–195.
 12. Beran D, Perone SA, Castellsague Perolini M, et al. Beyond the virus: ensuring continuity of care for people with diabetes during COVID-19. *Prim Care Diabetes* 2021; 15: 16–17.
 13. Hartmann-Boyce J, Morris E, Goyder C, et al. Diabetes care. Diabetes and COVID-19: risks, management, and learnings from other national disasters. *Diabetes Care* 2020; 43: 1695–1703.
 14. Tsai J, Szymkowiak D and Jutkowitz E. Developing an operational definition of housing instability and homelessness in veterans health administration's medical records. *PLoS One* 2022; 17: e0279973.
 15. Price LE, Shea K and Gephart S. The veterans affairs's corporate data warehouse: uses and implications for nursing research and practice. *Nurs Adm Q.* 2015; 39: 311–318.
 16. Meditz RW, Manberg CL and Rosner F. Improving access to a primary care medical clinic. *J Natl Med Assoc* 1992; 84: 361–364.
 17. Elixhauser A, Steiner C, Harris DR, et al. Comorbidity measures for use with administrative data. *Med Care* 1998; 36: 8–27.
 18. Weller SC, Vickers B, Bernard HR, et al. Open-ended interview questions and saturation. *PLoS One* 2018; 13: e0198606.
 19. Scientific Software Development. *Atlas.ti, Qualitative Data Analysis Software*. 2020 <https://atlasti.com/support/support-center/>
 20. Kinney RL, Haskell S, Relyea MR, et al. Coordinating women's preventive health care for rural veterans. *J Rural Health* 2022; 38: 630–638.
 21. Nowell B. *Public Administration Research and Theory*, 2019.
 22. Wagner C, Dichter ME and Mattocks K. Women Veterans' pathways to and perspectives on veterans affairs health care. *Womens Health Issues* 2015; 25: 658–665.
 23. Cordasco KM, Yuan AH, Rollman JE, et al. Veterans' use of telehealth for veterans health administration community care urgent care during the early COVID-19 pandemic. *Med Care* 2022; 60: 860–867.
 24. Darrat I, Tam S, Boulis M, et al. Socioeconomic disparities in patient use of telehealth during the coronavirus disease 2019 surge. *JAMA Otolaryngol Head Neck Surg* 2021; 147: 287–295.
 25. Lindsay JA, Caloudas A, Hogan J, et al. Getting connected: a retrospective cohort investigation of video-to-home telehealth for mental health care utilization among women veterans. *J Gen Intern Med* 2022; 37: 778–785.
 26. Moreau JL, Cordasco KM, Young AS, et al. The use of telemental health to meet the mental health needs of women using department of veterans affairs services. *Womens Health Issues* 2018; 28: 181–187.
 27. Goldstein KM, Zullig LL, Dedert EA, et al. Telehealth interventions designed for women: an evidence map. *J Gen Intern Med* 2018; 33: 2191–2200.
 28. Pandey KR, Yang F, Cagney KA, et al. The impact of marital status on health care utilization among medicare beneficiaries. *Medicine (Baltimore)* 2019;98:e14871.
 29. Mohammadi I, Wu H, Turkcan A, et al. Data analytics and modeling for appointment no-show in community health centers. *J Prim Care Community Health* 2018; 9: 215013271881169.
 30. Ferguson JM, Wray CM, Jacobs J, et al. Variation in initial and continued use of primary, mental health, and specialty video care among veterans. *Health Serv Res* 2023; 58: 402–414.
 31. Ryskina KL, Shultz K, Zhou Y, et al. Older adults' access to primary care: gender, racial, and ethnic disparities in telemedicine. *J Am Geriatr Soc* 2021; 69: 2732–2740.
 32. Guajardo E, Amspoker AB, Stanley MA, et al. Patterns of telehealth use for mental health treatment among hispanic veterans. *Telemed J E Health* 2023; 29: 788–792.
 33. Tai DBG, Sia IG, Doubeni CA, et al. Disproportionate impact of COVID-19 on racial and ethnic minority groups in the United States: a 2021 update. *J Racial Ethn Health Disparities* 2022; 9: 2334–2339.
 34. Hise N, Buckner J, Ince S, et al. Telemedicine at the VA: examining smartphone connectivity rates to VA video connect and doximity dialer video. *Ann Vasc Surg Brief Rep Innov* 2022; 2: 100147.
 35. Tsai J and Rosenheck RA. Use of the internet and an online personal health record system by US veterans: comparison of veterans affairs mental health service users and other veterans nationally. *J Am Med Inform Assoc* 2012; 19: 1089–1094.
 36. Slightam C, Gregory AJ, Hu J, et al. Patient perceptions of video visits using veterans affairs telehealth tablets: survey study. *J Med Internet Res* 2020; 22: e15682.
 37. Gordon HS, Solanki P, Bokhour BG, et al. "I'm not feeling like I'm part of the conversation" patients' perspectives on

communicating in clinical video telehealth visits. *J Gen Intern Med* 2020; 35: 1751–1758..

38. SY L. Smartphones prove a smart solution to combatting homelessness. October 10, 2023 <https://news.va.gov/111139/smartphones-solution-combattinghomelessness/#:~:text=Initially%2C%20the%20smartphones%20were%20provided,were%20considered%20%E2%80%9Ccleaner%20devices.%E2%80%9D>.
39. Zulman DM, Wong EP, Slightam C, et al. Making connections: nationwide implementation of video telehealth tablets to address access barriers in veterans. *JAMIA Open* 2019; 2: 323–329.
40. Wray CM, Van Campen J, Hu J, et al. Crossing the digital divide: a veteran affairs program to distribute video-enabled devices to patients in a supportive housing program. *JAMIA Open* 2022; 5: ooac027.
41. McInnes DK, Sawh L, Petrakis BA, et al. The potential for health-related uses of mobile phones and internet with homeless veterans: results from a multisite survey. *Telemed J E Health* 2014; 20: 801–809.
42. McInnes DK, Solomon JL, Shimada SL, et al. Development and evaluation of an internet and personal health record training program for low-income patients with HIV or hepatitis C. *Med Care* 2013; 51: S62–S66.

Appendix A: VA specialty care telehealth

Telehealth enables you to connect to the quality care you need in more than 50 specialty areas from your local VA clinic.

Clinical specialty

TeleCardiology	TeleChaplain	TeleDentistry
TeleDermatology	TeleEyeCare	TeleGenomics
TeleGIHepatology	TeleHematology	TeleICU
TeleInfectious Disease	TeleMental Health	TeleMOVE!
TeleNephrology	TeleNeurology	TeleNutrition
TelePathology	TelePharmacy	TelePodiatry

(continued)

TelePrimary Care	TelePulmonology	TeleRheumatology
TeleSCI/D	TeleSpirometry	TeleStroke
TeleSurgery	TeleTransplant	TeleWholeHealth
TeleWound Care	Virtual PACT	Womens Health

TeleRehabilitation

TeleAmputation	Telehealth Assistive Technology
TeleBlind Rehabilitation	TeleChiropractic Care
TeleKinesiotherapy	TeleOccupational Therapy TeleOrthotic Prosthetic Care
TelePhysical Medicine Physicia	TelePhysical Therapy
TeleProsthetic & Sensory Aids Service	TeleRecreational and Creative Arts Therapy
TeleSpeech Patholog	

Source: VA telehealth: In the Clinic | Telehealth VA, 2024.

Appendix B: List of acronyms

Acronym definition

CDW	Corporate Data Warehouse
COVID-19	Coronavirus Disease-2019
DSS	Decision Support System
HER	Electronic Health Record
HI	Housing Instability
IT	Information Technology
NCHAV	National Center on Homelessness for Veterans
NPCD	National Patient Care Database
PY	Pandemic Year
VA	Veterans Affairs
VHA	Veterans Health Administration
VISN	Veterans Integrated Services Network:
VVC	VA Video Connect