

Heparin stability: effects of diluent, heparin activity, container, and pH

KENNETH T GOODALL, CHEE-CHOONG CHOOI, AND
ALEXANDER S GALLUS

From the Department of Haematology, Flinders Medical Centre, Bedford Park, South Australia 5042, Australia

SUMMARY The effects have been studied of diluent, heparin activity after dilution, container, and pH on the stability of heparin solutions stored under conditions resembling those present during heparin infusion by intravenous drip or syringe pump. Heparin activity was measured by activated partial thromboplastin time and thrombin clotting time (and, in one set of studies, also by factor Xa inhibitor assay and protamine sulphate neutralisation).

Heparin activity was stable for 6 hours regardless of storage conditions. After 24 hours heparin activity was stable when the drug was diluted in 0.9% saline and stored in plastic, but a small loss of activity was observed in several studies after dilution in 5% dextrose or storage in glass. A more extensive comparison confirmed a 3-5% loss in heparin activity over 24 hours after dilution in 5% dextrose. Changing the pH to 3.5 or 10.0 had little effect on storage stability.

We conclude that heparin activity *in vitro* remains stable during short infusions but recommend dilution in 0.9% saline and a plastic container when a heparin solution is infused over 24 hours.

Heparin is now commonly given by continuous intravenous infusion rather than by intermittent injection. For this mode of administration heparin is diluted in one of several intravenous infusion fluids and may then remain at room temperature in the infusion bag, bottle, or syringe for 8-24 hours. It is, therefore, important to know if its anticoagulant activity remains stable *in vitro* for this time.

This question has been repeatedly investigated with varying results (Table). Thus, it has been reported that heparin activity is constant for 24 hours regardless of the diluting fluid or its pH over a wide range,¹⁻⁴ or that heparin activity is reduced by 40-50% within 1 hour of dilution except in 0.9% sodium chloride,⁵ or that heparin activity is reduced by 50-60% after 4-6 hours in all diluents but then increases again with further storage.⁶

Because of these contradictory and sometimes confusing reports about heparin stability *in vitro* we decided to re-investigate the question.

Further, because there is disagreement about the most appropriate test of heparin effect, we decided to measure heparin stability *in vitro* using four tests: the

activated partial thromboplastin time (APTT), the thrombin clotting time (TCT), factor Xa inhibitory activity (Xa time), and protamine sulphate neutralisation.

Methods

STUDY DESIGN

Sodium heparin was added to 500 ml 0.9% saline or 5% dextrose, supplied commercially for intravenous use in plastic bags or glass bottles, to give a heparin activity of 10, 20, or 40 IU/ml, and to 50 ml 0.9% saline or 5% dextrose drawn into a plastic syringe to give a heparin activity of 400 or 1000 IU/ml. These conditions were chosen to mimic those obtaining during heparin infusion by intravenous drip or syringe pump. Heparin activity was assayed immediately and after 3, 6, and 24 hours of room temperature storage. In addition, heparin stability was measured during storage at varying pH.

HEPARIN AND INTRAVENOUS FLUIDS

Mucous heparin BP (1000 or 5000 IU/ml) was obtained from Allen and Hanbury's Division of Glaxo Australia Pty Ltd (Boronia, Victoria). Intra-

Previously reported changes in heparin activity after dilution and storage

Authors	Heparin activity (U/ml)	Storage (hours)	Diluting fluid	pH	Assay	Conclusions
Stock and Warner (1971) ¹	964 200 40	24	5% Dextrose	7.0 6.8 6.6	Not described	No loss of activity
Hodby <i>et al.</i> (1972) ²	32	24	0.9% Sodium chloride 5% Dextrose Dextrose saline 0.9% Sodium chloride 0.9% Sodium chloride	6.9 7.2 6.9 3.2 6.9	APTT	No loss of activity
Mitchell <i>et al.</i> (1976) ³	20, 40	48	Normal saline 5% Dextrose	5.2-7.0 5.5-6.8	TCT	No loss of activity
Joy <i>et al.</i> (1979) ⁴	20	24	5% Dextrose 0.45% Saline 5% Dextrose	2.0-9.0	TCT	No loss of activity
Jacobs <i>et al.</i> (1973) ⁵	20	24	Normal saline 5% Dextrose Saline/Dextrose Ringer lactate Saline lactate	6.8 3.9 3.9 6.3 6.3	Modified TCT	No loss of activity ≈ 50% loss of activity ≈ 40% loss of activity ≈ 50% loss of activity ≈ 50% loss of activity
Okuno and Nelson (1975) ⁶	10	24	Normal saline 5% Dextrose 10% Dextrose 2½% Dextrose Hartmann's solution	5.5-6.9	Xa inhibitor TCT	Marked loss of activity within 4 hours, then complete recovery by Xa inhibitor assay but not by TCT

venous fluids were obtained in plastic bags from Travenol Laboratories Pty Ltd (Sydney) and in glass bottles from Abbott Laboratories Pty Ltd (Sydney). Plastic syringes were supplied by Terumo Australia Pty Ltd (Melbourne).

In some experiments the pH of 0.9% saline and 5% dextrose was adjusted to 3.5 or 10.0 with 0.1 N HCl or 0.1 N NaOH before the addition of heparin.

HEPARIN ACTIVITY MEASUREMENTS

Heparin solutions were subsampled into pooled normal platelet poor plasma for the measurement of heparin activity after each storage interval. After a preliminary dilution, 10 or 20 μ l of heparin solution were added to 1.0 ml plasma to yield a calculated final plasma activity of 0.2 IU/ml (assuming no change in activity with storage). All tests were done in duplicate at least five times by two workers using coded samples. Samples were stored on melting ice until tests had been done.

To prepare the normal plasma, blood was collected from volunteer blood donors into 3.8% tri-sodium citrate (9 parts blood:1 volume citrate) and centrifuged at 2000 *g* for 30 minutes at 4°C. Pooled normal plasma from 20 donors was then stored in small aliquots at -70°C. The same batch of pooled normal plasma was used for each set of studies.

Heparin activity was measured with the APTT and the TCT in all studies, and, in one series of experiments, heparin activity was also measured by a factor Xa inhibitor method (Xa time)⁷ and by protamine sulphate neutralisation.⁸ All tests were done using a clot-detecting device (Clotek, Hyland), except for the

Xa time, which was measured using a manual (hook) technique.

The observed clotting times for each test were converted to heparin activity (IU/ml) by the use of standard heparin sensitivity curves for each test (see below, and Figs 1-3).

Automated APTT reagent was obtained from General Diagnostics (Morris Plains, New Jersey) and used according to the manufacturer's instructions. The TCT was measured after diluting Topical




Fig. 1 Heparin sensitivity of APTT. The mean APTT \pm SEM is plotted for each heparin activity tested. The line of best fit was derived by least squares analysis.




Fig. 2 Heparin sensitivity of TCT. The mean TCT \pm SEM is plotted for each activity tested.




Fig. 3 Heparin sensitivity of factor Xa inhibitor assay (XaCT). The mean XaCT \pm SEM is plotted for each activity tested.

Thrombin (Parke Davis, Detroit, Michigan) to an activity of 5 units/ml in 0.1 M calcium chloride and adding 0.1 ml of this reagent to 0.2 ml platelet poor plasma.

The Xa time was measured using the Deci Unit range of method A of Yin *et al.*⁷ Reagents for this test were obtained from Sigma Chemicals Co (St Louis, Mo).

Protamine sulphate neutralisation was done as described by Refn and Vestergaard⁸ using dilutions of protamine sulphate (10 mg/ml) supplied by Boots Co Ltd (Nottingham, England).

HEPARIN SENSITIVITY CURVES AND HEPARIN RECOVERY FROM PLASMA

The sensitivities of the APTT, TCT, and Xa CT to heparin are shown in Figures 1-3. These curves were derived by adding appropriate dilutions of the heparin batch used in the stability studies to pooled normal platelet poor plasma and measuring clotting times in duplicate on five separate occasions.

To check heparin recovery from the pooled normal plasma used in these studies, protamine sulphate was used to neutralise a crystalline porcine sodium mucous heparin (169 IU/mg) obtained by courtesy of Dr Peter Forsell, Allen and Hanbury's, Boronia, Victoria. 100 IU of this heparin was neutralised by 1.045 mg of protamine sulphate. When 0.2 IU/ml of this standard heparin was added to pooled normal plasma, 0.19 IU/ml could be measured by protamine sulphate neutralisation, indicating substantially complete recovery.

STATISTICAL METHODS

Lines of best fit for heparin activity-test response curves were derived by the method of least squares. Change of heparin activity with time was tested by paired *t* test and analysis of variance. All results in text and figures are expressed as mean \pm standard error of mean (SEM).

Results

HEPARIN RECOVERY

At zero time, all heparin solutions showed the predicted heparin activity with all test systems used, so that subsequent results were expressed as a percent of the initial measured activity.

PLASTIC CONTAINERS

There was no significant loss of activity when heparin was diluted in 0.9% saline and stored for 24 hours in plastic containers (bag or syringe) (Figs 4, 5, 7). This occurred regardless of the heparin dilution, the storage pH, and whether heparin activity was measured by APTT, TCT, or (with the 10 IU/ml




Fig. 4 Heparin stability tested by APTT, TCT, Xa time (Xa), and protamine sulphate neutralisation after heparin was diluted to 10 IU/ml and stored in commercial solutions of 0.9% saline (above) or 5% dextrose (below) sold for intravenous use in plastic bags. Bars represent percent of initial activity \pm SEM after 3, 6, and 24 hours' storage. ** $p < 0.01$ when activity after storage was compared with initial activity.




Fig. 5 Heparin stability tested by APTT (clear bars) and TCT (shaded bars) after dilution to 20 IU/ml, 40 IU/ml, 400 IU/ml, and 1000 IU/ml in 0.9% saline (above) or 5% dextrose (below), and storage for 3, 6, and 24 hours in plastic bags (20 IU/ml and 40 IU/ml) or plastic syringes. Bars represent percent of initial activity \pm SEM. * $p < 0.05$ ** $p < 0.01$ (compared with initial activity).

solution) by Xa time or protamine sulphate neutralisation. Similarly, there was at most a small loss of activity when heparin was diluted in 5% dextrose solution and kept in plastic containers. There was a statistically significant decrease in activity after 24 hours when the 10 IU/ml solution was tested by Xa time (Fig. 4), the 40 IU/ml solution was tested by TCT (Fig. 5), the 1000 IU/ml solution was tested by APTT (Fig. 5), and the pH 10, 20 IU/ml solution was tested by APTT (Fig. 7), but the apparent loss of activity was small: 9-11% in all cases except for a 23% loss of activity measured by Xa time.

GLASS BOTTLES

In several experiments a statistically significant 10-14% loss of activity was recorded after 24 hours' storage in glass bottles. This was seen when activity was measured by APTT (10 IU/ml, 0.9% saline) or TCT (40 IU/ml, 0.9% saline; 20 IU/ml, 40 IU/ml, 5% dextrose) (Fig. 6).

CONFIRMATORY STUDIES

In a second series of studies, 10 IU/ml heparin was stored for 24 hours in 20 500-ml plastic bags of 0.9% saline and 20 500-ml bags of 5% dextrose, and




Fig. 6 Heparin stability tested by APTT (clear bars) and TCT (shaded bars) after dilution to 10 IU/ml, 20 IU/ml, and 40 IU/ml in commercial solutions of 0.9% saline (above) or 5% dextrose (below) sold for intravenous use in glass bottles. Bars represent percent of initial activity \pm SEM after 3, 6, and 24 hours' storage. * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$ (compared with initial activity).

heparin activity was measured by APTT and TCT at zero time and after 24 hours.

There was no change of heparin activity in 0.9% saline (24-hour APTT = $99.9 \pm 0.6\%$; 24-hour TCT = $100.3 \pm 0.6\%$ of initial activity), but a very small though statistically significant decrease of heparin activity in 5% dextrose (24-hour APTT = $97.3 \pm 0.9\%$, 24-hour TCT = $95.1 \pm 1.3\%$ of initial activity; $P < 0.01$ for both observations).

Discussion

Our results suggest that there is no loss of heparin activity when heparin is diluted in 0.9% saline and stored in plastic for 24 hours at room temperature. Heparin also appears to be stable for 6 hours under other storage conditions, but a small loss was observed in some studies after 24 hours when the drug was diluted in 5% dextrose or stored in glass; this loss was no greater than 10-15% of initial activity. Further studies comparing heparin stability in 0.9% saline and 5% dextrose in plastic bags, using a larger sample of 20, confirmed that heparin is stable for 24 hours in 0.9% saline, but showed a 3-5% loss of activity in 5% dextrose. Any loss of activity associated with storage at the pH extremes




Fig. 7 Effect of pH 3.5 and pH 10 on heparin stability tested by APTT (clear bars) and TCT (shaded bars) 6 and 24 hours after dilution of heparin to 20 IU/ml in 0.9% saline (above) or 5% dextrose (below) sold for intravenous use in plastic bags. Bars represent percent of initial activity \pm SEM * $P < 0.05$.

of 3.5 and 10.0 was also small.

These conclusions are similar to those of most previous reports¹⁻⁴ and are consistent with the stable anticoagulant response to a 12-hour infusion of heparin diluted in 5% dextrose observed by Chessels *et al.*⁹

Our results do not confirm the marked loss of heparin activity reported after dilution in 5% dextrose by Jacobs *et al.*,⁵ and after dilution in 5% dextrose or 0.9% saline by Okuno and Nelson⁶ (Table), and we are unable to explain these discrepancies. Our methods of measuring heparin activity differ from those used by Jacobs *et al.*⁵ but are similar to those of Okuno and Nelson.⁶

We conclude that heparin activity *in vitro* remains stable during a short infusion, but, because of the small loss in activity after 24 hours when heparin is diluted in 5% dextrose or stored in glass, we recommend dilution in 0.9% saline and storage in plastic containers when a heparin solution is to be infused over 24 hours.

We thank Miss J Tillett for able technical help, Dr Peter Forsell, of Allen and Hanbury's, for supplying crystalline heparin, and Mrs Jenny Riddle and Mrs Sheila Phillips for typing the manuscript.

References

- ¹ Stock SL, Warner N. Heparin in acid solution. *Br Med J* 1971;**3**:307.
- ² Hodby ED, Hirsh J, Adeniyi-Jones C. The influence of drugs upon the anticoagulant activity of heparin. *Can Med Assoc J* 1972;**106**:562-4.
- ³ Mitchell JF, Barger RC, Cantwell L. Heparin stability in 5% dextrose and 0.9% sodium chloride solutions. *Am J Hosp Pharm* 1976;**33**:540-2.
- ⁴ Joy RT, Hyneck ML, Bernardi RR, Ho NFH. Effect of pH on the stability of heparin in 5% dextrose solution. *Am J Hosp Pharm* 1979;**36**:618-21.
- ⁵ Jacobs J, Kletter D, Superstine E, Hill KR, Lynn B, Webb RA. Intravenous infusions of heparin and penicillins. *J Clin Pathol* 1973;**26**:742-6.
- ⁶ Okuno T, Nelson CA. Anticoagulant activity of heparin in intravenous fluids. *J Clin Pathol* 1975;**28**:494-7.
- ⁷ Yin ET, Wessler S, Butler JV. Plasma heparin: a unique, practical submicrogram-sensitive assay. *J Lab Clin Med* 1973;**81**:298-310.
- ⁸ Refn I, Vestergaard L. The titration of heparin with protamine. *Scand J Clin Lab Invest* 1954;**6**:284-7.
- ⁹ Chessels JM, Braithwaite TA, Chamberlain DA. Dextrose and sorbitol as diluent for continuous intravenous heparin infusion. *Br Med J* 1972;**2**:81-2.

Requests for reprints to: Dr AS Gallus, Department of Haematology, Flinders Medical Centre, Bedford Park, South Australia 5042.

Reports and Bulletins prepared by the Association of Clinical Biochemists

The following reports and bulletins are published by the Association of Clinical Biochemists. They may be obtained from The Publishing Department, British Medical Journal (ACB Technical Bulletins), BMA House, Tavistock Square, London WC1H 9JR. Overseas readers should remit by British Postal or Money Order.

SCIENTIFIC REVIEWS (price £1.00/\$2.00 each)

- 1** The assessment of thyroid function March 1971
FV FLYNN and JR HOBBS
- 2** Renal function tests suitable for clinical practice
January 1972 FL MITCHELL, N VEALL, and RWE WATTS
- 3** Biochemical tests for the assessment of fetoplacental
function May 1975 CE WILDE and RE OAKLEY
- 4** Test of exocrine pancreatic function March 1977
AH GOWENLOCK
- 5** Assay of cholinesterase in clinical chemistry March
1979 ELSIE SILK, J KING, and MARY WHITTAKER

TECHNICAL BULLETINS (price £1.00/\$2.00 each)

- 22** Bilirubin standards and the determination of bilirubin
by manual and technicon AutoAnalyzer methods
January 1971 BARBARA BILLING, RUTH HASLAM, and N
WALD
- 23** Interchangeable cells for spectrophotometers and
fluorimeters September 1971 SS BROWN and AH
GOWENLOCK
- 24** Simple tests to detect poisons March 1972 BW
MEADE *et al.*
- 25** Blood gas analysers May 1972 K DIXON
- 26** Kits for enzyme activity determination September
1972 SB ROSALKI and D TARLOW
- 27** Assessment of pumps suitable for incorporation into
existing continuous flow analytical systems November
1972 A FLECK *et al.*
- 28** Routine clinical measurements of transferrin in
human serum September 1973 K DIXON

- 29** Control materials for clinical biochemistry (5th
edition) September 1973 JF STEVENS
- 30** Notes on the quality of performance of serum
cholesterol assays September 1973 SS BROWN
- 31** Determination of uric acid in blood and in urine
July 1974 RWE WATTS
- 32** A survey of amino acid analysers readily available in
the United Kingdom September 1974 JE CARLYLE
and P PURKISS
- 33** Definitions of some words and terms used in auto-
mated analysis November 1974 A FLECK, R ROBINSON,
SS BROWN, and JR HOBBS
- 34** Measurement of albumin in the sera of patients
January 1975 LINDA SLATER, PM CARTER, and JR HOBBS
- 35** Investigation of the validity of temperature correction
factors for serum aspartate and alanine transaminases
March 1975 SB ROSALKI *et al.*
- 36** Factors influencing the assay of creatinine November
1975 JGH COOK
- 37** A survey of enzyme reaction rate analysers readily
available in the United Kingdom July 1977 RA
SAUNDERS and RF BURNS
- 38** Transport of specimens for clinical chemistry an-
alysis November 1977 P WILDING, JF ZILVA, and
CE WILDE
- 39** A scheme for the evaluation of diagnostic kits May
1978 PH LLOYD
- 40** A practical guide to gamma-counting in radio-
immunoassay January 1980 CE WILDE and D OTTEWELL
- 41** The use of biochemical tests [in the diagnosis of
disorders of calcium metabolism July 1980 ANGELA
FAIRNEY