



Research Article

Introducing palmfungi.org, an integrated fungal-host data platform

Yinru Xiong^{‡,§,||}, Manawasinghe S. Ishara^l, Kevin D. Hyde^{‡,§,¶,¶}, Joanne E. Taylor[¶], Alan Phillips[«], Diana Santos Pereira[«], Li Lu^{‡,§,»}, Sheng-Nan Zhang[^], Ausana Mapook[‡], Biao Xu^l

‡ Center of Excellence in Fungal Research, Chiang Rai, Thailand

§ School of Science, Mae Fah Luang University, Chiang Rai, Thailand

| Innovative Institute for Plant Health, Key Laboratory of Green Prevention and Control on Fruits and Vegetables in South China, Ministry of Agriculture and Rural Affairs, Zhongkai University of Agriculture and Engineering, Guangzhou, China

¶ CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Science, Kunming, China

Department of Botany and Microbiology, College of Science, Riyadh, Saudi Arabia

¶ Royal Botanic Garden Edinburgh, Edinburgh EH3 5LR, United Kingdom

« Biosystems and Integrative Sciences Institute, Faculdade de Ciências da Universidade de Lisboa, Lisbon, Portugal

» Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University, Qujing, China

^ School of Life Science and Technology, Center for Informational Biology, University of Electronic Science and Technology of China, Chengdu, China

Corresponding author: Manawasinghe S. Ishara (ishara9017@gmail.com),

Kevin D. Hyde (kdhhyde3@gmail.com)

Academic editor: Danny Haelewaters

Received: 30 Apr 2024 | Accepted: 20 Aug 2024 | Published: 02 Oct 2024

Citation: Xiong Y, Ishara MS, Hyde KD, Taylor JE, Phillips A, Pereira D, Lu L, Zhang S-N, Mapook A, Xu B (2024) Introducing palmfungi.org, an integrated fungal-host data platform. Biodiversity Data Journal 12: e126553. <https://doi.org/10.3897/BDJ.12.e126553>

Abstract

Palm fungi are a diverse and unique group mostly found on Arecaceae hosts. They have been studied for approximately 200 years resulting in a large number of known fungal species representing over 700 genera. The timeline of palm fungal studies could be roughly divided into three phases, based on the methods and frequency of reports. They are the “Historical palm fungi era”, “Classical palm fungi era” and “Molecular palm fungi era”. In the first two periods, the identification of palm fungi was based on morphology, which resulted in a considerable number of morphological species scattered across the data in books, monographs and papers. With the advancement of molecular techniques, studies on palm fungi accelerated. A large number of new species were introduced in the

molecular era, especially from Asia, including China and Thailand. However, there is a necessity to link these three generations of studies into a single platform combining data related to host factors, geography and utilisation. Herein, we introduce the palm fungi website: <https://palmfungi.org>, an integrated data platform for interactive retrieval, based on palm and fungal species. This website is not only a portal for the latest, comprehensive species information on palm fungi, but also provides a new platform for fungal researchers to explore the host-specificity of palm fungi. Additionally, this study uses palmfungi.org and related data to briefly discuss the current status of research on the distribution of palm fungi populations, showing how palmfungi.org links fungi with their palm hosts. Furthermore, the website will act as a platform for collaboration amongst taxonomists, plant pathologists, botanists, ecologists and those who are interested in palms and their relationship with ecological sustainability.

Keywords

Areaceae, Ascomycota, Basidiomycetes, website, fungal host interaction

Introduction

To document the diversity and distribution of different organismal taxa, it is important to set out recording and standardised monitoring programmes, often with the help of citizen scientists (Mueller et al. 2004, Halme et al. 2012, Haelewaters et al. 2024). fungi are an ancient and thriving group of organisms that play a vital role in natural ecosystems (Stubblefield et al. 1985, Taylor et al. 2014, Xie et al. 2023). As early as the beginning of the 20th century, evidence of fungi living in the Tertiary period was found in fossils and, since then, a large number have been documented (Kidston and Lang 1921, Edwards 1922). In the 21st century, the description of new species and recording of fungi have been occurring at a higher frequency (Hawksworth and Lücking 2017, Fisher et al. 2020). Over the centuries, mycologists have been working to study the diversity of fungi. The "*Sylloge Fungorum*" published by Pier Andrea Saccardo in 1882 stimulated extensive research and discussion on fungal diversity amongst modern mycologists (Bolman 2023). This discussion continues to this day, with Hawksworth (2001) suggesting that the number of fungi might be 1.5 million species, O'Brien et al. (2005) estimating there might be 3.5 to 5.1 million species, while Schmit and Mueller (2007) suggest that there may be as few as 712,000 species. According to a more recent study by Niskanen et al. (2023), the current estimates for global fungal richness ranges between 2 and 3 million species, with a "best guess" at 2.5 million of which only about 150,000 fungi have been described (Species Fungorum 2024; <http://www.speciesfungorum.org>). There are three major ecological strategies of fungi: saprotrophic, parasitic and mutualistic (Willis 2018). To adapt to different ecological setting, fungi have a variety of life modes, including endophytes, pathogens or saprobes and, depending on environmental conditions, they may be able to shift from one lifestyle to another, such as endophytes becoming pathogens (Hyde et al. 2014, Bhunjun et al. 2023). These diverse lifestyles allow fungi to have a wide range of hosts, from animals to plants. Amongst these, the relationship

between fungi and plants has been well studied (Senanayake et al. 2020, Jayawardena et al. 2021, Maharachchikumbura et al. 2021, Manawasinghe et al. 2021).

Arecaceae, which is commonly known as palms, ranks fifth in species richness amongst monocots families (Chen et al. 2022). According to APG IV (Angiosperm Phylogeny Group et al. 2016), Arecaceae contains five subfamilies, namely Arecoideae, Calamoideae, Ceroyloideae, Coryphoideae and Nypoideae. This family is one of the most morphologically diverse angiosperm groups, with 181 genera and approximately 2600 species (Dransfield et al. 2005, Dransfield et al. 2008, Baker and Dransfield 2016). To better record palm species data, Kissling et al. (2019) established the PalmTraits 1.0 database and a species information retrieval website palmweb (<https://palmweb.org>). An increasing number of palm fungi are also being continuously recorded and reported. As represented by monographs of Fröhlich and Hyde (2000), Hyde et al. (2000) and Taylor and Hyde (2003), a large number of palm fungi have been recorded, laying a solid foundation for subsequent research on fungal diversity on palm plants and exploring the diversity of the entire fungal population.

According to the database of Farr et al. (2021), there are about 9500 records of palm fungi, distributed within 99 palm genera. The diversity of palm fungi is extremely broad, covering almost all major fungal taxa (Taylor et al. 1999, Fröhlich and Hyde 2000). fungi on palms may be saprobes (Taylor et al. 1999, Fröhlich and Hyde 2000), pathogens (Douanla-Meli and Scharnhorst 2021, Xiong et al. 2022b), endophytes (Taylor et al. 1999) or epiphytes (Marasinghe et al. 2022, Marasinghe et al. 2023). Earlier focal studies by Pinruan et al. (2007) reported fungi on the *Licuala longicalycata*, while Pinnoi et al. (2006) reported those on the *Eleiodoxa conferta*, both from a peat swamp in southern Thailand. Recent introductions of new saprobic taxa on palms are by Konta et al. (Konta et al. 2020, Konta et al. 2021, Konta et al. 2023b) and Xiong et al. (2022a), who have expanded our understanding of palm-associated fungal diversity. Zheng et al. (2017), Pandian et al. (2021) and Xiong et al. (2022b) reported pathogenic fungi on palms and Guo et al. (1998) and Mahmoud et al. (Mahmoud et al. 2016, Mahmoud et al. 2017) reported endophytic fungi on palms. However, the research on palm fungi is scattered, lacking a comprehensive and up to date source of knowledge to precisely document them. Many asexual palm fungi have also not been linked to their sexual morphs (Hyde et al. 2011, Wijayawardene et al. 2017). To overcome these gaps, we proposed a website dedicated to palm fungi which will be an interactive platform for mycologists, as well the those who are interested in palms in general.

In recent years, web pages dedicated to fungal groups have become important resources to retrieve information. Other than the traditional fungal databases dedicated to fungal classification, several new databases were introduced as websites. Taxa depository databases include MycoBank (Crous et al. 2004) and Index Fungorum (Index Fungorum 2024, <https://www.indexfungorum.org>). Other generally important databases are Facesoffungi (Jayasiri et al. 2015, <https://www.facesoffungi.org>), Outlineoffungi (Wijayawardene et al. 2020, Wijayawardene et al. 2022b, <https://www.outlineoffungi.org>) and Fungalpedia (Hyde et al. 2023). In addition, there are several recently introduced fungal web pages which are dedicated to specific hosts, ecosystems and localities,

allowing researchers to easily access and cite relevant data. A few examples are dothideomycetes.org (Pem et al. 2019) dedicated to Dothideomycetes, botryosphaerales.org (Wu et al. 2021) dedicated to Botryosphaerales, coelomycetes.org (Huanraluek et al. 2021) dedicated to Coelomycetes, Italian microfungi.org (Wijesinghe et al. 2021) dedicated to fungi associated with Italian flora, Soilfun.org on soil-inhabiting Ascomycota (Yasanthika et al. 2023) and Beeltehangers.org (de Groot et al. 2024) dedicated to explore spatiotemporal trends of *Hesperomyces harmoniae*.

As a timely and significant addition to the palm fungal studies, herein we introduce palmfungi.org an online platform dedicated to fungi associated with various palm species worldwide. This study uses palmfungi.org and related data to briefly discuss the current status of research on the distribution of palm fungi populations, showing how palmfungi.org serves as a bridge between the available data on fungi and their palm hosts, providing researchers with a sustainable platform for fungal information. In addition, this website will be the global consortium for studies on various aspects of palm fungi.

Why do we need palmfungi.org?


Palm fungi research and records have a long history. Based on the language and frequency of reports, Pereira and Phillips (2023) divide the history of palm fungus research into three phases: 1880s – 1920s, 1920s – 1990s and 1990s – the present. However, we re-define the time ranges of these three phases after adding the standard based on the method of fungal identification. We named the first phase as “Historical palm fungi era”, which has been reported sporadically using short Latin paragraphs and relatively subjective identification methods, from the 1820s to 1990s. The second phase we named as “Classical palm fungi era”, which was led by K.D. Hyde and co-authors using standard morphological methods to identify fungi on palms, from the 1990s to around 2005. The third phase is named “Molecular palm fungi era”, which combined molecular analysis with morphology for standardised fungal identification, started around 2005 to continues today. Taylor and Hyde (2003) reported the classification of a large amount of palm fungi, prompting researchers to pay more attention to how to collect and integrate palm fungi. However, the decentralised reporting palm fungi means that data cannot be collected and integrated simultaneously, which might cause misjudgement of the diversity of palm fungi. Most of the fungal species introduced in the early 20th century lacked molecular data or living cultures. Furthermore, these data are mostly published in printed books and not frequently referred to or cited. Palmfungi.org as an integrated sustainable data platform will facilitate solving this issue. This website will provide researchers with concise, relatively complete information on palm fungal species and as a directory of palm fungal diversity and host-specificity (Zhou and Hyde 2001).

In addition to connecting the history of palm fungi with modern data, this website also correlates palm fungi and palm plant taxa by citing the identification and classification of palms by Kissling et al. (2019) and Yao et al. (2023). In recent years, the research on palms has not been limited to the classification and identification of species. Most people

have focused on the application and uses of palms (Khan et al. 2023). By establishing itself as a data bridge between available data on fungi and palms, Palmweb (<https://palmweb.org>) has the potential to become an important tool in applied research.

What is palmfungi.org?

The role of palmfungi.org is to establish a retrieval database, based on fungi species that use palms as hosts. The operation process of the website is shown in Fig. 1. In terms of host data, we listed the names of all known species of Arecaceae, according to Kissling et al. (2019) and Palmweb (2023) and the taxonomic status of Arecaceae species will be kept updated. In terms of fungal classification, we followed the “Outline of fungi” (Wijayawardene et al. 2022b, <https://www.outlineoffungi.org>). Meanwhile, we have added the vast majority of fungal species names from palms and dedicated entries for some of the species for which information is available. In addition, we implemented interactive searches for palm plant species and fungal species through hyperlinks and dual-function tags.



We will further refine the summary including the order, family, genus of fungi and other important data. Readers can click on relevant links from the palm fungi website which will redirect to the other databases including “Faces of fungi” (Jayasiri et al. 2015, <http://www.facesoffungi.org>); “Onestopshop fungi” (Jayawardena et al. 2019, <https://onestopshopfungi.org>); “Marine fungi” (Jones et al. 2019, <http://marinefungi.org>); “Freshwater fungi” (Calabon et al. 2020, <http://freshwaterfungi.org>); “Sordariomycetes” (Bundhun et al. 2020, <https://sordariomycetes.org>); “Fungal Genera” (Monkai et al. 2020, <https://www.fungalgenera.org>); “Outline of fungi” (Wijayawardene et al. 2020, <https://www.outlineoffungi.org>) and “gmsmicrofungi” (Chaiwan et al. 2021, <http://gmsmicrofungi.org>).

Association with plant classification

According to APG IV (Angiosperm Phylogeny Group et al. 2016), Arecaceae (Palm) is the largest family within the order Arecales. Kissling et al. (2019) reported that there are 181 genera and nearly 2,600 species of palms. Despite this, taxonomic studies on palm plants are still common in recent years (Bellot et al. 2020, Helmstetter et al. 2020, Hodel et al. 2021, Jiménez et al. 2021, Eiserhardt et al. 2022). For fungi, especially saprophytic fungi, confirming the host species is a key step in taxonomic identification (Senanayake et al. 2020). Palmfungi.org cites identification and classification information of palm trees by Kissling et al. (2019) and Yao et al. (2023) and keeps it updated. Thus, Palmfungi.org will be a useful platform to solve this challenge and, together with Palmweb (<https://palmweb.org>), it will build a data bridge between the fungi and plants.

Construction

Following the Outline of Ascomycetes (Wijayawardene et al. 2022b), all fungi using palm plants as hosts are included in the database. The database of palmfungi.org will be updated periodically as new information becomes available. Outlines, detailed descriptions and relevant information on each entry on the website will be carefully verified by the expert curators (Table 1).

Database interface and visualisation

Palmfungi.org is an online fungi-palm interactive retrieval platform that compiles published information based on the taxonomy of fungi that host palm plants. The functions of the website are diverse, the interface is simple and user-friendly. The website consists of eight tabs; home, hosts substrate, archives, curators, history, references, notes and contact details as each tab with different functions. In addition, there is a right toolbar for searching and displaying recent updates. Finally, the lower border for displaying copyright and entry content to display fungal species details.

Table 1.

List of expert curators with their contact information.

Position	Name	Address	Contact information
Head Curators	Kevin D. Hyde	Center of Excellence in Fungal Research, School of Science Mae Fah Luang University, Chiang Rai, Thailand 57100	kdhyde3@gmail.com
	Ishara S. Manawasinghe	Innovative Institute for Plant Health, Zhongkai University of Agriculture and Engineering, Guangzhou 510225, P.R. China	ishara9017@gmail.com
Managing curator	Yinru Xiong	Center of Excellence in Fungal Research, School of Science, Mae Fah Luang University, Chiang Rai, 57100, Thailand	richard_xyr@163.com

Right toolbar

The right toolbar consists of three sections, the search toolbar (a), recent genus and recent species (b), are fixed on the right side of the entire website (Fig. 3). The fungal genus or taxon of interest can be entered in the search toolbar. Then a pop-up window will prompt the target fungus, including its taxonomic level. Clicking on the corresponding entry name will lead you to the species entry interface.




Figure 3. doi

Right toolbar and search result: a Search toolbar; b Recently updated news, recent genera and species; c The entry details the interface of fungi species.

Lower border

This information shows contact details and copyright ownership (Fig. 4).



Figure 4. doi

Bottom border: a Contact details; b Publisher and copyright information.

Homepage

The homepage (Fig. 5) shows the objectives of the website and the general information of the web (the function menu includes the search toolbar, home page, host/substrate, archive, curator, history, references, notes, contact). In addition, in Fig. 5c, it shows users the reference materials when citing this website. In addition, the number of palm fungi (including each classification level) and the number of palm species (including each classification level) currently included on the website are also shown.

The screenshot shows the homepage of palmfungi.org. At the top, there is a navigation menu (a) with links for Home, Host/substrate, Archives, Curators, History, References, Notes, and Contact. Below the menu is a search bar (b) with a search icon. The main content area features a large image (c) of a palm leaf with a dark spot, likely a fungal infection. To the right of the image is a sidebar with sections for Recent Notes, Recent Genus, and Recent Species. Below the image, there are statistics (d, e, f, g) showing the number of palm fungal taxa, records, species entries, and species. The statistics are: Number of palm fungal taxa (Orders: 95, Families: 284, Genera: 755), Estimated number of palm fungal records in palmfungi.org (as of 14/06/2024): 1,126, Number of palm fungal species entries in palmfungi.org (as of 14/06/2024): 78, and Estimated number of palm species (Genera: 184, Species: 2,557).

Figure 5. [doi](#)

Homepage: **a** Headers; **b** Objectives of the website; **c** Citation of the website; **d** Number of palm fungi (including Order, family and genus); **e** Number of palm fungi records (Documented on the website after screening); **f** Number of palm fungi species entries; **g** Number of palm species (including each classification level).

Host/Substrate

By entering this tab, users can obtain information regarding all currently-known palm species based on their classification (<https://palmweb.org>) (Fig. 6). Users can intuitively find the fungal species and quantities reported on each palm species. At the same time, each palm species provides entry to the fungal species that host this palm species. In addition, there is also a search bar frozen at the top of the page, where you can enter the palm genus or taxon of interest and the fungus genus or taxon and the page will automatically lock to the target entry. Clicking on the corresponding entry name will guide users to the entry's detailed information interface.




Figure 6. [doi](#)
Host/Substrate: **a** Host/Substrate tag; **b** Frozen search bar; **c** Search bar; **d** Palm fungus species names with hyperlinks; **e** The entry details the fungal species that host this palm species.

Archives

Provides users with a relevant list of palm fungi at various classification levels (highest classification level is order, lowest classification level is species) (Fig. 7). By clicking on the relevant term, the user is presented with options for "Read more about this entry" and a list of sub-categories.




Figure 7. [doi](#)
Archive: **a** Archive tag; **b** Order list; **c** Family list; **d** Genus list; **e** Species list; **f** The entry details the fungal species.

Curators

Provides the contact information and affiliation of the website curators (Fig. 8).

The screenshot shows the 'Curators' page of the PALM FUNGI website. The page features a navigation bar at the top with links for Home, Host/substrate, Analysis, Curators (highlighted), History, References, Notes, and Contact. Below the navigation bar, there is a search bar and a list of curators. Each curator entry includes a small portrait photo, the curator's name, title, affiliation, and email address. The curators listed are: Head Curator Emerita Prof. Kevin D. Hyde, Head Curator Dr. Inês S. Matavola, Managing Curator Dr. Yuxia Xiong, Curator Dr. Joanne Taylor, Curator Prof. Alan J.L. Phillips, Curator Dr. Shengnan Zhang, and Curator Dr. Carlos Santos Pereira. To the right of the curators list, there are sections for 'Recent Notes', 'Recent Genus', and 'Recent Species'.

Figure 8. [doi](#)

Curators: Website curator information.

History

Shows the palm fungi collected, examined and recorded with a brief historical background (Fig. 9).

The screenshot shows the 'History' page of the PALM FUNGI website. The page features a navigation bar at the top with links for Home, Host/substrate, Analysis, Curators, History (highlighted), References, Notes, and Contact. Below the navigation bar, there is a search bar and a world map showing the distribution of palm fungi. Below the map, there is a section titled 'Palm: Importance, Diversity and Distribution' and another section titled 'Early studies on Palm fungi'. The text discusses the importance of palm fungi, their diversity, and the early studies on them.

Figure 9. [doi](#)

History: Website history information.

References

Assembly of the main literature (such as books, reviews, monographs and articles) and websites (Fig. 10).

Figure 10. [doi](#)

References: Website reference information.

Notes

The note section is dedicated to additional details of relevant to the palm fungi, with two sections (Fig. 11). One link is for Important News (a) and will link the recent reviews, publications and events or other updated news relevant to the palm fungi. The second is Updated Log (b) which shows the updated time and person who updated the website and includes the hyperlink of the update log.

Figure 11. [doi](#)

Notes: a Important News; b Updated Log.

Contact

Provides the contact information of the website and allows users provide any comments or suggestions (Fig. 12).




Figure 12. [doi](#)

Contact: Website contact information.

Use palmfungi.org to briefly discuss the current status of research on the population distribution of palm fungi

Palm species have many uses (Dransfield et al. 2008), and Cámara-Leret et al. (2017) reported that 208 species of palm plants can be used as cash crops. Data from FAOSTAT (Food and Agriculture Organization of the United Nations 2024, <https://www.fao.org/faostat/en/#data>) also show that palms, especially oil palms and coconuts, have huge economic benefits. Based on data from Index Fungorum (2024) and Farr et al. (2021), palmfungi.org reviewed and included 1,521 fungal species associated with palm.

Based on the above data, this study briefly analysed the reported distribution of fungi in the cash crops and non-cash crops of the palm. The results show in (Fig. 13a): 784 fungal species were reported from cash crop palm hosts, accounting for 51.55%; 421 fungal species were reported on non-cash crop palm hosts, accounting for 27.68%; 316 fungal species reported from unidentified palm species, accounting for 20.78%. Simultaneously, utilising the information from Host/Substrate in palmfungi.org, this study conducted a basic analysis to determine the number of palm species that have fungus records. The results showed as follows (Fig. 13b): there were 316 species of palm with fungi recorded, accounting for 12.36%, and 2241 species of Palm without fungi recorded, accounting for 87.64%.




Figure 13. [doi](#)

A diagram of numbers of palm fungi vs. number of palm species or Areaceae.

Leveraging palmfungi.org as a platform, this study successfully conducted a basic analysis of the population distribution of palm fungi. The findings suggest that, while there is widespread documentation of palm fungi, the study focuses on a limited number of economically valuable palm species and overlooks the vast majority of naturally-occurring palm species.

Discussion

The advancement of modern online platforms has greatly affected the ease of gathering information, which connects various databases about different biological information (Chaiwan et al. 2021). Basic information on related fungi can be found in several global fungal nomenclature and classification databases outlined in this paper. Those resources provides users with fungal host records and, as fungal species are continually being discovered, new host records and regional records are constantly being reported (Hyde et al. 2024). In recent years studies have recorded the fungal host-specificity (Zhou and Hyde 2001). Bubner (2013) reported the host-specificity of ectomycorrhizal fungi in pure and mixed stands of Scots pine (*Pinus sylvestris* L.) and beech (*Fagus sylvatica* L.), Miao et al. (2021) found that plant genetics are a deciding factor affecting *Nitraria tangutorum* endophytic fungal composition, Wang et al. (2019) pointed out that soil plant pathogenic fungi are specialised in low-level host taxa and Li et al. (2020) describe and discuss host-specific genes in most fungi at the molecular level. In addition, Hyde et al. (2007) pointed out that there are also differences in fungi from different locations on the same host.

Despite the fact that fungi are so closely associated with plants, the only fungal-plant database available is for fungi on rice (https://mycolab.pp.nchu.edu.tw/rice_fungi/contact.php). However, this website cannot be used to explore fungal host specificity and is not interlinked with other large databases. The brief discussion of the distribution of palm fungi populations in this study fully demonstrates the effective application of palmfungi.org after interconnection with other large databases. Therefore, the operation of palmfungi.org provides a platform for the exploration of the host-fungus relationship.

We have presently uploaded 50 species entries to palmfungi.org and the website content will be continuously updated with the assistance of all curators. In the future, the palm fungi website will establish links between large and small fungal classification websites to form an integrated and interactive data collection platform. More attention will be paid to links with palm-related taxonomic websites such as palm web (<https://palmweb.org>), where associations with palm species will further improve the understanding of the relationship between fungi and palm hosts. Palmfungi.org will also be the first retrieval platform to record and retrieve taxonomic data on fungi and their specifically corresponding palm hosts. This platform will enable a new direction for exploring the host specificity of palm fungi and even the whole fungal kingdom.

Acknowledgements

Yinru Xiong would like to thank Mae Fah Luang University for awarding a scholarship. This research was funded by the High-level Talents at Zhongkai University of Agriculture and Engineering, grant no: J2201080102. Ishara Manawasinghe would like to acknowledge Zhongkai University of Agriculture and Engineering, talent funding (grant number KA210319288) and the Guangzhou Science and Technology Plan Project (2023A04J1427). We would like to acknowledge the Innovative team programme of the Department of Education of Guangdong Province (2022KCXTD015 and 2022ZDJS020). The authors also extend their appreciation to the Researchers Supporting Project number (RSP2024R114), King Saud University, Riyadh, Saudi Arabia for funding this work.

Conflicts of interest

The authors have declared that no competing interests exist.

References

- Angiosperm Phylogeny Group, Chase MW, Christenhusz MJ, Fay MF, Byng JW, Judd WS, Soltis DE, Mabberley DJ, Sennikov AN, Soltis PS, Stevens PF (2016) An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of The Linnean Society* 181 (1): 1-20. <https://doi.org/10.1111/boj.12385>

- Baker WJ, Dransfield J (2016) Beyond genera palmarum: progress and prospects in palm systematics. *Botanical Journal of The Linnean Society* 182 (2): 207-233. <https://doi.org/10.1111/boj.12401>
- Bellot S, Odufuwa P, Dransfield J, Eiserhardt WL, Perez-Escobar OA, Petoe P, Usher E, Baker WJ (2020) Why and how to develop DNA barcoding for Palms? A case study of *Pinanga*. *Palms* 64 (3): 109-120.
- Bhunjun CS, Phukhamsakda C, Hyde KD, McKenzie EH, Saxena RK, Li Q (2023) Do all fungi have ancestors with endophytic lifestyles? *Fungal Diversity* 1-26 <https://doi.org/10.1007/s13225-023-00516-5>
- Bolman B (2023) What mysteries lay in spore: taxonomy, data, and the internationalization of mycology in Saccardo's *Sylloge Fungorum*. *The British Journal for the History of Science* 56 (3): 369-390. <https://doi.org/10.1017/s0007087423000158>
- Bubner B (2013) Host specificity and biodiversity of ectomycorrhizal fungi in pure and mixed stands of Scots pine (*Pinus sylvestris* L.) and beech (*Fagus sylvatica* L.). BTU Cottbus-Senftenberg [In English].
- Bundhun D, Maharachchikumbura SS, Jeewon R, Senanayake IC, Jayawardena RS, Hongsanan S (2020) <https://sordariomycetes.org/>, a platform for the identification, ranking and classification of taxa within Sordariomycetes. *Asian Journal of Mycology* 3: 13-21. <https://doi.org/10.5943/ajom/3/1/2>
- Calabon MS, Hyde KD, Jones EB, Chandrasiri S, Dong W, Fryar SC, Yang J, Luo ZL, Lu YZ, Bao DF, Boonmee S (2020) www.freshwaterfungi.org, an online platform for the taxonomic classification of freshwater fungi. *Asian Journal of Mycology* 3: 419-445. <https://doi.org/10.5943/ajom/3/1/14>
- Cámara-Leret R, Faurby S, Macía M, Balslev H, Gödel B, Svenning J, Kissling WD, Rønsted N, Salsis-Lagoudakis CH (2017) Fundamental species traits explain provisioning services of tropical American palms. *Nature Plants* 3 (2). <https://doi.org/10.1038/nplants.2016.220>
- Chaiwan N, Gomdola D, Wang S, Monkai J, Tibpromma S, Doilom M, Wanasinghe DN, Mortimer PE, Lumyong S, Hyde KD (2021) <https://gmsmicrofungi.org>: an online database providing updated information of microfungi in the Greater Mekong Subregion. *Mycosphere* 12 (1): 1513-1526. <https://doi.org/10.5943/mycosphere/12/1/19>
- Chen DJ, Landis JB, Wang HX, Sun QH, Wang Q, Wang HF (2022) Plastome structure, phylogenomic analyses and molecular dating of Arecaceae. *Frontiers in Plant Science* 13: 960588. <https://doi.org/10.3389/fpls.2022.960588>
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G (2004) MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology* 50 (1): 19-22.
- de Groot M, Christou M, Pan J, Adriaens T, Maes D, Martinou A, Roy H, Verbeke A, Haelewaters D (2024) Beetlehangars.org: harmonizing host-parasite records of *Harmonia axyridis* and *Hesperomyces harmoniae*. *Arthropod-Plant Interactions* 18: 665-679. <https://doi.org/10.1007/s11829-023-10037-2>
- Douanla-Meli C, Scharnhorst A (2021) Palm foliage as pathways of pathogenic Botryosphaeriaceae fungi and host of new *Lasiodiplodia* species from Mexico. *Pathogens* 10 (10): 1297. <https://doi.org/10.3390/pathogens10101297>
- Dransfield J, Uhl NW, Asmussen CB, Baker WJ, Harley MM, Lewis CE (2005) A new phylogenetic classification of the palm family, Arecaceae. *Kew Bulletin* 60: 559-569.

- Dransfield J, Uhl NW, Asmussen CB, Baker WJ, Harley MM, Lewis CE (2008) *Genera Palmarum - The Evolution and Classification of the Palms*. Royal Botanic Gardens, Kew, London, UK, 732 pp. [In English]. [ISBN 978-1-84246-182-2] <https://doi.org/10.34885/92>
- Edwards WN (1922) An Eocene microthyriaceous fungus from Mull, Scotland. *Transactions of the British Mycological Society* 8: 66-72. [https://doi.org/10.1016/S0007-1536\(22\)80008-5](https://doi.org/10.1016/S0007-1536(22)80008-5)
- Eiserhardt WL, Bellot S, Cowan RS, Dransfield J, Hansen LE, Heyduk K, Rabarijaona RN, Rakotoarinivo M, Baker WJ (2022) Phylogenomics and generic limits of *Dypsidinae* (Arecaceae): the largest palm radiation in Madagascar. *Taxon* 71 (6): 1170-1195. <https://doi.org/10.1002/tax.12797>
- Farr DF, Rossman AY, Castlebury LA (2021) United States National Fungus Collections Fungus-Host Dataset. Ag Data Commons URL: <https://doi.org/10.15482/USDA.ADC/1524414>
- Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, EH S, Stajich JE, Kahmann R, Boone C, Denning DW, Gow NA, Klein BS, Kronstad JW, Sheppard DC, Taylor JW, Wright GD, Heitman J, Casadevall A, Cowen LE (2020) Threats posed by the fungal kingdom to humans, wildlife, and agriculture. *MBio* 11 (3): 10.1128. <https://doi.org/10.1128/mBio.00449-20>
- Food and Agriculture Organization of the United Nations (2024) <https://www.fao.org/faostat/en/#home>. Accessed on: 2024-6-18.
- Fröhlich J, Hyde K (2000) *Palm microfungi*. Fungal Divers Press, The University of Hong Kong, Hong Kong.
- Guo LD, Hyde KD, Liew EC (1998) A method to promote sporulation in palm endophytic fungi. *Fungal Diversity* 1: 109-113.
- Haelewaters D, Quandt CA, Bartrop L, Cazabonne J, Crockatt M, Cunha S, De Lange R, Dominici L, Douglas B, Drechsler-Santos ER, Heilmann-Clausen J, Irga P, Jakob S, Lofgren L, Martin T, Muchane MN, Stallman J, Verbeken A, Walker A, Gonçalves S (2024) The power of citizen science to advance fungal conservation. *Conservation Letters* 17 (3): e13013. <https://doi.org/10.1111/conl.13013>
- Halme P, Heilmann-Clausen J, Rämä T, Kosonen T, Kunttu P (2012) Monitoring fungal biodiversity – towards an integrated approach. *Fungal Ecology* 5 (6): 750-758. <https://doi.org/10.1016/j.funeco.2012.05.005>
- Hawksworth D, Lücking R (2017) Fungal Diversity Revisited: 2.2 to 3.8 Million Species. *Microbiology Spectrum* 5 (4): 10.1128. <https://doi.org/10.1128/microbiolspec.funk-0052-2016>
- Hawksworth DL (2001) The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycological Research* 105 (12): 1422-1432. <https://doi.org/10.1017/S0953756201004725>
- Helmstetter AJ, Kamga SM, Bethune K, Lautenschläger T, Zizka A, Bacon CD, et al. (2020) Unraveling the phylogenomic relationships of the most diverse African palm genus *Raphia* (Calamoideae, Arecaceae). *Journal Plants* 9 (4): 549. <https://doi.org/10.3390/plants9040549>
- Hodel DR, Baker WJ, Bellot S, Pérez-Calle V, Cumberledge A, Barrett CF (2021) Reassessment of the Archontophoenicinae of New Caledonia and description of a new species. *Palms* 65 (3): 109-31.
- Huanraluek N, Padaruth OD, Jayawardena RS, Li WJ, Hongsanan S, Jeewon R, Maharachchikumbura SS, Senanayake IC, Phukhamsakda C, Wijayawardene NN,

- Shivas RG, Goonasekara ID, Norphanphoun C, Pem D, Calabon MS, Bhunjun CS, Hyde KD (2021) <https://www.coelomycetes.org>: Databank that contributes for the classification, identification and nomenclature of Coelomycetes. *Asian Journal of Mycology* 4 (2): 114-122. <https://doi.org/10.5943/ajom/4/2/8>
- Hyde KD, Taylor JE, Fröhlich J (2000) Genera of ascomycetes from palms. Vol. 2. *Fungal Divers* Press, The University of Hong Kong, Hong Kong.
 - Hyde KD, Bussaban B, Paulus B, Crous PW, Lee S, McKenzie EH, Photita W, Lumyong S (2007) Diversity of saprobic microfungi. *Biodiversity and Conservation* 16: 7-35. <https://doi.org/10.1007/s10531-006-9119-5>
 - Hyde KD, McKenzie EH, Koko TW (2011) Towards incorporating anamorphic fungi in a natural classification - checklist and notes for 2010. *Mycosphere* 2 (1): 1-88.
 - Hyde KD, Nilsson RH, Alias SA, Ariyawansa HA, Blair JE, Lei.Cai, Cock AWAMd, Dissanayake AJ, Glockling SL, Goonasekara ID, Gorczak M, Hahn M, Jayawardena RS, Kan JALv, Laurence MH, Lévesque CA, Li X, Liu J, Maharachchikumbura SSN, Manamgoda DS, Martin FN, McKenzie EHC, McTaggart AR, Mortimer PE, Nair PVR, Pawłowska J, Rintoul TL, Shivas RG, Spies CFJ, Summerel BA, Taylor PIWJ, Terhem RB, Udayanga D, Vaghef N, Walther G, Wilk M, Wrzosek M, Xu J, Yan J, Zhou N (2014) One stop shop: backbones trees for important phytopathogenic genera: I. *Fungal Diversity* 67: 1-105.
 - Hyde KD, Amuhene TB, Apurillo CC, Asghari R, Aumentado HD, Bahkali AH, Bera I, Bhunjun CS, Calabon MS, Chandrasiri S, Chethana KW, Doilom M, Dong W, Fallahi M, Kapov SA, Khyaju S, Le L, Li CJ, Li QR, Li YX, Lin CG, Linn MM, Liu JK, Liu NG, Luangharn T, Madagammana AD, Manawasinghe IS, Marasinghe DS, McKenzie EH, Meakala N, Meng QF, Mukhopadhyay S, Norphanphoun C, Pem D, Phukhamsakda C, Sarma VV, Selcuk F, Senanayake IC, Shah S, Shu YX, Silva HV, Su HL, Tavakol M, Thakshila SA, Thiyagaraja V, Thongklang N, Tian Q, Tibpromma S, Tun ZL, Ulukapi M, Wang Y, Wannasawang N, Wijayawardene NN, Wimalasena SD, Xiao Y, Xiong YR, Yasanthika WA, Li Q, Dai DQ (2023) Fungalpedia, an illustrated compendium of the fungi and fungus-like taxa. *Mycosphere* 14 (1): 1835-1959. <https://doi.org/10.5943/mycosphere/14/1/22>
 - Hyde KD, Wijesinghe SN, Afshari N, Aumentado HD, Bhunjun CS, Boonmee S, Camporesi E, Chethana KW, Doilom M, Dong W, Du TY, Farias AR, Gao Y, Jayawardena RS, Karimi O, Karunarathna SC, Kularathnage ND, Lestari AS, Li CJ, Li YX, Liao CF, Liu XF, Lu L, Lu YZ, Luo ZL, Ma J, Mamarabadi M, Manawasinghe IS, Mapook A, Mi LX, Niranjana M, Senanayake IC, Shen HW, Su HL, Tibpromma S, Xu RJ, Yan JY, Yang YH, Yang YY, Yu FQ, Kang JC, Zhang JY (2024) *Mycosphere Notes* 469-520. *Mycosphere* 15 (1): 1294-1454. <https://doi.org/10.5943/mycosphere/15/1/11>
 - Index Fungorum (2024) Index Fungorum. <http://www.indexfungorum.org/Names/Names.asp>. Accessed on: 2024-6-18.
 - Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai Y, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu J, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo J, Ghobad-Nejhad M, Nilsson H, Pang K, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen T, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsanan S, Jayawardena RS, Li W, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao R, Zhao

- Q, Kang J, Promputtha I (2015) The Faces of fungi database: fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74 (1): 3-18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jayawardena RS, McKenzie EH, Chen YJ, Phillips AJ, Hongsanan S, Norphanphoun C, Abeywikrama PD, Maharachchikumbura SS, Manawasinghe IS, McTaggart A, Shivas RG, Gentekaki E, Hyde KD (2019) <https://onestopshopfungi.org/>, a webpage to enhance identification of phytopathogenic genera. *Asian Journal of Mycology* 2: 281-286. <https://doi.org/10.5943/ajom/2/1/18>
 - Jayawardena RS, Hyde KD, Farias AR, Bhunjun CS, Fernandez HS, de Farias ARG, Bhunjun CS, Fernandez HS, Manamgoda DS, Udayanga D, Herath IS, Thambugala KM, Manawasinghe IS, Gajanayake AJ, Samarakoon BC, Bundhun D, Gomdola D, Huanraluek N, Sun Y, Tang X, Promputtha I, Thines M (2021) What is a species in fungal plant pathogens? *Fungal Diversity* 109 (1): 239-266. <https://doi.org/10.1007/s13225-021-00484-8>
 - Jiménez MFT, Prata EMB, Zizka A, Cohn-Haft M, de Oliveira AVG, Emilio T, Chazot N, Couvreur TLP, Kanga SM, Sonké B, Cano Á, Collevatti RG, Kuhnhäuser BG, Baker WJ, Antonelli A, Bacon CD (2021) Phylogenomics of the Palm Tribe Lepidocarpaceae (Calamoideae: Arecaceae) and Description of a New Species of *Mauritiella*. *Systematic Botany* 46 (3): 863-874. <https://doi.org/10.1600/036364421x16312067913543>
 - Jones EB, Pang KL, Abdel-Wahab MA, Scholz B, Hyde KD, Boekhout T, Ebel R, Rateb ME, Henderson L, Sakayaroj J, Suetrong S, Dayarathne MC, Kumar V, Raghukumar S, Sridhar KR, Bahkali AH, Gleason FH, Norphanphoun C (2019) An online resource for marine fungi. *Fungal Diversity* 96: 347-433. <https://doi.org/10.1007/s13225-019-00426-5>
 - Khan MI, Shanableha A, Manzoorb S, Rehman A, Shahidad S, Ahmade F (2023) Application of United Arab Emirates Arecaceae leaves biochar for adsorptive removal of Rhodamine B from an aqueous solution. *Desalination And Water Treatment* 283: 247-258. <https://doi.org/10.5004/dwt.2023.29217>
 - Kidston R, Lang WH (1921) On Old Sandstone plants showing structure, from the Rhyne Chert Bed, Aberdeenshire. Part V. The thallophyta occurring in the peat-bed; the succession of the plants through a vertical section of the bed, and the conditions of accumulation and preservation of the deposit. *Transactions of the Royal Society of Edinburgh* 52: 855-902. <https://doi.org/10.1017/S0080456800016045>
 - Kissling WD, Balslev H, Baker W, Dransfield J, Gödel B, Lim JY, Onstein R, Svenning J (2019) PalmTraits 1.0, a species-level functional trait database of palms worldwide. *Scientific Data* 6 (1): 178. <https://doi.org/10.1038/s41597-019-0189-0>
 - Konta S, Hyde KD, Eungwanichayapant PD, Doilom M, Tennakoon DS, Senwana C, Boonmee S (2020) *Fissuroma* (Aigialaceae: Pleosporales) appears to be hyperdiverse on Arecaceae: evidence from two new species from southern Thailand. *Acta Botanica Brasilica* 34: 384-393. <https://doi.org/10.1590/0102-33062020abb0021>
 - Konta S, Hyde KD, Eungwanichayapant PD, Karunarathna SC, Samarakoon MC, Xu J, Dauner LAP, Aluthwattha ST, Lumyong S, Tibpromma S (2021) Multigene Phylogeny Reveals *Haploanthostomella elaeidis* gen. et sp. nov. and Familial Replacement of *Endocalyx* (Xylariales, Sordariomycetes, Ascomycota). *Life* 11 (6): 486. <https://doi.org/10.3390/life11060486>
 - Konta S, Tibpromma S, Karunarathna SC, Samarakoon MC, Steven LS, Mapook A, Boonmee S, Senwana C, Balasuriya A, Eungwanichayapant PD, Hyde KD (2023) Morphology and multigene phylogeny reveal ten novel taxa in Ascomycota from terrestrial

- palm substrates (Arecaceae) in Thailand. *Mycosphere* 14 (1): 107-152. <https://doi.org/10.5943/mycosphere/14/1/2>
- Li J, Cornelissen B, Rep M (2020) Host-specificity factors in plant pathogenic fungi. *Fungal Genetics and Biology* 144: 103447. <https://doi.org/10.1016/j.fgb.2020.103447>
 - Maharachchikumbura SSN, Chen Y, Ariyawansa HA, Hyde KD, Haelewaters D, Perera RH, Samarakoon MC, Wanasinghe DN, Bustamante DE, Liu J, Lawrence DP, Cheewangkoon R, Stadler M (2021) Integrative approaches for species delimitation in Ascomycota. *Fungal Diversity* 109 (1): 155-179. <https://doi.org/10.1007/s13225-021-00486-6>
 - Mahmoud FM, Yekkour A, Boudffeur S, Errahmani MB, Krimi Z (2016) Root endophytic fungi from date palm (*Phoenix dactylifera* L.) grove of Algerian Sahara and screening of their growth promotion activities. *Advances in Environmental Biology* 10 (11): 18-27.
 - Mahmoud FM, Krimi Z, Maciá-Vicente J, Errahmani M, Lopez-Llorca L (2017) Endophytic fungi associated with roots of date palm (*Phoenix dactylifera*) in coastal dunes. *Revista Iberoamericana de Micología* 34 (2): 116-120. <https://doi.org/10.1016/j.riam.2016.06.007>
 - Manawasinghe IS, Phillips AJL, Xu J, Balasuriya A, Hyde KD, Stępień Ł, Harischandra DL, Karunarathna A, Yan J, Weerasinghe J, Luo M, Dong Z, Cheewangkoon R (2021) Defining a species in fungal plant pathology: beyond the species level. *Fungal Diversity* 109 (1): 267-282. <https://doi.org/10.1007/s13225-021-00481-x>
 - Marasinghe DS, Hongsanan S, Wanasinghe DN, Boonmee S, Lumyong S, Hyde KD, Ning X (2022) Morpho-molecular characterization of *Brunneofissuraceae* fam. nov., *Cirsosia mangiferae* sp. nov., and *Asterina neomangiferae* nom. nov. *Mycological Progress* 21 (1): 279-295. <https://doi.org/10.1007/s11557-021-01767-9>
 - Marasinghe DS, Hongsanan S, Zeng XY, Jones EB, Chomnunti P, Boonmee S, Hyde KD (2023) Taxonomic monograph of epifoliar fungi. *Fungal Diversity* 121 (1): 139-334. <https://doi.org/10.1007/s13225-023-00522-7>
 - Miao SM, Zhang YY, Cui JL, Zhang G (2021) Species and geographic specificity between endophytic fungi and host supported by parasitic *Cynomorium songaricum* and its host *Nitraria tangutorum* distributed in desert. *Archives of Microbiology* 203: 2511-2519. <https://doi.org/10.1007/s00203-021-02224-7>
 - Monkai J, McKenzie EH, Phillips AJ, Hongsanan S, Pem D, Liu JK, Chethana KW, Tian Q, Ekanayaka AH, Lestari AS, Zeng M, Zhao Q, Norphanphoun C, Abeywikrama PD, Maharachchikumbura SS, Jayawardena RS, Chen YJ, Zhao RL, He MQ, Raspé O, Kirk PM, Gentekaki E, Hyde KD (2020) <https://fungalgenera.org/>: a comprehensive database providing web-based information for all fungal genera. *Asian Journal of Mycology* 2: 297-304.
 - Mueller GM, Bills GF, Foster MS (2004) Biodiversity of fungi: Inventory and Monitoring Methods. 1. Academic Press [ISBN 0125095511]
 - Niskanen T, Lücking R, Dahlberg A, Gaya E, Suz L, Mikryukov V, Liimatainen K, Druzhinina I, Westrip JS, Mueller G, Martins-Cunha K, Kirk P, Tedersoo L, Antonelli A (2023) Pushing the Frontiers of Biodiversity Research: Unveiling the Global Diversity, Distribution, and Conservation of fungi. *Annual Review of Environment and Resources* 48 (1): 149-176. <https://doi.org/10.1146/annurev-environ-112621-090937>
 - O'Brien H, Parrent J, Jackson J, Moncalvo J, Vilgalys R (2005) Fungal community analysis by large-scale sequencing of environmental samples. *Applied And Environmental Microbiology* 71 (9): 5544-5550. <https://doi.org/10.1128/AEM.71.9.5544-5550.2005>

- Palmweb (2023) Palmweb: Palms of the World Online. <https://palmweb.org>. Accessed on: 2024-4-24.
- Pandian RTP, Thube SH, Bhavishya, Merinbabu, Chaithra, Santhoshkumar P, Nirmalkumar BJ, Hegde V (2021) First report of *Phytophthora palmivora* (E. J. Butler) E. J. Butler, 1919 causing fruit rot in *Areca triandra* Roxb. ex Buch.-Ham. from India. Australasian Plant Pathology 50 (4): 495-499. <https://doi.org/10.1007/s13313-021-00802-3>
- Pem D, Hongsanan S, Doilom M, Tibpromma S, Wanasinghe DN, Dong W, Ningguo L, Phookamsak R, Phillips AJ, Jeewon R, Hyde KD (2019) <https://www.dothideomycetes.org>: an online taxonomic resource for the classification, identification, and nomenclature of Dothideomycetes. Asian Journal of Mycology 2 (1): 287-297. <https://doi.org/10.5943/ajom/2/1/19>
- Pereira DS, Phillips AJ (2023) Palm fungi and their key role in biodiversity surveys: A review. Journal of fungi 9 (11): 1121. <https://doi.org/10.3390/jof911121>
- Pinnoi A, Lumyong S, Hyde KD, Jones EB (2006) Biodiversity of fungi on the palm *Eleiodoxa conferta* in Sirindhorn peat swamp forest, Narathiwat, Thailand. Fungal Diversity 22: 205-218.
- Pinruan U, Hyde KD, Lumyong S, McKenzie EH, Jones EB (2007) Occurrence of fungi on tissues of the peat swamp palm *Licuala longicalycata*. Fungal Diversity 25: 157-173.
- Schmit JP, Mueller GM (2007) An estimate of the lower limit of global fungal diversity. Biodiversity and Conservation 16 (1): 99-111. <https://doi.org/10.1007/s10531-006-9129-3>
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Jayawardena RS, Wanasinghe DN, Jeewon R, Bhat DJ, Xiang MM (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. Mycosphere 11 (1): 2678-2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Species Fungorum (2024) Species Fungorum. <http://www.speciesfungorum.org>. Accessed on: 2024-4-24.
- Stubblefield SP, Taylor TN, Beck CB (1985) Studies of Paleozoicfungi. IV. Wood decaying fungi in *Callixylon newberryi* from the Upper Devonian. American Journal of Botany 72 (11): 1765-1774. <https://doi.org/10.1002/j.1537-2197.1985.tb08449.x>
- Taylor JE, Hyde KD, Jones EB (1999) Endophytic fungi associated with the temperate palm, *Trachycarpus fortunei*, within and outside its natural geographic range. New Phytologist 142: 335-346. <https://doi.org/10.1046/j.1469-8137.1999.00391.x>
- Taylor JE, Hyde KD (2003) Microfungi of Tropical and Temperate palms. Fungal Diversity Press 108 (2): 223-223. <https://doi.org/10.1017/S0953756204249668>
- Taylor TN, Krings M, Taylor EL (2014) Fossil fungi. 1st Edition. Elsevier Science <https://doi.org/10.1016/C2010-0-68335-0>
- Wang Z, Jiang Y, Deane DC, He F, Shu W, Liu Y (2019) Effects of host phylogeny, habitat and spatial proximity on host specificity and diversity of pathogenic and mycorrhizal fungi in a subtropical forest. New Phytologist 223 (1): 462-474. <https://doi.org/10.1111/nph.15786>
- Wijayawardene NN, Hyde KD, Tibpromma S, Wanasinghe DN, Thambugala KM, Tian Q, Wang Y (2017) Towards incorporating asexual fungi in a natural classification: checklist and notes 2012-2016. Mycosphere 8 (9): 1457-1554. <https://doi.org/10.5943/mycosphere/8/9/10>

- Wijayawardene NN, Hyde KD, Al-Ani LK, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, Tokarev YS, Dai DQ, Letcher PM, Stephenson SL, Ertz D, Lumbsch HT, Kukwa M, Issi IV, Madrid H, Phillips AJ, Selbmann L, Pfliegler WP, Horváth E, Bensch K, Kirk PM, Kolaříková K, Raja HA, Radek R, Papp V, Dima B, Ma J, Malosso E, Takamatsu S, Rambold G, Gannibal PB, Triebel D, Gautam AK, Avasthi S, Suetrong S, Timdal E, Fryar SC, Delgado G, Réblová M, Doilom M, Dolatabadi S, Pawłowska J, Humber RA, Kodsueb R, Sánchez-Castro I, Goto BT, Silva DK, de Souza FA, Oehl F, da Silva GA, Silva IR, Błaszczowski J, Jobim K, Maia LC, Barbosa FR, Fiuza PO, Divakar PK, Shenoy BD, Castañeda-Ruiz RF, Somrithipol S, Lateef AA, Karunarathna SC, Tibpromma S, Mortimer PE, Wanasinghe DN, Phookamsak R, Xu J, Wang Y, Tian F, Alvarado P, Li DW, Kušan I, Matočec N, Maharachchikumbura SS, Papizadeh MPM, Heredia G, Wartchow F, Bakhshi M, Boehm E, Youssef N, Hustad VP, Lawrey JD, Santiago AL, Bezerra JD, Souza-Motta CM, Firmino AL, Tian Q, Houbraken J, Tanaka K, Dissanayake AJ, Monteiro JS, Grossart HP, Suija A, Weerakoon G, Etayo J, Tsurukau A, Vázquez V, Mungai P, Damm U, Li QR, Zhang H, Boonmee S, Lu YZ, Becerra AG, Kendrick B, Brearley FQ, Motiejūnaitė J, Sharma B, Khare R, Gaikwad S, Wijesundara DS, Tang LZ, He MQ, Flakus A, Rodríguez-Flakus P, Zhurbenko MP, McKenzie EH, Stadler M, Bhat DJ, Liu JK, Raza M, Jeewon R, Nassonova ES, Prieto M, Jayalal RG, Erdoğan M, Yurkov A, Schnittler M, Shchepin ON, Novozhilov YK, Silva-Filho AG, Liu P, Cavender JC, Kang Y, Mohammad S, Zhang LF, Xu RF, Li YM, Dayarathne MC, Ekanayaka AH, Wen TC, Deng CY, Pereira OL, Navathe S, Hawksworth DL, Fan XL, Dissanayake LS, Kuhnert E, Grossart HP, Thines M (2020) Outline of fungi and fungus-like taxa. *Mycosphere* 11: 1060-1456. <https://doi.org/10.5943/mycosphere/11/1/8>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğan M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EB, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SS, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJ, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkalčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DS, Wimalaseena SD, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang L, Stadler M, Yurkov A, Thines M (2022) Outline of fungi and fungus-like taxa - 2021. *Mycosphere* 13: 53-453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Wijesinghe SN, Camporesi E, Wanasinghe DN, Maharachchikumbura SS, Senanayake IC, Phookamsak R, Hongsanan S, Tibpromma S, Thambugala KM, Luangharn T, McKenzie EH, Hyde KD (2021) A dynamic online documentation of Italian ascomycetes with hosts and substrates: www.italianmicrofungi.org. *Asian Journal of Mycology* 4: 10-18. <https://doi.org/10.5943/ajom/4/1/2>
- Willis KJ (Ed.) (2018) State of the World's fungi 2018. Royal Botanic Gardens, Kew [ISBN 978-1-84246-678-0]
- Wu N, Dissanayake AJ, Manawasinghe IS, Rathnayaka AR, Liu JK, Phillips AJ, Promputtha I, Hyde KD (2021) <https://botryosphaerales.org/>, an online platform for up-to-date classification and account of taxa of Botryosphaerales. Database baab061.
- Xie A, Gee CT, Tian N (2023) Ancient Basidiomycota in an extinct conifer-like tree, *Xenoxylon utahense*, and a brief survey of fungi in the Upper Jurassic Morrison

Formation USA. *Journal of Paleontology* 97 (3): 754-763. <https://doi.org/10.1017/jpa.2023.12>

- Xiong YR, Manawasinghe IS, Liao CF, Hyde KD, Dong ZY (2022a) *Neodeightonia arengae* sp. nov., botryosphaeriaceous taxa on *Arenga tremula* (Arecaceae) from Guangdong, China. *Phytotaxa* 530 (2): 130-140. <https://doi.org/10.11646/phytotaxa.530.2.1>
- Xiong YR, Manawasinghe IS, Maharachchikumbura SS, Lu L, Dong ZY, Xiang MM, Xu B (2022b) Pestalotioid species associated with palm species from Southern China. *Current Research in Environmental & Applied Mycology (Journal of Fungal Biology)* 12 (1): 285-321. <https://doi.org/10.5943/cream/12/1/18>
- Yao G, Zhang YQ, Barrett C, Xue B, Bellot S, Baker WJ, Ge XJ (2023) A plastid phylogenomic framework for the palm family (Arecaceae). *BMC biology* 21 (1): 1-15.
- Yasanthika WA, Farias AR, Wanasinghe DN, Chethana KW, Zare R, Cai L, Maharachchikumbura SS, Tennakoon DS, Perera RH, Luangharn T, Chomnunti P (2023) <https://soilfun.org/>, a web-based platform for soil-inhabiting Ascomycota species. *Studies in fungi* 8: 16. <https://doi.org/10.48130/SIF-2023-0016>
- Zheng L, Xi P, Tu JJ, Chen X, Li J, Qin X, Shen H, Xie C (2017) First report of *Phoma herbarum* causing leaf spot of Oil Palm (*Elaeis guineensis*) in China. *Plant Disease* 101 (4): 629-630. <https://doi.org/10.1094/PDIS-05-16-0692-PDN>.
- Zhou DQ, Hyde KD (2001) Host-specificity, host-exclusivity, and host-recurrence in saprobic fungi. *Mycological Research* 105: 1449-1457. <https://doi.org/10.1017/S0953756201004713>