



www.bioinformation.net  
Volume 20(8)

Views

Received August 1, 2024; Revised August 31, 2024; Accepted August 31, 2024, Published August 31, 2024

DOI: 10.6026/973206300200898

BIOINFORMATION 2022 Impact Factor (2023 release) is 1.9.

**Declaration on Publication Ethics:**

The author's state that they adhere with COPE guidelines on publishing ethics as described elsewhere at <https://publicationethics.org/>. The authors also undertake that they are not associated with any other third party (governmental or non-governmental agencies) linking with any form of unethical issues connecting to this publication. The authors also declare that they are not withholding any information that is misleading to the publisher in regard to this article.

**Declaration on official E-mail:**

The corresponding author declares that lifetime official e-mail from their institution is not available for all authors

**License statement:**

This is an Open Access article which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. This is distributed under the terms of the Creative Commons Attribution License

**Comments from readers:**

Articles published in BIOINFORMATION are open for relevant post publication comments and criticisms, which will be published immediately linking to the original article without open access charges. Comments should be concise, coherent and critical in less than 1000 words.

**Disclaimer:**

The views and opinions expressed are those of the author(s) and do not reflect the views or opinions of Bioinformation and (or) its publisher Biomedical Informatics. Biomedical Informatics remains neutral and allows authors to specify their address and affiliation details including territory where required. Bioinformation provides a platform for scholarly communication of data and information to create knowledge in the Biological/Biomedical domain.

Edited by P Kanguane

Citation: Bathla *et al.* Bioinformation 20(8): 898-900 (2024)

# Nanobots: An endodontist saviour

Saurav Bathla<sup>1,\*</sup>, Manoj Kumar Hans<sup>1</sup>, Saurav Kumar Dutta<sup>1</sup>, Shyamalima Bhattacharyya<sup>2</sup>, Anindita Talukdar<sup>3</sup> & Sana Saifi<sup>1</sup>

<sup>1</sup>Department of Conservative Dentistry and Endodontics, Institute of Dental Sciences, Bareilly, Uttar Pradesh, India; <sup>2</sup>Department of Public Health Dentistry, Regional Dental College, SSUHS, Guwahati, India; <sup>3</sup>Department of Pediatric & Preventive Dentistry, Regional Dental College, SSUHS, Guwahati, India; \*Corresponding author

**Affiliation URL:**

<https://www.idsbareilly.in/>

<https://www.ssuhs.in/>

**Author contacts:**

Saurav Bathla - E-mail: saurav.bathla@yahoo.com; Phone: +919927770456

Manoj Kumar Hans - E-mail: hansie51@yahoo.com; +919829779528

Saurav Kumar Dutta - E-mail: sauravd.kumar@gmail.com; +919520523814

Shyamalima Bhattacharyya - E-mail: shyamalima113@gmail.com; +919639541544

Anindita Talukdar - E-mail: anin2228@gmail.com; +919916375198

Sana Saifi - E-mail: saifi.sana1804@gmail.com; +917906926233

**Abstract:**

The application of nanoparticles in the form of solution for irrigation, medication and as an additive for sealer/restorative material has been evaluated to improve the antibacterial efficacy in the field of endodontics. Recently developed nanobots are injected into the teeth to destroy pathogens and they are more effective in root canal therapy. They are helical shaped and composed of silicon dioxide with iron embedded into the silica body to provide magnetic properties. They are also able to move within the dentinal tubules and are manipulated peripherally through a low-intensity magnetic field. These nanobots can be administered into the cleaned central canal of a tooth samples which are immersed in deionized water and their motions are monitored using a near-infrared imaging technique to produce heat which is a hyperthermia-based bactericidal method which help to eradicate the surrounding pathogens.

**Keywords:** Dentinal tubules, *Enterococcus faecalis*, Magnetic field

**Description:**

Dental caries, trauma and tooth malformations frequently cause an infection of the dental pulp which results in pulpitis, pulp necrosis and finally apical periodontitis if left untreated. In that case, root canal treatment (RCT) is indicated, as the removal of necrotic pulp tissue and bacteria from the entire root canal system enables healing of periapical pathosis [1-3]. However, biomechanical preparation independent of the technique or type of irrigation used provides a significant reduction of the microbiota but only in the main root canal [4]. Even after completion of biomechanical preparation, the microorganisms remain persisted in dentinal tubules, isthmuses, lateral canals and accessory canals which may reinfect the root canal [5]. The

extent of penetration of microorganisms into the dentinal tubules of infected root canals reported in the literature varied from 200-1,500µm (Table 1) [6-17]. In fact, within oval canals, only 40% of the root canal wall can be contacted by instruments when a rotating technique is used. Therefore, irrigation is an essential part of a root canal treatment as it allows for cleaning beyond the root canal instruments [18]. The primary objective of chemo-mechanical preparation of root canal systems is elimination of pulpal tissue, inorganic and organic debris, bacteria and their toxic by-products through the use of instruments and irrigation devices [19].

**Table 1:** Depth of penetration of microorganisms into the dentinal tubules of infected root canals

S.No	Author(s)	Maximum depth of penetration	References
1)	Kouchi Y <i>et al.</i> (1980)	1050 - 1150 µm	[6]
2)	Haapasalo M & Orstavik D. (1987)	800 - 1000 µm	[7]
3)	Safavi KE <i>et al.</i> (1989)	50 - 300 µm	[8]
4)	Orstavik D & Haapasalo M. (1990)	600 - 1200 µm	[9]
5)	Horiba N <i>et al.</i> (1990)	300 - 800 µm	[10]
6)	Perez F <i>et al.</i> (1993)	Up to 800 µm	[11]
7)	Peters LB <i>et al.</i> (1995)	Up to 375 µm	[12]
8)	Berutti E <i>et al.</i> (1997)	Up to 300 µm	[13]
9)	Al-Nazhan S <i>et al.</i> (2014)	200 - 1,500 µm	[14]
10)	Vatkar NA <i>et al.</i> (2016)	>1000 µm	[15]
11)	Rosen E <i>et al.</i> (2020)	Up to 1480 µm	[16]
12)	Kumar PS <i>et al.</i> (2021)	800 to 1062 µm	[17]

**Table 2:** Penetration depth of root canal irrigants into dentinal tubules

S. No	Type of irrigants activation methods	Maximum depth of penetration	References
1	Conventional needle irrigation	31-280 µm	[21]
2	Sonic irrigation	110 - 337 µm	[21]
3	Ultrasound irrigation	160 - 330 µm	[21, 22]
4	Laser-PIPS (photon induced photoacoustic streaming)	Up to 900 µm	[23-25]
5	Laser-SWEEPS (shock-wave enhanced emission photoacoustic streaming)	650 - 800 µm	[26]
6	Magnetic nanobots	Up to 2000 µm	[27]

The success of endodontic therapy is based on the combination of adequate instrumentation, irrigation and three dimensional obturation of the root canal system. Approximately 2.3 billion individuals are affected by dental caries globally and about 24 to

50 million procedures are performed annually in the United States alone, of which more than 10% are reported to fail due to the persistence of microbes deep into the dentinal tubules. It is really challenging to remove bacteria from dentinal tubules using

traditional methods, because the dentinal tissue has a complicated and constrained morphology [20]. Ultrasonic or laser pulses were employed to produce shockwaves in the fluid used to flush away the pathogens and tissue debris to increase the effectiveness of root canal therapy. But the energy of these pulses quickly dissipated and they can only travel approximately up to 900 micro-meters (Table 2) [21-26]. Currently available nanobots can be injected into the teeth to destroy pathogen and they are more effective in root canal therapy. They are helical shaped and composed of silicon dioxide with iron embedded into the silica body to provide magnetic properties [27]. Moreover, the amount of silica used in the nanobots is negligible that it can be considered harmless for the human body. They are able to move within the dentinal tubules, manipulated extraneous through the low-intensity magnetic field. They are immersed in water or in a biocompatible medium that resembles water. A billion of them can fit in 0.5 ml of water drop. These nanobots can be administered into the cleaned central canal of a tooth samples which are immersed in deionized water, and their motions are monitored using a near-infrared imaging approach. The nanobots can reach up to 2,000 micro-meters and produce heat which is a hyperthermia-based bactericidal method which might eradicate the surrounding pathogens (Table 2) [27]. They were exposed to rotating magnetic fields, to retrieve them and reduce the chances of nanoparticle accumulation and toxicity. Thus, a hyperthermia-based bactericidal method provides a safer alternative to chemicals or antibiotics. *Enterococcus faecalis* is the most pronounced resistant bacteria isolated from the infected root canal. A nanobot with a rotating magnetic field into an infected tooth sample produced heat and within 15 minutes of exposure pathogen were destroyed [27].

#### Conclusion:

Available developed dental nanobots which is a hyperthermia-based bactericidal method provides a safer alternative to chemicals or antibiotics in root canal therapy. Further, the dental nanobots had been tested in animal models and proven to be safe and efficient.

#### Future clinical implications:

In the future, the nanobots can aid in surgical procedures, pulpal regeneration, inducing anesthesia, curing tooth hypersensitivity, non-vital tooth bleaching and targeted drug delivery.

**Conflicts of interest:** There are no conflicts of interest.

#### References:

- [1] Paque F *et al.* *J Endod.* 2009 **35**:1044. [PMID: 19567331]
- [2] Haapasalo M *et al.* *Br Dent J.* 2014 **216**:299. [PMID: 24651335]
- [3] Meto A *et al.* *Materials (Basel).* 2021 **14**:678. [PMID: 33540551]
- [4] Nangia D *et al.* *J Conserv Dent* 2020 **23**:185. [PMID: 33384493]
- [5] Souza CC *et al.* *J Conserv Dent* 2019 **22**:155. [PMID: 31142985]
- [6] Kouchi Y *et al.* *J Dent Res.*1980 **59**:2038. [PMID: 7019277]
- [7] Haapasalo M & Orstavik D. *J Dent Res* 1987 **66**:1375. [PMID: 3114347]
- [8] Safavi KE *et al.* *J Endodo* 1990 **16**:207. [PMID: 2074411]
- [9] Orstavik D & Haapasalo M. *Endod Dent Traumatol* 1990 **6**:142. [PMID: 2133305]
- [10] Horiba N *et al.* *J Endod* 1990 **16**:331. [PMID: 2081947]
- [11] Perez F *et al.* *Oral Surg Oral Med Oral Pathol* 1993 **76**:97. [PMID: 8351128]
- [12] Peters LB *et al.* *Int Endod J* 1995 **28**:95. [PMID: 7665208]
- [13] Berutti E *et al.* *J Endod* 1997 **23**:725. [PMID: 9487845]
- [14] Al-Nazhan S *et al.* *Restor Dent Endod* 2014 **39**:258. [PMID: 25383343]
- [15] Vatkar NA *et al.* *J Conserv Dent.* 2016 **19**:445. [PMID: 27656064]
- [16] Rosen E *et al.* *Applied Sciences.* 2020 **10**:6996. [https://doi.org/10.3390/app10196996]
- [17] Kumar PS *et al.* *Eur Endod J.* 2021 **6**:205. [PMID: 34650016]
- [18] Wu MK *et al.* *Int Endod J* 2003 **36**:218. [PMID: 12657148]
- [19] Cesario F *et al.* *J Conserv Dent* 2018 **21**:383. [PMID: 30122818]
- [20] Wong R. *Dent Clin North Am.* 2004 **48**:265. [PMID: 15066516]
- [21] Virdee SS *et al.* *Int Endod J.* 2020 **53**:986. [PMID: 32163598]
- [22] G. Faria *et al.* *Int Endod J.* 2019 **52**:385. [PMID: 30220093]
- [23] Galler KM *et al.* *Int Endod J.* 2019 **52**:1210. [PMID: 30828819]
- [24] De Moor RJ *et al.* *Lasers Surg Med.* 2009 **41**:520. [PMID: 19639621]
- [25] Blanken J *et al.* *Lasers Surg Med.* 2009 **41**:514. [PMID: 19639622]
- [26] Lukac N & Jezersek M. *Lasers Med Sci.* 2018 **33**:823. [PMID: 29327088]
- [27] Dasgupta D *et al.* *Adv Healthc Mater.* 2022 **11**:e2200232. [PMID: 35481942]