

A new species of *Bonnetia* Mart. (Bonnetiaceae) from the Pantepui of South America

Rafael G. Barbosa-Silva^{1,2}, Benjamin M. Torke², Pedro L. Viana^{1,3}

¹ Coordenação de Botânica, Museu Paraense Emílio Goeldi, Belém, Pará, 66077-830, Brazil

² Center for Biodiversity & Evolution, New York Botanical Garden, 2900 Southern Boulevard, Bronx, NY, 10458-5126, USA

³ Instituto Nacional da Mata Atlântica, Santa Teresa, Espírito Santo, 29650-000, Brazil

Corresponding author: Rafael G. Barbosa-Silva (rafa.g29@gmail.com)

Abstract

Bonnetia is the most representative genus of the Pantepui woody flora and is among the groups with the greatest endemism in the local flora. The genus has 28 currently recognized species in tropical America, 26 of them endemic to the Pantepui. Here we describe *Bonnetia ayangannensis* from the summit of Mount Ayanganna tepui in Guyana, providing a morphological description, illustrations, distribution maps, characterization of micromorphology under scanning electron microscopy and leaf venation, comments comparing the new species with closely related species, and a key for the identification of the species of *Bonnetia* occurring in Guyana. With its restricted distribution threatened by climate change, *Bonnetia ayangannensis* is assessed in the conservation threat category of Critically Endangered. Its description raises the number of endemic species of *Bonnetia* in the Pantepui to 27.

Key words: Amazonia, endemism, floristic inventory, Guayana Shield, herbarium, Malpighiales



Academic editor: John L. Clark

Received: 12 June 2024

Accepted: 15 September 2024

Published: 8 October 2024

Citation: Barbosa-Silva RG, Torke BM, Viana PL (2024) A new species of *Bonnetia* Mart. (Bonnetiaceae) from the Pantepui of South America. *PhytoKeys* 247: 55–65. <https://doi.org/10.3897/phytokeys.247.126950>

Copyright: © Rafael G. Barbosa-Silva et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Bonnetia Mart. is the largest genus of the Bonnetiaceae, comprising 28 species of trees and shrubs distributed across tropical America, with 26 species endemic to the Pantepui region of the Guayana Shield in northern South America (Weitzman et al. 2007). Species of *Bonnetia* often dominate woody landscapes of the tepui summits and upper slopes (Maguire 1972; Huber 1988, 2005, 2006; Barbosa-Silva et al. 2020), and the genus provides an excellent model for understanding vascular plant diversity and endemism within the Pantepui region (Riina et al. 2019). The seven species of *Bonnetia* occurring in Brazil were the focus of a recent taxonomic study (Barbosa-Silva 2020), but additional taxonomic work is needed, especially on the species in other parts of the Pantepui region.

Floral morphology holds significant value in the circumscription of genera of Bonnetiaceae. Within the genus *Bonnetia*, however, leaf venation patterns, particularly branching patterns, which are highly diverse in the genus, have been more important in the group's taxonomy (Maguire 1972; Dickison and Weitzman 1996). Such characteristics have even been used to support generic

segregates of *Bonnetia* (Maguire 1972; Steyermark 1984), but all these generic names are now treated as synonyms of *Bonnetia* (Barbosa-Silva 2020).

The most comprehensive advances in cataloging *Bonnetia* species diversity in the Pantepui resulted from several intensive floristic inventory programs in the region (see: Maguire 1972; Steyermark 1984; Barbosa-Silva et al. 2016; Kelloff et al. 2019). An example of these programs is the Smithsonian Institution's Biological Diversity of the Guiana Shield (**BDG**) Program, which began in 1983 as an initiative aimed at studying, documenting, and preserving the biological diversity of the Guayana Shield over 30 years. Here, we describe a new species of the genus based on collections gathered by the BDG (Kelloff et al. 2019). Its description marks the first addition of a new species to the genus (and the family Bonnetiaceae) in 37 years (Sastre 1987). It raises to 27 the number of species of *Bonnetia* endemic to the Pantepui region, and to six the number of species occurring in Guyana.

Materials and methods

All specimens of *Bonnetia* in the herbaria BM, COL, IAN, MG, MO, K, NY and US (acronyms follow Index Herbariorum, Thiers updated continuously) were examined. Specialized terminology for morphological structures follows Radford et al. (1974) and, for terms describing venation, Ellis et al. (2009). The description and illustration of reproductive structures was done from rehydrated material. Geographical data were sourced from specimen labels. The distribution map was generated using QGIS.org (2023). Assessment of the conservation status of the new species employed the categories and criteria of the IUCN Red List (IUCN 2012, 2024).

Samples of leaves, floral bracts, sepals and seeds from herbarium specimens deposited at NY were examined by Scanning Electron Microscopy (SEM), using a Hitachi model Su3500. Images were colored using Photoshop CS4 (Adobe). For the analysis of leaf venation, leaves were rendered transparent by soaking them in a 5% sodium hydroxide solution for two weeks at room temperature (Vasco et al. 2014). The transparent leaves were subsequently stained with 1% safranin and then immersed in ethanol to remove excess stain prior to imaging by an EPSON Perfection V750 PRO scanner.

Results

Taxonomic treatment

Bonnetia ayangannensis Barb.Silva, sp. nov.

urn:lsid:ipni.org:names:77349807-1

Type. GUYANA • Potaro-Siparuni, Mt Ayanganna, East slope, summit plateau; 5°23'18"N, 59°58'54"W; 1955m; fl.; 20 Mar 2014; A. Radosavljevic et al. 247 (holotype: NY 04078880, isotype US 03457967) (Figs 1–4).

Description. Shrub or small tree up to 1.5 m tall. Branchlets often with remnants of inflorescences below the congested leaves. Leaves alternate, sessile or with a petiole c. 1 mm long, leaf blade (1–)1.5–4 × (0.5–)1–1.8 cm, coriaceous, obtrullate, rarely rhombic or narrowly rhombic, base cuneate, apex slightly




Figure 1. *Bonnetia ayangannensis* **A** fertile branch, with details of the apex of a persistent inflorescence (at left) and the leaf abaxial surface and crenulate margin (at right) **B** immature inflorescence apex, showing bracts **C** flower bud **D** dimorphic sepals with glands, the detail showing the distal sepal margin **E** flower **F** stamens **G** closed capsule **H** transverse cut of the capsule **I** longitudinal cut of the capsule **J** seed (illustrated by: Bobbi Angell, from Radosavljevic et al. 247 and Clarke et al. 9524).

rounded to acuminate, margins crenulate, abaxial surface glabrous, with sparse black glands when dried, secondary veins scarcely visible, adaxial surface glabrous, rarely with black glands; venation pinnate with no naked basal veins, one basal vein, and no agrophic veins (Fig. 4), the major secondaries semicraspedodromous with irregular spacing, the secondary angle slightly increasing proximally, and with decurrent attachment to the midvein, the minor secondaries and interior secondaries absent, the intersecondaries parallel, spanning less than 50% of the length of the subjacent secondaries, the intercostal tertiary veins

irregularly reticulate, the epimedial tertiaries ramified with the admedial course parallel to the subjacent secondary and the exmedial course parallel to the intercostal tertiary, the exterior tertiaries terminating at the margin, the quaternary vein fabric irregularly reticulate, the quinternary vein fabric freely ramifying, areolation well developed, freely ending veinlets with two or more dendritic branches. Inflorescence axillary or occasionally terminal, once-paniculate, 2 cm long, with each branch bearing 3–4 flowers; bracts 2–3 × ca. 0.5 mm, oblanceolate to narrowly oblong, base truncate, rarely cuneate, apex rounded, margin ciliate; pedicels 2–3 mm long. Flowers 1.2–1.5 cm long, sepals 5, 2–4 × 1–2 mm, heteromorphic in size, oblong or ovate, apex rounded to acuminate, glabrous, margin ciliate, with sessile glands on the external surface; petals 5–6, 7–8 × 5–6 mm, broadly obtrullate, apex obcordate with asymmetric lobes, base cuneate, white with red margins; stamens 70–80, yellow, filaments 3 mm long, glabrous, anthers ca. 0.6 mm long; gynoecium ca. 5 × 2.5 mm, locules 3, carpels 3, green or yellow, stigma 3-branched in the upper fifth, the lobes reflexed. Capsules 7–8 × 3–3.5 mm, widely ovate to ovate. Seeds 1.5–2 mm long, linear.

Paratype. GUYANA • Potaro-Siparuni, Mt. Ayanganna, east face, edge of summit plateau; 5°23'18"N, 59°58'56"W; 2000 m elev.; fl., fr.; 24 June 2001; D. Clarke et al. 9524 (NY, US).

Notes. *Bonnetia ayangannensis* is morphologically most similar to *B. paniculata* Spruce ex Benth., but it differs from that species by the leaf blades (1–)1.5–4 cm long (vs. 6–18 cm in *B. paniculata*), the leaves with glandular punctuations on the abaxial surface (vs. glands absent), the bracts 2–3 mm long (vs. 6–7 mm), the sepals ciliate and glandular (vs. cilia and glands lacking) and 2–4 × 1–2 mm (vs. 10 mm × 6–7 mm). *Bonnetia tepuiensis* Kobuski & Steyerl. and *B. rubicunda* (Sastre) A.L. Weitzman & P.F. Stevens also occurs on the summit of Mount Ayanganna, however the new species can be differentiated by the leaves cuneate at base (vs. rounded in *B. tepuiensis*), the flowers pedicellate (vs. sessile), arranged in inflorescences (vs. solitary flowers) and by having leaf with venation pinnate and flowers with sepals and petals less than 8 mm long (vs. parallel leaf venation and sepals and petals more than 18 mm long in *B. rubicunda*). For better identification of the *Bonnetia* species of Guyana, see the identification key below.

The voucher Clarke et al. 9535 (NY 04067354) is a mixed specimen, consisting of three branches of *B. tepuiensis*, and a small branch to the right of *B. ayangannensis*.

Etymology. The epithet refers to the tepui Mount Ayanganna, where the new species was discovered.

Distribution and habitat. *Bonnetia ayangannensis* appears to be endemic to the summit area of Mount Ayanganna tepui, Potaro-Siparuni, in western Guyana. The species is known from only two collections gathered in close proximity from a single population on the eastern summit slope of Mount Ayanganna. The species occurs in scrub forest on sandstone, together with *B. tepuiensis*, *Clusia* spp. (Clusiaceae), and *Brocchinia* spp. (Bomeliaceae), among others, at elevations between 1900 and 2000 m.

Preliminary conservation status. We recommend that *Bonnetia ayangannensis* be assigned to the Critically Endangered category based on criterion B2ab(iv). The species is known from basically a single location (the two collection localities being separated by only c. 0.06 km), yielding an area of occupancy of 4 km², assuming the default settings in GeoCAT (Bachman et al. 2011;




Figure 2. SEM images of the *Bonnetia ayangannensis* **A** abaxial surface of the leaf showing the crenate margin, sometimes with deciduous spinulose projections (purple) **B** abaxial leaf surface with sessile glands (orange) and stomatal complexes (purple) **C** stomate immersed in a granular epidermis **D** bract **E** bract margin with short-stalked glands **F** sepals with sessile glands on the external surface **G** sepal margin with projections or short-stalked glands **H** seed (from Radosavljevic et al. 247 and Clarke et al. 9524).



Figure 3. Leaf venation of *Bonnetia ayangannensis* (from Radosavljevic et al. 247).

Bachman and Moat 2012). However, considering that the area above 1900 m elevation on Mount Ayanganna, where taller forest gives way to scrub forest, is only about 0.7 km², we suspect that the actual AOO is substantially less than 4 km². Moreover, it is likely that the population will undergo anthropogenic climate change-driven reduction, paralleling reduction in the extent of the summit scrub vegetation on Mount Ayanganna, the summit peak of which is only




Figure 4. **A** The map shows a topographic representation where warm colors indicate higher elevations, illustrating the geographical distribution of *Bonnetia ayangannensis* **B, C** view of Mount Ayanganna (photo by Adrian Warren/sasyimages.com).

2041 m elevation. This projection is based on modeling (Nogué et al. 2009) that suggests upward migration and potential habitat loss for many species with restricted tepui distributions. This, in turn, subjects these localities to threats according to the conditions of subcriterion b(iv). Other species occurring in the

eastern district of Pantepui, which are endemic to the summit of one or a few tepuis, are also threatened by rising temperatures, as is the case with *B. fasciculata* P.F.Stevens & A.L.Weitzman (Nogué et al. 2009) or are already listed on the IUCN Red List, such as *B. rubicunda* (Vulnerable) and *B. ptariensis* Steyerm. (Critically Endangered) (World Conservation Monitoring Centre 1998a, World Conservation Monitoring Centre 1998b).

Micromorphology. The leaves have a crenulate margin, sometimes bearing deciduous spinular projections (Fig. 2A). The abaxial leaf surface is rugose (i.e., not smooth and presenting different levels) and bears many sessile glands and stomata (Fig. 2B). At higher magnification, the surface takes on a warty-crustose appearance with granular projections (Fig. 2C). The margins of the bracts have short stalked glands with thin-walled elongate heads (Fig. 2D, E). The sepals have sessile glands on the external surface (Fig. 2F) and short stalked glands on the margins (Fig. 2F, G). The seeds bear longitudinal striations (Fig. 2H).

Key to the species of *Bonnetia* occurring in Guyana

- 1a Petals and capsules < 1 cm long..... 2
- 2a Leaf blade rounded at base ***B. tepuiensis* Kobuski & Steyerm.**
- 2b Leaf blade cuneate at base 3
- 3a Flowers arranged in inflorescences ***B. ayangannensis* Barb.Silva**
- 3b Flowers solitary ***B. roraimae* Oliv.**
- 1b Petals and capsules > 1 cm long..... 4
- 4a Flowers arranged in inflorescences ***B. paniculata* Spruce ex Benth.**
- 4b Flowers solitary 5
- 5a Leaf venation pinnate..... ***B. sessilis* Benth.**
- 5b Leaf venation parallel..... ***B. rubicunda* (Sastre) A.L. Weitzman & P.F. Stevens**

Discussion

Leaf characteristics hold significant taxonomic and historical value for the systematics of Bonnetiaceae. They provide support for the segregation of Bonnetiaceae as a distinct family (Dickison and Weitzman 1996) and have been used for the delimitation of genera now subsumed within *Bonnetia* (Maguire 1972). The micromorphological leaf features and leaf venation characters that we described for the new species may be of broader use in studies within the family, aiding in species delineation. For example, the warty-crustose leaf surface with a granular appearance that we observed in *B. ayangannensis* occurs in several other species of *Bonnetia*: *B. roraimae* Oliv., *B. steyermarkii* Kobuski, *B. wurdackii* Maguire (Dickison and Weitzman 1996). However, the spinulose projections on the leaf margin, which typically lack vascularization, are, as far as we know, shared only with *B. liesneri* Steyerm., although a similar structure is found in *B. ahogadoi* (Steyerm.) A.L. Weitzman & P.F. Stevens (Dickison and Weitzman 1996).

With respect to leaf venation, there is a fascinating array of branching patterns exhibited by different species of *Bonnetia*, ranging from pinnate to uniformly parallel (Maguire 1972; Dickison and Weitzman 1996). Our leaf clearing method showed that in *B. ayangannensis* the secondary veins are pinnate and

the higher order veins, which in uncleared leaves are almost imperceptible, are irregularly reticulate, with the quaternary veins delimiting well defined areolae. Comprehensive phylogenetic sampling of *Bonnetia* is necessary to test whether species with pinnate (most species) and parallel veins (i.e., *B. fasciculata* A.L. Weitzman & P.F. Stevens, *B. maguireorum* Steyerm., *B. multinervia* (Maguire) Steyerm., and *B. rubicunda*) form distinct clades.

As is the case for *B. ayangannensis*, many plant species, including species of herbs, epiphytes, and trees, appear to be endemics of single mountains in the Pantepui region (Riina et al. 2019). The conservation of these species is of grave concern, in light of ongoing and projected impacts of climate change on the vegetation of the tepuis (Nogué et al. 2009; Vieira et al. 2024). These projections suggest reductions in the areas covered by unique summit vegetation types and upward migration of species. In many cases, these trends may result in the total loss of narrowly endemic species as current habitats become unsuitable for their occupation. With many species teetering on the brink of losing their climatic suitability, urgent conservation efforts are warranted to safeguard the biodiversity of these fragile ecosystems.

Acknowledgements

We thank Bobbi Angell for preparing the illustration of the new species, Leeann Dabydeen and Tynisha Smalls for assistance in the laboratory at the New York Botanical Garden, Ingrid Gracielle Martins da Silva for assistance in the Microscopy and Microanalysis Laboratory of the Institute of Biological Sciences of the University of Brasilia (LMM-UnB), and Carol Kelloff and Ken Wurdack for providing additional details about collections at US.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study is part of the first author's PhD thesis research and was financially supported by the following Brazilian research agencies, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001 and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (03086/2022-1 and grants 473244/2017-4 and scholarship Pq-1A 307670/2022-8). RGBS also thanks EUROFINs, FUNBIO, HUMANIZE, Idea Wild, International Association for Plant Taxonomy, Neotropical Grassland Conservancy, American Society of Plant Taxonomists, The Linnean Society of London, and The Systematics Association for additional funding. PLV is supported by a research grant from CNPq (grant 312486/2020-0).

Author contributions

All authors have contributed equally.

Author ORCIDs

Rafael G. Barbosa-Silva  <https://orcid.org/0000-0002-0668-4840>

Benjamin M. Torke  <https://orcid.org/0000-0001-8823-3519>

Pedro L. Viana  <https://orcid.org/0000-0001-5044-0758>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bachman S, Moat J (2012) GeoCAT-an open source tool for rapid Red List assessments. *BGjournal* 9: 11–13.
- Bachman S, Moat J, Hill A, De La Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: Geospatial conservation assessment tool. *ZooKeys* 150: 117–126. <https://doi.org/10.3897/zookeys.150.2109>
- Barbosa-Silva RG (2020) Bonnetiaceae in Flora do Brasil 2020. Jardim Botânico do Rio de Janeiro. <http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB63> [accessed 30.03.2024]
- Barbosa-Silva RG, Labiak PH, Dos Santos Bragança Gil A, Goldenberg R, Michelangeli FA, Martinelli G, Nadruz Coelho MA, Zappi DC, Forzza RC (2016) Over the hills and far away: New plant records for the Guayana Shield in Brazil. *Brittonia* 68(4): 397–408. <https://doi.org/10.1007/s12228-016-9435-3>
- Barbosa-Silva RG, Bueno ML, Labiak PH, Coelho MAN, Martinelli G, Forzza RC (2020) The Pantepui in the Brazilian Amazon: Vascular Flora of Serra Do Aracá, a Cradle of Diversity, Richness and Endemism. *Botanical Review* 86(3–4): 359–375. <https://doi.org/10.1007/s12229-020-09235-x>
- Dickison WC, Weitzman AL (1996) Comparative Anatomy of the Young Stem, Node, and Leaf of Bonnetiaceae, Including Observations on a Foliar Endodermis. *American Journal of Botany* 83(4): 405–418. <https://doi.org/10.1002/j.1537-2197.1996.tb12722.x>
- Ellis B, Daly DC, Hickey LJ, Johnson KR, Mitchell JD, Wilf P, Wing SL (2009) Manual of leaf architecture. The New York Botanical Garden Press, 190 pp. <https://doi.org/10.1079/9781845935849.0000>
- Huber O (1988) Vegetacion y flora de Pantepui, Region Guayana. *Acta Botanica Brasílica* 1(2 suppl 1): 41–52. <https://doi.org/10.1590/S0102-33061987000300005>
- Huber O (2005) Diversity of vegetation types in the Guayana Region: An overview. *Biologiske Skrifter* 55: 169–188.
- Huber O (2006) Herbaceous ecosystems on the Guayana Shield, a regional overview. *Journal of Biogeography* 33(3): 464–475. <https://doi.org/10.1111/j.1365-2699.2005.01454.x>
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. Second edition. IUCN, Gland, Switzerland and Cambridge, UK, 32 pp.
- IUCN (2024) Guidelines for Using the IUCN Red List Categories and Criteria. Prepared by the Standards and Petitions Committee. <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Kelloff CL, Funk VA, Alexander SN (2019) Smithsonian Plant Collections, Guiana Shield: 1985–2014, Biological Diversity of the Guiana Shield Collections. Smithsonian Institution Scholarly Press, 361 pp. <https://doi.org/10.5479/si.1938-2812.110>

- Maguire B (1972) Bonnetiaceae. The botany of the Guayana Highland – Part IX. Memoirs of the New York Botanical Garden 23: 131–165.
- Nogué S, Rull V, Vegas-Vilarrúbia T (2009) Modeling biodiversity loss by global warming on Pantepui, northern South America: Projected upward migration and potential habitat loss. Climatic Change 94(1–2): 77–85. <https://doi.org/10.1007/s10584-009-9554-x>
- QGIS.org (2023) QGIS Geographic Information System. QGIS Association. <http://www.qgis.org>
- Radford AE, Dickison WC, Massey JR, Bell CR (1974) Vascular plant systematics. Harper & Row, New York.
- Riina R, Berry PE, Huber O, Michelangeli FA (2019) Vascular plants and bryophytes. Biodiversity of Pantepui. Elsevier, 121–147. <https://doi.org/10.1016/B978-0-12-815591-2.00006-9>
- Sastre C (1987) Studies on the Flora of the Guianas 27. Description d'un *Neblinaria* (Bonnetiaceae) nouveau de Guyana. Bulletin du Muséum National d'Histoire Naturelle. Section B. Adansonia 2: 129–131.
- Steyermark JA (1984) Flora of the Venezuelan Guayana – I. Annals of the Missouri Botanical Garden 71(1): 297–340. <https://doi.org/10.2307/2399074>
- Vasco A, Thadeo M, Conover M, Daly DC (2014) Preparation of samples for leaf architecture studies, a method for mounting cleared leaves. Applications in Plant Sciences 2(9): 1400038. <https://doi.org/10.3732/apps.1400038>
- Vieira TL, Barbosa-Silva RG, Acosta AL, Van Den Berg C (2024) Expanding the Distribution of *Prosthechea jauana* (Orchidaceae) in the Pantepui and Highlighting the Urgent Need for Conservation Strategies in the Region in Face of Climate Change. Plants 13(2): 222. <https://doi.org/10.3390/plants13020222>
- Weitzman AL, Kubitzki K, Stevens PF (2007) Bonnetiaceae. In: Kubitzki K (Ed.) The Families and Genera of Vascular Plants. Volume IX. Flowering Plants · Eudicots. Springer Berlin Heidelberg, Berlin, Heidelberg, 36–39. https://doi.org/10.1007/978-3-540-32219-1_8
- World Conservation Monitoring Centre (1998a) *Bonnetia ptariensis*. The IUCN Red List of Threatened Species 1998: e.T36186A9982305. <https://doi.org/10.2305/IUCN.UK.1998.RLTS.T36186A9982305.en> [Accessed on 05 September 2024]
- World Conservation Monitoring Centre (1998b) *Bonnetia rubicunda*. The IUCN Red List of Threatened Species 1998: e.T36463A9997059. <https://doi.org/10.2305/IUCN.UK.1998.RLTS.T36463A9997059.en> [Accessed on 05 September 2024]